

ARTICLE

Open Access

The NAM/ATAF1/2/CUC2 transcription factor PpNAC.A59 enhances *PpERF.A16* expression to promote ethylene biosynthesis during peach fruit ripening

Zhi-Hua Guo¹, You-Jia Zhang¹, Jia-Long Yao², Zhi-Hua Xie¹, Yu-Yan Zhang³, Shao-Ling Zhang¹ and Chao Gu¹✉

Abstract

Peach is a typical climacteric fruit that releases ethylene during fruit ripening. Several studies have been conducted on the transcriptional regulation of ethylene biosynthesis in peach fruit. Herein, an ethylene response factor, *PpERF.A16*, which was induced by exogenous ethylene, could enhance ethylene biosynthesis by directly inducing the expression of *1-aminocyclopropane-1-carboxylic acid synthase (PpACS1)* and *1-aminocyclopropane-1-carboxylic acid oxidase (PpACO1)* genes. Moreover, the NAM/ATAF1/2/CUC2 (NAC) transcription factor (TF) *PpNAC.A59* was coexpressed with *PpERF.A16* in all tested peach cultivars. Interestingly, *PpNAC.A59* can directly interact with the promoter of *PpERF.A16* to induce its expression but not enhance *LUC* activity driven by any promoter of *PpACS1* or *PpACO1*. Thus, *PpNAC.A59* can indirectly mediate ethylene biosynthesis via the NAC-ERF signaling cascade to induce the expression of both *PpACS1* and *PpACO1*. These results enrich the genetic network of fruit ripening in peach and provide new insight into the ripening mechanism of other perennial fruits.

Introduction

Fruit ripening is the last stage of the fruit developmental process and is accompanied by a series of physiological and metabolic changes in color, texture, flavor, and nutritional compounds¹. Fleshy fruits are classically defined as climacteric or nonclimacteric; the former exhibit a burst of ethylene production and respiration during ripening, while the latter do not². Ethylene is a major signaling molecule that controls fruit ripening in climacteric fruits³ and plays an important role in the fruit ripening of nonclimacteric fruits^{4,5}.

Ethylene biosynthesis and signal transduction have been widely studied in angiosperms^{6–8}. Ethylene biosynthesis is

catalyzed by two enzymes, 1-aminocyclopropane-1-carboxylic acid synthase (ACS), and oxidase (ACO). Synthesized ethylene activates the expression of ripening-related genes through transcription factors (TFs), ETHYLENE INSENSITIVE-like factors and ethylene responsive factors (ERFs). In tomato, LeERF1 promotes fruit ripening⁹, while SlERF6 delays fruit ripening¹⁰. In perennial fruit trees, MaERF9, MdERF3, and PbERF24 induce ethylene biosynthesis during fruit ripening in banana, apple, and pear, respectively^{11–13}. In contrast, MaERF11, MdERF2, and CpERF9 inhibit ethylene biosynthesis during fruit ripening in banana, apple, and papaya, respectively^{11,12,14}. Notably, apple MdERF3 is the downstream target of MdERF2, MdMYC2, and MdARF5^{12,15,16}. These findings suggest that ERFs are important in ethylene biosynthesis during fruit ripening.

In addition to ERF, other TFs, such as NAM/ATAF1/2/CUC2 (NAC), MADS-box, homeobox, apetala 2, SQUAMOSA promoter binding protein (SBP), auxin response

Correspondence: Chao Gu (guchao@njau.edu.cn)

¹College of Horticulture/State Key Laboratory of Crop Genetics and Germplasm Enhancement, Nanjing Agricultural University, 210095 Nanjing, China

²New Zealand Institute of Plant & Food Research Ltd, Private Bag 92169, Auckland 1142, New Zealand

Full list of author information is available at the end of the article

© The Author(s) 2021



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

factor (ARF), and zinc-finger proteins, are also involved in mediating ethylene biosynthesis in climacteric fruits^{8,17–20}. Among the NAC TFs, NOR, SINAC4, and NAC-like 1 are positive regulators of fruit ripening^{21–23}, while SINAC1 is a transcriptional repressor of ripening-related genes in tomato²⁴. AdNAC6 and AdNAC7 can bind to the *AdACS1* and *AdACO1* promoters to stimulate its expression and increase ethylene production in kiwifruit²⁵. MaNAC1 and MaNAC2 can directly interact with ethylene signaling components, such as ETHYLENE INSENSITIVE, in banana²⁶. CmNAC-NOR is located in a quantitative trait locus for climacteric ripening and is likely associated with ethylene biosynthesis and fruit ripening in melon²⁷. Moreover, a heterodimer of two NACs can activate the transcription of an MYB TF²⁸, indicating that the NAC TF is probably an upstream factor that regulates the expression of downstream TF(s). These findings suggest that NAC could be involved in regulating ethylene biosynthesis by directly and/or indirectly inducing ethylene biosynthesis genes. This suggestion is supported by previous reports, in which three MADS-box TFs, RIN, FUL1, and FUL2, can mediate the expression of ACS genes and their upstream TFs, including RIN, NOR, and CNR (an SBP TF)^{29,30}.

Peach (*Prunus persica*) emits a large amount of ethylene during fruit ripening^{31,32}. Exogenous ethylene accelerates fruit softening and induces the expression of softening-related genes³³, such as *endopolygalacturonase* (*PpendoPGM*) controlling the melting flesh phenotype in the clingstone melting flesh (CMF) peach fruit³⁴. Moreover, ethylene biosynthesis and signal transduction in peach can be interrupted by aminoethoxyvinylglycine and 1-methylcyclopropene (1-MCP), respectively, leading to the delay of fruit ripening^{35,36}. Therefore, ethylene is crucial for catalyzing fruit ripening in peach. The genes *PpACS1* and *PpACO1*, which encode the two catalytic enzymes for ethylene biosynthesis, have been identified in peach fruit^{37–39}. Two TFs, PpHB.G7 and PpTCP.A2, were identified to be involved in peach fruit ripening^{40,41}. PpHB.G7 can bind to the *PpACS1* and *PpACO1* promoters to induce the expression of both genes and increase ethylene production⁴⁰, while PpTCP.A2 can inhibit ethylene biosynthesis by negatively affecting *PpACS1* expression⁴¹. To broaden the transcription regulation network of ethylene biosynthesis in peach, two TFs, PpERF.A16 and PpNAC.A59 were isolated from peach fruit. Of these, PpERF.A16 could directly bind to the *PpACS1* and *PpACO1* promoters to positively regulate ethylene biosynthesis in peach fruit. *PpNAC.A59* was coexpressed with *PpERF.A16*, and PpNAC.A59 could directly bind to the promoter of *PpERF.A16* to indirectly promote ethylene biosynthesis in peach fruit. This study further elucidates the molecular network of ethylene-induced fruit ripening in peach and other perennial fruit trees.


Results

Identification of ACS, ACO, and ERF genes during fruit ripening

Members of the ACS, ACO, and ERF gene families were isolated from peach and other fruit trees. A total of 106 AP2/ERF genes (Table S1) were detected in the peach genome and could be divided into two groups, A and B (Fig. S1). Of the two groups, group A corresponded to subfamily ERF, which comprised 56 clusters (A1 → A56), and group B contained the AP2, RAV, and soloist subfamilies. Moreover, six members of the ACS gene family were detected in peach (Fig. S2) along with 32 possible members of the ACO (or ACO-like) gene family (Fig. S3). Based on the RNA-Seq data of fruit samples of the peach cultivars Nanshantiantao (NS) and Zhaohui (ZH) in a previous study³⁴, the ACS, ACO, and AP2/ERF genes, which had higher levels of expression in ripening fruit than in developing fruit (Fig. 1A and Table S2), were detected. Based on previous reports^{37,40}, the ACS gene *Ppa004774m* was identical to *PpACS1*, and the two ACO genes, *Pp008791m* and *Ppa009228m*, were identical to *PpACO1* and *PpACO3*, respectively. The three ERF genes belonged to the A16, A29, and A31 clusters and were thus designated *PpERF.A16* (Ppa008730m), *PpERF.A29* (Ppa021711m), and *PpERF.A31.1* (Ppa007193m), respectively (Table S1). However, the expression level of *PpACO3* in ripening fruit was similar to that in developing fruit of cv. Xiahui No. 5 (Fig. 1B) and cvs. Yuhualu, Chumei, and Hujimilu (Fig. 1C). Moreover, *PpERF.31.1* and *PpERF.A29* had similar levels of expression between developing and ripening fruits of some cultivars (Fig. 1C), although these two genes were more highly expressed in ripening fruit than in developing fruit of cv. Xiahui No.5 (Fig. 1B). These results indicate that *PpACO3*, *PpERF.31.1*, and *PpERF.A29* are not involved in peach fruit ripening. In contrast, *PpACS1*, *PpACO1*, and *PpERF.A16* were more highly expressed in ripening fruit than in developing fruit of all 12 tested cultivars (Fig. 1B, C) and thus are probably associated with peach fruit ripening.

Interaction of PpERF.A16 with the promoters of the *PpACS1*, *PpACO1*, and *PpendoPGM* genes

Similar to the previous study⁴⁰, to test whether PpERF.A16 could bind to the promoters of ripening-related genes, *PpACS1*, *PpACO1*, and *PpendoPGM*, dual-luciferase assays were conducted and revealed that the LUC activities that were individually driven by the *PpACS1*, *PpACO1*, or *PpendoPGM* promoters were significantly increased in *Arabidopsis* protoplasts overexpressing *PpERF.A16* (35S::PpERF.A16) compared to *Arabidopsis* protoplasts transformed with an empty vector or without any transformation (control; Fig. 2B). This result indicates that PpERF.A16 can interact with the promoters of *PpACS1*, *PpACO1*, and *PpendoPGM*.



Subsequently, yeast one-hybrid (Y1H) assays showed that PpERF.A16 binds to the *PpACS1*, *PpACO1*, and *PpendoPGM* promoters (Fig. 2C). Moreover, a PpERF.A16::HIS fusion protein was generated by inserting the whole sequences of *PpERF.A16* into the pCold-TF vector. Kinetic assay showed that binding values between PpERF.A16::HIS, and the promoters of the three ripening-related genes gradually increased with increasing concentrations of PpERF.A16::HIS (Fig. 2D). Furthermore, electrophoretic mobility shift assay (EMSA) showed that PpERF.A16::HIS could bind to the probes of the *PpACS1*,

PpACO1, and *PpendoPGM* promoters, and the binding signals weakened with increasing concentrations of the cold probe (Fig. 2D). Taken together, PpERF.A16 directly binds to the *PpACS1*, *PpACO1*, and *PpendoPGM* promoters to enhance their activities.

Effect of PpERF.A16 on ethylene biosynthesis

To unravel the role of PpERF.A16 in ethylene biosynthesis, overexpression and antisense vectors of *PpERF.A16* were individually constructed and then introduced to *Agrobacterium tumefaciens*. After introducing the





Fig. 2 PpERF.A16 directly interacts with the promoters of *PpACS1*, *PpACO1*, and *PpendoPGM*. **A** GCC-box element(s) were predicted in the promoters of *PpACS1*, *PpACO1*, and *PpendoPGM*. **B** PpERF.A16 activated *LUC* expression that was controlled by the promoters of *PpACS1*, *PpACO1*, or *PpendoPGM*. **C** Y1H assay of PpERF.A16 with the predicted GCC-box elements in the *PpACS1*, *PpACO1*, and *PpendoPGM* promoters. **D** Kinetic assay and EMSA of PpERF.A16 with the *PpACS1*, *PpACO1*, and *PpendoPGM* promoters. + and – indicate the presence and absence of recombinant PpERF.A16::HIS, biotin-labeled probe, or cold probe, respectively. Cold probe concentrations were tenfold (10x), 50-fold (50x), 100-fold (100x), 150-fold (150x), 200-fold (200x), or 250-fold (250x) those of the labeled probes

overexpression vector of *PpERF.A16* into tobacco and identifying the positive lines (Fig. S4), the ethylene production rate was measured using a gas chromatography system. The results showed that the ethylene production rate was higher in the transgenic lines of *PpERF.A16* (P value < 0.001) than in the nontransgenic regeneration lines (Fig. 3A), suggesting that PpERF.A16 can promote ethylene biosynthesis in transgenic tobacco. Furthermore, *Agrobacterium tumefaciens* containing overexpression and antisense vectors of *PpERF.A16* was independently injected into the fruits harvested at ~10 days before the commercial harvesting date, and ethylene production rates were measured at 3, 4, and 5 days after injection (DAI). At 5 DAI, the ethylene production rate was increased in the fruits overexpressing *PpERF.A16* but decreased in the fruits silencing *PpERF.A16* compared to the fruits infiltrated with the empty vector (Fig. 3B). Thus, PpERF.A16 can positively mediate ethylene biosynthesis during peach fruit ripening.

To test whether the expression of the *PpACS1*, *PpACO1*, and *PpendoPGM* genes was affected by PpERF.A16, qRT-PCR analyses were conducted on the transiently transformed peach fruits. As a result, both *PpACS1* and *PpACO1* were upregulated in fruits overexpressing *PpERF.A16* but downregulated in the fruits silencing *PpERF.A16* compared to the fruits infiltrated with the empty vector (Fig. 3C). Further qRT-PCR analysis showed that the transcript levels of *PpERFA15.1/15.2/15.3* and *PpERFA17.2/17.2* were most closely related to *PpERF.A16* and were not downregulated in the fruits with silenced *PpERF.A16* (Fig. S5). These results suggest that PpERF.A16 can positively regulate ethylene biosynthesis by inducing the expression of both *PpACS1* and *PpACO1*. In addition, PpERF.A16 upregulated the expression of *PpendoPGM* (Fig. 3C) and ripening-related genes (such as *PpCHI*, *PpPDS*, *PpZEP*, and *PpNECD*) (Fig. S6). Together with the results of the promoter analyses described above, PpERF.A16 was shown to be able to



directly affect the expression of the *PpACS1*, *PpACO1*, and *PpendoPGM* genes.

Identification of the NAC TF coexpressed with *PpERF.A16*

To mine the upstream TF(s) of *PpERF.A16* expression and ethylene biosynthesis, the potential *cis*-regulatory elements in the *PpERF.A16* promoter were predicted, revealing 12 elements (the sequences with red color) for the NAC TF (Fig. S7). A higher number of elements for the NAC TF indicated that this TF should play an important role in mediating *PpERF.A16* expression and ethylene production. Subsequently, 103 members of the NAC gene family were isolated from the peach genome. Together with the NAC genes derived from other fruit trees, the NAC gene family can be divided into two groups, A and B (Fig. S8 and Table S3). Of the two groups, group A comprised 69 clusters (A1 → A69), while group B contained only three clusters (B1 → B3; Fig. S8). Based on the RNA-Seq data of fruit samples of cvs. NS and ZH in a previous study³⁴, two NAC genes were more highly expressed in ripening fruit than in developing fruit of both cvs. NS and ZH (Fig. 4A and Table S2). These two genes, *Ppa007445m* and *Ppa008828m*, belong to clusters A32 and A59 and were thus designated *PpNAC.A32* and *PpNAC.A59*, respectively (Table S3). Of the two genes, however, *PpNAC.A32* had similar levels of expression between developing and ripening fruits of cv. Xiahui No.5

(Fig. 4B) and cvs. Xiahui No. 2, Yuhualu, Chunxue, and Wasesimizu (Fig. 4C), while *PpNAC.A59* was more highly expressed in ripening fruit than in developing fruit of all 12 tested cultivars (Fig. 4B, C). Obviously, *PpNAC.A59* had an identical expression profile to *PpERF.A16* in all tested peach cultivars, indicating that this TF may induce the expression of *PpERF.A16* and is probably associated with peach fruit ripening.

Interaction of *PpNAC.A59* with the *PpERF.A16* promoter

To test whether *PpNAC.A59* interacts with the *PpERF.A16* promoter, a dual-luciferase assay was conducted and showed an interesting result. The *LUC* gene driven by the *PpERF.A16* promoter had higher activity in *Arabidopsis* protoplasts overexpressing *PpNAC.A59* (35S::PpNAC.A59) than in *Arabidopsis* protoplasts transformed with an empty vector or without any transformation (control; Fig. 5A). In contrast, the *LUC* gene driven by any promoter of *PpACS1*, *PpACO1*, and *PpendoPGM* had similar activity in *Arabidopsis* protoplasts overexpressing *PpNAC.A59* to that in *Arabidopsis* protoplasts transformed with an empty vector (Fig. 5A). Therefore, *PpNAC.A59* interacts with the *PpERF.A16* promoter.

To determine which fragment(s) in the promoter of *PpERF.A16* could be bound by *PpNAC.A59*, the promoter sequences of *PpERF.A16* were fragmented into seven sections (I → VII). Only fragment V was bound by *PpNAC.A59*.





Fig. 4 Identification of the NAC genes probably associated with peach fruit ripening. **A** Transcriptome analysis showed that *PpNAC.A59* and *PpNAC.A32* were differentially expressed between developing (S1 to S3) and ripening (S4) fruits in peach cvs. NS and ZH. The color bar was calculated by the log₂ fold-change values of the ratio. **B** The expression profiles of *PpNAC.A59* and *PpNAC.A32* were tested by qRT-PCR in cv. Xiahui No. 5 fruits at S1, S2, S3, and S4. **C** The expression levels of *PpNAC.A59* and *PpNAC.A32* were tested in the fruits of 11 peach cultivars at S2 and S4

A59 in yeast cells (Fig. 5B). In fragment V, we detected two possible *cis*-regulatory elements (WNNYBTNNNNNN NAMGNHW and TTRCGT) for NAC TF^{42–44}, and these elements were used to synthesize the biotin-labeled probes (Fig. 5C). Subsequently, a recombinant PpNAC.A59::HIS fusion protein was generated by inserting the whole sequences of *PpNAC.A59* into the pCold-TF vector. Kinetic assay showed that the binding value between PpNAC.A59::HIS and each of the three probes gradually increased with increasing concentrations of PpNAC.A59::HIS, and the binding response between PpNAC.A59::HIS and the probe boxed with a yellow color peaked when the concentration reached 4500 nM (KD <1 μM; Fig. 5C). Moreover, EMSA showed that PpNAC.A59::HIS could bind to the probe boxed with a yellow color, and the binding signal weakened as the concentration of the cold probe increased (Fig. 5C). Taken together, PpNAC.A59 can directly bind to the promoter of *PpERF.A16*.

Effect of PpNAC.A59 on ethylene biosynthesis

To unravel the role of PpNAC.A59 in ethylene biosynthesis, overexpression and antisense vectors of *PpNAC.A59* were constructed and then introduced into *A. tumefaciens*. The resuspended *Agrobacterium* was injected into peach fruits, and ethylene production rates were measured at 3, 4, and 5 DAI. At 5 DAI, the ethylene production rate was increased in the fruits overexpressing *PpNAC.A59* but reduced in the fruits with silenced *PpNAC.A59* compared to the fruits injected with an empty vector (Fig. 6A). This result indicated that PpNAC.A59 can positively regulate ethylene biosynthesis during peach fruit ripening.

To test whether the expression of ethylene biosynthesis and ripening-related genes was affected by PpNAC.A59, the transiently transformed peach fruits were analyzed using qRT-PCR. The results showed that the expression levels of *PpACS1*, *PpACO1*, and *PpERF.A16* were



identified in climacteric fruits such as apple, banana, and pear^{11–13}. This study identified a similar gene *PpERF.A16* from previously reported RNA-Seq data from peach fruit. Overexpression of *PpERF.A16* could enhance the expression of *PpACS1* and *PpACO1* and thus increase ethylene production during peach fruit ripening (Fig. 3). Furthermore, *PpERF.A16* could also induce the expression of other ripening-related genes (Figs. 3C, S6) and is thus likely associated with peach fruit ripening. *PpERF.A16* showed a similar function to the previously reported *ERF* genes *MdERF3*, *PbERF24*, and *MaERF9* involved in regulating the ripening of climacteric fruits^{11–13}. However, a difference is that *PpERF.A16* binds to the promoters of both the *PpACS1* and *PpACO1* genes (Fig. 2), while *MdERF3*, *PbERF24*, and *MaERF9* interact with the promoters of an *ACS* or *ACO* gene but not both genes^{11–13}. Another difference is that these genes are clustered into different subgroups of the *ERF* family, i.e., *PpERF.A16*, *PbERF24*, *MdERF3*, and *MaERF9* were clustered into the A16, A17, A39 and A13 subgroups, respectively (Table S1).

It has been reported that exogenous auxin can effectively induce the expression of *PpACS1* and *PpERF.A16*^{37,48}. Ethylene production is associated with the TC microsatellite genotypes of *PpYUC11* in peach fruits with flesh melting and stony hard phenotypes^{49,50}. Recently, it was confirmed that auxin can promote ethylene biosynthesis in apple fruit by activating *MdARF5*¹⁶. *MdARF5* can bind to the *MdACS1*, *MdACS3a*, *MdACO1*, and *MdERF2* promoters to induce their expression in apple fruit. It is possible that auxin induces ethylene

biosynthesis in peach fruit by an ARF TF. However, exogenous auxin treatment cannot upregulate ARF TFs in peach fruit^{37,48}. Instead, the expression levels of *auxin/indole-3-acetic acid (auxin/IAA)* genes (*IAA11*, *IAA13*, and *IAA9_2*) were increased in the peach fruits with exogenous auxin treatment compared with those in the control fruits^{37,48}. It is more possible that these auxin/IAA TFs directly or indirectly induce the expression of *PpACS1* and *PpERF.A16* to promote ethylene biosynthesis in peach fruit.

PpNAC.A59 indirectly mediates ethylene biosynthesis during fruit ripening

Plant-specific NAC TFs of woody and grassy plants have been phylogenetically analyzed^{51,52} and are reported to regulate various physiological traits^{53–56}. *NOR* was the first NAC gene known to promote ethylene biosynthesis in a nonripening tomato mutant²². Subsequently, the *SINAC4* and *NOR-like 1* genes were proven to be associated with fruit ripening in tomato^{21,23}, similar to *AdNAC6* and *AdNAC7* in kiwifruit²⁵. In the present study, *PpNAC.A59* was shown to have a higher level of expression in ripening fruit than in developing fruit of all tested peach cultivars (Fig. 4). Interestingly, *PpNAC.A59* cannot interact with the promoters of *PpACS1* and *PpACO1* (Fig. 5A) but could induce the expression of both *PpACS1* and *PpACO1* to promote ethylene biosynthesis in transiently transformed fruits (Fig. 6). This result is different from previous studies of the NACs associated with ethylene biosynthesis during fruit ripening. For example, *NOR-like 1* of tomato interacts with the *SIACS2* and

SLACS4 promoters²¹, and AdNAC6/7 of kiwifruit interacts with the *AdACS1* and *AdACO1* promoters²⁵. In contrast, PpNAC.A59 directly interacts with the *PpERF.A16* promoter (Fig. 5) but not with the *PpACS1* or *PpACO1* promoter. Then, PpERF.A16 interacts with the promoters of *PpACS1* and *PpACO1*. Obviously, PpNAC.A59 indirectly mediates ethylene biosynthesis during peach fruit ripening. It is noteworthy that PpNAC.A59 was not clustered together with any of the previously reported NACs associated with ethylene biosynthesis during fruit ripening. PpNAC.A59, AdNAC6/7, NOR/NOR-like 1, SINAC1, and SINAC4 were individually clustered into subgroups A59, A21, A35, A25, and A24 of the NAC family, respectively (Table S3).


In previous studies, NACs were reported to be involved in abscisic acid (ABA) biosynthesis^{57,58}. Both *Oryza sativa* OsNAC2 and *Solanum lycopersicum* SINAP2 directly control the expression of the ABA biosynthesis gene *9-cis-epoxycarotenoid dioxygenase* and the ABA catabolic gene *ABA 8'-hydroxylase* to affect ABA production^{59,60}. It is noteworthy that ABA indirectly promotes fruit ripening by triggering ethylene biosynthesis in climacteric fruits^{61,62}. In this study, we found that PpNAC.A59 positively regulated the expression of ABA biosynthesis genes (*Zeaxanthin epoxidase* and *9-cis-epoxycarotenoid dioxygenase*; Fig. S10), indicating that PpNAC.A59 is likely involved in ABA biosynthesis in peach fruit. Therefore, ethylene biosynthesis may also be influenced by the PpNAC.A59-ABA-ethylene signal cascade during peach fruit ripening.

Molecular network of fruit ripening in peach

Controlling the ripening process of climacteric fruits is important to the commercialization of these fruit crops and has been widely studied at the molecular genetics level^{19,63}. In tomato, a genetic network regulating fruit ripening has been established and consists of TFs of different families, including RIN, FUL1, FUL2, HB-1, AP2a, ERF6, TAGL1, NOR, CNR, and TDR4^{29,30}. This network was further enriched by the most recently identified TFs, SINAC4, SINAC1, SIZFP2, and NOR-like 1^{20,21,23,24}, and epigenetic modifiers, such as DNA methylation^{64,65}, microRNA^{66,67}, and long noncoding RNA^{68,69}. Obviously, the understanding of the genetic network regulating fruit ripening in tomato is much advanced compared to our understanding of the network in tree fruits. During apple fruit ripening, the regulatory network of ethylene biosynthesis was constructed to include the TFs MYC2, ARF5, ERF2, and ERF3 and ethylene biosynthesis genes^{12,15,16}. However, in other perennial fruit trees, few TFs were concatenated into a network. In peach fruit, two TFs, PpHB.G7 and PpTCP.A2, were identified in the previous studies^{40,41}. Of these two TFs, the transcription activator PpHB.G7

can directly bind to the *PpACS1* and *PpACO1* promoters to increase ethylene production⁴⁰, while the transcription repressor PpTCP.A2 can inhibit the expression of *PpACS1* to hinder ethylene biosynthesis⁴¹.

In the present study, two TFs, PpERF.A16 and PpNAC.A59, were shown to be involved in ethylene biosynthesis during peach fruit ripening. These results could be integrated into a primary network of fruit ripening in peach (Fig. 7). First, PpNAC.A59 can induce the expression of *PpERF.A16* by directly binding to its promoter. Second, PpERF.A16 and PpHB.G7 enhances the expression of both *PpACS1* and *PpACO1* by directly interacting with their promoters, contributing to an increase in ethylene production. Finally, ethylene accelerates fruit ripening by mediating fruit metabolism, such as flavonoid and carotenoid metabolism. It is noteworthy that this conclusion is supported by a previous study in which the NAC TF was predicted to control ethylene biosynthesis during peach fruit ripening⁷⁰. The only difference is that the predicted NAC TF can induce the expression of ethylene biosynthesis genes by directly interacting with their promoters⁷⁰, while PpNAC.A59 indirectly induces the expression of both *PpACS1* and *PpACO1* by directly mediating the expression of *PpERF.A16*. Further work will focus on the enrichment of this network by identifying more TFs associated with fruit ripening.



Materials and methods

Plant materials

A total of 12 peach cultivars were used in this study. Of these cultivars, Wasesimizu and Xiahui No. 5 were maintained at the Jiangsu Academy of Agricultural Sciences (Nanjing, Jiangsu Province, China). The other ten cultivars (Chunxue, Hakuho, Hujinmilu, Xiahui No. 2, Yuhualu, Chiyuanmi, Chunmei, Xiahui No. 6, Baixianglu, and Zao-baifeng) were identical to those in a previous study⁴⁰. The cv. Baixianglu fruit presents a semifreestone melting flesh phenotype, and the fruits of the other ten peach cultivars present a clingstone melting flesh phenotype. Based on the dynamic investigation of fruit development and ripening⁷¹, the fruits of cv. Xiahui No. 5 were collected at the fruitlet (S1), stone hardening (S2), preripening (S3), and ripening (S4) stages, and the fruits of the other 11 cultivars were collected at the rapid enlargement (roughly corresponding to S2) and ripening stages (S4; Table S4). Grouping, biological replication, and storage of fruit samples were as described in a previous study⁴⁰.

Sequence and phylogenetic analyses of the ACS, ACO, AP2/ERF, and NAC genes

To isolate the ACS, ACO, AP2/ERF, and NAC genes in peach and other fruit trees, the nucleotide and amino acid sequences of the ACS, ACO, AP2/ERF, and NAC genes in peach were extracted from the peach genome version 1.0 (Genome Database for Rosaceae; <http://www.rosaceae.org/>). Then, these downloaded sequences were used as indexes to find their homologs in papaya (*Carica papaya* ASGPBv0.4), strawberry (*Fragaria vesca* v 1.1), orange (*Citrus sinensis* v1.1), grape (*Vitis vinifera* Genoscope.12X), apple (*Malus domestica* v1.0) (<https://phytozome.jgi.doe.gov/>), and pear (*Pyrus bretschneideri*; <http://peargenome.njau.edu.cn/>) genomes (Tables S1, S3). Sequence alignment and phylogenetic analyses were as described in a previous study⁴⁰.

Gene expression analysis in developing and ripening fruits

In a previous study, RNA-Seq was performed in the developing and ripening fruits of three peach cultivars, Nanshantiantao (NS), Zhaohui (ZH), and Myojo³⁴. Of these three cultivars, NS and ZH fruits presented a melting flesh phenotype, and the fruit hardness gradually decreased during ripening, while fruits of the Myojo cultivar did not soften during fruit ripening. For this reason, the transcriptome data fruits from cvs. NS and ZH were used to isolate the ACS, ACO, AP2/ERF, and NAC genes that may be associated with fruit ripening. The hierarchical clustering of transcriptome data was performed according to a previous study⁴⁰. By comparing the gene expression levels of the ripening fruit (S4) to those of the developing fruits (S1 to S3), we isolated the target genes *PpACS1*, *PpACO1*, *PpACO3*, *PpERF.A16*, *PpERF.A29*,

PpERF.A31.1, *PpNAC.A32*, and *PpNAC.A59*. Thus, these eight genes were further tested by qRT-PCR analysis. The details for qRT-PCR and data analysis were identical to those of a previous study⁴⁰.

Transformations in peach fruit and tobacco

The full length CDSs of *PpERF.A16* and *PpNAC.A59* were amplified with the PCR primers listed in Table S5 and inserted in a sense orientation downstream of the CaMV 35S promoter in the pCAMBIA1301 vector to generate gene overexpression vectors. An approximately 400-bp fragment of the CDS of *PpERF.A16* or *PpNAC.A59* was amplified using the primers listed in Table S5 and inserted in an antisense orientation downstream of the CaMV 35 S promoter in the pSAK277 vector to create a gene silencing construct. The 400 bp fragments were located in the nonconserved region and showed no significant similarity (defined as no >21 bp in length with 100% identity) to the sequences of the most closely related genes within the same gene family (Figs. S11–S20). pCAMBIA1301 contains a hygromycin resistance gene, and pSAK277 contains a kanamycin resistance gene as a selection marker in plant transformation. These vectors were transferred into *Agrobacterium tumefaciens* EHA 105 cells that were used to transiently transform peach fruit tissues with a previously reported *Agrobacterium* infiltration protocol⁴⁰. The fruit used for the transformation experiments was harvested from the peach cultivar Xiahui No. 5 at ~10 days before the commercial harvesting date. At three, four and five DAI, at least five fruits were sealed in a 5-L airtight jar, and six jars were used for each construct. Three DAI, ethylene production was measured every day. Fruit flesh tissues were sampled at 5 DAI for qRT-PCR analysis. All of the PCR primer sequences are listed in Supplementary Table S5.

Overexpression vectors of *PpERF.A16* were also introduced into tobacco (*Nicotiana tabacum*) based on *Agrobacterium*-mediated transformation as described previously⁷². The regenerated tobacco plants were tested by PCR for amplification of the *hygromycin* gene from the pCAMBIA1301 vector. PCR-positive plants and wild-type control plants were used for measuring ethylene production.

Ethylene measurements

The ethylene production rate was measured using a gas chromatography system (GC2010, Shimadzu, Japan) with a 1000 µl rheodyne injector (Gaoze, Shangshai, China) according to a previously reported protocol⁷³ with a few modifications. In brief, at room temperature, at least four fruits were weighed as a replicate and then placed into a 3-l airtight chamber. For tobacco, each plant was a replicate and individually weighed after collection from tissue culture medium and placed into a 15 ml centrifuge

tube. Six replicates were used for each gene construct and each control. A 0.5-ml gas sample was collected from the chambers and tubes and injected into a gas chromatograph fitted with a 30-cm glass column (3.2 mm ID). Gas flows for N₂, H₂, and air were 40, 40, and 500 ml/min, respectively. The oven, injector, and FID temperatures were 300, 60, and 260 °C, respectively.

Protoplast isolation and dual-luciferase assay

Protoplast isolation was as described in a previous report⁴⁰. Approximately 2000-bp sequences upstream of the initiation codons of *PpACS1*, *PpACO1*, *PpendoPGM*, and *PpERF.A16* were inserted into a pGreenII 0800-LUC vector for a dual-luciferase assay that was performed according to a previous study⁴⁰. At least six biological replicates were used in each assay. All of the primer sequences are listed in Table S5.

Yeast one-hybrid assay (Y1H)

It is well known that ERF TFs can bind to the GCC-box element^{74,75}. Thus, the possible GCC-box elements were surveyed from the promoters of *PpACS1*, *PpACO1*, and *PpEndoPGM* (Fig. 2A). The fragments containing the possible GCC-box element were amplified from the genomic DNA of cv. Xiahui No.5 and then inserted into the pAbAi vector. The whole sequences of *PpERF.A16* and *PpNAC.A59* genes were individually inserted into the pGADT7 vector. The Y1H assay was performed using the Matchmaker Gold Yeast One-Hybrid Library Screening System (Clontech, Palo Alto, CA). All of the primer sequences are listed in Table S5.

Recombinant proteins of both PpERF.A16 and PpNAC.A59 in *Escherichia coli*

The whole sequences of *PpERF.A16* and *PpNAC.A59* were individually inserted into the pCold-TF expression vector (including a HIS tag) to create recombinant proteins. Recombinant plasmids were introduced into *Escherichia coli* cells and incubated at 37 °C until an OD₆₆₀ of 0.6 was reached. Protein expression was induced at 16 °C for 24 h, and the recombinant protein was extracted and purified at 4 °C⁷⁶. SDS-PAGE was conducted to analyze the extracted proteins. All of the primer sequences are listed in Table S5.

Kinetic assay

Fragments containing possible protein-binding elements were used to synthesize probes biotinylated at the ends of both the N- and C-termini (Sangon, Shanghai, China). Protein-DNA interactions were determined by biolayer interferometry using a ForteBio Octet 96 Red Instrument (Menlo Park, CA, USA). The kinetic assay comprised the following five steps: (1) baseline acquisition was conducted by loading PBS buffer (1.4 mM KH₂PO₄,

2.7 mM KCl, 4.3 mM Na₂HPO₄, 137 mM NaCl, 0.02% (w/v) Tween-20; pH = 7.2) onto streptavidin-coated biosensors, (2) the biotin-labeled probe was loaded onto the sensor, (3) a second baseline acquisition was conducted with the same PBS buffer, and steps (4) and (5) were the respective association and dissociation between the recombinant protein and the biotin-labeled probe(s). Equilibrium binding affinities (KD) were measured using nanomole quantities of samples as well as rates of association and dissociation (KA/KD)⁷⁷.

Electrophoretic mobility shift assay (EMSA)

The recombinant proteins and biotin-labeled probes were also used for EMSAs. In brief, the binding reaction was performed in a 0.2 ml centrifuge tube for 20 min at room temperature, and the reaction mixtures contained 2 µl loading buffer, 1 mg poly(dI-dC), 5 mM MgCl₂, 2.5% glycerol, 0.05% NP-40, 200 nM probe, and 1 µg recombinant protein, with ultrapure water added for a total volume of 20 µl. For probe competition, 2, 10, 20, 30, 40, or 50 µM nonlabeled probe was individually mixed into the reaction mixtures. After running in a 6% polyacrylamide gel, the protein-DNA complexes were electrophoretically transferred to a nylon membrane (GE Healthcare, Danderyd, Sweden). SDS-PAGE imaging was performed by a Tanon 5200 chemiluminescence imaging system (Tanon, Shanghai, China) and scanned using a ChemiDoc™ XRS+ System (BioRad, Hercules, CA).

Statistical analysis

Means and standard errors were calculated from at least three biological replicates for qRT-PCR analysis, five biological replicates for ethylene measurement in tobacco, and at least six replicates for ethylene measurement in peach fruit. Analysis of variance was calculated by Student's *t* test. The lowercase letters a, b, and c indicate the level of significance at *P* < 0.05. Single, double and triple asterisks represent the levels of significance at *P* < 0.05, <0.01, and <0.001, respectively.

Acknowledgements

We thank LetPub (www.letpub.com) for its linguistic assistance during the preparation of this paper. This work was supported by the Natural Science Foundations of China (31672118 and 31471856) and the Natural Science Foundations of Jiangsu Province (BK20140756).

Author details

¹College of Horticulture/State Key Laboratory of Crop Genetics and Germplasm Enhancement, Nanjing Agricultural University, 210095 Nanjing, China. ²New Zealand Institute of Plant & Food Research Ltd, Private Bag 92169, Auckland 1142, New Zealand. ³Institute of Pomology, Jiangsu Academy of Agricultural Sciences/Jiangsu Key Laboratory for Horticultural Crop Genetic Improvement, 210014 Nanjing, China

Author contributions

C.G. and S.L.Z. conceived this study; C.G. conducted transcriptome-based data analyses, isolated the candidate genes by qRT-PCR, and drafted the paper. Z.H.G. contributed to most experiments with help from Y.J.Z. and Z.H.X. Y.Y.Z.

collected peach samples at the developing and ripening stages. J.L.Y. and S.L.Z. revised this paper.

Conflict of interest

The authors declare no competing interests.

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41438-021-00644-6>.

Received: 21 January 2021 Revised: 17 June 2021 Accepted: 14 July 2021

Published online: 01 October 2021

References

- Klee, H. J. & Giovannoni, J. J. Genetics and control of tomato fruit ripening and quality attributes. *Annu. Rev. Genet.* **45**, 41–59 (2011).
- Vrebalov, J. et al. A MADS-box gene necessary for fruit ripening at the tomato ripening-inhibitor (Rin) locus. *Science* **296**, 343–346 (2002).
- Pech, J. C., Purgatto, E., Bouzayen, M. & Latché, A. Ethylene and fruit ripening. *Annu. Plant Rev.* **44**, 275–304 (2012).
- Merchante, C. et al. Ethylene is involved in strawberry fruit ripening in an organ-specific manner. *J. Exp. Bot.* **64**, 4421–4439 (2013).
- Villarreal, N. M., Bustamante, C. A., Civello, P. M. & Martínez, G. A. Effect of ethylene and 1-MCP treatments on strawberry fruit ripening. *J. Sci. Food Agric.* **90**, 683–689 (2010).
- Gu, C. et al. Multiple regulatory roles of AP2/ERF transcription factor in angiosperm. *Bot. Stud.* **58**, 6 (2017).
- Guo, H. W. & Ecker, J. R. The ethylene signaling pathway: new insights. *Curr. Opin. Plant Biol.* **7**, 40–49 (2004).
- Seymour, G. B., Ostergaard, L., Chapman, N. H., Knapp, S. & Martin, C. Fruit development and ripening. *Annu. Rev. Plant Biol.* **64**, 219–241 (2013).
- Li, Y. C. et al. LeERF1 positively modulated ethylene triple response on etiolated seedling, plant development and fruit ripening and softening in tomato. *Plant Cell Rep.* **26**, 1999–2008 (2007).
- Lee, J. M. et al. Combined transcriptome, genetic diversity and metabolite profiling in tomato fruit reveals that the ethylene response factor SlERF6 plays an important role in ripening and carotenoid accumulation. *Plant J.* **70**, 191–204 (2012).
- Xiao, Y. Y. et al. Banana ethylene response factors are involved in fruit ripening through their interactions with ethylene biosynthesis genes. *J. Exp. Bot.* **64**, 2499–2510 (2013).
- Li, T. et al. Apple (*Malus domestica*) MdERF2 negatively affects ethylene biosynthesis during fruit ripening by suppressing *MdACS1* transcription. *Plant J.* **88**, 735–748 (2016).
- Hao, P. P. et al. Transcriptome analysis unravels an ethylene response factor involved in regulating fruit ripening in pear. *Physiol. Plant.* **163**, 124–135 (2018).
- Fu, C. C. et al. Papaya CpERF9 acts as a transcriptional repressor of cell-wall-modifying genes *CpPME1/2* and *CpPG5* involved in fruit ripening. *Plant Cell Rep.* **35**, 2341–2352 (2016).
- Yue, P. et al. The jasmonate-activated transcription factor MdMYC2 regulates ETHYLENE RESPONSE FACTOR and ethylene biosynthetic genes to promote ethylene biosynthesis during apple fruit ripening. *Plant Cell* **29**, 1316–1334 (2017).
- Yue, P. et al. Auxin-activated MdARF5 induces the expression of ethylene biosynthesis genes to initiate apple fruit ripening. *N. Phytol.* **226**, 1781–1795 (2020).
- Breitel, D. A. et al. AUXIN RESPONSE FACTOR 2 intersects hormonal signals in the regulation of tomato fruit ripening. *PLoS Genet.* **12**, e1005903 (2016).
- Hao, Y. W. et al. Auxin response factor SlARF2 is an essential component of the regulatory mechanism controlling fruit ripening in tomato. *PLoS Genet.* **11**, e1005649 (2015).
- Karlova, R. et al. Transcriptional control of fleshy fruit development and ripening. *J. Exp. Bot.* **65**, 4527–4541 (2014).
- Weng, L. et al. The zinc finger transcription factor SlZFP2 negatively regulates abscisic acid biosynthesis and fruit ripening in tomato. *Plant Physiol.* **167**, 931–949 (2015).
- Gao, Y. et al. A NAC transcription factor, NOR-like1, is a new positive regulator of tomato fruit ripening. *Hortic. Res.* **5**, 75 (2018).
- Osorio, S. et al. Systems Biology of tomato fruit development: combined transcript, protein, and metabolite analysis of tomato transcription factor (nor, rin) and ethylene receptor (Nr) mutants reveals novel regulatory interactions. *Plant Physiol.* **157**, 405–425 (2011).
- Zhu, M. et al. A new tomato NAC (NAM/ATAF1/2/CUC2) transcription factor, SINAC4, functions as a positive regulator of fruit ripening and carotenoid accumulation. *Plant Cell Physiol.* **55**, 119–135 (2014).
- Meng, C. et al. Suppression of tomato SINAC1 transcription factor delays fruit ripening. *J. Plant Physiol.* **193**, 88–96 (2016).
- Wang, W. Q. et al. Genome-wide analysis of coding and non-coding RNA reveals a conserved miR164-NAC regulatory pathway for fruit ripening. *N. Phytol.* **225**, 1618–1634 (2019).
- Shan, W. et al. Molecular characterization of banana NAC transcription factors and their interactions with ethylene signalling component EIL during fruit ripening. *J. Exp. Bot.* **63**, 5171–5187 (2012).
- Rios, P. et al. ETHQV6.3 is involved in melon climacteric fruit ripening and is encoded by a NAC domain transcription factor. *Plant J.* **91**, 671–683 (2017).
- Zhou, H. et al. Molecular genetics of blood-fleshed peach reveals activation of anthocyanin biosynthesis by NAC transcription factors. *Plant J.* **82**, 105–121 (2015).
- Fujisawa, M., Nakano, T., Shima, Y. & Ito, Y. A large-scale identification of direct targets of the tomato MADS box transcription factor RIPENING INHIBITOR reveals the regulation of fruit ripening. *Plant Cell* **25**, 371–386 (2013).
- Fujisawa, M. et al. Transcriptional regulation of fruit ripening by tomato FRUITFULL homologs and associated MADS box proteins. *Plant Cell* **26**, 89–101 (2014).
- Hayama, H., Tatsuki, M., Ito, A. & Kashimura, Y. Ethylene and fruit softening in the stony hard mutation in peach. *Postharvest Biol. Technol.* **41**, 16–21 (2006).
- Tonutti, P., Casson, P. & Ramina, A. Ethylene biosynthesis during peach fruit development. *J. Am. Soc. Hortic. Sci.* **116**, 274–279 (1991).
- Hayama, H., Shimada, T., Fujii, H., Ito, A. & Kashimura, Y. Ethylene-regulation of fruit softening and softening-related genes in peach. *J. Exp. Bot.* **57**, 4071–4077 (2006).
- Gu, C. et al. Copy number variation of a gene cluster encoding endopolygalacturonase mediates flesh texture and stone adhesion in peach. *J. Exp. Bot.* **67**, 1993–2005 (2016).
- Bregoli, A. M. et al. Peach (*Prunus persica*) fruit ripening: aminoethoxyvinylglycine (AVG) and exogenous polyamines affect ethylene emission and flesh firmness. *Physiol. Plant.* **114**, 472–481 (2002).
- Hayama, H., Tatsuki, M., Ito, A. & Kashimura, Y. Combined treatment of aminoethoxyvinylglycine (AVG) and 1-methylcyclopropene (1-MCP) reduces melting-flesh peach fruit softening. *Postharvest Biol. Technol.* **50**, 228–230 (2008).
- Tadiello, A. et al. On the role of ethylene, auxin and a GOLVEN-like peptide hormone in the regulation of peach ripening. *BMC Plant Biol.* **16**, 44 (2016).
- Tatsuki, M., Haji, T. & Yamaguchi, M. The involvement of 1-aminocyclopropane-1-carboxylic acid synthase isogene, *Pp-ACS1*, in peach fruit softening. *J. Exp. Bot.* **57**, 1281–1289 (2006).
- Tonutti, P., Bonghi, C., Ruperti, B., Tornielli, G. B. & Ramina, A. Ethylene evolution and 1-aminocyclopropane-1-carboxylate oxidase gene expression during early development and ripening of peach fruit. *J. Am. Soc. Hortic. Sci.* **122**, 642–647 (1997).
- Gu, C. et al. A HD-ZIP II HOMEBOX transcription factor, PpHB.G7, mediates ethylene biosynthesis during fruit ripening in peach. *Plant Sci.* **278**, 12–19 (2019).
- Guo, Z. H. et al. Expression analysis of *TCP* genes in peach reveals an involvement of *PpTCPA2* in ethylene biosynthesis during fruit ripening. *Plant Mol. Biol. Report.* **36**, 588–595 (2018).
- Ge, H. et al. E1NAC3 transcriptionally regulates chilling-induced lignification of loquat fruit via physical interaction with an atypical CAD-like gene. *J. Exp. Bot.* **68**, 5129–5136 (2017).
- Nieuwenhuizen, N. J. et al. Natural variation in monoterpene synthesis in kiwifruit: transcriptional regulation of terpene synthases by NAC and ETHYLENE-INSENSITIVE3-like transcription factors. *Plant Physiol.* **167**, 1243–1258 (2015).
- Zhong, R. Q., Lee, C. H. & Ye, Z. H. Global analysis of direct targets of secondary wall NAC master switches in arabidopsis. *Mol. Plant* **3**, 1087–1103 (2010).
- Nakano, T., Suzuki, K., Fujimura, T. & Shinshi, H. Genome-wide analysis of the ERF gene family in Arabidopsis and rice. *Plant Physiol.* **140**, 411–432 (2006).
- Zhang, C. H. et al. Genome-wide analysis of the AP2/ERF superfamily in peach (*Prunus persica*). *Genet. Mol. Res.* **11**, 4789–4809 (2012).

47. Licausi, F., Ohme-Takagi, M. & Perata, P. AP2/ERF transcription factors: mediators of stress responses and developmental programs. *N. Phytol.* **199**, 639–649 (2013).
48. Trainotti, L., Tadiello, A. & Casadoro, G. The involvement of auxin in the ripening of climacteric fruits comes of age: the hormone plays a role of its own and has an intense interplay with ethylene in ripening peaches. *J. Exp. Bot.* **58**, 3299–3308 (2007).
49. Pan, L. et al. PpYUC11, a strong candidate gene for the stony hard phenotype in peach (*Prunus persica* L. Batsch), participates in IAA biosynthesis during fruit ripening. *J. Exp. Bot.* **66**, 7031–7044 (2015).
50. Cirilli, M. et al. Integrative genomics approaches validate *PpYUC11-like* as candidate gene for the stony hard trait in peach (*P. persica* L. Batsch). *BMC Plant Biol.* **18**, 88 (2018).
51. Hu, R. et al. Comprehensive analysis of NAC domain transcription factor gene family in *Populus trichocarpa*. *BMC Plant Biol.* **10**, 145 (2010).
52. Nuruzzaman, M. et al. Genome-wide analysis of NAC transcription factor family in rice. *Gene* **465**, 30–44 (2010).
53. Jensen, M. K. & Skriver, K. NAC transcription factor gene regulatory and protein-protein interaction networks in plant stress responses and senescence. *IUBMB Life* **66**, 156–166 (2014).
54. Kim, H. J., Nam, H. G. & Lim, P. O. Regulatory network of NAC transcription factors in leaf senescence. *Curr. Opin. Plant Biol.* **33**, 48–56 (2016).
55. Olsen, A. N., Ernst, H. A., Lo Leggio, L. & Skriver, K. NAC transcription factors: structurally distinct, functionally diverse. *Trends Plant Sci.* **10**, 79–87 (2005).
56. Zhong, R. Q., Demura, T. & Ye, Z. H. SND1, a NAC domain transcription factor, is a key regulator of secondary wall synthesis in fibers of *Arabidopsis*. *Plant Cell* **18**, 3158–3170 (2006).
57. Shang, X. G. et al. A cotton NAC transcription factor GhirNAC2 plays positive roles in drought tolerance via regulating ABA biosynthesis. *Plant Sci.* **296**, 110498 (2020).
58. Ren, T. T. et al. Involvement of NAC transcription factor SiNAC1 in a positive feedback loop via ABA biosynthesis and leaf senescence in foxtail millet. *Planta* **247**, 53–68 (2018).
59. Mao, C. J. et al. A rice NAC transcription factor promotes leaf senescence via ABA biosynthesis. *Plant Physiol.* **174**, 1747–1763 (2017).
60. Ma, X. M. et al. The NAC Transcription Factor SINAP2 Regulates Leaf Senescence and Fruit Yield in Tomato. *Plant Physiol.* **177**, 1286–1302 (2018).
61. Zhang, M., Yuan, B. & Leng, P. The role of ABA in triggering ethylene biosynthesis and ripening of tomato fruit. *J. Exp. Bot.* **60**, 1579–1588 (2009).
62. Kou, X. H. et al. The interplay between ABA/ethylene and NAC TFs in tomato fruit ripening: a review. *Plant Mol. Biol.* **106**, 223–238 (2021).
63. Shinozaki, Y. et al. High-resolution spatiotemporal transcriptome mapping of tomato fruit development and ripening. *Nat. Commun.* **9**, 364 (2018).
64. Lang, Z. et al. Critical roles of DNA demethylation in the activation of ripening-induced genes and inhibition of ripening-repressed genes in tomato fruit. *Proc. Natl Acad. Sci.* **114**, E4511–E4519 (2017).
65. Liu, R. et al. A DEMETER-like DNA demethylase governs tomato fruit ripening. *Proc. Natl Acad. Sci.* **112**, 10804–10809 (2015).
66. Gao, C. et al. MicroRNA profiling analysis throughout tomato fruit development and ripening reveals potential regulatory role of RIN on microRNAs accumulation. *Plant Biotechnol. J.* **13**, 370–382 (2015).
67. Moxon, S. et al. Deep sequencing of tomato short RNAs identifies microRNAs targeting genes involved in fruit ripening. *Genome Res.* **18**, 1602–1609 (2008).
68. Zhu, B. et al. RNA sequencing and functional analysis implicate the regulatory role of long non-coding RNAs in tomato fruit ripening. *J. Exp. Bot.* **66**, 4483–4495 (2015).
69. Li, R., Fu, D., Zhu, B., Luo, Y. & Zhu, H. CRISPR/Cas9-mediated mutagenesis of lncRNA1459 alters tomato fruit ripening. *Plant J.* **94**, 513–524 (2018).
70. Lu, P. et al. Genome encode analyses reveal the basis of convergent evolution of fleshy fruit ripening. *Nat. Plants* **4**, 784–791 (2018).
71. Gu, C. et al. RNA-Seq analysis unveils gene regulation of fruit size cooperatively determined by velocity and duration of fruit swelling in peach. *Physiol. Plant.* **164**, 320–336 (2018).
72. Liu, Y. et al. A WRKY transcription factor PbrWRKY53 from *Pyrus betulaefolia* is involved in drought tolerance and AsA accumulation. *Plant Biotechnol. J.* **17**, 1770–1787 (2019).
73. Fan, X., Argenta, L. & Mattheis, J. P. Inhibition of ethylene action by 1-methylcyclopropene prolongs storage life of apricots. *Postharvest Biol. Technol.* **20**, 135–142 (2000).
74. Han, Y. C. et al. Banana transcription factor MaERF11 recruits histone deacetylase MaHDA1 and represses the expression of *MaACO1* and expansins during fruit ripening. *Plant Physiol.* **171**, 1070–1084 (2016).
75. Yin, X. R. et al. Involvement of an ethylene response factor in chlorophyll degradation during citrus fruit degreening. *Plant J.* **86**, 403–412 (2016).
76. Chen, J. et al. Phosphatidic Acid Counteracts S-RNase Signaling in Pollen by Stabilizing the Actin Cytoskeleton. *Plant Cell* **30**, 1023–1039 (2018).
77. Sultana, A. & Lee, J. E. Measuring protein-protein and protein-nucleic acid interactions by biolayer interferometry. *Curr. Protoc. Protein Sci.* **79**, 19.25.1–19.25.26 (2015).