



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Short Communication

## Health inequities in COVID-19 vaccination among the elderly: Case of Connecticut

Hui Wang<sup>a</sup>, Ran Xu<sup>b,c</sup>, Shijin Qu<sup>d</sup>, Marlene Schwartz<sup>e</sup>, Aaron Adams<sup>f</sup>, Xiang Chen<sup>c,f,\*</sup><sup>a</sup> Institute for Modeling Collaboration and Innovation, University of Idaho, Moscow, ID 83844, USA<sup>b</sup> Department of Allied Health Sciences, University of Connecticut, Storrs, CT 06269, USA<sup>c</sup> Institute for Collaboration on Health, Intervention, and Policy (InCHIP), University of Connecticut, Storrs, CT 06269, USA<sup>d</sup> Department of Land Resources Management, China University of Geosciences, Wuhan, Hubei 430074, China<sup>e</sup> Department of Human Development and Family Sciences and Rudd Center for Food Policy & Obesity, University of Connecticut, Hartford, CT 06103, USA<sup>f</sup> Department of Geography, University of Connecticut, Storrs, CT 06269, USA

## ARTICLE INFO

## Article history:

Received 13 May 2021

Received in revised form 17 July 2021

Accepted 19 July 2021

## Keywords:

COVID-19

Elderly

Vaccination

Health inequity

Social vulnerability

## ABSTRACT

**Background:** In the United States, distribution plans for the COVID-19 vaccination were established at the state level. However, some states, such as Connecticut, followed an age-based strategy without considering occupations or co-morbid conditions due to its simplicity in implementation. This strategy raised concerns about exacerbating health inequities because it did not prioritize vulnerable communities, specifically, minorities and low-income groups. The study aims to examine the vaccination inequities among different population groups for people aged 65+.

**Methods:** A cross-sectional analysis of quantile-based independent sample t-test was employed to examine the relationship between eight social vulnerability indices (SVIs, i.e., below poverty, unemployed, without high school diploma, disability, minority, speaks English less than well, no vehicle, and mobile homes) and vaccination rates at the town level in Connecticut during the second phase of the vaccine distribution plan when individuals aged 65 and over were eligible. Negative binomial regressions were employed to further justify the relationships between SVIs and vaccination rates.

**Results:** The report shows that the differences in vaccination rates were statistically significant between the most vulnerable and the least vulnerable towns with respect to six SVIs (i.e., below poverty, without high school diploma, disability, minority, speaks English less than well, and no vehicle). The vaccination gap was greater for people aged 75+ than people aged 65–74. Among the selected SVIs, below poverty was negatively correlated with the vaccination rate for 75+, and without high school diploma was negatively correlated with both rates.

**Conclusions:** This report reveals the significant health inequities in COVID-19 vaccination among the elderly population at the early vaccination phase. It can shed insights into health policy initiatives to improve vaccination coverage in the elderly communities, such as promoting onsite scheduling and increasing at-home vaccination services.

Published by Elsevier Ltd on behalf of King Saud Bin Abdulaziz University for Health Sciences. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Introduction


In the United States, distribution plans for the Coronavirus Disease 2019 (COVID-19) vaccination have been established at the state level. Like many states, Connecticut's first phase included vaccinating health care workers and nursing home residents, followed

by people aged 75 and older starting on January 18, 2021. However, unlike most states, Connecticut continued a primarily age-based strategy: people 65 and over were eligible on February 11, 2021, followed by other age groups in reverse chronological order [1].

The rationale provided by state leaders for this approach was its simplicity: it did not require proof of a particular occupation (e.g., front-line worker) or co-morbid condition. However, this strategy raised concerns about exacerbating health inequities because it did not prioritize communities that have been disproportionately affected by COVID-19, specifically, minorities and low-income groups [2,3]. These population groups were found to lack access to

\* Corresponding author at: Department of Geography, University of Connecticut, Storrs, CT 06269, USA.

E-mail addresses: [huiwang@uidaho.edu](mailto:huiwang@uidaho.edu) (H. Wang), [ran.2.xu@uconn.edu](mailto:ran.2.xu@uconn.edu) (R. Xu), [qusj@cug.edu.cn](mailto:qusj@cug.edu.cn) (S. Qu), [marlene.schwartz@uconn.edu](mailto:marlene.schwartz@uconn.edu) (M. Schwartz), [aaron.adams@uconn.edu](mailto:aaron.adams@uconn.edu) (A. Adams), [xiang.chen@uconn.edu](mailto:xiang.chen@uconn.edu) (X. Chen).



**Fig. 1.** Spatial distribution of vaccination rates for population (a) aged 65–74, and (b) aged 75+ in Connecticut towns.

**Table 1**  
Vaccination rates for the least (Q1) and most (Q4) vulnerable towns.

SVI <sup>a</sup>	Quartile <sup>b</sup>	SVI range <sup>a</sup>	Rate for 65–74 (std.) <sup>c</sup>	Rate for 75+ (std.) <sup>c</sup>
Below poverty	Q1	0.80–3.45	79.58 (9.99)	98.01 (14.92)
	Q4	37.57–28.51	69.12 (12.37)	73.87 (13.70)
		Difference estimate <sup>d</sup>	10.46***	24.14***
		Difference 95% CI <sup>e</sup>	5.59–15.35	17.92–30.35
Unemployed	Q1	0.22–2.01	75.60 (9.39)	88.75 (15.64)
	Q4	3.57–7.15	70.99 (12.43)	83.72 (19.24)
		Difference estimate <sup>d</sup>	4.61	5.03
		Difference 95% CI <sup>e</sup>	–0.14 to 9.36	–2.54 to 12.58
Without high school diploma	Q1	0.91–2.95	79.77 (8.19)	98.62 (13.53)
	Q4	6.09–15.72	66.86 (10.14)	74.47 (14.91)
		Difference estimate <sup>d</sup>	12.91***	24.15***
		Difference 95% CI <sup>e</sup>	8.91–16.92	17.97–30.33
Disability	Q1	5.01–8.76	78.54 (9.47)	96.80 (15.31)
	Q4	12.57–17.15	70.69 (11.59)	80.52 (17.03)
		Difference estimate <sup>d</sup>	7.85***	16.28***
		Difference 95% CI <sup>e</sup>	3.25–12.45	9.25–23.31
Minority	Q1	3.28–7.39	78.81 (11.05)	95.80 (17.39)
	Q4	20.16–85.23	70.18 (11.49)	74.80 (10.88)
		Difference estimate <sup>d</sup>	8.63***	21.00***
		Difference 95% CI <sup>e</sup>	3.74–13.53	14.70–27.30
Speaks English less than well	Q1	0–0.29	78.07 (10.33)	94.38 (15.05)
	Q4	1.92–12.21	70.28 (13.24)	74.46 (13.23)
		Difference estimate <sup>d</sup>	7.79**	19.92***
		Difference 95% CI <sup>e</sup>	2.63–12.95	13.76–26.07
No vehicle	Q1	0–0.82	79.30 (8.94)	98.84 (13.78)
	Q4	2.4–11.75	70.50 (11.84)	73.65 (12.02)
		Difference estimate <sup>d</sup>	8.80***	25.19***
		Difference 95% CI <sup>e</sup>	4.24–13.36	19.58–30.80
Mobile homes	Q1	0–0	76.36 (11.53)	89.94 (16.57)
	Q4	0.55–4.82	76.14 (10.32)	87.49 (12.69)
		Difference estimate <sup>d</sup>	0.22	2.45
		Difference 95% CI <sup>e</sup>	–4.54 to 4.96	–3.96 to 8.87

\*\* p < 0.01.

\*\*\* p < 0.001.

<sup>a</sup> Town-level SVI was rescaled from the census tract-level SVI based on the total census tract population in a town. The variable in percentage (%) was normalized with respect to town population. A higher SVI score means that the town has higher levels of social vulnerability.

<sup>b</sup> To correspond to the census tract-level SVI data, we merged the towns of Canaan and Norfolk in the county of Litchfield, thus leading to a total of 168 towns. Each quartile had 42 towns.

<sup>c</sup> The vaccination rates are mean values in percentage (%); std. is the standard deviation.

<sup>d</sup> A positive number in the difference estimate means the Q1 rate was higher than Q4 rate.

<sup>e</sup> CI is the confidence interval.

quality health care because of the under-coverage of health insurance, language barriers, and distrust of medical professionals [4].

This short report aims to examine the relationship between social vulnerability and vaccination rates at the town level during the second phase of the vaccine distribution plan when individuals aged 65 and over were eligible. Existing studies identified that low socioeconomic status US counties had lower vaccination rates, and such vaccination disparities enlarged as the vaccine eligibility expanded [5,6]. However, few studies have focused on the elderly population at a small geographic scale, such as the town level. Findings from this report can help reveal the vaccination gap in current

vaccination practices for the elderly population, eventually providing evidence to increase the vaccination coverage and develop equitable health policy.

### Methods

We performed a cross-sectional analysis to employ the association of COVID-19 vaccination rates with selected social vulnerability variables in Connecticut towns (n = 168). We collected town-level vaccination rates for those having received at least one dose from the Connecticut Department of Public Health (CTDPH)

as of March 8, 2021 [1]. The vaccination data included two population groups: those aged 65–74 and those aged 75+. To characterize the towns, we selected eight social vulnerability variables from the 2018 United States Centers for Disease Control (CDC)'s Social Vulnerability Index (SVI). These variables were: below poverty, unemployed, without high school diploma, disability, minority, speaks English less than well, no vehicle, and mobile homes [7].

Specifically, to explore the relationship between social vulnerability and vaccination rates among the elderly population, we examined the vaccination inequities between the towns in the lower SVI quartile (Q1) and those in the upper SVI quartile (Q4). We applied an independent sample t-test to identify if a significant difference existed in the vaccination rates of those aged 65–74 and those aged 75+ between the towns in Q1 and Q4 with respect to each SVI. As a robustness check, we also included all SVIs in the same model using negative binomial regression analysis (see details in Appendix in Supplementary material). The statistical analysis was completed in IBM SPSS 18. The data visualization was completed in ESRI ArcGIS Pro 2.6.

## Results

As of March 8, the vaccination rates in Connecticut were 72.89% and 80.17% for age groups 65–74 and 75+, respectively. Both rates were comparatively higher in the eastern part of the state (Fig. 1). The differences in vaccination rates were statistically significant between Q1 and Q4 with respect to six SVIs (i.e., below poverty, without high school diploma, disability, minority, speaks English less than well, and no vehicle; Table 1). Specifically, the difference in vaccination rate for 65–74 was the largest with respect to lack of education (i.e., difference = 12.91%), and the difference in vaccination rate for 75+ was the largest with respect to no vehicle (i.e., difference = 25.19%). Our secondary analysis employing negative binomial regressions (see details in Appendix in Supplementary material) shows that among the selected SVIs, poverty was negatively correlated with the vaccination rate for 75+ (coefficient:  $-0.00725$ ,  $p < 0.05$ ), and lack of education was negatively correlated with both rates (coefficient for 65–74:  $-0.0251$ ,  $p < 0.001$ ; coefficient for 75+:  $-0.0249$ ,  $p < 0.001$ ).

## Discussion

Past research shows that people residing in socioeconomically disadvantaged areas were more vulnerable to COVID-19 diagnosis and death [3,8]. This report further strengthens that such health inequities existed in the COVID-19 vaccination within the two elderly age groups by comparing the most and least vulnerable towns in Connecticut. It also reveals that the vaccination gap was greater among people aged 75+ than people aged 65–74. The differences in vaccination rates were further correlated with two social vulnerability variables: below poverty and without high school diploma.

Although the US and many other countries have generally adopted an age-based criterion for vaccine prioritization, it has been found that age was negatively correlated with the willingness to receive COVID-19 vaccines [9]. The finding suggests that although the elderly population was prioritized in health care, they were more reluctant to take vaccines and thus more vulnerable to the infection. Such vaccination reluctance within the elderly population needs to be further explored, such as which age group had a wider vaccination gap and which socioeconomic variable most contributed to the vaccination disparities. By revealing the relationships between vaccination rates and SVI-based social vulnerability

variables among people aged 65+ in Connecticut, the study sheds further insights into health policy initiatives to improve vaccination coverage among the elderly in the state and beyond.

Specifically, the study justifies that the socioeconomically disadvantaged elderly groups, especially those aged 75+, had a lower vaccination rate. Also, poverty and under-education contributed most to the lack of vaccination. The relative lack of vaccination among the socially vulnerable elderly group, especially those in poverty and undereducated, could result from the scarcity of transportation resources to access vaccination clinics and a lack of computer literacy in scheduling appointments [10]. Thus, promoting onsite scheduling and increasing at-home vaccination services in the elderly communities in the US and other countries will be a necessity to reducing health inequities.

In this report, we focused on Connecticut towns, and thus the conclusion cannot be extended to another region or at a different geographical scale. The cross-sectional study does not account for individual-level inferences or consider the temporality in vaccination phases. Since the state government administers the COVID-19 vaccination with largely different timelines and eligibility criteria, it is advisable to fully examine the geographical heterogeneity in vaccination policy among states and even across countries in order to justify health inequities.

## Funding

No funding sources.

## Conflict of interest

None declared.

## Ethical approval and consent to publish

The study does not require ethical approval.

## Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.jiph.2021.07.013>.

## References

- [1] CTDPH. COVID-19 vaccine distribution in Connecticut; 2021 [Accessed 6 April 2021] <https://data.ct.gov/stories/s/CoVP-COVID-Vaccine-Distribution-Data/bhcd-4mnnv/>.
- [2] NASEM. Framework for equitable allocation of COVID-19 vaccine. National Academies of Sciences, Engineering, Medicine; 2020.
- [3] Karaye IM, Horney JA. The impact of social vulnerability on COVID-19 in the US: an analysis of spatially varying relationships. *Am J Prev Med* 2020;59(3):317–25.
- [4] Szczepura A. Access to health care for ethnic minority populations. *Postgrad Med J* 2005;81(953):141–7.
- [5] Hughes MM, Wang A, Grossman MK, Pun E, Whiteman A, Deng L, et al. County-level COVID-19 vaccination coverage and social vulnerability—United States, December 14, 2020–March 1, 2021. *Morb Mortal Wkly Rep* 2021;70(12):431.
- [6] Barry V, Dasgupta S, Weller DL, Kriss JL, Cadwell BL, Rose C, et al. Patterns in COVID-19 Vaccination Coverage, by Social Vulnerability and Urbanicity—United States, December 14, 2020?May 1, 2021. *Morb Mortal Wkly Rep* 2021;70(22):818.
- [7] CDC. CDC Social Vulnerability Index; 2021 [Accessed 6 April 2021] <https://www.atsdr.cdc.gov/placeandhealth/svi/index.html>.
- [8] Khazanchi R, Beiter ER, Gondi S, Beckman AL, Bilinski A, Ganguli I. County-level association of social vulnerability with COVID-19 cases and deaths in the USA. *J Gen Intern Med* 2020;35(9):2784–7.
- [9] Kreps S, Prasad S, Brownstein JS, Hswen Y, Garibaldi BT, Zhang B, et al. Factors associated with US adults' likelihood of accepting COVID-19 vaccination. *JAMA Network Open* 2020;3(10), e2025594–e2025594.
- [10] Ramsetty A, Adams C. Impact of the digital divide in the age of COVID-19. *J Am Med Inform Assoc* 2020;27(7):1147–8.