

Cardiomyocytes recruit monocytes upon SARS-CoV-2 infection by secreting CCL2

Liuliu Yang, Benjamin E. Nilsson-Payant, Yuling Han, Fabrice Jaffré, Jiajun Zhu, Pengfei Wang, Tuo Zhang, David Redmond, Sean Houghton, Rasmus Møller, Daisy Hoagland, Lucia Carrau, Shu Horiuchi, Marisa Goff, Jean K. Lim, Yaron Bram, Chanel Richardson, Vasuretha Chandar, Alain Borczuk, Yaoxing Huang, Jenny Xiang, David D. Ho,* Robert E. Schwartz,* Benjamin R. tenOever,* Todd Evans,* and Shuibing Chen**

*Correspondence: dh2994@cumc.columbia.edu (D.D.H.), res2025@med.cornell.edu (R.E.S.), benjamin.tenoever@mssm.edu (B.R.t.), tre2003@med.cornell.edu (T.E.), shc2034@med.cornell.edu (S.C.)
<https://doi.org/10.1016/j.stemcr.2021.09.008>

(Stem Cell Reports 16, 2274–2288; September 14, 2021)

In the experimental procedures of the main text and in the supplemental information, the authors have now corrected the source of Golden Syrian hamsters to “Charles River” instead of “Jackson Laboratory” (the latter of which mistakenly appeared in the original version). The authors apologize for the confusion caused.

