



Published in final edited form as:

Br J Haematol. 2020 December ; 191(5): e124–e128. doi:10.1111/bjh.17090.

A Sardinian founder mutation in glycoprotein Ib platelet subunit beta (*GP1BB*) that impacts thrombocytopenia

Fabio Busonero^{1,†}, Maristella Steri^{1,†}, Valeria Orrù¹, Gabriella Sole¹, Stefania Olla¹, Michele Marongiu¹, Andrea Maschio¹, Carlo Sidore¹, Sandra Lai¹, Antonella Mulas¹, Magdalena Zoledziewska¹, Matteo Floris^{1,2}, Mauro Pala¹, Paola Forabosco¹, Isadora Asunis¹, Maristella Pitzalis¹, Francesca Deidda¹, Marco Masala¹, Cristian Antonio Caria¹, Susanna Barella³, Goncalo R. Abecasis⁴, David Schlessinger⁵, Serena Sanna¹, Edoardo Fiorillo¹, Francesco Cucca^{1,2}

¹Istituto di Ricerca Genetica e Biomedica, Consiglio Nazionale delle Ricerche (CNR), Monserrato (Cagliari)

²Dipartimento di Scienze Biomediche, Università degli Studi di Sassari, Sassari

³Ospedale Pediatrico Microcitemico 'Antonio Cao' (A.O.Brotzu), Cagliari, Italy

⁴Center for Statistical Genetics, University of Michigan, Ann Arbor, MI

⁵Laboratory of Genetics and Genomics, National Institute on Aging, US National Institutes of Health, Baltimore, MD, USA.

Keywords

platelet count; genetic analysis; *GP1BB*; Bernard-Soulier syndrome; platelet glycoproteins; functional studies

fabio.busonero@irgb.cnr.it .

[†]These authors contributed equally to this work.

Author contributions

Fabio Busonero, Maristella Pitzalis and Francesca Deidda collected samples and extracted genomic DNA from blood; Fabio Busonero, Andrea Maschio, Antonella Mulas and Magdalena Zoledziewska performed genotyping; Fabio Busonero and Andrea Maschio, performed sequencing; Valeria Orrù and Edoardo Fiorillo designed flow cytometric panels; Valeria Orrù, Edoardo Fiorillo and Sandra Lai performed cytometric analysis; Maristella Steri, Gabriella Sole, Carlo Sidore, Matteo Floris, Mauro Pala, Paola Forabosco, Marco Masala and Serena Sanna performed statistical analyses; Maristella Steri, Michele Marongiu and Serena Sanna performed bioinformatic analyses; Fabio Busonero and Serena Sanna performed region-specific analysis and selected candidate genes; Stefania Olla performed *in silico* analyses; Isadora Asunis and Cristian Antonio Caria performed functional evaluation; Francesco Cucca, Goncalo R. Abecasis and David Schlessinger provided funds and supervised the work; Susanna Barella provided clinical support; Fabio Busonero, Maristella Steri and Francesco Cucca wrote the paper; Valeria Orrù, Gabriella Sole, Stefania Olla, Andrea Maschio, Carlo Sidore, Matteo Floris, David Schlessinger and Serena Sanna revised the paper. All authors read the paper and contributed to its final form.

Conflict of interest

The authors declare no competing interests.

Web resources

The URLs for data presented herein are as follows: Online Mendelian Inheritance in Man, <https://www.omim.org/>; SardinIA Project, <https://sardinia.irp.nia.nih.gov/>; 1000 Genomes Project data repository, <ftp://ftp.1000genomes.ebi.ac.uk/>; GoNL, Genome of the Netherlands, <http://www.nlgenome.nl/>; GnomAD, <http://gnomad.broadinstitute.org/>; Exome Sequencing Project, <https://esp.gs.washington.edu/drupal/>; NHLBI TOPMed Programme, <https://www.nhlbiwgs.org/>; GWAS catalogue, <https://www.ebi.ac.uk/gwas/>.

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Inherited platelets disorders can be severe, especially after trauma or surgical procedures in some monogenic disorders, as in Bernard–Soulier syndrome [BSS; Mendelian Inheritance in Man (MIM) #231200]. BSS is a rare autosomal recessive macrothrombocytopenia (incidence of about one per million). Its hallmark is a defective adhesion of platelets to the sub-endothelium, resulting from quantitative or qualitative defects in the glycoprotein Ib (GPIb)-IX-V complex, a platelet receptor for von Willebrand Factor (VWF), which is composed of four subunits: GPIb α , GPIb β , GPIX and GPV.¹ Laboratory diagnosis is based on prolonged bleeding time, moderate-to-severe thrombocytopenia (platelet count typically ranges from 20 to $100 \times 10^9/l$), giant platelets and deficient ristocetin-dependent platelet agglutination.² Very little is known about the biochemical and clinical features of heterozygous carriers of the mutations causing BSS, and about the impact in general population individuals of variation in genes encoding the GPIb-IX-V complex when present in heterozygosity. In fact, family members with only one mutated allele are generally asymptomatic, with sub-normal platelet count, slightly enlarged platelets and marginally reduced levels of glycoproteins expression.

Here, to dissect the impact of genetic variability on platelet count, a sequencing-based whole-genome association study was performed in 6528 volunteers included in the SardiNIA general population cohort.³ Six signals were identified (Table SI, Data S1 for description), including a novel nonsynonymous variant [22:19711445:C/T; minor allele frequency (MAF) = 0.0045; $P = 1.172 \times 10^{-16}$], mapping in the second exon (c.C79T, p.P27S) of the *GPIBB* gene (Figure S1). Completely independent of previously reported associations in the same genomic region (Data S1), p.P27S is Sardinian-specific, being completely missing in large sequencing datasets such as 1000 Genomes Project,⁴ Genome of the Netherlands (GoNL) project,⁵ Genome Aggregation Database (gnomAD),⁶ the Exome Sequencing Project in the National Heart, Lung, and Blood Institute's (NHLBI's) Trans-Omics for Precision Medicine (TOPMed) programme.⁷ No homozygous and 57 carriers for the rare 22:19711445-T allele were found. The platelet count in wild-type homozygous was $242.87 \pm 117.05 \times 10^9/l$ (mean $\pm 1.96 \times SD$), whereas in p.P27S carriers was $174.17 \pm 91.51 \times 10^9/l$, corresponding to a reduction of $70.13 \times 10^9/l$ for each copy of the minor allele (Fig 1A). With this large effect, the novel founder mutation explains ~1.05% of phenotypic variance for platelet count, representing the largest phenotypic effect among all the independent variants reported to date in the genome-wide association studies (GWAS) Catalogue (Tables SII and SIII).⁸ Moreover, in a subset of 2000 individuals, whose mean platelet volume was measured, this variant was associated with notably larger platelets ($P = 2.13 \times 10^{-10}$), consistently with evidence of morphologically enlarged platelets in patients with BSS (Fig 1B). To assess platelet functionality, a seven-colour flow cytometry panel (Table SIV) was set up in 24 of 57 p.P27S carriers (42.1%) and in an equal number of matched unrelated controls. Monoclonal antibodies directed against the GPIIb-IIIa complex (CD41a and CD61), and the VWF receptor complex (CD42a and CD42b) were used to investigate the basal receptor expression in resting platelets. The p.P27S carriers showed increased levels of GPIIb (CD41a, +22.36%; $P = 1.61 \times 10^{-4}$; $N = 48$) and GPIIIa (CD61, +16.20%; $P = 6.61 \times 10^{-4}$; $N = 48$) a typical finding in the presence of enlarged platelets (Figure S2). The expression of GPIX and GPIb α glycoproteins and their correct assembly into the GPIb-IX-V complex are known to be impaired by a defective

GPIIb β peptide⁹. Indeed, despite carrying only one mutated allele, the p.P27S heterozygous showed appreciably lower basal expression levels of both GPIX (-24.69% , $P=2.66 \times 10^{-6}$, $N=46$; Fig 1C) and GPIba (-26.51% , $P=3.66 \times 10^{-8}$, $N=48$; Fig 1D), and consequently less of the entire complex, compared to controls. This is far more than the normal expression levels of GPIX and GPIba in carriers of other missense mutations in *GPIBB*, as recently reported.¹⁰ Pre-activation and reactivity changes in p.P27S platelets were investigated after exposure to the agonist adenosine diphosphate (ADP). Indeed, activated α IIb β 3 was prominently induced in the p.P27S carriers, as shown by the extent of procaspase-activating compound 1 (PAC-1) binding to resting and activated platelets ($+41.94\%$, $P=4.84 \times 10^{-3}$, $N=48$, Fig 1E). Notably, no variation in the response of platelets after ADP stimulation was recently reported in patients with BSS and carriers.⁹ Remarkably, platelet reactivity turned out to be differentially regulated: no changes were observed in surface exposure of neo P-selectin (CD62P, $+35.22\%$, $P=0.138$, $N=48$; Fig 1F) and neo granulophysin (CD63, $+1.86\%$, $P=0.658$, $N=46$; Fig 1G), markers of granule content release. The unique functional effects of the p.P27S led us to examine its possible consequences on the molecular structure and conformational changes of GPIIb β by molecular modelling analysis based on the X-ray crystal structure.¹¹ Proline–Serine substitution falls in the leucine-rich repeat *N*-terminal (LRRNT) domain of the 206 amino acid long protein encoded by *GPIBB* (Fig 2A, B). Proline residues are expected to be disruptive of structure; and indeed, in that highly conserved region and close to cysteine residues involved in the Cys26–Cys32 disulphide bridge, p.P27S could thus modify the stability and consequently the conformation of GPIIb β . To test this hypothesis, we first performed *in silico* molecular dynamic simulations, observing an increased conformational mobility of the amino acid backbone close to p.P27S (Figure S3), suggesting the instability of the GPIIb β glycoprotein in accordance with the observed reduction in the expression of GPIX and GPIba. Strikingly, a greater fluctuation of the amino acids in loop 2 of the p.P27S protein was also recorded, as indicated by root-mean-square-fluctuation (Fig 2C).

In summary, all typical findings of macrothrombocytopenias (i.e. BSS) were observed in the p.P27S obligate carriers characterised in the present study: low levels of large platelets and low expression of GPIX and GPIba glycoproteins, as shown by flow cytometry. As one might anticipate, the most severe cases are caused by deletions and nonsense mutations, but some missense mutations are disabling enough to be clinically significant. In one of the reported cases,¹² a charge difference is introduced (p.Asn89Asp); in the other,¹³ as in this case, the Proline residue is replaced (p.Pro27Leu), which is expected to disrupt secondary structure in the protein. That p.P27S influences conformational changes and stability of GPIIb β , in turn affecting GPIb–IX–V complex function, is further clearly supported by the *in silico* molecular dynamic analyses. Noteworthy, a critical interaction of GPIIb β with GPIX involves *N*-terminal residues 15 through 32 of GPIIb β , precisely including Proline 27.¹⁴ According to Hardy–Weinberg expectation, at least four p.P27S homozygous individuals, most likely with BSS, are expected in Sardinia, but none have been reported to date: this may suggest that BSS is likely underdiagnosed in Sardinia, consistent with other reports.¹⁵ Thus, clinicians should be aware of the novel p.P27S mutation in the molecular characterisation of Sardinian-origin patients with a clinical picture of platelet macrocytosis and platelet count of $<100 \times 10^9/l$.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank all the volunteers who generously participated in this study; we are grateful to Mr Mario Lovicu and Mr Nazario Olla for the logistic support provided and helpful suggestions. Supported by contracts N01-AG-1-2109 and HHSN2712011 00005C from the Intramural Research Program of the National Institute on Aging, National Institutes of Health (NIH).

References

1. Savoia A, Pastore A, De Rocco D, Civaschi E, Di Stazio M, Bottega R, et al. Clinical and genetic aspects of Bernard-Soulier syndrome: searching for genotype/phenotype correlations. *Haematologica*. 2011;96:417–23. [PubMed: 21173099]
2. Berndt MC, Andrews RK. Bernard-Soulier syndrome. *Haematologica*. 2011;96:355–9. [PubMed: 21357716]
3. Sidore C, Busonero F, Maschio A, Porcu E, Naitza S, Zoledziewska M, et al. Genome sequencing elucidates Sardinian genetic architecture and augments association analyses for lipid and blood inflammatory markers. *Nat Genet*. 2015;47:1272–81. [PubMed: 26366554]
4. Genomes Project Consortium, Abecasis GR, Auton A, Brooks LD, DePristo MA, Durbin RM, et al. An integrated map of genetic variation from 1,092 human genomes. *Nature*. 2012;491:56–65. [PubMed: 23128226]
5. Boomsma DI, Wijmenga C, Slagboom EP, Swertz MA, Karssen LC, Abdellaoui A, et al. The Genome of the Netherlands: design, and project goals. *Eur J Hum Genet*. 2014;22:221–7. [PubMed: 23714750]
6. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al. Variation across 141,456 human exomes and genomes reveals the spectrum of loss-of-function intolerance across human protein-coding genes. *bioRxiv*. 2019:531210. 10.1101/531210.
7. Taliun D, Harris DN, Kessler MD, Carlson J, Szpiech ZA, Torres R, et al. Sequencing of 53,831 diverse genomes from the NHLBI TOPMed Program. *BioRxiv*, 06 3 2019. 2019. 10.1101/563866. PPR:PPR72371.
8. MacArthur J, Bowler E, Cerezo M, Gil L, Hall P, Hastings E, et al. The new NHGRI-EBI Catalog of published genome-wide association studies (GWAS Catalog). *Nucleic Acids Res*. 2017;45:D896–901. [PubMed: 27899670]
9. Hadjkacem B, Elleuch H, Gargouri J, Gargouri A. Bernard-Soulier syndrome: novel nonsense mutation in GPIIb gene affecting GPIIb-IX complex expression. *Ann. Hematol* 2009;88:465–72. [PubMed: 18825380]
10. Bragadottir G, Birgisdottir ER, Gudmundsdottir BR, Hilmarisdottir B, Vidarsson B, Magnusson MK, et al. Clinical phenotype in heterozygote and biallelic Bernard-Soulier syndrome—a case control study. *Am J Hematol*. 2015;90:149–55. [PubMed: 25370924]
11. McEwan PA, Yang W, Carr KH, et al. Quaternary organization of GPIIb-IX complex and insights into Bernard-Soulier syndrome revealed by the structures of GPIIb β and a GPIIb β /GPIX chimera. *Blood*. 2011;118:5292–301. [PubMed: 21908432]
12. Fiore M, De Thoré C, Ranjatoelina HR, Baas MJ, Jacquemont ML, Dreyfus M, et al. High prevalence of the natural Asn89Asp mutation in the GPIIb gene associated with Bernard-Soulier syndrome in French patients from the genetic isolate of Reunion Island. *Br J Haematol*. 2020;189:e67–e71. [PubMed: 31997307]
13. Bastida JM, Lozano ML, Benito R, Janusz K, Palma-Barqueros V, Del Rey M, et al. Introducing high-throughput sequencing into mainstream genetic diagnosis practice in inherited platelet disorders. *Haematologica*. 2018;103:148–62. [PubMed: 28983057]

14. Kenny D, Morateck PA, Montgomery RR. The cysteine knot of platelet glycoprotein Ib (GPIb-beta) is critical for the interaction of GPIb-beta with GPIX. *Blood*. 2002;99:4428–33. [PubMed: 12036872]
15. Noris P, Balduini CL. Inherited thrombocytopenias in the era of personalized medicine. *Haematologica*. 2015;100:145–8. [PubMed: 25638803]
16. Savoia A, Kunishima S, De Rocco D, Zieger B, Rand ML, Pujol-Moix N, et al. Spectrum of the mutations in Bernard-Soulier syndrome. *Hum Mutat*. 2014;35:1033–45. [PubMed: 24934643]
17. Sivapalaratnam S, Westbury SK, Stephens JC, Greene D, Downes K, Kelly AM, et al. Rare variants in GP1BB are responsible for autosomal dominant macrothrombocytopenia. *Blood*. 2017;129:520–4. [PubMed: 28064200]
18. Qiao J, Davis AK, Morel-Kopp MC, Ward CM, Gardiner EE, Andrews RK. Low levels of CD9 coincidental with a novel nonsense mutation in glycoprotein Ib β in a patient with Bernard-Soulier syndrome. *Ann Hematol*. 2015;94:2069–71. [PubMed: 26275786]
19. Kunishima S, Naoe T, Kamiya T, Saito H. Novel heterozygous missense mutation in the platelet glycoprotein Ib beta gene associated with isolated giant platelet disorder. *Am J Hematol*. 2001;68:249–55. [PubMed: 11754414]
20. Ferrari S, Lombardi AM, Cortella I, Businaro MA, Bertomoro A, Di Pasquale I, et al. New heterozygous variant in GP1BB gene is responsible for an inherited form of macrothrombocytopenia. *Br J Haematol*. 2019;184:855–8. [PubMed: 29527674]




Fig 1. Effects of chr22:19711445 genotype on platelet-related phenotypes. (A) Platelet (PLT) count distribution stratified on 57 heterozygous carriers and 6471 homozygous wild types. (B) Mean platelet volume (MPV) distribution stratified on 28 heterozygous carriers and 1972 homozygous wild types. Basal expression levels of the main GPIb-IX-V receptor glycoproteins on resting platelets: (C) GPIX on 23 carriers and 23 controls, and (D) GPIIb on 24 carriers and 24 controls. (E–G) Expression levels of the most relevant platelet activation-dependent markers (activated $\alpha IIb\beta 3$, P-selectin and granulophysin), in basal conditions and after stimulation with ADP. Violin plots represent the distribution of the data; the boxplots inside report the median value as a dot, the interquartile range (IQR) as a box, the 1st quartile -1.5 IQR and the 3rd quartile $+1.5$ IQR as whiskers. ** $P < 0.01$, *** $P < 0.001$. MFI, median fluorescence intensity.




Fig 2. GPIIb/IIIa amino acid sequence with BSS-causing mutations and molecular modelling analyses. (A) Positions of the mutations within the coding regions of platelet glycoprotein (GP)Ib/IIIa according to National Center for Biotechnology Information (NCBI) Reference Sequence, NP_000398.1. The different domains are indicated with different patterns. Different types of mutation are colour-coded; highlighted in yellow is the Proline27 to Serine27 substitution (P27S) described here first. Known mutations were obtained from: Savoia *et al.*, 2014¹⁶; Sivapalaratnam *et al.*, 2017¹⁷; Bragadottir *et al.*, 2015¹⁰; Qiao *et al.*, 2015¹⁸; Kunishima *et al.*, 2001¹⁹; Ferrari *et al.*, 2018²⁰; and Bastida *et al.*, 2018¹³. (B) Three-dimensional structure of GPIIb/IIIa sequence, colour-coded according to the schematic representation in (A). (C) X-ray structure of protein (26–143 aa code 3RFE) showing the impact of p.P27S on GPIIb/IIIa glycoprotein conformation; in particular, the superposition between the first (*teal*) and last (*yellow*) frame of the molecular dynamics for the p.P27S protein (*left*), and the superposition between the first (*pink*) and last (*green*) frame for the wild-type (WT) protein (*right*) are reported.