



Published in final edited form as:

Trop Med Int Health. 2021 October ; 26(10): 1248–1255. doi:10.1111/tmi.13647.

Catastrophic costs among tuberculosis affected households in Zimbabwe: a national health facility-based survey

Collins Timire, BSc, MPH^{1,2,3}, Mkhokheli Ngwenya, MBChB, MPH⁴, Joconiah Chirenda, MBChB, MPH, PhD⁵, John Z Metcalfe, MD, MPH, PhD⁶, Katharina Kranzer, MSc, MSc, PhD³, Debora Pedrazzoli, MSc, PhD^{7,8}, Kudakwashe C Takarinda, MSc, PhD^{1,2}, Peter Nguhiu, BSc, MSc⁹, Geshem Madzingaidzo, MSc¹, Kwenzikweyinkosi Ndlovu, MSc¹, Tawanda Mapuranga, MBChB¹, Morna Cornell, MPH, PhD¹⁰, Charles Sandy, MBChB, MPH¹

¹Ministry of Health and Child Care, AIDS & TB Control Programme, Harare, Zimbabwe

²International Union Against Tuberculosis and Lung Disease (The Union), Paris, France

³Faculty of Infectious & Tropical Diseases, London School of Hygiene & Tropical Medicine (LSHTM), UK

⁴World Health Organisation, Zimbabwe Country Office, Harare, Zimbabwe

⁵University of Zimbabwe, College of Health Sciences, Harare, Zimbabwe

⁶Division of Pulmonary and Critical Care Medicine, Zuckerberg San Francisco General Hospital and Trauma Center, University of California, San Francisco, CA, USA

⁷Faculty of Epidemiology & Population Health, LSHTM, UK

⁸World Health Organisation, Geneva, Switzerland

⁹Health Economics Research Unit, Kenya Medical Research Institute (KEMRI), Wellcome Trust Research Programme, Nairobi, Kenya

¹⁰Centre for Infectious Disease Epidemiology and Research, School of Public Health and Family Medicine, University of Cape Town, South Africa

Abstract

Corresponding Author: Charles Sandy, Ministry of Health and Child Care; National TB Control Programme, Harare, Zimbabwe, 5th Floor Kaguvi Building; Cnr 4th/Central Avenue, Harare, Zimbabwe; Mobile: (+263) 774 358476; dr.c.sandy@gmail.com.

Authors' contributions

Drafted the study protocol: CT, MN,CS,KN,TM, JC,GM

Collected data: CT,MN,CS,KN,TM,GM

Analysed and interpreted data: CT,MN,CS,KN,TM,JC,PN,DP,KK,MC,KCT

Drafted the paper:CT,CS,MC

Critically reviewed the draft paper: KK,DP,JZM,JC,PN,MC,KCT

All authors read and approved the final version of the manuscript

Ethics approval and consent to participate

Written informed consent and/or assent was obtained from patients and their guardians prior the interviews. Ethics approval was obtained from the Medical Research Council of Zimbabwe (MRCZ/A/2290).

Consent for publication

Not applicable

Competing interests

The authors declare they have no competing interests

Objectives: To determine the incidence and major drivers of catastrophic costs among TB-affected households in Zimbabwe.

Methods: We conducted a nationally representative health facility-based survey with random cluster sampling among consecutively enrolled drug susceptible (DS-TB) and drug resistant TB (DR-TB) patients. Costs incurred and income lost due to TB illness were captured using an interviewer administered standardised questionnaire. We used multivariable logistic regression to determine the risk factors for experiencing catastrophic costs.

Results: A total of 841 patients were enrolled and were weighted to 900 during data analysis. There were 500 (56%) males and 46 (6%) DR-TB patients. Thirty-five (72%) DR-TB patients were HIV co-infected. Overall, 80% (95% CI:77-82) of TB patients and their households experienced catastrophic costs. The major cost drivers pre-TB diagnosis were direct medical costs. Nutritional supplements were the major cost driver post-TB diagnosis, with a median cost US\$360 (IQR: 240-600). Post-TB median diagnosis costs were three-times higher among DR-TB (US\$1,659 [653-2,787]) versus drug DS-TB affected households (US\$537 [204-1,134]). Income loss was five-times higher among DR-TB versus DS-TB patients. In multivariable analysis, household wealth was the only covariate that remained significantly associated with catastrophic costs: the poorest households had sixteen times the odds of incurring catastrophic costs compared to wealthiest households (adjusted odds ratio [aOR]:15.7 95% CI:7.5-33.1).

Conclusion: The majority of TB-affected households, especially those affected by DR-TB experienced catastrophic costs. Since the major cost drivers fall outside the healthcare system, multi-sectoral approaches to TB control and linking TB patients to social protection may reduce catastrophic costs.

Keywords

patient cost; tuberculosis; financial protection; Social protection; Zimbabwe; universal health coverage

Introduction

In 2019 approximately 10 million people developed tuberculosis (TB) globally and 1.6 million of them died, making TB one of the top ten killer diseases worldwide and the leading cause from a single infectious agent. (1) In the same year, Africa accounted for 25% of the global TB notifications. (1) The prevalence of human immune-deficiency virus (HIV) is high in countries of sub-Saharan Africa (SSA), and many of them have generalised HIV epidemics. Zimbabwe, a low-and middle income country (LMIC) in SSA is among the high TB, TB/HIV and multi-drug resistant TB (MDR-TB) burdened countries. In 2019, Zimbabwe had an estimated TB incidence of 199 per 100,000 population (95% CI: 147–258). The prevalence of MDR-TB/rifampicin resistant TB is around 3.9% among new and 14% among previously treated TB cases. (2,3)

Traditionally TB control has been the responsibility of the health care sector with a focus on TB diagnoses and curative treatment. Until recently, socioeconomic determinants of TB disease have received limited attention. Addressing these determinants has not been an integral part of TB control and would require a multi-sectoral response. The End TB

Strategy has encouraged thinking beyond the biomedical model by including one milestone specifically relating to costs (no TB patients or households should experience catastrophic costs by 2020). (4) Total TB-related costs are defined as catastrophic if they exceed 20% of a household's annual income.

Tuberculosis-related catastrophic costs are a public health challenge requiring urgent attention. (5) Such costs may plunge households into poverty and financial catastrophes. (6) For this reason, countries were encouraged to set baseline measurements of incident catastrophic costs by 2020. The measurements are based on three types of costs: direct medical, direct non-medical and indirect costs such as income loss. Direct medical costs include money spent on consultations, laboratory tests and hospitalisation. Direct non-medical costs are money spent on transport and food during health seeking. Income loss is money foregone by the patient or carers during illness.

Global efforts to ameliorate TB-related catastrophic costs lie in i) provision of free TB treatment ii) decentralisation of TB services to ensure equity of access and iii) advocacy for social protection and universal health coverage by the United Nations, national and international stakeholders. Despite interventions aimed at cushioning TB patients against direct medical costs, surveys in Africa and Asia have shown high incidence of TB-related catastrophic costs especially among i) drug resistant TB (DR-TB) patients, ii) poorest households, iii) cases where the patients were breadwinners and iv) those co-infected with HIV.(7–10)

In 2016, Zimbabwe embraced the End TB targets of eliminating TB-related catastrophic costs by 2020. The national TB control programme (NTP) has decentralised TB services, provided cash transfers to DR-TB patients and adopted active case finding to detect TB cases early. However, the country had no baseline measure of TB-related catastrophic costs or the major drivers of such costs. We therefore aimed to determine among TB-affected households in Zimbabwe, the incidence of catastrophic costs and its risk factors, and the major drivers of costs incurred because of accessing TB-related services.

Methods

Study design

A nationally representative, health facility-based survey with random cluster sampling among TB patients across Zimbabwe.

Setting

Zimbabwe is a southern African country with an estimated population of 14.9 million in 2020. (11) Classified as a low income country, it has suffered from an economic and humanitarian crisis for much of the last decade. Zimbabwe belongs to the 14 countries with a triple-burden of TB, TB/HIV and multi-drug resistant TB. (12)

Study population

Patients of all age groups who were on treatment for drug susceptible TB (DS-TB) or DR-TB for any type of TB (pulmonary, extra-pulmonary and/or disseminated), and who attended

their scheduled appointments within the sampled health facilities from 23 July to 31 August 2018 were eligible for inclusion in the study. Patients were recruited consecutively when they attended their scheduled appointments. The study included patients who had been on treatment for at least two weeks in their current treatment phase (intensive or continuation phase). For patients who had been on treatment for <2 weeks at the time the facility was visited, interviews were rescheduled to a time when the patients had been on treatment for at least two weeks. Patients who had completed treatment were ineligible for inclusion in the study. This was to minimise recall bias and to ease logistics during recruitment.

Study procedures

Sample size calculation, sampling and patient enrolment—The sample was based on 26,677 TB patients notified in Zimbabwe in 2016. We assumed an absolute precision of 5% and a *priori* estimate of 50% for the incidence of households experiencing TB-related catastrophic costs. We used the standard formula for sample size calculation for a cluster sampled TB prevalence survey.⁽¹³⁾ After factoring a design effect of 2.0, the sample size was 780 patients across 60 clusters, each contributing 13 patients. The sample size was adjusted to 900 assuming a non-response rate of 10%.

Cluster sampling was used to select health facilities (clusters). First, a list of clusters and their corresponding 2016 notifications was compiled. The number of TB notifications per cluster was used as a proxy for the size of clusters. Clusters that notified <10 patients were merged with adjacent clusters. Second, cumulative notifications were compiled and 60 clusters were selected by a probability proportional to size sampling method using a randomly defined starting point and sampling interval.

The study questionnaire was adapted from the WHO generic instrument and was created in CSPro® (Census Bureau, USA). Data collectors, one per facility, were trained by the NTP and partner organisations. They comprised TB focal nurses and Environmental Health Technicians. All clinical and economic data were collected for the respective phase only. In case of minors (<18 years), costs were obtained from their parents and legal guardians. Interviews were conducted after obtaining assent from minors and consent from their parents and/or legal guardians. While indirect costs like loss of income were not applicable for this group, costs for diagnosis, treatment and food were obtained from the guardians. Data on dissavings (use of savings and sale of assets) were also obtained from guardians. At the end of each day, data from tablets were synchronised electronically with a central server at central level. Zimbabwe National Statistics Agency staff monitored the server and promptly highlighted errors for clarification. Periodic data quality checks and support visits were conducted by the steering committee. A WhatsApp group for data collectors and steering committee members was created to aid in addressing operational challenges, mostly related to patient recruitment and syncing electronic records in real-time.

Data variables, source of data and data collection

Socio-demographic and clinical data (age, HIV status, type of TB patient) were extracted from treatment registers and patient treatment booklets prior to the interview. Data on hospital visits, costs incurred, household assets and coping strategies were collected by

trained data collectors during face-to-face interviews with patients. All the interviews were conducted in separate rooms within health facilities to ensure confidentiality.

Data analysis

Anonymised data were exported to Stata version 15.0 (StataCorp, College Station, TX, USA) for cleaning and analysis. Categorical variables were summarised using frequencies (proportions), while continuous variables were summarised using medians and inter-quartile ranges (IQR) stratified by DR status. We summed up the direct medical costs (consultations fees, laboratory tests) and direct non-medical costs (transport, food). Costs that were in South African Rand were converted to USD using the prevailing conversion rate obtained from the Oanda currency converter (<http://www.oanda.com>). Productivity losses due to TB treatment were estimated using the output approach, where the difference in monthly income before and after TB diagnosis was extrapolated over the treatment period. Household income was based on self-reports. A sensitivity analysis of indirect costs estimation was done using valuation of the time lost by the patient in each phase of treatment. To estimate patient costs for the entire TB episode, including costs for all phases of treatment, we extrapolated costs based on data from patients in other phases of illness. We used the approach recommended by WHO, whereby we replaced missing cost data with median costs of the phase of illness among those in that phase with available data. Comparisons between categorical variables were done using the chi-square test. Multivariable logistic regression was used to determine risk factors for patients experiencing TB-related catastrophic costs after adjusting for sex, age, DR status; treatment phase, HIV status, breadwinner status, household income and location of health facility. The level of significance was set at $P < 0.05$.

Results

A total of 860 patients (96% of the target sample size) were reached and consented (Figure 1). Of these, 19 records were excluded from analysis for being on treatment for <14 days and not being able to reschedule an appointment (17) and for failure to complete the interview (two). Overall, the response rate was 841 (93%). The figure was 900 after factoring in non-response weights.

Of the 900 patients, 851 (94%) had DS-TB (Table 1). The mean age was 36.9 years and 500 (56%) were men. A greater proportion of DR-TB than DS-TB patients were in the continuation phase (66% vs 56%) and were HIV-positive (72% vs 61%). Almost all patients were new, rather than retreatment. DR-TB patients were more likely to live in urban locations than DS-TB patients (69% vs 59%).

The proportion of households who experienced TB-related catastrophic costs was 80% (95% CI: 77-82) (Figure 2). The incidence of TB-related catastrophic costs was 90% among DR-TB patients and 79% among DS-TB patients, $P=0.06$ (Table 2). Overall, 95% of patients in the poorest income quintile experienced TB-related catastrophic costs. Income quintile was strongly associated with catastrophic costs in a dose-response relationship after adjusting for other variables. Compared to the wealthiest group, the poorest households had higher odds of incurring TB-related catastrophic costs, (adjusted odds ratio (aOR): 15.7 [7.5-33.1]). Sex, age, type of TB, treatment phase, treatment delay (four weeks), HIV status, being a

breadwinner and location of health facility were not associated with TB-related catastrophic costs in either univariable or multivariable analysis.

The major cost drivers in the pre-diagnosis phase were direct medical costs with a median of US\$25 (IQR:6-58) followed by food US\$18 (IQR:2.2-27) (Table 3). During the pre-diagnostic phase, DS-TB patients incurred higher direct costs than DR-TB patients (median US\$54 vs. US\$35). The median direct costs post-TB diagnosis were US\$555 (IQR:220-600), and three times higher among DR-TB (US\$1659 [IQR:653–2787]) vs DS-TB patients US\$537 (IQR:204–1134). The major cost drivers post-TB diagnosis were nutritional supplements (US\$360 [IQR:240-528]) vs (960 [IQR:640-1680]), for DR-TB vs DS-TB patients respectively, medical and travel costs. Median travel costs were five times higher among DR-TB compared to DS-TB patients.

Overall, the total median cost per TB episode was US\$1,247 (IQR: 545-2405) (Table 3). The median total costs incurred by DR-TB patients were three times higher than those of DS-TB patients. Costs as a proportion of total costs were: non-medical costs (51%); indirect costs (36%) and medical costs (13%). During treatment, patients lost productive time with a median value of US\$249.1 (IQR: 128.3-486.1). The losses were greater for DR-TB patients with a median of US\$1,827.2 (IQR: 432.6-3,819.6) as compared to DS-TB patients US\$238.3 (IQR: 113.3–441). Only 1.2% of households reported that they had received social protection.

Discussion

We found a high incidence of catastrophic costs among TB-affected households. The poorest households experienced the highest incidence of catastrophic costs. A higher proportion of DR-TB households incurred catastrophic costs compared to DS-TB household, albeit not reaching statistical significance. The major drivers of catastrophic costs were direct non-medical and indirect costs related to productivity loss. Indirect costs were five-times higher among DR-TB compared to DS-TB patients.

Previous surveys done in Africa and Asia have shown that TB patients incur huge catastrophic costs despite free TB treatment.(7,8,10,13) Our study provides additional evidence to this finding. However, catastrophic costs were not homogeneous among the different groups as the poorest households were disproportionately affected and risk of catastrophic costs increased as wealth quintile decreased. (6–8,10) This makes intuitive sense, and may be attributed to reduced resilience to external shocks such as TB. Unlike studies done elsewhere, we did not find a significant difference in catastrophic costs by DR status. We had low numbers of DR-TB patients, and our study may not have been sufficiently powered to detect the difference. Also, in our context even DS-TB patients experienced higher catastrophic costs than overall costs reported from other low-and middle income countries. (8, 9,14) High catastrophic costs may negatively impact on both access and adherence to TB treatment.

The major drivers of catastrophic costs lay outside the healthcare sector, a consistent finding with studies wherein non-medical costs were reported to account for up to 80% of

universal health coverage. A multi-sectoral approach to TB control holds great promise to reducing catastrophic costs due to TB in Zimbabwe.

Acknowledgements

We would like to thank the participants who took part in this survey, the survey steering committee members and staff who collected the data.

Funding

This study was funded by the US Agency for International Development (USAID) Challenge TB through the World Health Organisation, Zimbabwe Country Office (Grant number *AID-OAA-A-14-00029*). Publication costs for this article were supported by the U.S. National Institutes of Health's National Institute of Allergy and Infectious Diseases, under award Number *U01AI069924*. The content is solely the responsibility of the authors and does not necessarily represent the official views of the funders.

Availability of data and materials

The datasets are available from the corresponding author on reasonable request.

List of abbreviations

TB	Tuberculosis
HIV	human immune-deficiency virus
DR-TB	drug resistant TB
DS-TB	drug susceptible TB
MDR-TB	multi-drug resistant TB
OOP	out of pocket
IQR	inter-quartile range
aOR	adjusted odds ratio
WHO	World Health Organisation
NTP	National Tuberculosis Control Programme
SD	standard deviation
US\$	United States Dollar

References

1. World Health Organisation. Global Tuberculosis Report 2020 [Internet]. Geneva, Switzerland; 2020. Available from: <https://www.who.int/publications/i/item/9789240013131>
2. Timire C, Metcalfe JZ, Chirenda J, Scholten JN, Manyame-murwira B, Ngwenya M, et al. International Journal of Infectious Diseases Prevalence of drug-resistant tuberculosis in Zimbabwe: A health facility-based cross-sectional survey. *Int J Infect Dis* [Internet]. 2019;87:119–25. Available from: 10.1016/j.ijid.2019.07.021 [PubMed: 31357057]
3. World Health Organisation. Global Tuberculosis Report 2018 [Internet]. Geneva, Switzerland; 2018. Available from: WHO/CDS/TB/2018.20

4. STOP TB Partnership. The paradigm shift 2016-2020: Global plan to End TB [Internet]. Geneva, Switzerland; 2015. Available from: <http://www.stoptb.org/global/plan/plan2/>
5. Wingfield T, Tovar MA, Huff D, Boccia D, Montoya R, Ramos E, et al. The economic effects of supporting tuberculosis-affected households in Peru. *Eur Radiol* [Internet]. 2016;48:1396–410. Available from: 10.1183/13993003.00066-2016
6. Tanimura T, Jaramillo E, Weil D, Raviglione M, Lönnroth K. Financial burden for tuberculosis patients in low- and middle-income countries: a systematic review. *Eur Respir J*. 2014;43(6): 1763–75. [PubMed: 24525439]
7. Fuady A, Houweling TAJ, Mansyur M, Richardus JH. Catastrophic total costs in tuberculosis-affected households and their determinants since Indonesia ' s implementation of universal health coverage. *Infect Dis Poverty*. 2018;7(3): 1–14. [PubMed: 29335021]
8. Ministry of Health Kenya. The First Kenya Tuberculosis Patient Cost Survey, 2017. Nairobi, Kenya; 2017.
9. Pedrazzoli D, Siroka A, Boccia D, Bonsu F, Nartey K, Houben R, et al. How affordable is TB care? Findings from a nationwide TB patient cost survey in Ghana. *Trop Med Int Heal*. 2018;23(8):870–8.
10. Ukwaja KN, Alobu I, Abimbola S, Hopewell PC. Household catastrophic payments for tuberculosis care in Nigeria : incidence , determinants , and policy implications for universal health coverage. *Infect Dis Poverty* [Internet]. 2013;2(21):1–9. Available from: <http://www.idpjournals.com/content/2/1/21> [PubMed: 23849433]
11. Worldometer. Zimbabwe population (live) [Internet]. 2020 [cited 2020 Sep 21]. Available from: <https://www.worldometers.info/world-population/zimbabwe-population/>
12. World Health Organisation. Global Tuberculosis Report 2019 [Internet]. Geneva, Switzerland; 2019. Available from: WHO/CDS/TB/2019.15
13. World Health Organization. Tuberculosis patient cost surveys: a handbook. Geneva, Switzerland; 2018.
14. Nhung NV, Hoa NB, Anh NT, Anh LTN, Siroka A, Lönnroth K, et al. Measuring catastrophic costs due to tuberculosis in Viet Nam. *Int J Tuberc Lung Dis* [Internet]. 2018;22(9):983–90. Available from: 10.5588/ijtld.17.0859. [PubMed: 30092862]
15. Fuady A, Houweling TAJ, Mansyur M, Burhan E, Richardus JH. Effect of financial support on reducing the incidence of catastrophic costs among tuberculosis-affected households in Indonesia : eight simulated scenarios. *Infect Dis Poverty* [Internet]. 2019;8(10):1–14. Available from: 10.1186/s40249-019-0519-7 [PubMed: 30626428]
16. Rudgard WE, das Chagas NS, Gayoso R, Barreto ML, Boccia D, Smeeth L, et al. Uptake of governmental social protection and financial hardship during drug-,resistant tuberculosis treatment in Rio de Janeiro, Brazil. *Eur Respir J* [Internet]. 2018;51(100274): 1–5. Available from: 10.1183/13993003.00274-2018
17. Ministry of Health and Child Care. Availability and coverage of social protection services for TB patients in Zimbabwe. Harare, Zimbabwe; 2019.




Figure 1:
Flow of patients who were enrolled in the Zimbabwe patient cost survey, 2018

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 2: Proportion of tuberculosis-affected households who experienced TB-related catastrophic costs during the Zimbabwe tuberculosis patient cost survey 2018

Table 1.

Socio-demographic and clinical profile of patients with DR-TB and DS-TB who were enrolled in the Zimbabwe patient cost survey, 2018

Characteristic	DR-TB		DS-TB		Total	
	N	(%)	n	(%)	n	(%)
	49	(6)	851	(94)	900 [‡]	
<i>Sex</i>						
Male	28	(56)	472	(56)	500	(56)
Female	21	(44)	379	(44)	400	(44)
<i>Age group (years)</i>						
<15	0	(0)	52	(6)	52	(6)
15-24	4	(8)	99	(12)	103	(12)
25-34	21	(42)	209	(24)	230	(26)
35-44	15	(31)	277	(33)	292	(33)
45-54	7	(14)	124	(15)	131	(15)
55-64	0	(0)	53	(6)	53	(6)
65	2	(5)	37	(4)	39	(4)
Mean (SD)	36.4	(11.9)	36.9	(14.8)	36.9	(14.7)
<i>Treatment phase</i>						
Intensive	17	(34)	376	(44)	392	(44)
Continuation	32	(66)	475	(56)	508	(56)
<i>HIV Status</i>						
Positive	35	(72)	522	(61)	557	(62)
Negative	14	(28)	321	(38)	335	(37)
Unknown	0	(0)	8	(1)	8	(1)
<i>Type of TB patient</i>						
New	45	(91)	796	(94)	841	(94)
Retreatment	4	(9)	55	(6)	59	(6)
<i>Facility location</i>						
Urban	34	(69)	506	(59)	540	(60)
Rural	15	(31)	345	(41)	360	(40)

[‡]=weighted sample size;

DR-TB=drug resistant TB; *DS-TB*=drug susceptible TB; *SD*=Standard deviation; HIV = Human Immunodeficiency Virus

Table 2:

Factors associated with catastrophic costs among tuberculosis-affected households enrolled in the Zimbabwe tuberculosis patient cost survey, 2018

Characteristic	Total	Number (%) who incurred catastrophic costs		OR (95% CI)	aOR (95% CI)
	900	720	(80)		
Sex					
Male	500	392	(78)	0.8 (0.6 - 1.1)	1.0 (0.7 - 1.5)
Female	400	328	(82)	Reference	Reference
Age group					
<15	52	47	(90)	Reference	Reference
15-24	103	79	(77)	0.4 (0.1 - 1.1)	0.5 (0.2 - 1.4)
25-34	230	171	(74)	0.3 (0.1 - 0.9)	0.4 (0.1 - 1.2)
35-44	292	240	(82)	0.5 (0.2 - 1.3)	0.6 (0.2 - 1.8)
45-54	131	102	(78)	0.4 (0.1 - 1.1)	0.4 (0.1 - 1.2)
55-64	53	45	(85)	0.6 (0.2 - 2.0)	0.6 (0.2 - 2.0)
>65	39	37	(95)	2.0 (0.3 - 11.9)	1.7 (0.3 - 10.7)
Type of TB					
DR-TB	49	44	(90)	2.2 (0.7 - 6.7)	0.3 (0.1 - 1.0)
DS-TB	851	676	(79)	Reference	Reference
Treatment phase					
Intensive	392	313	(80)	1.0 (0.7 - 1.4)	0.9 (0.6 - 1.3)
Continuation	508	407	(80)	Reference	Reference
Treatment delay (>4 weeks)					
Yes	232	192	(83)	1.3 (0.8 - 1.9)	1.3 (0.9 - 1.9)
No	668	528	(79)	Reference	Reference
HIV status					
Positive	557	450	(81)	1.1 (0.8 - 1.7)	1.4 (0.8 - 2.2)
Negative	343	270	(79)	Reference	Reference
Bread winner					
Yes	457	364	(80)	1.1 (0.8 - 1.5)	1.1 (0.7 - 1.9)
No	443	356	(80)	Reference	Reference
Income quintile					
Poorest	191	181	(95)	14.5 (7.1 - 29.6)	15.7 (7.5 - 33.1)
Less poor	199	180	(90)	7.0 (3.9 - 12.4)	7.2 (3.9 - 13.1)
Average	159	125	(79)	2.8 (1.6 - 4.7)	2.8 (1.6 - 4.8)
Less wealthy	174	133	(76)	2.4 (1.5 - 3.9)	2.5 (1.5 - 4.2)
Wealthiest	177	101	(57)	Reference	Reference
Location of health facility					
Rural	360	295	(82)	1.2 (0.6 - 2.5)	0.9 (0.4 - 1.8)
Urban	540	425	(79)	Reference	Reference

DR-TB=drug resistant TB; DS-TB=drug susceptible TB; OR=Odds ratio; aOR=adjusted odds ratio; CI=confidence interval; HIV = Human Immunodeficiency Virus.

Median costs (US\$) incurred by TB patients enrolled in Zimbabwe patient cost survey (2018)

Table 3:

Phase	Type of cost	Median	DR-TB (IQR)	Median	DS-TB (IQR)	Median	Total (IQR)
Pre-diagnosis	Medical	13 (4-39)	25 (6.5-60)	25 (6-58)			
	Travel	5 (2-15)	5 (2-10.4)	5 (2-10.7)			
	Accommodation	0 (0-0)	0 (0-0)	0 (0-0)			
	Food	9 (0-18)	18 (3.6-36)	18 (2.2-27)			
	Nutritional supplements	2 (0-5)	0 (0-2)	0 (0-2)			
	Total median direct costs	35 (22-70)	54 (23-116)	52 (23-111)			
Post-diagnosis	Medical	207 (129 - 295)	91 (61.8- 134)	91.2 (63.4-151.2)			
	Travel	152 (11.3-552)	32 (3.1- 178)	33.6 (3.3- 193)			
	Accommodation	0 (0-0)	0 (0-0)	0 (0-0)			
	Food	100 (0-480)	25 (0-169.3)	28.1 (0-180)			
	Nutritional supplements	960 (640-1,680)	360 (240-528)	360 (240-600)			
	Total median direct costs	1,659 (653 -2,787)	537 (204-1,134)	555 (220- 1,197)			
Total costs	Medical costs	207 (136-295)	103 (62-173)	109 (63- 194)			
	Non-medical costs	1545 (461- 2,477)	411 (120-948)	434.6 (121-1,018)			
	Indirect costs*	1200 (100-3,000)	240 (0-1,080)	300 (0-1,200)			
		3,569.2 (1,692-5,859)	1185 (523-2,222)	1247 (543-2,405)			

* =calculated based on the output approach.