



Published in final edited form as:

Trop Med Int Health. 2021 November ; 26(11): 1481–1493. doi:10.1111/tmi.13654.

HIV care and treatment models and their association with medication possession ratio among treatment-experienced adults in three African countries

Sharon Tsui¹, Caitlin E. Kennedy¹, Lawrence H. Moulton¹, Larry W. Chang^{1,2}, Jason E. Farley^{2,7,8,9}, Kwasi Torpey⁵, Eric van Praag⁶, Olivier Koole^{3,4}, Nathan Ford¹⁰, Fred Wabwire-Mangen¹¹, Julie A. Denison¹

¹Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, USA

²Department of Medicine – Infectious Diseases, Johns Hopkins University School of Medicine, Baltimore, USA

³Clinical Sciences Department, Institute of Tropical Medicine, Antwerp, Belgium

⁴Department of Clinical Research, London School of Hygiene and Tropical Medicine, UK

⁵School of Public Health, University of Ghana College of Health Sciences, Accra, Ghana

⁶FHI 360 (retired), Dar es Salaam, Tanzania

⁷The REACH Initiative of The Johns Hopkins School of Nursing, Baltimore, USA

⁸University of KwaZulu Natal, Durban, South Africa

⁹Association of Nurses in AIDS Care, Akron, USA

¹⁰Dept HIV, World Health Organization, Geneva, Switzerland

¹¹Department of Epidemiology and Biostatistics, Makerere University School of Public Health, Kampala, Uganda

Abstract

Objective: How clinics structure the delivery of antiretroviral therapy (ART) services may influence patient adherence. We assessed the relationship between models of HIV care delivery and adherence as measured by medication possession ratio (MPR) among treatment-experienced adults in Tanzania, Uganda, and Zambia.

Methods: Eighteen clinics were grouped into three models of HIV care. Model 1-Traditional and Model 2-Mixed represented task-sharing of clinical services between physicians and clinical officers, distinguished by whether nurses played a role in clinical care; in Model 3-Task Shifted, clinical officers and nurses shared clinical responsibilities without physicians. We assessed MPR among 3,419 patients and calculated clinic-level MPR summaries. We then calculated the

Correspondence: Sharon Tsui, Department of International Health, Johns Hopkins Bloomberg School of Public Health, 615 N. Wolfe Street, Baltimore, Maryland, U.S.A. Phone: +256 758-948-177, sharonstui@gmail.com.

Sustainable Development Goal: Good health and well-being

mean differences of percentages and adjusted residual ratio (aRR) of the association between models of care and incomplete adherence, defined as a MPR<90%, adjusting for individual-level characteristics.

Results: In the adjusted analysis, patients in Model 1-Traditional were more likely than patients in Model 2-Mixed to have MPR<90% (aRR=1.60, 95% CI 1–2.48). Patients in Model 1-Traditional were no more likely than patients in Model 3-Task-Shifted to have a MPR<90% (aRR=1.58, 95% 0.88–2.85). There was no evidence of differences in MPR<90% between Model 2-Mixed and Model 3-Task-Shifted (aRR=0.99, 95% CI 0.59–1.66).

Conclusion: Non-physician-led ART programs were associated with adherence levels as good as or better than physician-led ART programs. Additional research is needed to optimize models of care to support patients on life-long treatment.

Keywords

antiretroviral therapy; HIV/AIDS; Tanzania; Uganda; Zambia; medication adherence

Introduction

Country adoption of WHO's recommendation to task-shift antiretroviral therapy (ART) services has contributed to the rapid scale-up of HIV care and treatment in sub-Saharan Africa. In eastern and southern Africa, the number of ART patients had increased from 625,500 in 2005(1) to 15 million by 2019(2). Task-shifting involves the redistribution of health tasks by extending the scope of practice for existing health workers (e.g., allowing nurses to prescribe ART), or creating auxiliary cadres to substitute for health professionals (e.g., creation of adherence support workers to provide clinic-based adherence counselling, or expert patients to relieve nurses of administrative tasks, such as patient file retrieval and clinic navigation(3). Another form of redistributing tasks is task-sharing, whereby a team of health cadres provide differentiated care for patients depending on the severity of illness(4). For example, clinical officers and nurses may be tasked with monitoring stable patients while physicians manage patients with complex opportunistic infections or chronic diseases. Task-shifting and task-sharing allows for limited human resources for health to be strategically deployed for differentiated service delivery of ART for people living with HIV(5–7).

Previous studies have examined the impact of discrete aspects of task-shifting/task-sharing of ART on virologic failure(8–12) and mortality(8, 9, 11), but just two of these examined the impact on patient adherence. One study in Zambia compared how patients counselled by lay or professional staff varied in self-reported adherence in the past 3 days(13). Another study in a clinical trial context in Uganda compared ART pill counts 6–12 months after ART initiation among patients in a nurse-peer counsellor model of care (including home visits) versus a physician-centered model of care(10). Neither study found evidence of significance difference in adherence outcomes.

While existing studies provide a strong evidence base for task-shifting and task-sharing of ART services, they often focus narrowly on a limited range of services, including ART

prescription, management, and adherence counselling. To more comprehensively identify models of care, we previously conducted a cluster analysis and Delphi survey to describe healthcare staffing of comprehensive ART services in 19 health facilities in Tanzania, Uganda, and Zambia(14). The analysis identified three models of care: Model 1-Traditional, where major clinical responsibilities were shared between physicians and clinical officers; Model 2-Mixed, where major clinical responsibilities were shared between physicians, clinical officers, and nurses; and Model 3-Task-Shifted, where clinical officers and nurses completed all major clinical duties, while lay health workers facilitated ancillary services. These models of care were further characterized by environmental factors (e.g., health facility level, urban/rural setting, government/non-profit status) and program characteristics (e.g., ART refill schedule, adherence support strategies, and alternatives to clinic-based ART distribution).

Objective

Here, we assess the relationship between these three models of care and ART adherence using the ART medication possession ratio (MPR) measure – a validated adherence measure predictive of virologic failure(15–18). Our objectives were to assess the association between task-shifting/task-sharing models of care and the MPR in the past six months, and to determine if any association remains after adjusting for individual patient-level factors.

Methods

Design and Setting

We used cross-sectional data from an ART retention and adherence study conducted in Tanzania, Uganda, and Zambia led by FHI 360 (2008–2012). The parent study's purpose was to characterize retention and adherence rates in 19 ART clinics, and to examine programmatic and individual factors related to adult retention and adherence (16). Clinics were purposefully selected with country stakeholders to include those with 300 patients from different urban-rural locations with varying characteristics, including public/private/faith-based organizations, primary/secondary/tertiary-levels, and different ART adherence and provision strategies(16, 19). Eligible patients were at least 18 years old at ART initiation, had initiated ART at least 6 months prior to data collection, and spoke one of the study languages. Participants were systematically sampled, and if a patient was ineligible, unwilling, or unavailable, the study team selected the next ART patient attending the clinic. All participants underwent a screening and consent process by trained interviewers. Interviewers also abstracted data from the patient's medical, pharmacy, and laboratory records using structured data abstraction forms. Data collection from patients and medical chart abstractions took place in 2011. Information on the 19 ART clinics' task-shifting/task-sharing characteristics was collected in a cross-sectional survey with ART clinic managers in 2010–2011.

Measures

Participant adherence, the dependent variable, was assessed by medication possession ratio (MPR). MPR is based on pharmacy refill data and has been shown to significantly

correlate with virologic failure among adults(16). The MPR summarized the number of pills dispensed to participant in the six months prior to the interview divided by the total number of pills the participant should have received during that time(16). The MPR was dichotomized into <90% or ≥90%; this cut-off was selected based on a previous analysis relating adherence measures to virologic failure and receiver operating characteristic analysis with HIV RNA at least 1000 copies/ml as the reference standard(16).

Task-shifting/task-sharing models of care, the independent variable, was constructed by cluster analysis described elsewhere(14). Other variables included individual-level factors relevant to patient adherence to ART: demographic variables including sex, marital status, household wealth index; psychosocial variables including internalized stigma, stigma against HIV disclosure, depressive symptoms, social support assistance, alcohol abuse, HIV-related traditional healer/herbalist visits, average cost and average time to reach the HIV clinic; and clinical variables including current ART regimen, time on ART, daily pill burden, self-reported HIV symptoms in the past four weeks, pre-ART WHO stage, and pre-ART CD4 cell count(16). We did not include country as a covariate because we accounted for clustering at the health facility level and this should sufficiently cover higher-level clustering, particularly when country-level differences are experienced at the health facility level.

Cluster Level Analysis

We assessed the association between task-shifting model of care and patient MPR using generalized estimating equations to account for intra-cluster correlation in the data at the health facility level. The mean differences of percentages of MPR<90% and 95% confidence intervals (CI) were computed for each model of care compared to each other model (e.g., model-1 minus model-2). Each difference was assessed with a Student *t* test, and the corresponding 95% CI for the mean difference.

Adjusted residual ratios (aRR) were calculated by generating standardized clinic level summaries. aRR were calculated as the ratio of observed to expected outcomes predicted by fitting a logistic regression model on individual level data with the MPR<90% as the dependent variable(20). The final logistic regression included all independent variables that were significantly associated with MPR<90% ($p<0.05$) in the bivariate analysis (Table 2). Independent variables included age, household wealth, internalized stigma against HIV, potential depression, time to clinic, current ART regimen, time on ART, pill burden, and pre-ART CD4 cell count. Statistical significance was assessed with the Student *t* test on the sets of logarithms of the aRRs of the two models being compared, and corresponding 95% CIs, and then exponentiating to obtain an aRR.

Subgroup analyses were performed by ART duration (<5.3 or ≥5.3 years). Significance tests of interaction were conducted between ART duration and task-shifting/task-sharing model of care. Statistical analyses were performed using SAS/STAT Version 9 (SAS Institute, Cary, NC, USA).

All patients and ART managers provided written informed consent prior to data collection, and the study was approved by seven Institutional Review Boards.

Results

Characteristics of ART clinics and participating patients by task-shifting/task-sharing model and by MPR<90% outcome are summarized in Table 1 and Table 2, respectively.

Characteristics of ART clinics

Eighteen ART facilities had the MPR outcome and were included in the analysis. Of these, seven were in Tanzania, six in Uganda, and five in Zambia. The 18 facilities were diverse in health facility level (4 national referral hospitals, 8 provincial or district hospitals, and 5 primary health centers), management (9 government, 5 faith-based missions, and 4 non-profit, non-religious organizations), and size (5 sites with <1000 ART patients, 3 with 1000–2000 patients, 7 with 2000–4000 patients, and 3 with >4000 patients). Most facilities were based in urban locales (14 urban, 4 rural) (Table 1a).

Staffing patterns for each model are summarized in Table 1b. Only nurse/midwife and ART counsellors were assigned to provide adherence counselling in Model 1-Traditional as compared to Model 2-Mixed and Model 3-Task-Shifted where a wider range of staff, including medical officer, clinical officer, nurse/midwife, ART counsellor, dispenser, and lay or expert patients, were engaged to provide adherence counselling. For tracing of patients who have missed appointments or defaulted, all three models primarily relied on lay or expert patients to provide this service (Table 1b).

Programmatic details for each model are summarized in Table 1c. Notable differences in programmatic factors included number of counselling sessions required before ART initiation, routine pill counts during ART adherence counselling, community-based distribution of ART, and frequency of ART refill schedule. 40% of clinics in Model 1-Traditional required 3 pre-ART counselling sessions vs. 90% of clinics in Model 2-Mixed and 75% clinics in Model 3-Task-Shifted (Table 1c). 60% of clinics in Model 1-Traditional conducted routine pill counts for patients on ART vs. 89% of clinics in Model 2-Mixed and 75% of clinics in Model 3-Task-Shifted (Table 1c). 30% of clinics in Model 2-Mixed and 25% of clinics in Model 3-Task-Shifted offered community-based distribution of ART vs. 0% in Model 1-Traditional (Table 1c). Finally, 40% clinics in Model 1-Traditional offered 2 or more months of ARV drug refills for stable patients who had been on ART for 6 months or longer vs. 78% of clinics in Model 2-Mixed and 50% clinics in Task-Shifted (Table 1c).

Characteristics of study participants

Overall, 3,419 participants were eligible for analysis. Of these, 73% (2496/3419) were 35 years or older at the time of the interview; 67% (2282/3419) were female. Only 25% (859/3419) had been on ART for less than 2.2 years (Table 2). Overall, 25% of patients (842/3411) attended Model 1-Traditional clinics, 54% (1843/3411) attended Model 2-Mixed clinics, and 21% (726/3411) attended Model 3-Task-Shifted clinics. Patients with <90% MPR were significantly younger, wealthier, and needed more time to travel to the clinic than patients who achieved better MPR (Table 2). Patients with <90% MPR also reported significantly greater levels of internalized or self-stigma living with HIV, positive screening for depression, higher pill burden, and were more likely to have missing pre-CD4 cell count

than patients with higher MPR (Table 2). Finally, significantly fewer patients with <90% MPR were on AZT/3TC/EFV regimen and had more than 5.3 years of experience on ART than patients with higher MPR (Table 2).

Unadjusted proportions and residual ratio of MPR<90% by models of care

Cluster-level analysis found that 57% of patients had MPR<90% (SD 17.59) in Model 1-Traditional, 35% (SD 12.92) for Model 2-Mixed, and 34% (SD 17.37) for Model 3-Task-Shifted (Table 3). Patients in Model 1-Traditional were significantly more likely than patients in Model 2-Mixed to have an MPR<90% (difference in mean = -22.54, 95% CI -42.07, -3.01) and a similar trend was observed comparing patients in Model 1-Traditional to Model 3-Task-Shifted (difference in mean = -23.42, 95% CI -55.17, 8.33). There was no evidence of statistical difference in MPR<90% between patients in Model 2-Mixed and Model 3-Task-Shifted (difference in mean = -0.88, 95% CI -21.64, 19.87) (Table 3).

Adjusted residual ratio of MPR<90% by models of care

Adjusted results from the cluster-level analysis found that patients in Model 1-Traditional were more likely than patients in Model 2-Mixed to have MPR<90% (aRR=1.60, 95% CI 1.03, 2.48). Patients in Model 1-Traditional also showed a trend toward being more likely than patients in Model 3-Task-Shifted to have an MPR<90%, but this difference was not statistically significant (aRR=1.58, 95% CI 0.88, 2.85) (Table 3). There was no evidence of a difference in proportions of patients with an MPR<90% between Model 2-Mixed and Model 3-Task-Shifted (aRR=0.99, 95% CI 0.59, 1.66) (Table 3).

Discussion

Patients in Model 1-Traditional were 60% more likely to have incomplete ART adherence with an MPR<90% than patients in Model 2-Mixed. Similarly, there was some indication that patients in Model 1-Traditional were more likely to have an MPR<90% than patients in Model 3-Task-Shifted, but this difference was not statistically significant, possibly due to the small number of clinics in Model 3 (n=4). These results suggest that differences in ART service delivery are related to varying patient adherence.

Differences in ART service delivery may result from different cadres providing health services, or from differences in how ART and related tasks are implemented. Our findings indicate that non-physician-led ART programs supported adherence levels better than physician-led ART programs, and task-shifting and task-sharing of ART services were not associated with poorer patient adherence. These findings are congruous with other research that has demonstrated comparable standards of care in HIV care and treatment by trained clinical officers and nurses compared to physicians(21–24), and similar mortality, mean CD4 cell count, and virologic failure in effectiveness studies(8, 9, 11, 12).

Differences in staffing for ART service delivery may result in different implementation approaches. Recognizing the complexity of this subject and the fact that observed differences in models of care cannot fully explain the differences observed in MPR, we noted three main aspects distinguishing the three models of care which may have helped patients pick-up their pills.

First, Model 1-Traditional clinics were less likely to require three adherence counselling sessions before ART initiation than Model 2 and 3 clinics. Adherence may have been greater among patients in Models 2 and 3 because these patients had more pre-ART counselling sessions and were more prepared for lifelong HIV treatment. However, research on the value of more pre-ART counselling sessions for adherence is inconclusive(25). Benefits of pre-ART counselling must be balanced by risks of early mortality and greater morbidity associated with delayed ART initiation(26). Given the emphasis on immediate ART initiation within the test and treat approach, it is important to strengthen adherence support for patients at ART initiation and throughout their life on ART(27). An example strategy to strengthen post-initiation adherence includes task-sharing of adherence counselling to lay or peer health workers(13).

Second, Model 1-Traditional clinics were less likely to practice routine pill counts for patients during ART refill visits in contrast to Model 2-Mixed and Model 3-Task-Shifted clinics. While most literature discusses pill counts as a measure of adherence, some have hypothesized that routine pill counts serve as an intervention shaping patient adherence behavior. This research suggests that the repeated process of expecting to have pills counted, organizing medication for pill counts, and receiving increased attention from the person conducting the pill count can have a “reactive effect” for patients, resulting in improved ART adherence(28). This hypothesis is supported by a Kenyan study which found a dose-response relationship between clinician pill count and adherence – the greater the number of pill counts conducted, the more adherent patients were(29).

Finally, some clinics in Models 2 and 3 provided patients with alternatives to clinic-based pill pick-up, which may have helped overcome common barriers to non-adherence by making ART more accessible and affordable(30–32). Models of decentralized ART delivery varied from distribution of ART from mobile, satellite ART clinics within the community to home-based ART distribution by trained lay workers, and are coupled with multi-month dispensing of ART for stable patients (at the time of study stable was defined as patients who have been on ART for six months or longer, without VL results). Altogether, these models can allow patients to travel shorter distances, spend less on transport, and reduce time spent for medication pick-up(31, 33–35), overcoming key deterrents to treatment interruption.

Strengths and limitations

A strength of our study was the inclusion of 18 ART programs in three countries – a considerably larger and more diverse sample than past research, which has generally described task-shifting/task-sharing practices of a few clinics in one country (10, 13, 36–41). While our evaluation offered more information than previously available, the sample of 18 clinics still left us inadequately powered to examine all the associations between models of care and ART adherence. The sites were not randomly selected so data may not be generalizable to ART clinics in these countries or elsewhere in sub-Saharan Africa. Clinics were purposefully selected with national partners to be diverse on a range of characteristics; however, while diverse, they were not randomly selected and therefore not representative of all ART clinics in these countries. In particular, because of our requirements to have

a minimum of 300 ART clients the selected clinics may have been larger and have more established ART programs. For these reasons, our findings may not be generalizable to all ART clinics in the three countries or elsewhere in sub-Saharan Africa.

Another strength was our comprehensive measure of task-shifting/task-sharing practices to characterize models of HIV care. Prior research has described task-shifting/task-sharing models of care broadly as “doctor-centered”, “nurse-centered”, “non-physician care”, “peer-led”, or “community-supported care”(8, 10–12, 21, 27, 38, 39, 42–45) while the model of care is based on a few discrete ART-related tasks, such as non-physician ART initiation or clinical monitoring of patients(10, 11, 38, 40, 45–48), or lay health worker adherence counselling and support(13, 49). In contrast, our measure included healthcare staffing patterns from patient registration, triage, ART prescription, adherence counselling, dispensing, clinical monitoring, phlebotomy for laboratory testing, to tracing of patients lost to follow-up and medical records management, and added clarity to what the model of care entails. This comprehensive measure better considers the many facets that contribute to patient outcomes. A limitation to this measure is that we were unable to consider other important details, such as intensity and coverage of services per patient (and thus workload), or service quality.

Another strength of our analysis was our inclusion of a comprehensive range of factors affecting adherence, including socio-demographic, psychosocial, clinical, and structural characteristics. However, country was not included as a covariate in the analysis. This may limit the accuracy of the model as it will not account for country level factors, such as national policies on differentiated models of ART. Also, the cross-sectional design of the study restricted interpretations to associations rather than temporal relationships and causations.

Using MPR to measure adherence meant that we cannot know if patients actually ingested their drugs or achieved viral suppression. However, MPR is not as vulnerable as self-reported measures to social desirability or recall bias. Another advantage of MPR that it captures both individuals’ ability to access drugs and the system’s ability to dispense drugs to facilitate maximum possible adherence. Interestingly, MPR was not correlated to ART stock-outs in the past 6 months in this study. Our findings suggest a relationship between model of care and MPR – highlighting the importance of access to ART and support for adherence. However, it is not possible to isolate whether it is the health cadres or the ART program that is related to MPR, and further research is needed to elucidate these complex relationships.

Conclusions

Patient data from 18 routinely implemented ART programs in Tanzania, Uganda, and Zambia provide evidence that non-physician clinicians can support favorable levels of ART adherence and result in non-inferior adherence outcomes compared to physician-based models of care. Data on task-shifting/task-sharing models of care suggest further room to optimize the implementation of ART services to improve patient adherence, and additional research to enhance post-ART initiation adherence support is needed.

Acknowledgments

Funding

This research was funded in part and facilitated by the infrastructure and resources provided by the President's Emergency Plan for AIDS Relief (PEPFAR) through the Centers for Disease Control and Prevention (CDC) and the Health Resources & Services Administration (HRSA) under the terms of the contract number 2006-N-08428 with FHI 360, the National Institute of Health National Research Service Award under the terms of the fellowship number F31MH095665, the U.S. Department of Education Fulbright-Hays Doctoral Dissertation Research Abroad Award under the fellowship number P022A150076, and the Johns Hopkins University Center for AIDS Research, an NIH-funded program (P30AI094189), which is supported by the following NIH Co-Funding and Participating Institutes and Centers: NIAID, NCI, NICHD, NHLBI, NIDA, NIMH, NIA, FIC, NIGMS, NIDDK, and OAR. The content is solely the responsibility of the authors and does not necessarily represent the official views of CDC, HRSA, NIH, or any other federal agency or office.

References

1. UNAIDS. Getting to Zero Regional Report: HIV in Eastern & Southern Africa Geneva, Switzerland: UNAIDS; 2013 2013.
2. UNAIDS. Fact Sheet: Global HIV Statistics Geneva, Switzerland: UNAIDS; 2019.
3. Callaghan M, Ford N, Schneider H. A systematic review of task- shifting for HIV treatment and care in Africa. *Hum Resour Health* 2010;8:8. [PubMed: 20356363]
4. Olson D Task Sharing, Not Task Shifting: Team Approach Is Best Bet for HIV Care 2012 3 14, 2016. Available from: http://www.intrahealth.org/blog/task-sharing-not-task-shifting-team-approach-best-bet-hiv-care#.VuaAy_197Dc
5. Ehrenkrantz P, Grimsrud A, Rabkin M. Differentiated service delivery: navigating the path to scale. *Current opinion in HIV and AIDS* 2019;14(1):60–5. [PubMed: 30394947]
6. Hagey JM, Li X, Barr-Walker J, Penner J, Kadima J, Oyaro P, et al. Differentiated HIV care in sub-Saharan Africa: a scoping review to inform antiretroviral therapy provision for stable HIV-infected individuals in Kenya. *AIDS care* 2018;30(12):1477–87. [PubMed: 30037312]
7. Roy M, Bolton Moore C, Sikazwe I, Holmes CB. A Review of Differentiated Service Delivery for HIV Treatment: Effectiveness, Mechanisms, Targeting, and Scale. *Current HIV/AIDS reports* 2019;16(4):324–34. [PubMed: 31230342]
8. Fairall L, Bachmann MO, Lombard C, Timmerman V, Uebel K, Zwarenstein M, et al. Task shifting of antiretroviral treatment from doctors to primary-care nurses in South Africa (STRETCH): a pragmatic, parallel, cluster-randomised trial. *Lancet* 2012;380(9845):889–98. [PubMed: 22901955]
9. Jaffar S, Amuron B, Foster S, Birungi J, Levin J, Namara G, et al. Rates of virologic failure in patients treated in a home-based versus facility-based HIV care model in Jinja, Southeast Uganda: a cluster randomized equivalence trial. *Lancet* 2009;374(9707):2080–9. [PubMed: 19939445]
10. Kiweewa FM, Wabwire D, Nakibuuka J, Mubiru M, Bagenda D, Musoke P, et al. Noninferiority of a task-shifting HIV care and treatment model using peer counselors and nurses among Ugandan women initiated ART: evidence from a randomized trial. *J Acquir Immune Defic Syndr* 2013;63(4):e125–32. [PubMed: 23807157]
11. Sanne I, Orrell C, Fox MP, Conradie F, Ive P, Zeinecker J, et al. Nurse versus doctor management of HIV-infected patients receiving antiretroviral therapy (CIPRA-SA): a randomised non-inferiority trial. *Lancet* 2010;376(9734):33–40. [PubMed: 20557927]
12. Selke HM, Kimaiyo S, Sidle JE, Vedanthan R, Tierney WM, Shen C, et al. Task-shifting of antiretroviral delivery from health care workers to persons living with HIV/AIDS: clinical outcomes of a community-based program in Kenya. *J Acquir Immune Defic Syndr* 2010;55(4):483–90. [PubMed: 20683336]
13. Torpey KE, Kabaso ME, Mutale LN, Kamanga MK, Mwango AJ, Simpungwe J, et al. Adherence support workers: a way to address human resource constraints in antiretroviral treatment programs in the public health setting in Zambia. *PLoS One* 2008;3(5):e2204. [PubMed: 18493615]
14. Tsui S, Denison JA, Kennedy CE, Chang L, Koole O, Torpey K, et al. Identifying Models of HIV Care and Treatment Service Delivery in Tanzania, Uganda, and Zambia Using Cluster Analysis and Delphi Survey. *BMC Health Serv Res* 2017;17(1).

15. Ahonkhai AA, Banigbe B, Adeola J, Bassett IV, Idigbe I, Okonkwo P, et al. High Medication Possession Ratios Associated with Greater Risk of Virologic Failure Among Youth Compared to Adults in a Nigerian Cohort. *J Acquir Immune Defic Syndr* 2018.
16. Denison JA, Koole O, Tsui S, Menten J, Torpey KE, Van Praag E, et al. Incomplete adherence among treatment-experienced adults on antiretroviral therapy in Tanzania, Uganda, and Zambia. *AIDS* 2015;29:361–71. [PubMed: 25686684]
17. Goldman JD, Cantrell RA, Mulenga LB, Tambatamba BC, Reid SE, Levy JW, et al. Simple adherence assessments to predict virologic failure among HIV-infected adults with discordant immunologic and clinical responses to antiretroviral therapy. *AIDS research and human retroviruses* 2008;24(8):1031–5. [PubMed: 18724803]
18. Wu P, Johnson BA, Nachega JB, Wu B, Ordonez CE, Hare AQ, et al. The combination of pill count and self-reported adherence is a strong predictor of first-line ART failure for adults in South Africa. *Current HIV research* 2014;12(5):366–75. [PubMed: 25426940]
19. Koole O, Tsui S, Wabwire-Mangen F, Kwesigabo GP, Menten J, Mulenga M, et al. Retention and risk factors for attrition among adults in antiretroviral treatment programmes in Tanzania, Uganda, and Zambia. *Trop Med Int Health* 2014;19(12):1397–410. [PubMed: 25227621]
20. Bennett S, Parpia T, Hayes R, Cousens S. Methods for the analysis of incidence rates in cluster randomized trials. *International Journal of Epidemiology* 2002;31:839–46. [PubMed: 12177032]
21. Brentlinger PE, Assan A, Mudender F, Ghee AE, Vallejo Torres J, Martinez Martinez P, et al. Task shifting in Mozambique: cross-sectional evaluation of non-physician clinicians' performance in HIV/AIDS care. *Hum Resour Health* 2010;8(23):23. [PubMed: 20939909]
22. Monyatsi G, Mullan PC, Phelps BR, Tolle MA, Machine EM, Genarri FF, et al. HIV management by nurse prescribers compared with doctors at a paediatric centre in Gaborone, Botswana. *S Afr Med J* 2012;102(1):34.
23. Vasan A, Mugisha N, Seung KJ, Achieng M, Banura P, Lule F, et al. Agreement between physicians and non-physician clinicians in starting antiretroviral therapy in rural Uganda. *Hum Resour Health* 2009;7:75. [PubMed: 19695083]
24. Weigel R, Feldacker C, Tweya H, Gondwe C, Chiwoko J, Gumulira J, et al. Managing HIV-infected children in a low-resource, public clinic: a comparison of nurse vs. clinical officer practices in ART refill, calculation of adherence and subsequent appointments. *Journal of the International AIDS Society* 2012;15(2):17432. [PubMed: 22905359]
25. Myer L, Zulliger R, Pienaar D. Diversity of patient preparation activities before initiation of antiretroviral therapy in Cape Town, South Africa. *Trop Med Int Health* 2012;17(8):972–7. [PubMed: 22809271]
26. Cohen MS, Smith MK, Muessig KE, Hallett TB, Powers KA, Kashuba AD. Antiretroviral treatment of HIV-1 prevents transmission of HIV-1: where do we go from here? *Lancet* 2013;382(9903):1515–24. [PubMed: 24152938]
27. van Griensven J, De Naeyer L, Uwera J, Asimwe A, Gazille C, Reid T. Success with antiretroviral treatment for children in Kigali, Rwanda: experience with health center/nurse-based care. *BMC pediatrics* 2008;8:39. [PubMed: 18831747]
28. Kalichman SC, Amaral C, Swetsze C, Eaton L, Kalichman MO, Cherry C, et al. Monthly unannounced pill counts for monitoring HIV treatment adherence: tests for self-monitoring and reactivity effects. *HIV clinical trials* 2010;11(6):325–31. [PubMed: 21239360]
29. Achieng L, Musangi H, Billingsley K, Onguit S, Ombegoh E, Bryant L, et al. The use of pill counts as a facilitator of adherence with antiretroviral therapy in resource limited settings. *PLoS One* 2013;8(12):e67259. [PubMed: 24339861]
30. Bemelmans M, Baert S, Goemaere E, Wilkinson L, Vandendyck M, van Cutsem G, et al. Community-supported models of care for people on HIV treatment in sub-Saharan Africa. *Trop Med Int Health* 2014;19(8):968–77. [PubMed: 24889337]
31. Decroo T, Rasschaert F, Telfer B, Remartinez D, Laga M, Ford N. Community-based antiretroviral therapy programs can overcome barriers to retention of patients and decongest health services in sub-Saharan Africa: a systematic review. *International health* 2013;5(3):169–79. [PubMed: 24030268]

32. Mukherjee JS, Barry D, Weatherford RD, Desai IK, Farmer PE. Community-Based ART Programs: Sustaining Adherence and Follow-up. *Current HIV/AIDS reports* 2016;13(6):359–66. [PubMed: 27739018]
33. Fox MP, Pascoe S, Huber AN, Murphy J, Phokojoe M, Gorgens M, et al. Adherence clubs and decentralized medication delivery to support patient retention and sustained viral suppression in care: Results from a cluster-randomized evaluation of differentiated ART delivery models in South Africa. *PLoS Med* 2019;16(7):e1002874. [PubMed: 31335865]
34. Fatti G, Ngorima-Mabhena N, Mothibi E, Muzenda T, Choto R, Kasu T, et al. Outcomes of Three- Versus Six-Monthly Dispensing of Antiretroviral Treatment (ART) for Stable HIV Patients in Community ART Refill Groups: A Cluster-Randomized Trial in Zimbabwe. *J Acquir Immune Defic Syndr* 2020;84(2):162–72. [PubMed: 32097252]
35. Prust ML, Banda CK, Nyirenda R, Chimbwandira F, Kalua T, Jahn A, et al. Multi-month prescriptions, fast-track refills, and community ART groups: results from a process evaluation in Malawi on using differentiated models of care to achieve national HIV treatment goals. *Journal of the International AIDS Society* 2017;20(Suppl 4):21650. [PubMed: 28770594]
36. Alamo S, Wabwire-Mangen F, Kenneth E, Sunday P, Laga M, Colebunders RL. Task-shifting to community health workers: evaluation of the performance of a peer-led model in an antiretroviral program in Uganda. *AIDS Patient Care STDS* 2012;26(2):101–7. [PubMed: 22224411]
37. Bedelu M, Ford N, Hilderbrand K, Reuter H. Implementing antiretroviral therapy in rural communities: the Lusikisiki model of decentralized HIV/AIDS care. *J Infect Dis* 2007;196 Suppl 3(Suppl 3):S464–8. [PubMed: 18181695]
38. Brennan AT, Long L, Maskew M, Sanne I, Jaffray I, MacPhail P, et al. Outcomes of stable HIV-positive patients down-referred from a doctor-managed antiretroviral therapy clinic to a nurse-managed primary health clinic for monitoring and treatment. *AIDS* 2011;25(16):2027–36. [PubMed: 21997488]
39. Chang L, Alamo S, Guma S, Christopher J, Suntoko T, Omasete R, et al. Two-year virologic outcomes of an alternative AIDS care model: evaluation of a peer health worker and nurse-staffed community-based program in Uganda. *J Acquir Immune Defic Syndr* 2009;50(3):276–82. [PubMed: 19194316]
40. Gimbel-Sherr SO, Micek MA, Gimbel-Sherr KH, Koepsell T, Hughes JP, Thomas KK, et al. Using nurses to identify HAART eligible patients in the Republic of Mozambique: results of a time series analysis. *Hum Resour Health* 2007;5:7. [PubMed: 17328804]
41. Wools-Kaloustian KK, Sidle JE, Selke HM, Vedanthan R, Kemboi EK, Boit LJ, et al. A model for extending antiretroviral care beyond the rural health centre. *Journal of the International AIDS Society* 2009;12:22. [PubMed: 19788755]
42. Davies NE, Homfray M, Venables EC. Nurse and manager perceptions of nurse initiated and managed antiretroviral therapy (NIMART) implementation in South Africa: a qualitative study. *BMJ open* 2013;3(11):e003840.
43. Ford N, Kranzer K, Hilderbrand K, Jouquet G, Goemaere E, Vlahakis N, et al. Early initiation of antiretroviral therapy and associated reduction in mortality, morbidity and defaulting in a nurse-managed, community cohort in Lesotho. *AIDS* 2010;24(17):2645–50. [PubMed: 20980868]
44. Grimsrud A, Kaplan R, Bekker LG, Myer L. Outcomes of a nurse-managed service for stable HIV-positive patients in a large South African public sector antiretroviral therapy programme. *Trop Med Int Health* 2014;19(9):1029–39. [PubMed: 25041716]
45. Shumbusho F, Van Griensven J, Lowrance D, Turate I, Weaver MA, Price J, et al. Task shifting for scale-up of HIV care: evaluation of nurse-centered antiretroviral treatment at rural health centres in Rwanda. *PLoS Med* 2009;6(10):e1000163. [PubMed: 19823569]
46. Bemelmans M, van den Akker T, Ford N, Philips M, Zachariah R, Harries A, et al. Providing universal access to antiretroviral therapy in Thyolo, Malawi through task shifting and decentralization of HIV/AIDS care. *Trop Med Int Health* 2010;15(12):1413–20. [PubMed: 20958897]
47. Long L, Brennan A, Fox MP, Ndibongo B, Jaffray I, Sanne I, et al. Treatment outcomes and cost-effectiveness of shifting management of stable ART patients to nurses in South Africa: an observational cohort. *PLoS Med* 2011;8(7):e1001055. [PubMed: 21811402]

48. Sherr K, Pfeiffer J, Mussa A, Vio F, Gimbel S, Micek M, et al. The role of nonphysician clinicians in the rapid expansion of HIV care in Mozambique. *J Acquir Immune Defic Syndr* 2009;52 Suppl 1(Suppl 1):S20–3. [PubMed: 19858931]
49. Chang LW, Kagaayi J, Nakigozi G, Sempijja V, Packer AH, Serwadda D, et al. Effect of peer health workers on AIDS care in Rakai, Uganda: a cluster-randomized trial. *PLoS One* 2010;5(6):e10923. [PubMed: 20532194]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1a:

Contextual characteristics of the ART clinics in 2011, by the 3 task-shifting/task-sharing models of care

Model of Service Delivery	Country	Facility Level	Managing Authority	Number of Current ART Patients	Urban or Rural	Number of Providers on a Typical Clinic Day				
						# Medical Officer	# Clinical Officer	# Nurse / Midwife	# Lay worker	# Total
1-Traditional (n=5 clinics)	Tanzania	Nat Ref	Mission	1000–2000	Urban	3	3	2	0	8
	Tanzania	Nat Ref	Mission	<1000	Urban	5	0	3	0	8
	Tanzania	District	Government	2000–4000	Urban	3	2	10	2	17
	Uganda	Nat Ref	NPNR	>4000	Urban	7	0	8	4	17
	Zambia	Nat Ref	Government	2000–4000	Urban	3	1	2	2	8
2-Mixed (n=9 clinics)	Tanzania	Provincial	Mission	1000–2000	Urban	1	2	3	2	8
	Tanzania	District	Government	<1000	Rural	2	2	1	1	6
	Uganda	PHC	NPNR	2000–4000	Urban	1	2	3	5	11
	Uganda	PHC	Mission	2000–4000	Urban	4	1	12	0	17
	Uganda	PHC	NPNR	2000–4000	Urban	2	5	5	4	16
	Uganda	PHC	NPNR	2000–4000	Urban	1	1	3	0	5
	Zambia	District	Mission	>4000	Urban	1	2	3	3	9
	Zambia	District	Government	<1000	Rural	3	5	1	2	11
	Zambia	Provincial	Government	>4000	Urban	1	2	5	1	9
3-Task-Shifted (n=4 clinics)	Tanzania	District	Government	<1000	Rural	0	1	3	0	4
	Tanzania	Provincial	Government	1000–2000	Urban	0	2	9	1	12
	Uganda	District	Government	<1000	Rural	0	1	6	8	15
	Zambia	PHC	Government	2000–4000	Urban	0	2	4	1	7

Table 1b:

Staffing patterns of the ART clinics in 2011 by the 3 task-shifting/task-sharing models of care

Tasks, by cadre	Model 1: Traditional (n=5 clinics)	Model 2: Mixed (n=9 clinics)	Model 3: Task-Shifted (n=4 clinics)
Registration			
Medical officer	0 (0%)	1 (11%)	0 (0%)
Clinical officer	0 (0%)	1 (11%)	1 (25%)
Nurse/ midwife	3 (60%)	2 (22%)	2 (50%)
ART counsellor	0 (0%)	4 (44%)	0 (0%)
Phlebotomist	0 (0%)	1 (11%)	0 (0%)
Dispenser	0 (0%)	1 (11%)	1 (25%)
Records clerk	2 (40%)	2 (22%)	1 (25%)
Lay or expert patient	1 (20%)	5 (56%)	1 (25%)
Initial ART Prescription			
Medical officer	5 (100%)	8 (89%)	0 (0%)
Clinical officer	2 (40%)	8 (89%)	4 (100%)
Nurse/ midwife	0 (0%)	4 (44%)	3 (75%)
ART Monitoring & Management			
Medical officer	5 (100%)	9 (100%)	0 (0%)
Clinical officer	2 (40%)	8 (89%)	4 (100%)
Nurse/ midwife	2 (40%)	3 (33%)	3 (75%)
Dispenser	0 (0%)	0 (0%)	1 (25%)
Laboratory technician	0 (0%)	0 (0%)	1 (25%)
Lay or expert patient	1 (20%)	0 (0%)	0 (0%)
Adherence Counselling * (n=4)			
Medical officer	0 (0%)	3 (33%)	0 (0%)
Clinical officer	0 (0%)	3 (33%)	1 (25%)
Nurse/ midwife	2 (50%)	7 (78%)	2 (50%)
ART counsellor	2 (50%)	6 (67%)	2 (50%)
Dispenser	0 (0%)	3 (33%)	1 (25%)
Lay or expert patient	0 (0%)	3 (33%)	0 (0%)
ART Dispensing * (n=4)			
Nurse/ midwife	1 (25%)	4 (44%)	4 (100%)
Pharmacist/ dispenser	4 (100%)	7 (78%)	1 (25%)
Lay or expert patient	1 (25%)	1 (11%)	1 (25%)
Phlebotomy * (n=4)			
Clinical officer	0 (0%)	0 (0%)	1 (25%)
Nurse/ midwife	0 (0%)	4 (44%)	1 (25%)
ART counsellor	0 (0%)	1 (11%)	0 (0%)

Tasks, by cadre	Model 1: Traditional (n=5 clinics)	Model 2: Mixed (n=9 clinics)	Model 3: Task-Shifted (n=4 clinics)
Lab technician	4 (100%)	7 (78%)	3 (75%)
Lay or expert patient	0 (0%)	1 (11%)	0 (0%)
Patient Tracing on missed appointments and defaulters * (n=4)			
Medical officer	0 (0%)	1 (11%)	0 (0%)
Clinical officer	0 (0%)	2 (22%)	0 (0%)
Nurse/ midwife	0 (0%)	3 (33%)	1 (25%)
ART counsellor	0 (0%)	5 (56%)	0 (0%)
Records clerk	1 (25%)	1 (11%)	1 (25%)
Lay or expert patient	3 (75%)	6 (67%)	3 (75%)

The percentage is calculated for each cadre to enable comparison across each model of care.

* The question was not answered by one of sites grouped in Model 1: Traditional; therefore, percentages are calculated out of a total denominator of 4 ART clinics.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1c:

Programmatic characteristics of the ART clinics in 2011 by the 3 task-shifting/task-sharing models of care

Programmatic Characteristics, by cadre	Model 1: Traditional (n=5 clinics)	Model 2: Mixed (n=9 clinics)	Model 3: Task-Shifted (n=4 clinics)
ART Initiation Preparedness			
Counselling required to initiate ART (yes)	5 (100%)	9 (100%)	4 (100%)
Treatment supporter required to initiate ART (yes)	4 (80%)	8 (89%)	3 (75%)
# Counselling Pre-ART Counselling Sessions			
1 session	1 (20%)	0 (0%)	1 (25%)
2 sessions	2 (40%)	1 (11%)	0 (0%)
3 sessions	2 (40%)	8 (89%)	3 (75%)
Methods of Pre-ART Counselling			
Individual	5 (100%)	8 (89%)	4 (100%)
Individual with treatment supporter	5 (100%)	9 (100%)	4 (100%)
Group	3 (60%)	6 (67%)	3 (75%)
Pill count	3 (60%)	5 (56%)	2 (50%)
Total number of counselling methods used			
2 methods	1 (20%)	0 (0%)	1 (25%)
3 methods	2 (40%)	8 (89%)	1 (25%)
4 methods	2 (40%)	1 (11%)	2 (50%)
Clinic has referral linkage to CHW trained in adherence support (yes)	3 (60%)	8 (89%)	3 (75%)
ART Adherence Counselling after Initiation			
Methods of Counselling			
Individual	4 (80%)	7 (78%)	4 (100%)
Individual with treatment supporter	5 (100%)	7 (78%)	3 (75%)
Group	5 (100%)	5 (56%)	3 (75%)
Pill count practice	3 (60%)	8 (89%)	3 (75%)
Total number of counselling methods used			
2 methods	0 (0%)	2 (22%)	1 (25%)
3 methods	3 (60%)	5 (56%)	1 (25%)
4 methods	2 (40%)	2 (22%)	2 (50%)
Frequency of ART Refill			
In the first month on ART			
Every 2 weeks	5 (100%)	9 (100%)	4 (100%)
In the second to sixth months on ART			
Every month	5 (100%)	8 (89%)	4 (100%)
Every 2 months	0 (0%)	1 (11%)	0 (0%)
After six months on ART			
Every month	3 (60%)	2 (22%)	2 (50%)
Every 2 months	1 (20%)	6 (67%)	1 (25%)
Every 3 months	1 (20%)	1 (11%)	1 (25%)

Programmatic Characteristics, by cadre	Model 1: Traditional (n=5 clinics)	Model 2: Mixed (n=9 clinics)	Model 3: Task-Shifted (n=4 clinics)
Pharmacy Support for ART Adherence			
Methods			
Patient self-report adherence assessment	3 (60%)	6 (67%)	4 (100%)
Dispenser adherence assessment	4 (80%)	6 (67%)	3 (75%)
Verify refill dates	4 (80%)	7 (78%)	4 (100%)
Total number of pharmacy-based methods used			
None (zero method)	1 (20%)	2 (22%)	0 (0%)
3 methods	1 (20%)	4 (44%)	2 (50%)
4 methods	3 (60%)	3 (33%)	2 (50%)
Social Support for ART Adherence			
PLHIV support group	5, 100%	7, 78%	4, 100%
Adherence support worker	4, 80%	7, 78%	3, 75%
Home-based care worker	3, 60%	5, 56%	2, 50%
Community-based ART Adherence Services			
Methods			
Distribution of ARV drugs	0, 0%	3, 33%	1, 25%
Home based care	3, 60%	6, 67%	3, 75%
Adherence support	3, 60%	7, 78%	4, 100%
Emotion/ social support	3, 60%	8, 89%	3, 75%
Follow-up of missed appointments	3, 60%	8, 89%	3, 75%
Nutritional support	2, 40%	5, 56%	3, 75%
Number of community-based methods (out of 7)			
No methods	1, 20%	1, 11%	0, 0%
1–4 methods	1, 20%	1, 11%	1, 25%
5–7 methods	3, 60%	7, 78%	3, 75%
Stock outs of ART in the last 6 months			
	1, 20%	0, 0%	4, 100%
Provided viral load testing as needed			
	3, 60%	2, 22%	1, 25%

Table 2:

Summary of analysis population by task-shifting/task-sharing model and adherence outcome

	<i>N</i> =3411	Model 1 Traditional % (<i>n</i> =842)	Model 2 Mixed % (<i>n</i> =1843)	Model 3 Task- Shifted % (<i>n</i> =726)	<i>P</i>	Total %	MPR <90% (<i>n</i> =1997)	MPR 90% (<i>n</i> =1414)	<i>On</i> MPR<90% Odds Ratio (95% CI)	<i>P</i>
Demographics										
Age (in years)										
<35	921	23.16	30.66	22.18	<.0001	27.00	29.99	24.89	1.293 (1.110, 1.505)	0.0010
35 years	2490	76.84	69.34	77.82		73.00	70.01	75.11	1	
Female sex										
	2277	66.51	65.76	69.56	0.1814	66.75	68.46	65.55	0.877 (0.758, 1.014)	0.0756
Marital status										
Single	334	12.59	9.39	7.58	0.0121	9.79	10.25	9.46	3.836 (0.443, 33.187)	0.4631
Separated/ divorced/ widowed	1199	34.20	34.56	37.74		35.15	35.79	34.70	3.650 (0.425, 31.339)	
Married or cohabitating	1872	53.09	55.94	54.27 0.41		54.88	53.89	55.58	3.432 (0.400, 29.433)	
Missing	6	0.12	0.11			0.18	0.07	0.25	1	
DHS Wealth Index										
Low	1107	7.96	41.35	38.29	<.0001	32.45	28.15	35.50	0.638 (0.540, 0.755)	<.0001
Middle	1135	28.86	34.35	35.67		33.27	33.17	33.35	0.801 (0.679, 0.944)	
High	1169	63.18	24.31	26.03		34.27	38.68	31.15	1	
Psychosocial Factors										
Stigma internalized										
High (>median)	1156	37.17	34.89	27.55	0.0001	33.89	37.69	31.20	1.334 (1.156, 1.540)	<.0001
Low	2255	62.83	65.11	72.45		66.11	62.31	68.80	1	
Stigma disclosure										
High (>median)	951	28.15	29.73	22.87	0.0022	27.88	29.63	26.64	1.160 (0.997, 1.349)	0.0549
Low	2460	71.85	70.27	77.13		72.12	70.37	73.36	1	
Potential depression										

	<i>N</i> =3411	Model 1 Traditional % (<i>n</i> =842)	Model 2 Mixed % (<i>n</i> =1843)	Model 3 Task- Shifted % (<i>n</i> =726)	<i>P</i>	Total %	MPR <90% (<i>n</i> =1997)	MPR 90% (<i>n</i> =1414)	<i>On</i> <i>MPR</i> <90% Odds Ratio (95% CI)	<i>P</i>
Positive screen	437	7.84	15.46	11.85	<.0001	12.81	14.71	11.47	1.332 (1.089, 1.629)	0.0053
Negative screen (REF)	2974	92.16	84.54	88.15		87.19	85.29	88.53	1	
Ever-disclosed HIV status										
Yes	3250	96.67	94.47	95.73	0.0353	95.28	95.62	95.04	0.879 (0.635, 1.217)	0.4374
No	161	3.33	5.53	4.27		4.72	4.38	4.96	1	
Social support care										
Lower 10 th percentile	427	10.45	14.76	9.23	<.0001	12.52	11.53	13.22	0.855 (0.694, 1.053)	0.1415
Higher	2984	89.55	85.24	90.77		87.48	88.47	86.78	1	
Social support help										
Lower 10 th percentile	461	9.62	15.95	11.85	<.0001	13.52	12.94	13.92	0.919 (0.752, 1.123)	0.4101
Higher	2950	90.38	84.05	88.15		86.48	87.06	86.08	1	
CAGE alcohol abuse										
Positive 2	796	23.63	24.04	21.21	0.3045	23.34	24.47	22.53	1.114 (0.949, 1.307)	0.1880
Negative <2 (REF)	2615	76.37	75.96	78.79		76.66	75.53	77.47	1	
Traditional healer										
Ever consulted	224	15.20	4.23	2.48	<.0001	6.57	6.86	6.35	1.085 (0.825, 1.426)	0.5589
Never consulted	3187	84.80	95.77	97.52		93.43	93.14	93.64	1	
Cost to clinic										
1 USD	1486	42.28	39.77	54.68	<.0001	43.56	43.49	43.62	0.995 (0.867, 1.142)	0.9437
<1 USD	1925	57.72	60.23	45.32		56.44	56.51	56.38	1	
Time to clinic										
30 minutes	2739	80.17	79.00	83.75	0.0244	80.30	78.43	81.62	0.891 (0.691, 0.970)	0.0210
<30 minutes	672	19.83	21.00	16.25		19.70	21.57	18.38	1	
Clinical Characteristics										

	N=3411	Model 1 Traditional % (n=842)	Model 2 Mixed % (n=1843)	Model 3 Task- Shifted % (n=726)	P	Total %	MPR <90% (n=1997)	MPR 90% (n=1414)	On MPR<90% Odds Ratio (95% CI)	P
ART Regimen (rcurrartfinal2)										
1=D4T,3TC,NVP	570	12.95	15.57	23.97	<.0001	16.71	15.63	17.48	0.814 (0.639, 1.038)	<.0001
2=AZT,3TC,EFV	659	26.72	15.52	20.39		19.32	14.92	22.43	0.606 (0.477, 0.769)	
3=AZT,3TC,NVP	1227	26.84	49.75	34.44		35.97	39.46	33.50	1.072 (0.871, 1.320)	
4=Other regimen	443	12.83	12.75	13.77		12.99	14.14	12.17	1.058 (0.819, 1.367)	
5=TDF,3TC/ FTC,EFV (ref)	512	20.67	15.41	7.44		15.01	15.84	14.42	1	
Time on ART										
<2.2	859	19.60	29.19	21.49	<.0001	25.18	23.55	26.34	1.147 (0.942, 1.397)	<.0001
2.2–5.3	1714	47.39	49.81	54.68		50.25	55.37	46.62	1.524 (1.285, 1.807)	
>5.3	838	33.02	21.00	23.83		24.57	21.07	27.04	1	
Pill burden (self-report)										
<4	2766	86.22	81.55	73.97	<.0001	81.09	78.15	83.17	0.723 (0.609, 0.859)	0.0002
4	645	13.78	18.45	26.03		18.91	21.85	16.83	1	
HIV symptoms index (median)										
1808	50.00	56.81	46.83	<.0001	53.00	54.17	52.18	0.923 (0.805, 1.058)	0.2506	
Pre-ART WHO stage										
Missing	290	15.80	5.81	6.89	<.0001	8.50	8.63	8.41	1.040 (0.804, 1.345)	0.9214
Stage IV	408	9.98	13.56	10.19		11.96	12.38	11.67	1.076 (0.860, 1.346)	
Stage III	1351	39.67	35.32	50.41		39.61	39.39	39.76	1.005 (0.862, 1.171)	
Stage I and II	1362	34.56	45.31	32.51		39.93	39.60	40.16	1	
Pre-ART CD4⁺ cell count (cells/μl)										
Missing	671	17.22	18.77	24.79	<.0001	19.67	21.78	18.18	1.242 (1.045, 1.477)	0.0288

	<i>N</i> =3411	Model 1 Traditional % (<i>n</i> =842)	Model 2 Mixed % (<i>n</i> =1843)	Model 3 Task- Shifted % (<i>n</i> =726)	<i>P</i>	Total %	MPR <90% (<i>n</i>=1997)	MPR 90% (<i>n</i>=1414)	<i>On MPR<90%</i> Odds Ratio (95% CI)	<i>P</i>
>250	424	10.81	14.27	9.64		12.43	11.74	12.92	0.942 (0.762, 1.164)	
250	2316	71.97	66.96	65.56		67.90	66.48	68.90	1	

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3:

Percentages, mean difference, and adjusted relative risks of non-adherence between the three task-sharing/
task-sharing models of care in Antiretroviral Therapy.

	Model 1: Traditional	Model 2: Mixed	Model 3: Task- Shifted	Effect Estimates	P
Number of ART clinics	5	9	4		
Percentages of non-adherence based on cluster summaries (SD)	57.30 (17.59)	34.77 (12.92)	33.88 (17.37)		
Mean Difference and 95% CI					
Model 1-Traditional minus 2-Mixed				-22.54 (-42.07, -3.01)	0.0272
Model 1-Traditional minus 3-Task-Shifted				-23.42 (-55.17, 8.33)	0.1246
Model 2-Mixed minus 3-Task-Shifted				-0.88 (-21.64, 19.87)	0.9270
<i>Analyses based on ratio-residuals</i>					
Adjusted *Relative Risk and 95% CI					
Model 1-Traditional vs. 2-Mixed				1.60 (1.04, 2.46)	0.0354
Model 1-Traditional vs. 3-Task-Shifted				1.59 (0.88, 2.85)	0.1054
Model 2-Mixed vs. 3-Task-Shifted				0.99 (0.59, 1.65)	0.9720

* Adjusted for age, household wealth, internalized stigma against HIV, potential depression, time to clinic, ART regimen, time on ART, pill burden, and pre-ART CD4 cell count.