



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Original article

Female gender is associated with long COVID syndrome: a prospective cohort study

Francesca Bai^{1,*}, Daniele Tomasoni¹, Camilla Falcinella¹, Diletta Barbanotti¹, Roberto Castoldi¹, Giovanni Mulè¹, Matteo Augello¹, Debora Mondatore¹, Marina Allegrini¹, Andrea Cona¹, Daniele Tesoro¹, Gianmarco Tagliaferri¹, Ottavia Viganò¹, Elisa Suardi¹, Camilla Tincati¹, Tomaso Beringheli¹, Benedetta Varisco¹, Chiara Luridiana Battistini², Kyrie Piscopo², Elena Vegni², Alessandro Tavelli¹, Stefano Terzoni³, Giulia Marchetti¹, Antonella d'Arminio Monforte¹

¹ Clinic of Infectious Diseases, San Paolo Hospital, ASST Santi Paolo e Carlo, Department of Health Sciences, University of Milan, Italy

² Unit of Clinical Psychology, San Paolo Hospital, ASST Santi Paolo e Carlo, Department of Health Sciences, University of Milan, Italy

³ Department of Nursing Sciences, University of Milan, Italy

ARTICLE INFO

Article history:

Received 24 March 2021

Received in revised form

27 October 2021

Accepted 1 November 2021

Available online 9 November 2021

Editor: A. Huttner

Keywords:

COVID-19

Gender

Long COVID

Post acute sequelae SARS COV-2 infection

(PASC)

Post COVID-19 condition

Post-COVID syndrome

ABSTRACT

Objective: We explored the association between female gender and long COVID syndrome, defined as persistence of physical and/or psychological symptoms for more than 4 weeks after recovery from acute COVID-19 disease. The secondary aim was to identify predictors of long COVID syndrome by multivariable logistic regression analysis.

Methods: This was a single-centre prospective cohort study conducted at San Paolo Hospital in Milan, Italy. We enrolled adult patients who were evaluated at the post-COVID outpatient service of our Infectious Diseases Unit between 15 April 2020 and 15 December 2020. Participants were individuals who had clinically recovered from COVID-19 and in whom virological clearance had occurred. Previous infection by SARS-CoV-2 was microbiologically documented by positivity using a reverse-transcriptase polymerase chain reaction (RT-PCR) assay of nasopharyngeal swab. All enrolled patients underwent blood tests and a comprehensive medical examination at follow-up. Individuals were interviewed about resolved and persisting symptoms and were asked to fill in two questionnaires to allow assessment of the Hospital Anxiety and Depression symptoms (HADS) score and of the Impact of Event Scale–Revised (IES-R) score.

Results: A total of 377 patients were enrolled in the study. The median time from symptom onset to virological clearance was 44 (37–53) days. A diagnosis of long COVID syndrome was made in 260/377 (69%) patients. The most common reported symptoms were fatigue (149/377, 39.5%), exertional dyspnoea (109/377, 28.9%), musculoskeletal pain (80/377, 21.2%) and “brain fog” (76/377, 20.2%). Anxiety symptoms were ascertained in 71/377 (18.8%) individuals, whereas 40/377 (10.6%) patients presented symptoms of depression. Post-traumatic stress disorder (defined by a pathological IES-R score) was diagnosed in one-third of patients (85/275, 31%). Female gender was independently associated with long COVID syndrome at multivariable analysis (AOR 3.3 vs. males, 95% CI 1.8–6.2, $p < 0.0001$). Advanced age (adjusted (A)OR 1.03 for 10 years older, 95% CI 1.01–1.05, $p 0.01$) and active smoking (AOR 0.19 for former smokers vs. active smokers, 95% CI 0.06–0.62, $p 0.002$) were also associated with a higher risk of long COVID, while no association was found between severity of disease and long COVID (AOR 0.67 for continuous positive airway pressure (CPAP)/non-invasive mechanical ventilation (NIMV)/orotracheal intubation (OTI) vs. no O2 therapy, 95% CI 0.29–1.55, $p 0.85$).

Discussion: Factors that were found to be associated with a higher risk of developing “long COVID” syndrome were female gender, older age and active smoking, but not severity of the acute disease.

* Corresponding author. Francesca Bai, Clinic of Infectious Diseases, San Paolo Hospital, ASST Santi Paolo e Carlo, Department of Health Sciences, University of Milan, Via A. di Rudini, 8 - 20142, Milan, Italy.

E-mail address: francesca.bai@unimi.it (F. Bai).

Individuals affected by SARS-CoV-2 infection with the aforementioned features should be early identified and involved in follow-up programmes. **Francesca Bai, *Clin Microbiol Infect* 2022;28:611.e9–611.e16**
© 2021 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved.

Introduction

Over half of individuals affected by COVID-19 report persisting symptoms after recovery from acute illness. These symptoms can endure up to 6–7 months and even longer [1–5]. Since the persistence of symptoms represents a health concern, this phenomenon was given the name of “long COVID” syndrome. In the UK, the National Institute for Health and Care Excellence (NICE) defined long COVID as the persistence of symptoms for more than 4 weeks after the onset of acute illness [6–8]. Ongoing physical symptoms include fatigue, muscle weakness, shortness of breath or cough, joint or chest pain, as well as difficulty in concentration and memory disorders; the last two symptoms re-appear in a broader state of diminished mental capacity that may be described as “brain fog” [1,4,9–11]. As far as psychological manifestations are concerned, the most frequent are post-traumatic stress disorder (PTSD), depression, anxiety, obsessive–compulsive traits and insomnia [1,12–15]. Previous studies have not distinctly identified predictors of long COVID syndrome; factors found to have a correlation with long COVID so far were increased length of acute illness, longer time from symptom onset to virological clearance and comorbidities [11,16]. Increased COVID-19 severity and female gender as predicting factors are instead not equivocal findings [1,4,5,8,13,17–20]. Nevertheless, what emerges from preliminary data is the negative impact of long COVID on quality of life and its effects on patient work ability and social relationships [1,4,19]. Given the extent of such phenomenon, healthcare public resources will be utilized to face the issue even when the COVID-19 emergency status will end. It is therefore necessary to better understand the characteristics of long COVID, its predictors and possible long-term sequelae. Furthermore, the likelihood of presenting ongoing symptoms might differ according to several factors (e.g. socioeconomic status or cultural background) that should therefore be identified.

We thus investigated the incidence of physical and/or psychological symptoms characterising the “long COVID” syndrome in female gender and we studied the possible predictors of long COVID.

Materials and methods

Study design and ethics

This was a single-centre prospective cohort study conducted at San Paolo Hospital in Milan, Italy. Informed consent from all participants was obtained. The study was approved by the Ethics Committee Area 1, Milan (2020/ST/049 and 2020/ST/049_BIS, 11 March 2020).

Study population

We enrolled adult patients who were evaluated at the post-COVID outpatient clinic, which had been set up in April 2020, in our Infectious Diseases Unit between 15 April 2020 and 15 December 2020. Participants were patients who had been hospitalized for COVID-19 at San Paolo and San Carlo Hospitals in Milan

and had been discharged between 1 March 2020 and 1 November 2020. Previous infection by SARS-CoV-2 was microbiologically documented by positivity of a reverse-transcriptase polymerase chain reaction (RT-PCR) assay for a nasopharyngeal swab. At the time of hospital discharge, patients were given an appointment for a follow-up visit in the outpatient clinic.

Not included in the study were patients who died during hospitalization or after discharge, those who were not given a follow-up appointment because they declined to participate or were unable to reach the hospital, and those who missed their appointment and were therefore lost to follow-up.

Patients who had been hospitalized for other medical issues and had a positive SARS-CoV-2 nasopharyngeal swab but remained asymptomatic for COVID-19 were also excluded from the study.

Follow-up implied

- repetition of nasopharyngeal swab for patients who were discharged with the indication to self-isolate since their upper respiratory swab was still positive at the time of discharge;
- blood tests and comprehensive medical visit at 1–3 months and later at 6 months after virological clearance (namely a negative nasopharyngeal swab).

This study reports data about the first follow-up assessment at 1–3 months after virological clearance.

A trained nurse was in charge of taking blood samples, whereas a team of Infectious Diseases specialists and clinical psychologists were involved in visiting and interviewing participants. Patients were asked to self-report a detailed list of symptoms and signs, specifying whether they had resolved or were still ongoing. Moreover, they were invited to fill in two questionnaires: the Hospital Anxiety and Depression Symptoms (HADS) [21] and the Impact of Event Scale–Revised (IES-R) [22]. A comprehensive physical examination was performed by trained Infectious Diseases specialists. Clinical psychologists revised the questionnaires and identified symptoms of anxiety or depression and PTSD by evaluating HADS and IES-R scores respectively.

Aims of the study

The primary aim of the study was to assess the incidence of long COVID syndrome in females. The secondary aim was to identify predictors of long COVID by multivariable logistic regression analysis.

Study procedures

Data of the acute phase of illness were collected. Disease severity was defined by the highest level of respiratory support required during hospitalization (no oxygen therapy required, low- or high-flow oxygen therapy, continuous positive airway pressure (CPAP), non-invasive mechanical ventilation (NIMV) and orotracheal intubation (OTI)). Treatments for COVID-19 included antiviral agents (remdesivir (RDV) or HIV protease inhibitors), hydroxychloroquine alone or in association with azithromycin,

corticosteroids and immunomodulatory drugs (IL-6 receptor antagonists and JAK-STAT inhibitors). At follow-up, all patients underwent routine blood tests and a medical evaluation. All patients were asked about the following signs and symptoms: anosmia, dysgeusia, fever, gastrointestinal symptoms, rest or exertional dyspnoea, fatigue, musculoskeletal pain, muscle weakness, “brain fog” manifestations (namely difficulties in attention or concentration and memory disorders). HADS [21] was intended to measure anxiety and depression symptoms, whereas IES-R was used as a screening tool of PTSD [22]. A total HADS score higher than 8 denoted considerable symptoms of anxiety and depression, while a IES-R score above 33 was interpreted as highly suggestive for PTSD. Long COVID was defined as the persistence of ≥ 1 physical and/or psychological symptoms at follow-up [7].

Statistical analyses

Data are presented as median and interquartile range (IQR) for quantitative parameters and absolute numbers and percentages for categorical variables. A Venn diagram was used to show possible correlations among physical and psychological symptoms characterizing long COVID. Comparison between the group of patients diagnosed with long COVID and that of patients who did not and between females and males was investigated by Mann–Whitney test and chi-squared or Fisher's exact test. The association between female gender and long COVID was analysed by fitting a multivariable logistic regression analysis, adjusting for possible confounders (age, severity of disease, length of hospital stay [LOS], comorbidities, body mass index [BMI], smoking and time to

virological clearance). Missing data were handled by the missing-indicator method. p values have been adjusted by Bonferroni's correction. Statistical analyses were performed with STATA software, version 14.0.

Results

Fig. 1 depicts the study flowchart. A total of 492 patients discharged from San Paolo and San Carlo Hospitals in the aforementioned time period were given an appointment for a follow-up evaluation at the post-COVID outpatient clinic. In total 115/492 (23%) patients missed their visit, thus the study includes a total of 377/492 individuals, namely 77% of those who were given an appointment. Demographic and clinical characteristics of patients who missed the follow up appointment are presented in the Table S1.

The follow-up examination was done at a median of 102 (IQR 86–126) days from acute symptom onset, a median of 79 (IQR 69–102) days from clinical recovery and a median of 56 (IQR 47–74) days from virological clearance. Baseline demographic and clinical features of the enrolled patients are shown in Table 1. Females accounted for 137/377 (36%) of all patients; median age was 57 (IQR 49–68) years. The most frequent acute symptoms were fever (280/377, 74%), cough (198/377, 53%), dyspnoea (161/377, 43%) and fatigue (59/377, 16%). In total, 346/377 (92%) presented interstitial pneumonia at chest X-ray or CT scan at the time of hospital admission. One-third of patients received CPAP or NIMV (115/377, 31%) during hospitalization, whereas only 30/377 (8%) patients underwent OTI.




Fig. 1. Study flow chart.

Women were less frequently active smokers than men. Furthermore, overall patients of female gender presented a milder form of disease, estimated as a lower proportion of OTI and a shorter LOS (Table 2). Blood tests at follow-up showed no significant alterations: C-reactive protein (CRP) and complete blood count (CBC) were normal for all patients, while D-dimer

was still out of range in 28/377 (7.4%) patients without however any correlation with organ dysfunction (median pathological D-dimer was 399, IQR 310.7–579.2 ng/mL). In all 28 patients with elevated D-dimer at follow-up, levels were lower than the acute phase of illness; however, two of these patients presented persisting dyspnoea and therefore underwent chest

Table 1
Demographic and clinical characteristics of enrolled patients according to “long COVID” syndrome

Parameters	Total population (N 377)	No long COVID (N 117)	Long COVID (N 260)	p
Age, years	57 (49–68)	56 (42–64)	58 (51–69)	0.01
Gender, males	240 (63.7)	92 (78.6)	148 (56.9)	<0.0001
Comorbidities	164 (43.5)	45 (38.5)	119 (45.8)	0.241
Smoking:				0.046
Never	216 (57.3)	71 (60.7)	145 (55.8)	
Ex-smokers	20 (5.3)	9 (7.7)	11 (4.2)	
Active-smokers	131 (34.7)	30 (25.7)	101 (38.8)	
Unknown	10 (2.7)	7 (5.9)	3 (1.2)	
Ethnicity:				0.579
White/Caucasian	252 (66.8)	82 (70.1)	170 (65.4)	
Arabic	16 (4.2)	6 (5.1)	10 (3.8)	
Black	2 (0.5)	0	2 (0.8)	
Asian	7 (1.9)	0	7 (2.7)	
Hispanic	33 (8.7)	9 (7.7)	24 (9.2)	
Unknown	67 (17.8)	20 (17.1)	47 (18.1)	
During hospitalization				
Symptoms at admission:				
Fever	280 (74.3)	96 (82.1)	184 (70.8)	0.023
Cough	198 (52.5)	67 (57.3)	131 (50.4)	0.360
Sore throat	7 (1.9)	3 (2.6)	4 (1.5)	0.501
Chest pain	16 (4.2)	6 (5.1)	10 (3.8)	0.593
Myalgia-arthralgia	28 (7.4)	10 (8.5)	18 (6.9)	0.422
Fatigue	59 (15.6)	20 (17.1)	39 (15)	0.732
Headache	11 (2.9)	4 (3.4)	7 (2.7)	0.754
Dyspnoea	161 (42.7)	46 (39.3)	115 (44.2)	0.458
Vomiting/nausea	25 (6.6)	9 (7.7)	16 (6.2)	0.526
Anosmia/dysgeusia	24 (6.4)	9 (7.7)	15 (5.8)	0.478
Pneumonia	346 (91.8)	108 (92.3)	238 (91.5)	0.372
Oxygen therapy:				0.879
No O ₂ therapy	35 (9.3)	13 (11.1)	22 (8.4)	
Low/high flows systems	197 (52.2)	60 (51.3)	137 (52.7)	
CPAP, NIV	115 (30.6)	35 (29.9)	80 (30.8)	
OTI	30 (7.9)	9 (7.7)	21 (8.1)	
Length of hospital days (LOS)	11 (6–20)	10 (6–16)	12 (7–21)	0.232
Treatments for COVID-19:				
Antivirals (RDV or PI)	92 (22.4)	30 (25.6)	62 (23.8)	0.746
HQC ± azithromycin	260 (68.9)	83 (70.9)	177 (68.1)	0.661
Steroid	42 (11.1)	14 (11.9)	28 (10.8)	0.798
Immunomodulators	28 (7.4)	11 (9.4)	17 (6.5)	0.334
Obesity:				0.021
BMI <30	227 (60.2)	79 (67.5)	148 (56.9)	
BMI ≥30	136 (36.1)	31 (26.5)	105 (40.4)	
Unknown	14 (3.7)	7 (6)	7 (2.7)	
Follow-up visit				
Time since clinical recovery, days	79 (69–102)	81 (69–108)	79 (69–97)	0.599
Time since virological clearance, days	56 (47–74)	56 (46–79)	56 (47–74)	0.981
Time since symptoms onset, days	102 (86–126)	101 (85–131)	103 (86–124)	0.664
Time from symptoms onset to virological clearance, days	44 (37–53)	43 (35–50)	44 (38–55)	0.081
Blood exams at follow-up				
WBC, × 10 ³ /mm ³	6.07 (5.11–7.07)	5.89 (5.19–7.22)	6.08 (5.1–6.99)	0.770
Hb, g/dL	14.1 (13.2–15)	14.6 (13.7–15.5)	13.9 (12.8–14.8)	<0.0001
PLT, × 10 ³ /mm ³	243 (203–281)	237 (208–272)	246 (202–288)	0.325
Lymphocytes, cell/mm ³	2.2 (1.81–2.6)	2.24 (1.87–2.7)	2.18 (1.79–2.57)	0.274
Creatinine, mg/dL	0.8 (0.67–0.9)	0.8 (0.7–1)	0.8 (0.6–0.9)	0.129
GOT, UI/L	27 (24–31)	25 (21–30)	26 (23–31)	0.198
GPT, UI/L	21 (19–31)	20 (16–29)	20 (16–27)	0.571
LDH, UI/L	196 (177–213)	188 (173–209)	196 (180–219)	0.247
CRP, mg/L	5 (5–5.2)	5 (5–5.1)	5 (5–5.1)	0.484
D dimer, mg/L	120 (81–199)	110 (78–177)	126 (87–204)	0.04

CPAP, continuous positive airway pressure; OTI, orotracheal intubation; NIV, non-invasive ventilation. PI, protease inhibitor (darunavir/cobicistat, darunavir/ritonavir, lopinavir/ritonavir); RDV, remdesivir; HQC, hydroxychloroquine; WBC, white blood cell; Hb, haemoglobin; PLT, platelet count; CRP, C-reactive protein; GOT, serum glutamic oxalacetic transaminase; GPT, serum glutamic pyruvic transaminase; LDH, lactate dehydrogenase.

Mann Whitney and Chi-square for comparison between patients diagnosed with long COVID and patients without long COVID.

Table 2
Comparison between males and females

Parameters	Females (N 137)	Males (N 240)	p
Age, years ^a	58 (50–68)	57 (49–67)	0.931
Comorbidities	57 (41.6)	107 (44.6)	0.672
Smoking:			0.008
Never	94 (68.6)	122 (50.8)	
Ex-smokers	10 (7.3)	10 (4.2)	
Active smokers	31 (22.6)	100 (41.7)	
Unknown	2 (1.5)	8 (3.3)	
Ethnicity:			0.691
White/Caucasian	90 (65.6)	162 (67.5)	
Arabic	6 (4.4)	10 (4.2)	
Black	2 (1.5)	0	
Asian	2 (1.5)	5 (2.1)	
Hispanic	12 (8.8)	21 (8.7)	
Unknown	25 (18.2)	42 (17.5)	
During hospitalization			
Symptoms at admission:			
Fever	96 (70.1)	184 (76.7)	0.215
Cough	61 (44.5)	137 (57.1)	0.02
Sore throat	3 (2.2)	4 (1.7)	0.91
Chest pain	10 (7.3)	6 (2.5)	0.03
Myalgia-arthralgia	16 (11.7)	12 (5)	0.392
Fatigue	21 (15.4)	38 (15.8)	0.914
Headache	4 (2.9)	7 (2.9)	0.999
Dyspnoea	54 (39.4)	107 (44.6)	0.452
Vomiting/nausea	11 (8.1)	14 (5.8)	0.49
Anosmia/dysgeusia	12 (8.8)	12 (5)	0.237
Pneumonia	126 (91.9)	220 (91.7)	0.771
Oxygen therapy:			0.009
No O ₂ therapy	18 (13.1)	17 (7.1)	
Low/high flows systems	79 (57.7)	118 (49.2)	
CPAP, NIV	37 (27)	78 (32.5)	
OTI	3 (2.2)	27 (11.2)	
Length of hospital days (LOS) ^a	10 (6–16)	12 (7–21)	0.033
Treatments for COVID-19:			
Antivirals (RDV or PI)	33 (24.1)	59 (24.5)	0.892
HQC ± azithromycin	90 (65.7)	170 (70.8)	0.356
Steroid	13 (9.5)	29 (12.1)	0.576
Immunomodulating therapies	8 (5.8)	20 (8.3)	0.351
BMI, categories:			0.003
<18.5	5 (3.6)	0	
18.5–24.99	31 (22.6)	45 (18.7)	
25–29.99	39 (28.5)	107 (44.6)	
≥30	56 (40.9)	80 (33.3)	
Unknown	6 (4.38)	8 (3.3)	
Follow-up visit			
Time since virological clearance, days ^a	59 (49–77)	55 (46–72)	0.05
Time from symptoms onset to virological clearance, days ^a	45 (38–55)	44 (37–52)	0.471

Antivirals include: darunavir/cobicistat, darunavir/ritonavir, lopinavir/ritonavir, remdesivir. CPAP, continuous positive airway pressure; OTI, orotracheal intubation; NIV, non-invasive ventilation; HQC, hydroxychloroquine.

^a Quantitative variables are presented as median, (Interquartile Range); categorical variables are presented as absolute numbers (%).

CT scan with contrast medium that excluded pulmonary thromboembolism.

Long COVID was observed in 260/377 (69%) patients; interestingly, 112/137 (81.7%) females presented long COVID syndrome. Within long COVID patients, 97/260 (37.3%) participants had only one persisting symptom, 84/260 (32.3%) had two persisting symptoms and 79/260 (30.4%) had three or more persisting symptoms. More specifically, a significant number of patients (142/260, 55%) reported ongoing physical symptoms only, 100/260 (38%) complained both physical and psychological symptoms, while 18/260 (7%) presented psychological distress solely at follow-up (Fig. S1a). Physical and psychological manifestations were similarly represented in both genders (Fig. S1b,c).

The most common physical symptoms characterising long COVID were fatigue (149/377, 39.5%), exertional dyspnoea (109/377, 28.9%), musculoskeletal pain (80/377, 21.2%) and “brain fog” (76/377, 20.2%) (Table 3: physical symptoms).

As far as psychological sequelae are concerned, manifestations of anxiety were the most frequent (71/377, 18.8%), while depression symptoms were present in 40/377, 10.6% of patients (although 38 patients did not complete HADS questionnaire being non-Italian speakers) (Table 3: psychological symptoms and PTSD). IES-R was available in an unselected sub-group of patients (275/377, 73% patients): in one-third (85/275, 31%) of cases the IES-R score resulted pathological (according to the aforementioned definition), possibly suggesting the presence of PTSD (Table 3: psychological symptoms and PTSD). Interestingly, women were characterized by a higher proportion of most physical symptoms and all psychological symptoms than men (Table 3).

At multivariable logistic regression analysis, female gender was associated with a threefold higher risk of long COVID, also after adjustment for age, severity of illness, LOS, comorbidities, smoking, BMI and time from symptoms onset to virological clearance (Table 4). Other independent predictors of long COVID were

Table 3
Physical and psychological symptoms at follow-up

Parameter	Study population N = 377	Males N = 240	Females N = 137	p
Physical symptoms				
Anosmia				
No, ever	179 (47.5%)	127 (52.9%)	52 (37.9%)	0.017
Ongoing	18 (4.8%)	8 (3.3%)	10 (7.3%)	
Resolved	174 (46.2%)	104 (43.4%)	70 (51.1%)	
Unknown	6 (1.6%)	1 (0.4%)	5 (3.6%)	
Dysgeusia				
No, ever	167 (44.3%)	120 (50%)	47 (34.3%)	0.005
Ongoing	20 (5.3%)	8 (3.3%)	12 (8.7%)	
Resolved	184 (48.8%)	111 (46.2%)	73 (53.3%)	
Unknown	6 (1.6%)	1 (0.4%)	5 (3.6%)	
GI symptoms				
No, ever	201 (53.3%)	143 (59.6%)	58 (42.3%)	<0.001
Ongoing	6 (1.6%)	1 (0.4%)	5 (3.7%)	
Resolved	162 (43%)	91 (37.9%)	71 (51.8%)	
Unknown	8 (2.1%)	5 (2.1%)	3 (2.2%)	
Fever				
No, ever	22 (5.8%)	14 (5.9%)	8 (5.8%)	0.966
Ongoing	0	0	0	
Resolved	348 (92.3%)	223 (92.9%)	125 (91.2%)	
Unknown	7 (1.9%)	3 (1.2%)	4 (2.9%)	
Joint pain or myalgia				
No, ever	215 (57%)	156 (65%)	59 (43.1%)	0.548
Ongoing	80 (21.2%)	30 (12.5%)	50 (36.4%)	
Resolved	74 (19.6%)	50 (20.8%)	24 (17.5%)	
Unknown	8 (2.1%)	4 (1.7%)	4 (2.9%)	
Rest dyspnoea				
No, ever	103 (27.3%)	70 (29.2%)	33 (24.1%)	0.548
Ongoing	24 (6.4%)	16 (6.7%)	8 (5.8%)	
Resolved	242 (64.2%)	150 (62.5%)	92 (67.1%)	
Unknown	8 (2.1%)	4 (1.6%)	4 (2.9%)	
Exertional dyspnoea				
No, ever	87 (23.1%)	61 (25.5%)	26 (19%)	0.036
Ongoing	109 (28.9%)	59 (24.6%)	50 (36.5%)	
Resolved	175 (46.4%)	117 (48.7%)	58 (42.3%)	
Unknown	6 (1.6%)	3 (1.2%)	3 (2.2%)	
Fatigue				
No, ever	67 (17.8%)	50 (20.9%)	17 (12.5%)	<0.001
Ongoing	149 (39.5%)	74 (30.8%)	75 (54.7%)	
Resolved	154 (40.8%)	113 (47.1%)	41 (29.9%)	
Unknown	7 (1.9%)	3 (1.2%)	4 (2.9%)	
Brain fog				
No, ever	274 (72.7%)	190 (79.2%)	84 (61.3%)	<0.001
Ongoing	76 (20.2%)	35 (14.6%)	41 (29.9%)	
Resolved	13 (3.4%)	6 (2.5%)	7 (5.1%)	
Unknown	14 (3.7%)	9 (3.7%)	5 (3.7%)	
Other				
No, ever	300 (79.5%)	198 (82.6%)	102 (74.5%)	0.003
Ongoing	58 (15.4%)	28 (11.7%)	30 (21.9%)	
Resolved	19 (5%)	14 (5.7%)	5 (3.6%)	
Psychological symptoms and PTSD				
IES-R, PTSD				
Normal	190 (50.4%)	133 (55.4%)	57 (41.6%)	0.002
Pathological	85 (22.5%)	43 (17.9%)	42 (30.6%)	
Unknown	102 (27.1%)	64 (26.7%)	38 (27.8%)	
HADS, depression symptoms				
Normal	299 (79.3%)	198 (82.5%)	101 (73.7%)	0.009
Pathological	40 (10.6%)	18 (7.5%)	22 (16.1%)	
Unknown	38 (10.1%)	24 (10%)	14 (10.2%)	
HADS, anxiety symptoms				
Normal	268 (71.1%)	185 (77.1%)	83 (60.6%)	<0.001
Pathological	71 (18.8%)	31 (12.9%)	40 (29.2%)	
Unknown	38 (10.1%)	24 (10%)	14 (10.2%)	

Parameters are presented as absolute numbers (%).

GI, gastrointestinal symptoms; IES-R, impact of event scale - revised for diagnosis of post-traumatic stress disorders; HADS, Hospital Anxiety and Depression Scale for diagnosis of anxiety and depression symptoms; other, chest pain; headache; constipation; tinnitus; insomnia; palpitations; NSTEMI, cough; sore throat.

Chi-square or Fisher's exact test for comparison between males and females.

advanced age and active smoking. Severity of disease, LOS and time to virological clearance were not associated with long COVID at univariable and multivariable analyses (Table 4).

Discussion

The American Center for Disease Control (CDC) and the British NICE have identified symptoms that could persist for weeks or even months after recovery from COVID-19. How long these symptoms could endure for, possible risk factors for their persistence and the predisposing patient features are all aspects to be further elucidated [6–8,17]. Our study reveals that:

1. The most common symptoms characterizing long COVID were both physical and psychological.
2. Females had a 3-fold higher risk of being diagnosed with long COVID.
3. Severity of disease and time to virological resolution were not associated with long COVID.

The high incidence of long COVID within our cohort was similar to that previously reported: more than half of patients reported symptoms lasting more than 2 months after symptom onset [4,5,16,18,19]. Some symptoms are also commonly seen in other viral illnesses and both psychical and psychological sequelae have been described in MERS and SARS [12,23–29].

Literature data are not equivocal about the association between females and long COVID: some preliminary studies have shown an increased prevalence of fatigue [24] or other symptoms among women [1,5,13,16], while in other studies no gender association was found [4,18–20]. Differences in ethnicity, living country and possibly socio-economic status might explain such contrasting results. Hormones may play a role in perpetuating the hyper-inflammatory status of the acute phase even after recovery [30,31] and a stronger IgG antibodies production in females in the early phase of disease has been reported; this could turn out in a more favourable outcome in women [32], but might play a role as well in perpetuating disease manifestations. Furthermore, we might

hypothesise that women are in general more attentive to their body and related distress.

Advanced age was associated with ongoing fatigue and musculoskeletal pain, or impairment in pulmonary functions, reflecting a decline in organ function and a slower ability to recover [1,16]. As far as our finding that reported a weak association between obesity and long COVID is concerned, this was not confirmed in a previous study that however focused on persisting fatigue solely [24]. Increased inflammation, defective adaptive immune responses, endothelial dysfunction and coagulation-related disorders are all phenomena that have been well described in obesity and might represent a plausible explanation to the link we observed between high BMI and long COVID [33]. Finally, the relationship between smoking and long COVID has not been previously explored; the most frequent symptom reported by smokers was however shortness of breath, which possibly reflects an underlying pulmonary dysfunction.

We were expecting to find an association between severity of disease and residual symptoms, as reported by previous studies [1]. However, long COVID has been previously described also in not hospitalised patients diagnosed with a mild, self-limiting disease [6,16–18,24].

Our study has some limitations: a possible selection bias due to losses to follow-up; the limited sample size for patients with severe disease and females; the lack in validated tools to assess dyspnoea and fatigue and in information about the presence of the symptoms before the onset of acute infection. The possible association between female gender and long COVID should be confirmed in larger populations and by means of a longer follow-up. The evaluation of our patients at six months still needs to be completed and this will certainly allow us to collect further data and build stronger evidence.

In conclusion, in our setting long COVID was a frequent long-term complication of COVID-19 and was diagnosed more frequently in women than in men. It was also commonly seen in patients who recovered from a mild disease form. Follow-up outpatient services are therefore needed in order to manage this syndrome and to better understand the possible association

Table 4
Factors associated with long COVID syndrome by fitting univariable and multivariable logistic regression analyses

Parameters	OR (95%CI)	p	AOR (95%CI)	p
Gender:				
Male	1		1	
Female	2.78 (1.68–4.62)	<0.0001	3.32 (1.78–6.17)	<0.0001
Age, 10 years older	1.03 (1.01–1.04)	0.001	1.03 (1.01–1.05)	0.01
O ₂ therapy:				
No O ₂	1		1	
O ₂ therapy low–high flows	0.67 (0.38–1.19)	0.17	0.39 (0.19–0.82)	0.44
CPAP/NIV/IOT	0.97 (0.55–1.71)	0.91	0.67 (0.29–1.55)	0.85
LOS, each day more	1.01 (0.99–1.03)	0.28	0.998 (0.97–1.03)	0.92
Comorbidities:				
No	1		1	
Yes	1.35 (0.86–2.11)	0.19	1.05 (0.597–1.84)	0.87
Smoking:				
Active	1		1	
Unknown	0.13 (0.03–0.52)	0.004	0.16 (0.04–0.75)	0.31
Never	0.61 (0.37–0.997)	0.05	0.56 (0.31–1.01)	0.41
Former	0.36 (0.14–0.96)	0.04	0.19 (0.06–0.62)	0.002
BMI:				
≥30	1		1	
Unknown	0.29 (0.096–0.91)	0.03	0.13 (0.30–0.53)	0.03
<30	0.55 (2.27–5.06)	0.02	0.55 (0.31–0.98)	0.28
Time from symptoms onset to virological clearance, each day more	1.01 (0.99–1.02)	0.64	0.99 (0.98–1.01)	0.47

Each variable is mutually adjusted.

Outcome, long COVID syndrome. AOR, adjusted odds ratio; CPAP, continuous positive airway pressure; OTI, orotracheal intubation; NIV, non-invasive ventilation; LOS, length of hospital stay; BMI, body mass index.

between symptoms and residual organ impairment, and their impact on patient quality of life.

Author contributions

F. Bai wrote the manuscript. D. Tomasoni created the dataset, D. Tomasoni and F. Bai performed statistical analyses. F. Bai, D. Tomasoni, C. Falcinella, D. Barbanotti, R. Castoldi, G. Mulè, M. Augello, D. Mondatore, M. Allegrini, A. Cona, D. Tesoro, G. Tagliaferri, O. Viganò, E. Suardi, C. Tincati, B. Varisco and T. Beringheli followed up patients, contributed to acquisition of data and revised the article. A. Tavelli and S. Terzoni contributed to analyses and interpretation of data. G. Marchetti and A. d'Arminio Monforte contributed to the design of the study and revised the article. C. Luridiana, K. Piscopo, E. Vegni performed psychological evaluations and revised the final version of the manuscript. All the authors have read and approved the final version of the paper.

Transparency declaration

Authors do not have conflicts of interests to declare for this work. Prof. G. Marchetti and Prof. A. d'Arminio Monforte report grants from Gilead, ViiV and Janssen outside this research. Funding: No financial support was obtained for this study.

Acknowledgements

Preliminary data were presented as oral communication at the 31st European Congress of Clinical Microbiology and Infectious Diseases (ECCMID), 9–12 July 2021. We are grateful to all patients and their families for their kind participation in the study. We would like to acknowledge all the staff of the Clinic of Infectious Diseases, Department of Health Sciences, San Paolo Hospital, ASST Santi Paolo e Carlo, University of Milan, Italy, for the precious help in the study.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.cmi.2021.11.002>.

References

- [1] Huang C, Huang L, Wang Y, Li X, Ren L, Gu X, et al. 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study. *Lancet* 2021;397:220–32.
- [2] Editorial Long COVID: let patients help define long-lasting COVID symptoms. *Nature* 2020;586:170.
- [3] Davis AE, Assaf GS, McCorkell L, Wei H, Low RJ, Re'em Y, et al. Characterizing long COVID in an international cohort: 7 months of symptoms and their impact. *EClinicalMedicine* 2021;38:101019.
- [4] Carfi A, Bernabei R, Landi F. Gemelli against COVID-19 post-acute care study group. Persistent symptoms in patients after acute COVID-19. *JAMA* 2020;324:603–5.
- [5] Venturelli S, Benatti SV, Casati M, Binda F, Zuglian G, Imeri G, et al. Surviving COVID-19 in Bergamo Province: a post-acute outpatient re-evaluation. *Epidemiol Infect* 2021;1–25.
- [6] Venkatesan P. NICE guideline on long COVID. *Lancet Respir Med* 2021;9:129.
- [7] Shah W, Hillman T, Playford ED, Hishmeh L. Managing the long term effects of COVID-19: summary of NICE, SIGN, and RCGP rapid guideline. *BMJ* 2021;372:n136.
- [8] Sivan M, Taylor S. NICE guideline on long COVID. *BMJ* 2020;371:m4938.
- [9] Mahase E. COVID-19: what do we know about "long COVID". *BMJ* 2020;370:m2815.
- [10] Mandal S, Barnett J, Brill SE, Brown JS, Denny EK, Hare SS, et al. Long-COVID: a cross-sectional study of persisting symptoms, biomarker and imaging abnormalities following hospitalisation for COVID-19. *Thorax* 2021;76:396–8.
- [11] Carvalho-Schneider C, Laurent E, Lemaignan A, Beauflis E, Bourbao-Tournois C, Laribi S, et al. Follow-up of adults with noncritical COVID-19 two months after symptom onset. *Clin Microbiol Infect* 2020;27:258–63.
- [12] Tomasoni D, Bai F, Castoldi R, Barbanotti D, Falcinella C, Mulè G, et al. Anxiety and depression symptoms after virological clearance of COVID-19: a cross-sectional study in Milan, Italy. *J Med Virol* 2020;93:1175–9.
- [13] Mazza MG, De Lorenzo R, Conte C, Poletti S, Vai B, Bollettini I, et al. Anxiety and depression in COVID-19 survivors: role of inflammatory and clinical predictors. *Brain Behav Immun* 2020;89:594–600.
- [14] Troyer EA, Kohn JN, Hong S. Are we facing a crashing wave of neuropsychiatric sequelae of COVID-19? Neuropsychiatric symptoms and potential immunologic mechanisms. *Brain Behav Immun* 2020;87:34–9.
- [15] Kaseda ET, Levine AJ. Post-traumatic stress disorder: a differential diagnostic consideration for COVID-19 survivors. *Clin Neuropsychol* 2020;34:1498–514.
- [16] Xiong Q, Xu M, Li J, Liu Y, Zhang J, Xu Y, et al. Clinical sequelae of COVID-19 survivors in Wuhan, China: a single-centre longitudinal study. *Clin Microbiol Infect* 2021;27:89–95.
- [17] Centers for Disease Control and Prevention. Long-term effects of COVID-19. Available at: <https://www.cdc.gov/coronavirus/2019-ncov/long-term-effects.html>; 2020.
- [18] Moreno-Pérez O, Merino E, Leon-Ramirez JM, Andres M, Ramos JM, Arenas-Jiménez J, et al. Post-acute COVID-19 Syndrome. Incidence and risk factors: a Mediterranean cohort study. *J Infect* 2021;82:378–83.
- [19] Chopra V, Flanders SA, O'Malley M, Malani AN, Prescott HC. Sixty-day outcomes among patients hospitalized with COVID-19. *Ann Intern Med* 2020;174:576–8.
- [20] Daher A, Balfanz P, Cornelissen C, Muller A, Bergs I, Marx N, et al. Follow up of patients with severe coronavirus disease 2019 (COVID-19): pulmonary and extrapulmonary disease sequelae. *Respir Med* 2020;174:106197.
- [21] Zigmund AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361–70.
- [22] Mak IW, Chu CM, Pan PC, Yiu MG, Chan VL. Long-term psychiatric morbidities among SARS survivors. *Gen Hosp Psychiatry* 2009;31:318–26.
- [23] Ahmed H, Patel K, Greenwood DC, Halpin S, Lewthwaite P, Salawu A, et al. Long-term clinical outcomes in survivors of severe acute respiratory syndrome and Middle East respiratory syndrome coronavirus outbreaks after hospitalisation or ICU admission: a systematic review and meta-analysis. *J Rehabil Med* 2020;52:jrm00063.
- [24] Townsend L, Dyer AH, Jones K, Dunne J, Mooney A, Gaffney F, et al. Persistent fatigue following SARS-CoV-2 infection is common and independent of severity of initial infection. *PLoS One* 2020;15:e0240784.
- [25] Moldofsky H, Patcai J. Chronic widespread musculoskeletal pain, fatigue, depression and disordered sleep in chronic post-SARS syndrome: a case-controlled study. *BMC Neurol* 2011;11:37.
- [26] Hickie I, Davenport T, Wakefield D, Vollmer-Conna U, Cameron B, Vernon SD, et al. Post-infective and chronic fatigue syndromes precipitated by viral and non-viral pathogens: prospective cohort study. *BMJ* 2006;333:575.
- [27] Lee SH, Shin HS, Park HY, Kim JL, Lee JJ, Lee H, et al. Depression as a mediator of chronic fatigue and post-traumatic stress symptoms in Middle East respiratory syndrome survivors. *Psychiatry Investig* 2019;16:59–64.
- [28] Vindegaard N, Benros ME. COVID-19 pandemic and mental health consequences: systematic review of the current evidence. *Brain Behav Immun* 2020;89:531–42.
- [29] Murata S, Rezeppa T, Thoma B, Marengo L, Krancevich K, Chiyka E, et al. The psychiatric sequelae of the COVID-19 pandemic in adolescents, adults, and health care workers. *Depress Anxiety* 2020;38:233–46.
- [30] Mohamed MS, Moulin TC, Schiöth HB. Sex differences in COVID-19: the role of androgens in disease severity and progression. *Endocrine* 2021;71:3–8.
- [31] Bienvenu LA, Noonan J, Wang X, Peter K. Higher mortality of COVID-19 in males: sex differences in immune response and cardiovascular comorbidities. *Cardiovasc Res* 2020;116:2197–206.
- [32] Zeng F, Dai C, Cai P, Wang J, Xu L, Li J, et al. A comparison study of SARS-CoV-2 IgG antibody between male and female COVID-19 patients: a possible reason underlying different outcome between sex. *J Med Virol* 2020;92:2050–4.
- [33] Drucker DJ. Diabetes, obesity, metabolism, and SARS-CoV-2 infection: the end of the beginning. *Cell Metab* 2021;33:479–98.