



Published in final edited form as:

*Otol Neurotol.* 2021 October 01; 42(9): 1329–1333. doi:10.1097/MAO.0000000000003247.

## Tinnitus and Subjective Hearing Loss are More Common in Migraine: A Cross-Sectional NHANES Analysis

Khodayar Goshtasbi, MS<sup>1,\*</sup>, Mehdi Abouzari, MD, PhD<sup>1,\*</sup>, Adwight Risbud, BS<sup>1</sup>, Navid Mostaghni, BS<sup>1</sup>, Ethan Muhonen, MD<sup>1</sup>, Elaine Martin, MD<sup>1</sup>, Hamid R. Djalilian, MD<sup>1,2</sup>

<sup>1</sup>Department of Otolaryngology–Head and Neck Surgery, University of California, Irvine, USA

<sup>2</sup>Department of Biomedical Engineering, University of California, Irvine, USA

### Abstract

**Objectives:** To investigate whether migraine is independently associated with tinnitus and subjective hearing loss (HL) in a large national database.

**Methods:** The de-identified 1999-2004 National Health and Nutrition Examination Survey database was retrospectively queried for subjects aged 18-65. HL and tinnitus were subjectively reported by subjects.

**Results:** A total of 12,962 subjects (52.9% female) with a mean age of 38.1±14.6 years were included. This consisted of 2,657 (20.5%), 2,344 (18.1%), and 2,582 (19.9%) subjects who had migraine, subjective-HL, and tinnitus, respectively. In patients with tinnitus or subjective-HL, migraine was reported in 35.6% and 24.5%, respectively. Migraineurs were more likely to have subjective-HL (25.0% vs. 16.6%,  $p<0.001$ ) and tinnitus (34.6% vs. 16.9%,  $p<0.001$ ) compared to the non-migraineurs. This corresponded to migraine having an odds ratio of 1.5 (95% CI 1.3–1.7,  $p<0.001$ ) and 2.2 (95% CI 2.0–2.4,  $p<0.001$ ) for subjective-HL and tinnitus, respectively. After adjusting for confounders, subjective-HL (OR=1.2, 95% CI 1.1–1.4,  $p=0.003$ ), tinnitus (OR=2.1, 95% CI 1.9–2.3,  $p<0.001$ ), and neck pain (OR=4.0, 95% CI 3.6–4.5,  $p<0.001$ ) were more common in migraineurs. Among migraineurs, a higher proportion of those with tinnitus also had subjective-HL compared to those without tinnitus (40.0% vs. 15.3%,  $p<0.001$ ), and a higher proportion of those with subjective-HL also had tinnitus compared to those without HL (58.1% vs. 27.3%,  $p<0.001$ ).

**Conclusions:** This study suggests an independent association between migraine with subjective-HL and tinnitus. Otologic migraine, which is the effects of migraine on the ear, may be partly responsible for the link between HL, tinnitus, neck pain, and migraine.

### Keywords

Migraine; Tinnitus; Hearing loss; NHANES; Otologic migraine

---

**Corresponding Author:** Hamid R. Djalilian, MD, Division of Neurotology and Skull Base Surgery, Department of Otolaryngology–Head and Neck Surgery, University of California, Irvine, 19182 Jamboree Road, Otolaryngology-5386, Irvine, CA 92697, Phone: (714) 456-5753, Fax: (714) 456-5747, [hdjalili@hs.uci.edu](mailto:hdjalili@hs.uci.edu).

\*K.G. and M.A. contributed equally to this work.

**Conflicts of Interest:** Hamid R. Djalilian holds equity in MindSet Technologies and Cactus Medical LLC, and is a consultant to NXT Biomedical.

## Introduction

Migraine affects approximately 12% of U.S. individuals with significant quality of life consequences.<sup>1</sup> The pathophysiology of migraine is not fully elucidated, but its association with related neuro-otological conditions has been known for years.<sup>2</sup> Two such conditions are tinnitus and hearing loss (HL), which are debilitating chronic conditions experienced by approximately 25%<sup>3</sup> and 20%<sup>4</sup> of the adult population, respectively. An association between migraine and tinnitus among French young individuals was recently reported.<sup>5</sup> Similarly, a study of Australian patients found migraine to increase the risk of developing mild tinnitus.<sup>6</sup> Migraine has also been found to be a risk factor for developing sudden sensorineural HL.<sup>7,8</sup> These possibly secondary conditions have significant effects on patients' health and well-being,<sup>9,10</sup> warranting continued investigation of their relationship with migraine. Although the aforementioned studies utilized French, Taiwanese, and Korean national databases, these associations have yet to be evaluated using U.S. population-based data.

Recent suggestions that migraine may be a risk factor for developing HL and tinnitus have warranted testing this hypothesis on the general U.S. population.<sup>5-8</sup> Investigating these relationships via different approaches and populations will encourage future research into the underlying pathophysiologies and shared otologic principles between them. This is important because if migraine is independently related to tinnitus and HL, comprehensive migraine prevention/treatment protocols may improve the outcomes of these associated otologic conditions. The objective of this study was to investigate the National Health and Nutrition Examination Survey (NHANES) U.S. national database to elucidate whether migraine was associated with subjective HL and tinnitus after adjusting for confounders.

## Methods

This study did not require institutional review board approval because of the publicly available nature of the database and lack of identifying patient information. The NHANES database has been conducted by the National Center for Health Statistics, which is part of the Centers for Disease Control and Prevention since 1971. Since 1999, NHANES has gathered extensive health and diet-related information from thousands of subjects through interviews, physical examinations, and laboratory tests.<sup>11</sup> The 1999-2004 NHANES database was queried for this study, as this was the only timeline that included a migraine question. This is in-line with a 2016 study by Pogoda *et al.* which used the same 1999-2004 NHANES cohort to investigate the association between sodium intake and migraine.<sup>12</sup> Since NHANES provides data categories (e.g., demographic, laboratory, physical, etc.) as separate files, subject-specific sequence numbers were used to merge data into a compiled file. Since migraine was the independent variable, patients with unknown headache status were excluded.

Only patients aged 18-65 were included to exclude subjects with otitis media-related, age-related HL, or a low-reliability test. The upper cutoff for age was set as 65 because prior U.S. population studies have reported a high prevalence of age-related HL in people older than 65,<sup>13,14</sup> which is in-line with several population-based hearing loss studies that have similarly excluded elderly patients.<sup>15-17</sup> Migraine and tinnitus sufferers were

identified from the questions “During the past 3 months, did you have severe headache or migraine?” and “In the past 12 months, have you ever had ringing, roaring, or buzzing in your ears?”. Subjective hearing quality was evaluated by “Which best describes your hearing?” with answers “good”, “little trouble”, “lots of trouble”, and “deaf”. Subjects choosing any of the latter three answers were categorized as having subjective HL. PASW Statistics 18.0 software (SPSS Inc., Chicago, IL) was used for statistical analysis. Two-tailed unpaired t-test and Chi-squared analyses were used to compare continuous and categorical variables, respectively, with a threshold of  $p<0.05$  for significance. Multivariate binary logistic regression was performed to investigate independent associations between migraine and other variables while adjusting for confounders. For logistic regression analysis, the cohort’s median age and body mass index (BMI) were used as threshold for binarizing age/weight variables.

## Results

A total of 12,962 patients (52.9% female) with a mean age of  $38.1\pm 14.6$  years were included. Migraine headache was reported by 2,657 (20.5%) subjects. Patients with migraine tended to be women (68.5% vs. 48.9%), slightly older ( $39.4\pm 12.5$  vs.  $37.8\pm 15.1$  years), with higher BMI ( $29.1\pm 7.1$  vs.  $27.8\pm 6.4$ ), and experience neck pain (38.4% vs. 11.8%) (all  $p<0.001$ ) (Table 1). Tinnitus and HL were reported by 2,582 (19.9%) and 2,344 (18.1%) subjects, respectively. A higher proportion of subjects with any reported tinnitus (34.6% vs. 16.9%,  $p<0.001$ ) or constant tinnitus (5.5% vs. 3.2%,  $p=0.018$ ) existed among migraineurs compared to non-migraineurs (Table 1). There was also a higher proportion of subjects with subjective HL among migraineurs compared to non-migraineurs (25.0% vs. 16.6%,  $p<0.001$ ).

In patients with tinnitus or subjective HL, migraine was reported in 919 (35.6%) and 574 (24.5%), respectively. Among patients with tinnitus, migraineurs tended to be younger than non-migraineurs ( $41.5\pm 12.7$  vs.  $44.4\pm 13.8$  years,  $p<0.001$ ). However, patients reporting migraine, tinnitus, and HL ( $n=353$ , 2.7%) were significantly older than the rest of the cohort ( $45.0\pm 12.2$  vs.  $37.9\pm 14.6$  years,  $p<0.001$ ). Among migraineurs, a higher proportion of those with tinnitus also had subjective HL compared to those without tinnitus (40.0% vs. 15.3%,  $p<0.001$ ). In migraineurs with HL, tinnitus was reported significantly more than those without HL (58.1% vs. 27.3%,  $p<0.001$ ).

Multivariate logistic regression demonstrated that patients with age  $\geq 39$  years (OR=1.16), BMI  $\geq 28.1$  (Odds Ratio [OR]=1.30), female gender (OR=2.38), neck pain (OR=4.04), HL (OR=1.21), and tinnitus (OR=2.09) were all more likely to have migraine (all  $p<0.01$ ) (Table 2). After controlling for age, gender, race, BMI, and neck pain as possible confounders, patients with migraine were significantly more likely to have subjective HL (OR=1.46, 95% confidence interval [CI] 1.30-1.65,  $p<0.001$ ) and tinnitus (OR=2.17, 95% CI 1.95-2.43,  $p<0.001$ ) compared to non-migraineurs. These results remained statistically significant for patients aged 18-60 after excluding patients aged 61-65 ( $n=1,213$ ). Lastly, 2,200 patients had reported answers for presence or absence of positive diagnosis for depression, panic disorder, or generalized anxiety disorder. After adjusting for age, gender, race, BMI, neck pain, depression, panic, and anxiety for these patients, those with migraine were

significantly more likely to have subjective HL (OR=1.42, 95% CI 1.04-1.92,  $p=0.026$ ) and tinnitus (OR=2.17, 95% CI 1.70–2.76,  $p<0.001$ ).

## Discussion

This study of approximately 13,000 U.S. adults demonstrated that migraineurs were more likely to have subjective HL and tinnitus than non-migraineurs even after adjusting for confounders. The observed relationships are not new, yet many otolaryngologists are unaware of this association due to a paucity of literature on the topic. We believe that this paper adds to the current literature on the topic of migraine's association with tinnitus and HL.

HL and tinnitus can be seen during migraine episodes.<sup>18,19</sup> In addition to any subjective HL, major HL (subjectively described as lots of trouble hearing or being deaf) was also significantly more common among migraineurs, although it is difficult to infer that major HL signifies sudden HL with the current data limitation. In-line with our findings, previous analyses have reported a significantly greater risk of developing sudden HL among subjects with migraine<sup>7,8</sup> or non-migraine headache.<sup>20</sup> Specifically, Chu *et al.*'s study of 10,280 Taiwanese subjects reported a 1.8 times greater risk of developing sudden HL among migraineurs compared to those without migraine headache.<sup>7</sup> Likewise, Kim *et al.*'s study of 44,714 Korean subjects found that the adjusted hazard ratio of migraine for sudden HL was 1.3.<sup>8</sup> Among a cohort of sudden HL patients evaluated by Arslan *et al.*, 40% fulfilled migraine criteria as well.<sup>21</sup> It has been suggested that sudden HL in patients with migraine can be attributed to ischemia from migraine-associated vasospasm.<sup>22</sup> A recent study reported improvement in sudden HL outcomes when adjunctive migraine medication were used in addition to standard oral and intratympanic steroids therapy.<sup>23</sup> The improved outcome of sudden HL patients with migraine therapy is likely due to the role of migraine in sudden HL in some of the patients.

In 1996, it was hypothesized that idiopathic HL in pre-existing migraine may be related to migraine-induced vasospasm of cochlear vasculature.<sup>19</sup> This shared epidemiology between migraine and HL or tinnitus may suggest related underlying pathophysiologies. The precise details regarding how migraine occurs are unknown, but cortical spreading depression resulting in release of neuropeptides particularly from the trigeminal nerve may play an important role.<sup>24-26</sup> Trigeminal nerve fibers innervate the spiral modiolary artery and stria vascularis, both important components of the cochlear vascular network.<sup>27,28</sup> When the ophthalmic branch of the trigeminal nerve has been stimulated in an animal model, fluid extravasation within the cochlea was observed, which increased with a greater degree of stimulation.<sup>29</sup> Therefore, it has been proposed that the trigeminal innervation of the cochlear vasculature and stria vascularis likely links migraine and HL.<sup>30</sup>

In addition to subjective HL, tinnitus was also independently associated with migraine in the studied cohort. The high prevalence of tinnitus in different types of migraine has been previously reported.<sup>5,31,32</sup> Tinnitus can also worsen in intensity with migraine episodes,<sup>33,34</sup> or only appear as a concurrent symptom and subside after the headache.<sup>18</sup> For instance, a study by Langguth *et al.* comparing tinnitus patients with (n=193) or without

(n=765) headache reported that those with both tinnitus and headache had higher tinnitus severities.<sup>34</sup> A significant relationship between headache laterality and tinnitus laterality has also been reported.<sup>35</sup> Additionally, HL and tinnitus are prominent features of Meniere's disease (MD), which is strongly associated with migraine.<sup>36,37</sup> MD can also be described as "migraine of the ear" with HL occurring secondary to vasospasm of the inner ear microvasculature.<sup>36,38,39</sup> It has been reported that migraine is common in MD and that migraine prophylaxis medications improves the quality of life in MD, indicating a link between the two conditions.<sup>40</sup> This study further supports the association between migraine and HL/tinnitus. As such, the authors suggest the umbrella term "otologic migraine" to describe migraine-induced ear (otalgia<sup>41</sup> and pressure<sup>42</sup>), cochlear, and vestibular symptoms.

Although we took diligent care in the analysis and interpretation of data, this study contains several limitations. First, the main variables of this study (i.e., migraine, tinnitus, and HL) were qualitatively reported by the patients without objective measures, and further details (e.g., migraine with/without aura, sudden/gradual or conductive/sensorineural HL, unilateral/bilateral tinnitus) could not be elucidated. As such, future institutional studies with more controlled and objective measures are warranted to further evaluate the cross-sectional relationship between migraine with hearing loss and tinnitus. Also, although the migraine cohort may include chronic-severe-headache patients without migraine, a relationship between non-migraine headache and both tinnitus and HL has already been demonstrated.<sup>20</sup> Furthermore, it is difficult to infer that minor or major HL within these patients signify sudden HL due to database limitations. NHANES is a national de-identified database which makes it susceptible to selection bias, recall/reporting bias, inaccurate coding, missing values, and variable limitations. Therefore, the results should be interpreted with caution and future in-depth studies to investigate these relationships are warranted. Despite these limitations, this is the first study utilizing U.S. population-based data to identify significant and independent relationships between migraine, tinnitus, and HL. This encourages future studies to investigate whether migraine prevention/treatment protocols can prevent or decrease these potentially secondary otologic conditions.

## Conclusions

In a large representative population of U.S. adults, migraine was independently associated with an increased likelihood of subjective tinnitus and HL. Patients with subjective HL or tinnitus were also more likely to have migraine after adjusting for various confounders. Female patients, those with a higher than median age, higher BMI, and neck pain had significantly higher representations in the migraine group.

## Financial Disclosure:

Mehdi Abouzari is supported by the National Center for Research Resources and the National Center for Advancing Translational Sciences, National Institutes of Health, through Grant TL1TR001415.

## References

1. Lipton RB, Stewart WF, Diamond S, Diamond ML, Reed M. Prevalence and burden of migraine in the United States: data from the American Migraine Study II. *Headache*. 2001; 41:646–57. [PubMed: 11554952]

2. Kayan A, Hood JD. Neuro-otological manifestations of migraine. *Brain*. 1984; 107:1123–42. [PubMed: 6334543]
3. Shargorodsky J, Curhan GC, Farwell WR. Prevalence and characteristics of tinnitus among US adults. *Am J Med*. 2010; 123:711–718. [PubMed: 20670725]
4. Lin FR, Niparko JK, Ferrucci L. Hearing loss prevalence in the United States. *Arch Intern Med*. 2011; 171:1851–1853. [PubMed: 22083573]
5. Guichard E, Montagni I, Tzourio C, Kurth T. Association Between Headaches and Tinnitus in Young Adults: Cross-Sectional Study. *Headache*. 2016; 56:987–94. [PubMed: 27197786]
6. Sindhusake D, Golding M, Wigney D, Newall P, Jakobsen K, Mitchell P. Factors predicting severity of tinnitus: a population-based assessment. *J Am Acad Audiol*. 2004; 15:269–80. [PubMed: 15239201]
7. Chu CH, Liu CJ, Fuh JL, Shiao AS, Chen TJ, Wang SJ. Migraine is a risk factor for sudden sensorineural hearing loss: a nationwide population-based study. *Cephalalgia*. 2013; 33:80–6. [PubMed: 23197354]
8. Kim SY, Kim MK, Lim J-S, Kong IG, Choi HG. Migraine increases the proportion of sudden sensorineural hearing loss: A longitudinal follow-up study. *Auris Nasus Larynx*. 2019; 46:353–9. [PubMed: 30389313]
9. Kennedy V, Wilson C, Stephens D. Quality of life and tinnitus. *Audiol Med*. 2004; 2:29–40.
10. Arlinger S. Negative consequences of uncorrected hearing loss—a review. *Int J Audiol*. 2003; 42:2S17–12S20. [PubMed: 12918624]
11. Johnson CL, Paulose-Ram R, Ogden CL et al. National health and nutrition examination survey. Analytic guidelines. [CDC.gov](https://www.cdc.gov). 1999-2010 2013.
12. Pogoda JM, Gross NB, Arakaki X, Fonteh AN, Cowan RP, Harrington MG. Severe headache or migraine history is inversely correlated with dietary sodium intake: NHANES 1999–2004. *Headache*. 2016; 56:688–698. [PubMed: 27016121]
13. Ries PW. Prevalence and characteristics of persons with hearing trouble: United States, 1990-91. *Vital Health Stat* 10. 1994; 188:1–75.
14. Homans NC, Metselaar RM, Dingemans JGet al. Prevalence of age-related hearing loss, including sex differences, in older adults in a large cohort study. *Laryngoscope*. 2017; 127:725–730. [PubMed: 27377351]
15. Agrawal Y, Platz EA, Niparko JK. Prevalence of hearing loss and differences by demographic characteristics among US adults: data from the National Health and Nutrition Examination Survey, 1999-2004. *Arch Intern Med*. 2008; 168:1522–1530. [PubMed: 18663164]
16. Glass LE, Elliot H. Work place success for persons with adult-onset hearing impairment. *Volta Rev*. 1993; 95:403–415.
17. Hoffman HJ, Dobie RA, Losonczy KG, Themann CL, Flamme GA. Declining prevalence of hearing loss in US adults aged 20 to 69 years. *JAMA Otolaryngol Head Neck Surg*. 2017; 143:274–285. [PubMed: 27978564]
18. Evans RW, Ishiyama G. Migraine with transient unilateral hearing loss and tinnitus. *Headache*. 2009; 49:756–758. [PubMed: 19472451]
19. Viirre ES, Baloh RW. Migraine as a cause of sudden hearing loss. *Headache*. 1996; 36:24–28. [PubMed: 8666532]
20. Chen Y-C, Tsai S-J, Chen J-C, Hwang J-H. Risks of tinnitus, sensorineural hearing impairment, and sudden deafness in patients with non-migraine headache. *PloS One*. 2019; 14:e0222041. [PubMed: 31490980]
21. Arslan Y, Arslan IB, Aydin H, Yagiz Ö, Tokuçoglu F, Çukurova I. The etiological relationship between migraine and sudden hearing loss. *Otol Neurotol*. 2017; 38:1411–1414. [PubMed: 29076927]
22. Lee H, Lopez I, Ishiyama A, Baloh RW. Can migraine damage the inner ear? *Arch Neurol*. 2000; 57:1631–1634. [PubMed: 11074796]
23. Abouzari M GK, Chua JT, Tan D, Sarna B, Saber T, Lin HW, Djalilian HR. Adjuvant Migraine Medications in the Treatment of Sudden Sensorineural Hearing Loss. *Laryngoscope*. 2020 [Online Ahead of Print]

24. Charles AC, Baca SM. Cortical spreading depression and migraine. *Nat Rev Neurol*. 2013; 9:637–44. [PubMed: 24042483]
25. Harriott AM, Takizawa T, Chung DY, Chen S-P. Spreading depression as a preclinical model of migraine. *J Headache Pain*. 2019; 20:45. [PubMed: 31046659]
26. Malhotra R. Understanding migraine: Potential role of neurogenic inflammation. *Ann Indian Acad Neurol*. 2016; 19:175. [PubMed: 27293326]
27. Vass Z, Shore S, Nuttall A, Jancsó G, Brechtelsbauer P, Miller J. Trigeminal ganglion innervation of the cochlea—a retrograde transport study. *Neuroscience*. 1997; 79:605–615. [PubMed: 9200743]
28. Brain SD, Grant AD. Vascular actions of calcitonin gene-related peptide and adrenomedullin. *Physiol Rev*. 2004; 84:903–934. [PubMed: 15269340]
29. Vass Z, Steyger P, Hordichok A, Trune D, Jancso G, Nuttall A. Capsaicin stimulation of the cochlea and electric stimulation of the trigeminal ganglion mediate vascular permeability in cochlear and vertebro-basilar arteries: a potential cause of inner ear dysfunction in headache. *Neuroscience*. 2001; 103:189–201. [PubMed: 11311800]
30. Vass Z, Shore S, Nuttall A, Miller J. Direct evidence of trigeminal innervation of the cochlear blood vessels. *Neuroscience*. 1998; 84:559–567. [PubMed: 9539226]
31. Farri A, Enrico A, Lacilla M, Sartoris A. Tinnitus during headache: clinical-instrumental evaluation. *Acta otorhinolaryngol Ital*. 1999; 19:70–75. [PubMed: 10434437]
32. Kirchmann M, Thomsen LL, Olesen J. Basilar-type migraine: clinical, epidemiologic, and genetic features. *Neurology*. 2006; 66:880–886. [PubMed: 16567706]
33. Volcy M, Sheftell FD, Tepper SJ, Rapoport AM, Bigal ME. Tinnitus in migraine: an allodynic symptom secondary to abnormal cortical functioning? *Headache*. 2005; 45:1083–1087. [PubMed: 16109127]
34. Langguth B, Hund V, Landgrebe M, Schecklmann M. Tinnitus patients with comorbid headaches: the influence of headache type and laterality on tinnitus characteristics. *Front Neurol*. 2017; 8:440. [PubMed: 28894434]
35. Langguth B, Hund V, Busch V et al. Tinnitus and headache. *Biomed Res Int*. 2015; 2015:797416. [PubMed: 26583133]
36. Ghavami Y, Mahboubi H, Yau AY, Maducdoc M, Djalilian HR. Migraine features in patients with Meniere's disease. *Laryngoscope*. 2016; 126:163–168. [PubMed: 26109273]
37. Radtke A, Lempert T, Gresty M, Brookes G, Bronstein A, Neuhauser H. Migraine and Meniere's disease: is there a link? *Neurology*. 2002; 59:1700–1704. [PubMed: 12473755]
38. Sarna B, Abouzari M, Lin HW, Djalilian HR. A hypothetical proposal for association between migraine and Meniere's disease. *Med Hypotheses*. 2020; 134:109430. [PubMed: 31629154]
39. Atkinson M. Migraine and Meniere's disease. *Arch Otolaryngol*. 1962; 75:220–225. [PubMed: 13863111]
40. Ghavami Y, Haidar YM, Moshtaghi O, Lin HW, Djalilian HR. Evaluating quality of life in patients with Meniere's disease treated as migraine. *Ann Otol Rhinol Laryngol*. 2018; 127:877–887. [PubMed: 30198300]
41. Teixido M, Seymour P, Kung B, Lazar S, Sabra O. Otagia associated with migraine. *Otol Neurotol*. 2011; 32:322–325. [PubMed: 21178808]
42. Paparella MM, Djalilian HR. Etiology, pathophysiology of symptoms, and pathogenesis of Meniere's disease. *Otolaryngol Clin North Am*. 2002; 35:529–545. [PubMed: 12486838]

**Table 1.**

Cohort's demographics and the prevalence of tinnitus and hearing loss.

Variable	Migraineurs (N=2,657)	Non-Migraineurs (N=10,305)	<i>p</i> value
Age	39.4 ± 12.5	37.8 ± 15.1	<b>&lt;0.001</b>
Body mass index	29.1 ± 7.1	27.8 ± 6.4	<b>&lt;0.001</b>
Gender: Female	1821 (68.5%)	5038 (48.9%)	<b>&lt;0.001</b>
Race: White Hispanic/Mexican Black	1169 (44.0%) 818 (30.8%) 559 (21.0%)	4443 (43.1%) 3136 (30.4%) 2285 (22.2%)	0.440
Neck pain	1019 (38.4%)	1214 (11.8%)	<b>&lt;0.001</b>
Subjective hearing loss	663 (25.0%)	1711 (16.6%)	<b>&lt;0.001</b>
Major subjective hearing loss *	84 (3.2%)	190 (1.8%)	<b>&lt;0.001</b>
Tinnitus: Any duration	920 (34.6%)	1737 (16.9%)	<b>&lt;0.001</b>
Tinnitus: Constant duration	147 (5.5%)	327 (3.2%)	<b>0.018</b>

\* Major hearing loss was defined as answering "lots of trouble" or "deaf" to the hearing question.

Statistically significant *P* values are bolded.



**Table 2.**

Multivariate binary logistic regression of factors predictive of an existence of migraine headache while accounting for confounders. All variables were binarized, and odds ratios (OR) were calculated for the second option compared to the first option labeled as “[reference]”. For instance, compared to patients <39 years of age (reference), patients ≥ 39 years old were 1.16 times (95% CI 1.05-1.28) more likely to experience migraine headache

Covariate	Multivariate OR (95% CI)	<i>p</i> value
Age, years		
<39	1 [Reference]	
≥ 39	1.16 (1.05 – 1.28)	<b>0.003</b>
Body mass index		
<28.1	1 [Reference]	
≥ 28.1	1.30 (1.18 – 1.43)	<b>&lt;0.001</b>
Gender		
Male	1 [Reference]	
Female	2.38 (2.15 – 2.63)	<b>&lt;0.001</b>
Subjective hearing loss		
No	1 [Reference]	
Yes	1.21 (1.07 – 1.37)	<b>0.003</b>
Tinnitus		
No	1 [Reference]	
Yes	2.09 (1.87 – 2.34)	<b>&lt;0.001</b>
Neck pain		
No	1 [Reference]	
Yes	4.04 (3.62 – 4.50)	<b>&lt;0.001</b>

OR: odds ratio; CI: confidence interval.

Age and body mass index cutoffs were according to the mean values in the cohort. Statistically significant *P* values are bolded