



EPA Public Access

Author manuscript

Hydro Earth Syst Sci. Author manuscript; available in PMC 2021 November 17.

About author manuscripts

Submit a manuscript

Published in final edited form as:

Hydro Earth Syst Sci. 2018 March 15; 22(3): 1851–1873. doi:10.5194/hess-22-1851-2018.

Wetlands inform how climate extremes influence surface water expansion and contraction

Melanie K. Vanderhoof^{1,*}, Charles Lane², Michael McManus³, Laurie Alexander⁴, Jay Christensen⁵

¹U.S. Geological Survey, Geosciences and Environmental Change Science Center, P.O. Box 25046, DFC, MS980, Denver, CO 80225

²U.S. Environmental Protection Agency, Office of Research and Development, National Exposure Research Laboratory, 26 W. Martin Luther King Dr., MS-A110, Cincinnati, OH 45268

³U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Assessment, 26 W. Martin Luther King Dr., MS-A110, Cincinnati, OH 45268

⁴U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Assessment, 1200 Pennsylvania Ave. NW (8623-P), Washington, DC 20460

⁵U.S. Environmental Protection Agency, Office of Research and Development, National Exposure Research Laboratory, Environmental Science Division, 944 E. Harmon Ave., Las Vegas, NV 89119

Abstract

Effective monitoring and prediction of flood and drought events requires an improved understanding of how and why surface-water expansion and contraction in response to climate varies across space. This paper sought to (1) quantify how interannual patterns of surface-water expansion and contraction vary spatially across the Prairie Pothole Region (PPR) and adjacent Northern Prairie (NP) in the United States, and (2) explore how landscape characteristics influence the relationship between climate inputs and surface-water dynamics. Due to differences in glacial history, the PPR and NP show distinct patterns in regards to drainage development and wetland density, together providing a diversity of conditions to examine surface-water dynamics. We mapped surface-water extent across eleven Landsat path/rows representing the PPR and NP (images spanning 1985–2015). The PPR not only experienced a 2.6-fold increase of surface-water extent under median conditions relative to the NP, but also showed a 3.4-fold greater difference in surface-water extent between drought and deluge conditions. The relationship between surface-water extent and accumulated water availability (precipitation minus potential evapotranspiration) was quantified per watershed and statistically related to variables representing hydrology-related landscape characteristics (e.g., infiltration capacity, surface storage capacity, stream density). To investigate the influence stream-connectivity has on the rate at which surface water leaves a given location, we modeled stream-connected and stream-disconnected surface water separately. Stream-connected surface water showed a greater expansion with wetter climatic conditions in landscapes with greater total wetland area. Disconnected surface water showed a greater expansion

* mvanderhoof@usgs.gov, phone: 303-236-1411.

with wetter climatic conditions in landscapes with higher wetland density, lower infiltration and less anthropogenic drainage. From these findings, we can expect that shifts in precipitation and evaporative demand will have uneven effects on surface-water quantity. Accurate predictions regarding the effect of climate change on surface-water quantity will require consideration of hydrology-related landscape characteristics including wetlands.

Keywords

Drought; evapotranspiration; Landsat; prairie pothole region; precipitation; surface water

1. Introduction

Surface-water dynamics have strong implications for ecosystem functioning and human land use including biogeochemical balances (Hoffmann et al., 2009), species distribution (Boschilia et al., 2008; Calhoun et al., 2017), hydrologic connectivity (Heiler et al., 1995; Pringle, 2001), and agricultural productivity (Mokrech et al., 2008; Gornall et al., 2010). Yet natural variability in surface-water extent poses a basic challenge to gathering timely, accurate information (Poff et al., 1997; Beeri and Phillips, 2007). While satellite imagery can be used to map variability in surface-water extent over time, predicting future changes in surface-water extent (e.g., in response to changes in climate, land use, or natural disasters) requires improving our understanding of how the landscape influences surface-water extent over time and space. The relative importance of hydrologic processes and flowpaths across a landscape (e.g., surface storage, infiltration, evapotranspiration, runoff) can be expected to influence the timing, duration and extent of surface water for a given location (Euliss and Mushet, 1996; LaBaugh et al., 1996, van der Kamp et al., 1999)

Winter (2001) presented the concept of hydrologic landscapes as a means to classify landscape units based on their hydrologic attributes (land-surface form, geology and climate). These attributes, it is argued, could then be used to predict the partitioning of water into storage, infiltration, evapotranspiration and runoff (Wagener et al., 2007). In many landscapes when rainfall intensity is greater than both the rate of soil infiltration and the soil moisture deficit it is assumed that overland flow, subsurface flow, and groundwater flow will dominate, contributing to increased stream discharge (Eamus et al., 2006). These landscapes could be described as exhibiting a lower potential for surface-water expansion. Alternatively, in landscapes with low topographic gradients and poorly developed drainage networks, runoff events rarely deplete available surface storage, meaning that although stream discharge may elevate, much of the surplus water remains as surface water (Shaw et al., 2012; Kuppel et al., 2015). These landscapes show a higher potential for surface-water expansion with evapotranspiration often the primary mechanism for water loss (Winter and Rosenberry, 1998). Landscapes with a tendency to accumulate surface-water are relatively common across the globe and include former glacial landscapes including the Prairie Pothole Region (PPR) (Sass and Creed, 2008; Shaw et al., 2012), and parts of China (Yao et al., 2007) and Russia (Stokes et al., 2007), permafrost regions (Smith et al., 2007), as well as low gradient landscapes including the Argentine Pampas (Kuppel et al., 2015); the Pantanal in Brazil (Hamilton, 2002), and the Orinoco Llanos in Columbia and Venezuela (Hamilton,

2004). Although such landscapes have previously been shown to experience surface-water expansion in response to increased precipitation (Huang et al., 2011; Kuppel et al., 2015; Vanderhoof et al., 2016) or melting ice (Stokes et al., 2007; Yao et al., 2007), we are unaware of studies that have explicitly compared surface-water expansion and contraction between landscapes of differing surface-water expansion potential.

The PPR and adjacent Northern Prairie (NP), which span the upper mid-west of the United States, occur within and beyond the last glacial maximum, respectively, and together represent a range in the potential for surface-water expansion. The PPR is characterized by a high density of depression wetland and lake features (Zhang et al., 2009), a relic of glacial retreat (Flint, 1971). Most wetlands are relatively small (< 0.5 ha) depressions, underlain by glacial till with low permeability, and occur within a landscape matrix of natural grassland and agriculture (Winter and Rosenberry, 1995; Zhang et al., 2009; Cohen et al., 2016). This is in contrast to the adjacent NP such as the Northwestern Great Plains (Montana, western North and South Dakota) and the Central Irregular Plains (southern Iowa and northern Missouri), which lack the high density of small wetlands and show a well-developed drainage network due to its occurrence outside of the last maximum glacial extent (USGS, 2013). The NP and PPR are also characterized by substantial spatial and interannual variability in air temperature and precipitation (Bryson and Hare 1974). Variations in temperature and moisture content of competing air masses results in a strong north-south temperature and east-west precipitation gradient. The precipitation-evaporation deficit is least in the east (i.e., Minnesota and Iowa), and increases to the west (i.e., Montana) (Kantrud et al., 1989; Millet et al., 2009). This variability in climate has a strong influence on water levels across the region. In the PPR in spring, wetland depressions receive water from both precipitation and snowmelt. In the summer, water level is controlled by direct precipitation, evaporation and wetland vegetation transpiration (Winter and Rosenberry, 1995; LaBaugh et al., 1998; Carroll et al., 2005), with evapotranspiration typically dominating water loss (Rosenberry et al., 2004).

Monitoring variation in water levels across the PPR has been of high interest as it is a key factor in flood abatement, water quality, biodiversity, carbon management and aquifer recharge (Gleason et al., 2008). Water level data at Devils Lake, North Dakota, for example, have been collected as far back as 1867 and provide a regional indicator of hydrological conditions (Wiche, 1996; LaBaugh et al., 1996). Efforts have been expanded to map interannual changes in surface-water extent across the PPR at a landscape scale using remotely sensed imagery (Kahara et al., 2009; Niemuth et al., 2010; Vanderhoof et al., 2016). However, while substantial interannual variation in water level has been documented across the PPR (Huang et al., 2011; Vanderhoof et al., 2016), and primarily attributed to interannual variation in temperature and precipitation (Johnson et al., 2005; Zhang et al., 2009), such surface-water patterns have to date been minimally characterized for the remainder of the NP. In addition to interannual patterns of temperature and precipitation, we would also expect that surface-water extent will depend on landscape parameters such as infiltration capacity, storage capacity, and drainage characteristics (Euliss and Mushet, 1996; LaBaugh et al., 1996; van der Kamp et al., 1999). Spatial models incorporating some of these factors can provide additional insights into the risk of flood and drought events across the PPR (Niemuth et al., 2010).

The PPR, in conjunction with adjacent NP, provides an ideal physiographic example in which to analyze the influence of landscape characteristics on surface-water expansion and contraction. Although the interaction between water level and climate has been studied extensively at select locations within the PPR (e.g., Cottonwood Lake) (Winter and Rosenberry, 1998; Huang et al. 2011), minimal research has sought to understand spatial variability in the relationship between climate and surface-water extent. Our research questions addressed in this study are:

1. How do interannual patterns of surface-water expansion and contraction vary spatially across the Prairie Pothole Region and adjacent Northern Prairie of the United States?
2. How do landscape characteristics influence the relationship between climate inputs and surface-water dynamics?

The successful exploration of this spatial patterning and landscape-scale statistical functions will inform hydrologic/hydraulic and biogeochemical modeling and has implications for biodiversity/habitat modeling and management (e.g., Allen et al., 2016; Golden et al., 2017)

2. Methods

As detailed below, we used Landsat imagery to map surface-water extent under dry, average, and wet conditions across portions of the PPR and adjacent NP. We compared the expansion and contraction of surface-water extent between the PPR and adjacent NP. As stream-connected surface water can leave a location easily as stream flow, stream-connected and disconnected surface water were analyzed separately. We then used a two-level modeling approach to investigate the influence of landscape variables on surface-water dynamics. In the first stage, surface-water extent per watershed was statistically related to accumulated water availability, defined as precipitation minus potential evapotranspiration. This first stage produced the dependent variable for the second model, the slope of the relationship between surface-water extent and climate inputs per hydrological unit (a watershed) or the Surface Water Climate Variable (SWCV). The SWCV was then regressed against independent variables representing landscape characteristics (e.g., infiltration capacity, surface storage capacity, stream density, long-term climate normals). This approach allowed us to explore what landscape characteristics drive spatial variability in the relationship between surface-water extent and climate.

2.1 Study Area

Our study area consisted of eleven Landsat path/rows (total area = 308,439 km²) in the U.S. portion of the PPR and adjacent NP (Figure 1). The PPR across North and South Dakota, western Minnesota, northern Iowa and northern Nebraska, is dominated by the North and Northwest Glaciated Plains. This ecoregion is characterized by landscape features formed during its recent glacial history. Drift plains, large glacial lake basins and shallow river valleys support row crop agriculture. Grasslands and livestock grazing dominate areas where glaciers left deposits of uneven glacial till (Sayler et al., 2015). The PPR is dominated by cultivated crops (59%), herbaceous (18%) and hay/pasture (10%) (Homer et al., 2015). Adjacent to the PPR, the Northwestern Great Plains, across western North and South

Dakota, is a semiarid unglaciated plain which tends to have shallow soils with a clay texture not conducive to growing crops and instead dominated by livestock grazing across grasslands (Sayler et al., 2015). To the southeast of the North Glaciated Plains lies the Western Corn Belt and the Central Irregular Plains in Iowa and Nebraska. Glacial till forms the parent material for most of the soil in Western Corn Belt and the northern part of the Central Irregular Plains, within the study area. Level and gently rolling hills and fertile soils support agriculture (Sayler et al., 2015). The NP is dominated by herbaceous land cover (47%) with cultivated crops (28%) and hay/pasture (9%) also common (Homer et al., 2015). Using the precipitation averages (1981–2010) defined by the Parameter-elevation Regressions on Independent Slopes Model (PRISM, Daly et al., 2008), the PPR study area receives 6% more precipitation on average than the NP study area (626 mm yr⁻¹ relative to 592 mm yr⁻¹, respectively) and 1.5% less evaporative demand or potential evapotranspiration (PET) (603 mm yr⁻¹ relative to 594 mm yr⁻¹, respectively). These differences were not found to be statistically different using the Wilcoxon rank sum test.

Our regression analysis used eight-digit Hydrologic Unit Codes (HUC8s; USDA NRCS, 2015) as the unit of analysis (n=150) across all eleven Landsat path/rows (Figure 1). HUC8s were used instead of smaller watersheds such as HUC10s or HUC12s to ensure that patterns in surface-water expansion and contraction represented landscape patterns, not individual or small groups of water features. HUC8s that occurred at the edge of a Landsat path/row with an area of < 50 ha were excluded from further regression analysis to limit the inclusion of incompletely characterized watersheds. The threshold of 50 ha was selected as it was a natural break in the distribution of HUC8 sizes. Patterns of surface-water expansion and contraction were compared between the PPR and NP. We note that one path/row (p37r26) in northern Montana was technically within the most western section of the PPR, but was found to behave dissimilarly from the PPR and similarly to the NP in terms of both its landscape characteristics (e.g., stream density, wetland density) and surface-water expansion and contraction. Because of this, p37r26 was included in the adjacent NP for analyses where findings were organized by PPR and NP.

2.2 Landsat Image Processing

2.2.1 Path-Row and Image Selection—Surface-water extent was mapped for a series of images across 11 Landsat path/rows (Figure 1). These path/rows were selected to represent the PPR and adjacent NP and were intentionally selected to represent a range of ecoregions, climate conditions (west to east and north to south) and densities of wetlands and streams. Snow-free images (acquired approximately from April through October) containing less than 10% cloud cover from the Landsat 4–5 TM, Landsat 7 ETM+ (prior to failure of the scan-line corrector in 2003) and Landsat 8 OLI sensors were selected between 1985 and 2015. The number of images processed within each path/row averaged 14 (range: 9 to 17 acceptable images) and were intentionally selected to document interannual variability in surface-water extent, by selecting images from wet, average and dry years (Table 1). The terms “wet”, “average” and “dry” were defined in reference to local norms, using the Palmer Hydrological Drought Index (PHDI) and the 12-month Standardized Precipitation Index (SP12) (NOAA, NCDC, 2014). The range of conditions captured by the time series within each path/row in relation to the historical climate conditions (1895–2015)

are shown in Table 1. The PHDI is based on a monthly water balance accounting approach that considers precipitation, evapotranspiration, runoff and soil moisture. The indices rely on weather station data and are interpolated at 5 km (NOAA NCDC, 2014). A complete list of images included in the analysis is presented in the Appendix (Table A1).

2.2.2 Image Processing—Images were atmospherically corrected and converted to surface reflectance values using the Landsat Ecosystem Disturbance Adaptive Processing System (Masek et al., 2006). A minimum noise fraction transformation was applied to reduce within-image noise (Green et al., 1988). The per-pixel water fraction was estimated using the Matched Filtering algorithm, a partial unmixing method in the ENVI software package (Exelis Visual Information Solutions, Inc, Herndon, Va) (Turin, 1960; Vanderhoof et al., 2016). This algorithm is trained on a water spectral signature, which was derived from open-water polygons manually selected within each path/row, resulting in a water signature specific to each image. The water fraction output was linearly stretched to maximize our ability to separate water from non-water. CFmask, a quality control layer provided with Landsat images, was used to mask out clouds and cloud shadows (Zhu and Woodcock, 2014), while the National Land Cover Database (NLCD) (2011) was used to mask out impervious surfaces, defined as low, medium and high density development (Homer et al., 2015), which can show spectral confusion with surface water. Each surface-water image was visually inspected for quality using visual interpretation as well as ancillary datasets (e.g., National Agricultural Imagery Program (NAIP) imagery, National Wetland Inventory (NWI) dataset (USFWS, 2010)). Select images were removed or edited primarily due to spectral confusion between water and bare rock or shadowed vegetation.

2.2.3 Surface-Water Extent Validation—The surface-water extent maps were validated using 1-m resolution NAIP imagery. Landsat images were selected for validation based on the temporal coincidence of the Landsat and NAIP imagery collections (Table 2). Because terrestrial surface water is a relatively rare cover type, it is difficult to generate enough inundated reference points through a simple random-point generation. Therefore, random points were generated in reference to NWI polygons overlapping with the NAIP and Landsat imagery. Points were then visually identified as inundated or non-inundated using the NAIP imagery. To account for the scale difference between a random point and a 900 m² Landsat pixel, the Landsat pixel boundaries for each random point were identified. The point was classified as the majority class (inundated or non-inundated) identified by NAIP within the Landsat pixel boundary surrounding each random point. Reference points were generated per Landsat/NAIP pair (500 or 1000), with the number of reference points varying depending on the amount of NAIP imagery available within the Landsat path/row extent, and the number of random points that occurred within Landsat NA pixels. Metrics presented included overall accuracy, omission error, commission error, dice coefficient, and relative bias. Omission and commission errors were calculated for the category “water”. The dice coefficient is the conditional probability that if one classifier (product or reference data) identifies a pixel as water, the other one will as well, and therefore integrates omission and commission errors (Fleiss, 1981; Forbes, 1995). The relative bias provides the proportion that water is under (negative) or overestimated (positive).

The Landsat per-pixel fraction water was binned into inundated (≥ 0.3) and non-inundated (< 0.3) classes. This threshold was selected as it best balanced errors of omission and commission. Overall accuracy for the Landsat surface-water maps across the 11 path/rows was 93.9% with errors of omission for surface water averaging 8.5% and errors of commission for surface water averaging 8.2% (Table 3). The surface-water maps showed no relative bias and a dice coefficient of 92%. Errors of omission and commission can be primarily attributed to mixed Landsat pixels occurring over small wetlands (a few pixels in size) or at the edge of larger wetlands or open water features. In some images parts of or entire agricultural fields were classified as water. It is common in both the spring months, when crops need to be planted, and fall months, when crops are being harvested, for fields to experience wet conditions (Fausey et al., 1987; King et al., 2014). In addition, poorly drained soil is common across this region (Skaggs et al., 1994) and wetland depressions often occur within agricultural fields. Consequently, subsurface tile drainage has become increasingly popular across the region to speed up the removal of excess soil water (Blann et al., 2009). It is often unclear to what extent surface-water mapped within agricultural fields represents historical or current wetlands, poorly drained fields, or misclassified pixels. Lastly, a close match in acquisition date between the Landsat and NAIP images is essential for the NAIP imagery to accurately represent ground conditions. Variability in the date match can be considered one potential source of error, as the occurrence of a rain event or seasonal variability can change surface-water conditions over even short time periods.

2.3 Surface-Water Extent Analysis

Surface-water abundance (ha km^{-2}) was calculated per HUC8 with HUC8 area being adjusted for each image based on the abundance of not applicable (NA) pixels (e.g., cloud cover, cloud shadow) in each image. We used the high-resolution National Hydrography Dataset (NHD, 1:24,000) to classify surface water as (1) continuous connected with the stream network, or (2) disconnected from the stream network. The NHD line dataset was buffered by 14 m, the reported digital horizontal accuracy of the dataset (USGS, 2000) and NHD area was added to account for the width of large rivers. Surface-water polygons that intersected the stream network in a given image were classified as continuously connected water (CCW). Surface-water polygons that did not intersect the stream network in a given image were classified as discontinuous water (DCW) or discontinuous from the stream network. We acknowledge that the NHD is known to be incomplete (e.g., lacking short and ephemeral stream lines) and that some stream lines within the NHD are disconnected from downstream waters (Heine et al., 2004). However, the NHD is the most complete nationally-available stream dataset.

Processed images within each path/row were ranked from least-to-most amount of surface water per area. Median condition was defined as the image or two images representing the median amount of surface-water extent, estimated from all images within a path/row. Drought and deluge conditions were defined as the average of the two end-member images showing the least and most amount of total surface-water extent for each path/row, respectively. Surface-water extent was then summed across the PPR and NP path/rows and divided by the total area to calculate the hectares of surface-water extent per km^2 for each

region. The NP portion of path 27, row 30 (p27r30) and p30r30 were deleted, as was the PPR portion of p26r30 to avoid double counting overlapped path/rows.

2.4 Stage One – Derivation of the Surface Water Climate Variable (SWCV)

In stage one, surface-water extent in each HUC8 was related, using linear regression, to water availability, defined as precipitation minus PET summed over a time interval. Water availability provided an estimate of the amount of water in each watershed available to either (1) runoff, (2) infiltrate to shallow or deep groundwater sources, or (3) be stored as surface-water. Surface water was again partitioned into CCW and DCW using its spatial relationship to the NHD. Precipitation data were compiled using the Parameter-elevation Regressions on Independent Slopes Model (PRISM, Daly et al., 2008). PET, or the atmospheric demand for evaporation and transpiration in the absence of water limitations, which can be expected over open surface water, was compiled using gridded surface meteorological data PRISM and the North American Land Data Assimilation System Phase 2 (Abatzoglou et al., 2011). PET was calculated using the Penman-Monteith equation that required inputs of minimum and maximum temperature, daily average dewpoint temperature (equivalently, vapor pressure or vapor pressure deficit), wind speed and downward shortwave radiation (Abatzoglou et al., 2011, Mitchel et al., 2004). The datasets were resampled to 125 m using cubic convolution and summarized for each HUC8. Water availability was summed for a series of monthly periods preceding each image date (3, 6, 9, 12, 18, 24, 30 and 36 months) to identify the accumulation period for which the greatest number of HUC8's showed a significant ($p < 0.05$) slope between water availability and surface-water extent. This logic was meant to reduce the probability that a zero slope resulted from surface water responding more strongly to climate drivers at a different time interval. This first stage produced surface water climate variables (SWCV), our dependent variables for stage 2, i.e., the slope of the relationship between CCW and DCW surface-water extent to accumulated water availability (Figure 2). The slope or stage 2 dependent variable is referred to as the surface water climate variable (SWCV) from this point forward.

Cloud cover makes it challenging to restrict analysis of Landsat imagery to a specific season, while including imagery that covers more than one season potentially conflates seasonal surface-water dynamics with interannual surface-water dynamics. The influence of seasonal change in surface-water extent within our analysis contributed to the uncertainty (primarily through sampling error) in the SWCV. For example, if we included an image from June 1993 and one from August 1993 and related both images to the last nine months of precipitation and PET (Sept 1992 - May 1993 and November 1992 – July 1993, respectively), greater seasonal dynamics or variation in surface-water extent between the two dates can be expected to show up as greater uncertainty in the slope, defined by the standard error of the slope. This becomes more evident as the accumulated period becomes larger (e.g., 36 months). By explicitly considering the uncertainty of the slope in the regression analysis, as described below in the Stage 2 Analysis (Section 2.6), we can, to the extent possible, account for seasonally induced variation in surface-water extent.

2.5 Landscape Variables for Stage 2 Analysis

The independent variables summarized for each HUC8 and included in the analysis were selected to characterize mechanisms through which water can leave the landscape (e.g., infiltration, runoff, tile drainage), mechanisms through which water can remain and expand on the landscape (e.g., wetland density, wetland size, topography), as well as other potential influences on surface water dynamics (e.g., climate norms, land cover). The National Wetland Inventory (USFWS, 2010) and NHD stream dataset (USGS, 2013) were used to calculate wetland and stream characteristics including stream density, wetland count and areal density, and proportion of total wetland area attributed to large (>8 ha) features. A threshold of 8 hectares was selected as this is the size threshold used by USFWS to define a lacustrine system (Cowardin et al., 1979). We do not refer to these features as lakes, however, as water depth and associated vegetation are also important features to defining lacustrine systems, and were not evaluated. We did not include distance variables, which were previously found to be highly correlated with simpler variables already in the analyses: mean wetland-to-wetland distance was previously found to be highly correlated with wetland density ($r = -0.95$, $p < 0.01$) and mean wetland-to-stream distance highly correlated with stream density ($r = 0.88$, $p < 0.01$) (Vanderhoof et al., 2017). Surface topography can influence the capacity for surface water to expand and was quantified as the weighted averaged slope gradient, as defined by the U.S. Department of Agriculture's Soil Survey Geographic (SSURGO) Database (Soil Survey Staff, 2017). Topographic Wetness Index was not included because of the relative weakness of such indices in landscapes with little relief (e.g., Schmidt and Persson, 2003) and the data intensive nature of calculating TWI with a 10 m DEM across such a large study area. Additional variables derived from the SSURGO database to characterize infiltration capacity include available water storage (0 – 150 cm), annual minimum depth to water table, and saturated hydraulic conductivity (Ksat). Human influence was quantified as the abundance of agricultural activities, or the percent of each HUC8 classified as agriculture, defined as the NLCD (2011) cover categories hay/pasture and row crop. Anthropogenic modifications to drainage systems, or the percent land cover artificially drained, was estimated as the percent of each HUC8 where row crop cover type (NLCD 2011) and very poorly drained or poorly drained soils as defined by the National Resources Conservation Service's SSURGO database were collocated following Christensen et al., (2013). The climate normals per HUC8 (1989–2013) were calculated to represent the Landsat image range. The precipitation averages are provided as part of the PRISM dataset (Daly et al., 2008). PET was calculated as a function of monthly mean PRISM temperature and day length following Hamon (1961). The Moisture Index (MI) was calculated as the ratio of precipitation and PET where, if PET exceeded precipitation, $MI = \text{precipitation}/PET - 1$, and if precipitation exceed or equaled PET, then $MI = 1 = PET/\text{precipitation}$. Values range from -1 (dry) to 1 (wet) (Willmott and Feddema, 1992; Feddema, 2005). The climate averages were resampled to 1 km from 4 km using inverse-distance weighting, prior to being averaged per HUC8. The distribution of values within each of the independent variables are shown in Table 4. Spearman rank correlations with a Bonferroni correction (Dunn, 1961) were calculated for the independent variables (Table 5).

2.6 Stage 2 - Analysis - Landscape Mechanisms Explaining Variability in SWCV

In stage two, CCW and DCW SWCVs, or the slope of the relationship between CCW and DCW and accumulated water availability, were related to landscape variables using feasible generalized least-squares (FGLS) regression, with HUC8s ($n=150$) as the unit of analysis. FGLS allowed us to estimate the heteroscedastic structure of the residuals (Lewis and Linzer, 2005) and has been previously applied within landscape ecology contexts (e.g., Acharya, 2000; Villalobos-Jiménez and Hassall, 2017). The SWCVs were found to be significant for the largest number of HUC8s using a 9-month period of accumulation for both CCW and DCW, which was therefore used as the accumulation period for further analyses (Table 6). The SWCVs were found to be spatially autocorrelated using Global Moran's I (spatial relationship conceptualized using inverse distance) (DCW SWCV, 9 months, $z\text{-score}=7.8$, $p<0.01$, CCW SWCV, 9 month, $z\text{-score}=4.1$, $p<0.01$), violating the assumption of independence between samples. To account for spatial autocorrelation in the SWCVs, we calculated an autocovariate in ArcGIS 10.3, Geostatistical Analyst (ESRI, Redmond CA) which uses adjacent HUC8s to create a neighbor value. By including a spatial autocovariate in the ordinary least-squares (OLS) regression model, we controlled for how much the response variable reflected response values of adjacent HUCs, before identifying additional significant explanatory variables (Dormann et al., 2007; Betts et al., 2009). The autocovariate was automatically retained while only significant independent variables ($p<0.05$) were additionally retained. The dependent variable was normalized using a Box-Cox power transformation (R package MASS, Venables and Ripley, 2002). Multicollinearity was formally assessed using the regression collinearity diagnostics described by Belsley et al. (1980) and implemented in the R package perturb (Hendrickx, 2012). Collinearity may affect parameter estimation when a condition index greater than 10 is associated with variance decomposition proportions greater than 0.5 for two or more explanatory variables (Belsley, 1991). Both models complied with collinearity requirements.

Having an estimated dependent variable (e.g., SWCV) does not necessarily present a problem for a regression analysis, but we must recognize that the regression model error term contains two components: (1) the expected random error resulting from sources of variation not accounted for in the model, and (2) the difference between the true value of the dependent variable and the estimated value (sampling error). In this study, the uncertainty around the dependent variable (SWCV) was not constant across observations. Instead, the dependent variable showed a strong positive correlation with its standard error (DCW SWCV, $R^2 = 0.59$, $p<0.05$; CCW SWCV, $R^2 = 0.70$, $p<0.05$) (Figure 3). FGLS allowed us to estimate both components of the error. To do so, we: (1) calculated the logarithm of squared residuals from the OLS model, (2) regressed the log-residuals on the independent variables included in the OLS model, (3) calculated the exponential of fitted values from that regression, which estimates the variance of the regression residual that is not due to sampling of the dependent variable, z , and (4) estimated the basic model again now including weights ($1/z^2$) (Hanushek, 1974; Lewis and Linzer, 2005). We found the final model residuals to be random using the studentized Breusch-Pagan test (Breusch and Pagan, 1979).

To help add confidence regarding which landscape variables were more or less important, we also fit random forest models in R using the package `randomForest` (Liaw and Wiener, 2015). The random forests were run with the SWCVs as the dependent variable and landscape characteristics as independent variables. We derived 500 binary trees or bootstrap iterations using out of bag (OOB) samples (70% of samples to train and 30% of samples to validate). Variable importance was calculated as the change in node impurity (i.e., Gini importance). Random forest models are generally insensitive to collinearity among metrics, however the inclusion of correlated variables can deflate variable importance as well as the overall variation explained by the model (Murphy et al., 2010). We implemented random forest model selection to select the smallest number of non-redundant variables (`varSelRF` R package) (Murphy et al., 2010).

3 Results

3.1 Surface-Water Extent

Median surface-water extent as well as the amount of water added and lost from the surface between wet and dry periods was found to vary considerably across the study area (Figure 4 and 5). Analysis of the median total surface-water extent between the PPR and the NP demonstrated that the PPR had 2.6 times greater surface-water extent than the NP (Table 7). The PPR also showed greater variability in total surface-water extent, adding 5.7 ha km⁻² during very wet conditions and losing 2.8 ha km⁻² during very dry conditions, for a maximum net difference of 8 ha km⁻². This can be compared to the NP which gained 1.6 ha km⁻² during very wet conditions and lost 0.8 ha km⁻² during very dry conditions, a net difference of 2.4 ha km⁻² (Table 7). DCW, or water that was discontinuous with the stream network, showed greater expansion and contraction in extent in both the PPR and NP, relative to CCW which intersected the stream network. Consequently, DCW increased as a percent of total surface water during wet periods and decreased as a percent of total surface water in dry periods. This suggests that across the study area, surface water that was disconnected from the stream network disproportionately served a surface water storage function during wet periods, reducing the amount of water contributing to downstream flooding. Similarly, DCWs disproportionately experienced loss during dry periods.

3.2 Relationship between Surface-Water Extent and Water Availability

Including PET instead of using precipitation alone tended to increase the percentage of HUC8s showing a statistically significant relationship between surface-water extent and water availability across the different accumulation periods that we tested, although this was not true for all time periods. For instance, the percent change from precipitation to precipitation minus PET ranged from -1.4 to 38% for DCW and -6.3 to 24.3% for CCW. For DCW there was a jump in the percentage of HUC8s showing a significant relationship between six and nine months, but the percentage of HUC8s stabilized after this time period out to 36 months. CCW showed a similar but smaller jump in the percentage of HUC8s with a significant relationship between six and nine months (Table 6). At nine months, all images, regardless of being collected in the spring, summer or fall, would include winter precipitation. We observed substantial spatial variability in the statistical relationship between surface-water extent and water availability. Using nine months as the accumulation

period, we observed a strong spatial pattern in DCW. PPR HUC8s tended to show a steep slope, exhibited by a substantial increase in surface-water extent with increased water availability, while HUC8s across the NP tended to show a weaker slope, exhibited by minor to no increases in surface-water extent with increased water availability (Figure 6 and 7). For CCW, the spatial pattern was less consistent within the PPR or ecoregion boundaries. Instead, HUC8s with a steep slope tended to be HUC8s with large lakes or floodplains (Figure 6 and 7).

3.3 Landscape Variables Explaining Variability in Surface-Water

For DCW SWCV, when independent variables were assessed individually using Spearman's rank correlation, the SWCV was greater in locations with fewer streams ($R = -0.64$, $p < 0.05$), lower slope gradient ($R = -0.59$, $p < 0.05$), higher wetland density ($R = 0.52$, $p < 0.05$) and total wetland area ($R = 0.51$, $p < 0.05$), deeper minimum depth to water table ($R = 0.59$, $p < 0.05$) and where a greater proportion of the total surface water was disconnected from the stream network ($R = 0.42$, $p < 0.05$) (Table 8). When the relative importance of the variables was tested using random forest, variables found to be the most important included, wetland density, stream density, annual minimum depth to water table and the slope gradient (Table 8). However, after accounting for the spatial autocorrelation in the DCW SWCV and the significance of the variables, the DCW SWCV increased in the final feasible generalized least-squares model (adjusted $R^2 = 0.66$, F-statistic = 73.6) with (1) greater wetland density, (2) deeper depth to groundwater, and (3) less anthropogenic drainage (Table 9). The variable most consistent identified across statistical approaches was wetland density.

For CCW SWCV, fewer independent variables showed a significant Spearman rank correlation. The SWCV for stream-connected water increased in locations with a greater total wetland area ($R = 0.48$, $p < 0.05$) and less average precipitation ($R = -0.33$, $p < 0.05$) (Table 8). Using random forest, the total wetland area and proportion of total water from large features were found to be the most important variables in explaining variation. The final feasible generalized least-squares model (adjusted $R^2 = 0.54$, F-statistic = 37.4) also found the relationship between CCW and surface-water availability (i.e., SWCV) was stronger with greater total wetland area, but also found that it decreased with greater wetland density (Table 9).

4. Discussion

Surface-water extent, and in particular surface water within well-studied portions of the PPR, has been previously shown to exhibit seasonal and interannual patterns (Poff et al., 1997; Beeri and Phillips, 2007; Vanderhoof et al., 2016) that can, in turn, influence the cumulative hydrologic response of a watershed (Golden et al. 2016; Evenson et al. 2016; Ali and Creed 2017). What has been less commonly quantified is how surface-water dynamics vary across diverse landscapes. This is particularly relevant when we consider the need for communities and local agencies to plan ahead for expected changes in the precipitation regime associated with climate change (Dore, 2005; Johnson et al., 2005; Millett et al., 2009). Our results demonstrated that the relationship between surface-water extent and water availability (SWCV) is a function of both climate and landscape variables

and that the density of depressional wetlands, in particular, played a key explanatory role in the observed landscape response to increased climate inputs. Given our findings, we expect that changes in net precipitation from climate change or other climatic forcings will disproportionately affect surface-water extent across the PPR relative to the adjacent NP, and that these effects will be most evident in disconnected wetland systems (DCWs). Surface waters that are disconnected from the stream network showed a larger change in extent in response to wetter conditions in landscapes with higher wetland densities. That is to say that landscapes with a larger number of depressional features were found to show a greater increase in surface-water extent in response to a wetter climate, relative to landscapes with fewer depressional features. In landscapes with greater total wetland area, on the other hand, surface waters connected to the stream network showed more substantial expansion with increased water availability. This finding suggests that the presence of stream-connected lakes within large flat basins may be an important factor influencing surface-water expansion. Previous research found lakes within the PPR to be important features that commonly experience extensive surface-water expansion, subsuming adjacent wetlands during wet periods (Vanderhoof and Alexander, 2016). These findings suggest that if climate conditions within the U.S. portion of the PPR continue to get wetter, as predicted (e.g., Millett et al. 2009), then both small wetland depressions and larger features, such as lakes and floodplains, will both serve critical roles in storing increased inputs of surface water, which could prevent downstream flooding.

Our study area was intentionally selected to encompass a large area with a wide range of landscape conditions in regards to wetland and stream density and capacity for infiltration. Across the study area, variation in the values of many of the variables (e.g., stream density, wetland density) can be attributed to landscape age or the time since the last glacial retreat, and corresponding variability in drainage development across the region (Ahnert, 1996). The Wisconsin glacier retreated from the PPR by 11,300 BP, meaning the drainage system is still developing and surface water is being stored in glacially formed depressions (Winter and Rosenberry, 1998; Stokes et al., 2007). In contrast, west and south of the PPR, the landscape is much older (>20,000 BP) with a well-developed drainage network (Clayton and Moran, 1982).

In addition to extensive human-induced wetland loss across the region (Miller et al., 2009; Van Meter et al., 2015), the drainage network across the region is also increasingly modified with the expansion of ditch networks and tile drainage in association with agricultural activities (McCauley et al., 2015). Ditches, pipes and field tiles on the glacial till can hasten the speed with which water leaves a location and lower the water table through increased water withdrawal (De Laney, 1995; Blann et al., 2009; McCauley et al., 2015). We found in the FGLS model, the expansion of disconnected water was inversely related to the abundance of estimated anthropogenic drainage. Because anthropogenic drainage increases the rate at which water leaves a location, it results in the loss or reduction of landscape-scale functions of wetlands and other natural water storage features in the PPR (McCauley et al. 2015), and shifts the hydrologic behaviors of watersheds towards those more commonly seen in the NP.

Evapotranspiration is known to be a primary mechanism for water loss in the PPR (Winter and Rosenberry, 1998). By explicitly incorporating this value into the SWCV, we could better isolate the effects of landscape-based components such as surface storage, stream density, and topography. One challenging component to characterize was the capacity for water to infiltrate through soil horizons. Depth to bedrock SSURGO data was found to be too patchy (i.e., too much missing data) to be useful. A variable that instead was found to correlate significantly with the expansion of disconnected water was annual minimum depth to groundwater. The PPR tended to have a deeper minimum depth while the NP had a shallower minimum depth, on average. A reduction in infiltration due to the low permeability of glacial till (Sloan, 1972; Winter and Rosenberry, 1995), would reduce the potential for increased water table elevations. Concomitantly, with less infiltration, pulses of snowmelt or precipitation in the PPR would instead be transported as overland flow and fill wetlands with available storage.

We must also consider that we may be missing key landscape variables that could explain variability in the spatial response of surface-water extent to climate inputs. For example, major landscape characteristics required for stream-connected surface water to expand include (1) large, stream-connected water bodies such as lakes and (2) hydrologically-connected floodplains. The influence of large water bodies was considered by including total wetland area and the portion of water from larger (>8 ha) features, however we did not explicitly consider the presence/absence of active floodplains beyond including stream density as a variable. Floodplain activity typically exhibits strong seasonal patterns; however, the goal of our analysis was to focus on patterns of surface-water extent that occurred on longer-time scales (i.e., interannual variability). Because of this, we excluded two Landsat path/rows from the analysis that were originally included because strong seasonal flooding outweighed interannual patterns in climate as evidenced by a lack of a relationship between climate indices (e.g., Standardized Precipitation Index (12 months) and Palmer Hydrologic Drought Index) and surface-water extent. These path/rows included p30r27 which straddles North Dakota and Minnesota and exhibits strong seasonal flooding of the Red River and p28r32 in the southeastern corner of Nebraska, which exhibits strong seasonal flooding of the Missouri River. However, even with the exclusion of these two path/rows, the importance of floodplains is still evident in Figure 6B where we observed higher slopes in areas with an abundance of lakes or floodplain systems. Because complete floodplain maps across the study area are lacking, we were not able to explicitly identify the role of floodplains in the CCW models.

In addition to decision points regarding study area extent, other decision points may have influenced our findings. For example, the period of time for which the greatest number of HUC8s showed a significant slope was used as the climate accumulation period. This logic was meant to avoid, to the extent possible, a HUC8 showing a zero slope because surface water responded at a time period different than the one selected. However, its usage meant that the study results are limited to interpreting the relationship of surface-water extent to same year climate inputs (or the previous 9 months) and may be less applicable to understanding the relationship of surface-water extent to shorter (seasonal) or longer (multi-year) time periods. In addition, decisions regarding image inclusion may have also influenced the analysis. Although the Landsat images used in the analysis were selected

strategically to represent historically dry, average, and wet conditions, because the Landsat images were processed individually we were ultimately limited in the number of Landsat images we could process. As more remotely sensed products become available, such as the U.S. Geological Survey's Dynamic Surface Water Extent (DSWE) Product, which plans to utilize the entire Landsat archive (1984 to present) (Jones, 2015), we could utilize many more images and reduce the uncertainty in estimates of the slope value or watershed-specific response to available water. Although decision points regarding the data included or excluded from the analysis are important to consider, this study provides an improved understanding of how the relationship between surface-water extent and climate may vary spatially across different landscapes.

5. Conclusion

Shifts in climate patterns and the frequency of extreme climate events will influence surface-water extent. This has implications for habitat availability (Boschilia et al., 2008; Calhoun et al., 2017), agricultural productivity (Mokrech et al., 2008; Gornall et al., 2010) and hydrologic connectivity (Golden et al. 2016; Ali and Creed 2017). This study demonstrated that not only is surface-water extent variable across landscapes, but shifts in climate patterns will have an uneven effect on surface-water extent across these different landscapes. The PPR experienced a 2.6 fold greater surface-water extent than the adjacent NP under average conditions and a 3.4 fold larger range in surface-water extent between drought and deluge conditions. To move from ecoregion boundaries to a clearer characterization of the spatial distribution of surface water on the landscape, we used a statistical approach to explore potentially significant landscape variables that could explain the spatially variable change in surface water to climate inputs (precipitation minus evapotranspiration). Landscapes with higher wetland density and less anthropogenic drainage showed a greater expansion of disconnected (from the stream network) surface water (e.g., depressional wetlands) and wetter climatic conditions relative to landscapes with fewer wetlands and more anthropogenic drainage. Landscapes with fewer wetlands but more total surface water area (e.g., lakes, large river systems) showed a greater expansion of stream-connected surface water and wetter climatic conditions relative to landscapes with less total surface water area. Enhancing our knowledge of spatial and temporal variability in the relationship between surface-water extent and climate inputs can advance efforts to predict the hydrological effects of climate change, including drought and floods, on water resources and improve hydrological modeling in low gradient landscapes.

Acknowledgements

This research was funded by the Drought Resilience Initiative through an interagency agreement with the U.S. Environmental Protection Agency, Office of Research and Development (DW-014-92454401 – 0). We thank Tedros Berhane, Hayley Distler and Marena Gilbert for their assistance in processing the Landsat imagery. We thank Maliha Nash for providing data on climate averages. The views expressed in this manuscript are solely those of the authors and do not necessarily reflect the views or policies of the U.S. EPA. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Appendix

Table 1.

A complete list of Landsat TM images used in the analysis and the corresponding Palmer Hydrological Drought Index (PHDI).

Landsat path/row	Date	PHDI	Landsat path/row	Date	PHDI	Landsat path/row	Date	PHDI
p26r30	1989 170	-4.29	p30r30	1990 121	-4.70	p31r29	1989 109	-1.62
p26r30	1989 186	-4.29	p30r30	1989 294	-4.66	p31r29	2003 196	-1.22
p26r30	1988 296	-4.15	p30r30	1989 110	-3.47	p31r29	2004 279	2.52
p26r30	1996 222	-0.24	p30r30	1991 236	-2.79	p31r29	1999 121	5.19
p26r30	1987 117	0.06	p30r30	1988 148	-1.23	p31r29	2011 154	6.55
p26r30	1996 142	0.30	p30r30	2002 122	-1.12	p31r29	2010 167	6.94
p26r30	2010 148	1.10	p30r30	2013 184	-0.94	p31r29	2010 279	8.63
p26r30	2006 153	1.17	p30r30	2003 141	0.26	p33r28	1988 249	-5.68
p26r30	2008 95	2.82	p30r30	2003 285	0.88	p33r28	1990 254	-3.87
p26r30	1993 133	3.95	p30r30	1993 161	5.40	p33r28	2008 112	-2.86
p26r30	1993 277	6.92	p30r30	2011 211	6.49	p33r28	1988 137	-2.47
p26r32	1988 264	-4.18	p30r30	2011 179	6.87	p33r28	2005 135	-2.35
p26r32	2000 105	-3.03	p30r30	2010 288	8.93	p33r28	2003 146	-1.78
p26r32	2003 145	-2.98	p30r31	2002 250	-4.62	p33r28	2005 263	-0.62
p26r32	1989 266	-2.92	p30r31	2000 269	-3.75	p33r28	1998 148	0.22
p26r32	1991 288	-1.88	p30r31	2000 173	-2.66	p33r28	2006 106	0.36
p26r32	1991 96	0.55	p30r31	1990 105	-2.63	p33r28	1998 260	0.70
p26r32	2007 108	0.74	p30r31	2003 141	-2.46	p33r28	1995 188	4.09
p26r32	2002 158	1.59	p30r31	1990 297	-2.45	p33r28	1997 129	5.11
p26r32	1994 136	2.76	p30r31	1990 137	-2.43	p33r28	2015 67	5.37
p26r32	1993 133	3.66	p30r31	2003 221	-2.41	p33r28	2014 160	5.61
p26r32	1994 104	3.79	p30r31	2000 221	-2.38	p33r28	2014 256	9.15
p26r32	2010 100	4.06	p30r31	2000 125	-2.05	p36r28	1988 222	-6.07
p26r32	2008 271	5.07	p30r31	2002 122	-1.84	p36r28	2002 212	-5.14
p26r32	2010 228	5.90	p30r31	2005 178	1.58	p36r28	2004 154	-4.72
p27r30	1988 239	-4.52	p30r31	1986 174	2.19	p36r28	2004 282	-4.29
p27r30	1989 161	-4.34	p30r31	1994 148	3.63	p36r28	2003 135	-2.38
p27r30	2003 280	-1.32	p30r31	1994 260	4.12	p36r28	1985 149	-2.04
p27r30	2002 141	-1.25	p30r31	2011 179	5.22	p36r28	1989 112	-1.94
p27r30	2003 104	1.44	p30r31	2009 173	5.29	p36r28	2013 178	-0.91
p27r30	2008 182	3.03	p31r27	1990 160	-4.12	p36r28	1993 91	-0.89
p27r30	1992 266	3.22	p31r27	2006 252	-3.32	p36r28	2013 242	-0.42
p27r30	1992 122	4.29	p31r27	1991 163	-2.45	p36r28	1998 121	1.67
p27r30	1993 172	6.52	p31r27	1992 118	-1.93	p36r28	2008 181	1.70
p29r29	1990 130	-3.55	p31r27	1999 121	2.01	p36r28	1996 244	2.06
p29r29	2003 118	-2.01	p31r27	2007 255	2.41	p36r28	1996 100	3.81
p29r29	2002 323	-1.69	p31r27	1997 195	2.72	p36r28	1993 235	5.17

Landsat path/row	Date	PHDI	Landsat path/row	Date	PHDI	Landsat path/row	Date	PHDI
p29r29	1991 133	-0.69	p31r27	2005 169	3.06	p36r28	1994 142	
p29r29	1992 136	1.35	p31r27	2009 244	3.28	p37r26	1988 213	-5.70
p29r29	2006 286	2.30	p31r27	2004 279	4.38	p37r26	2006 246	-3.41
p29r29	1998 120	2.77	p31r27	2001 190	4.46	p37r26	1994 261	-2.54
p29r29	2005 91	3.15	p31r27	1995 270	5.97	p37r26	2008 108	-2.37
p29r29	2006 94	4.20	p31r27	2010 279	6.43	p37r26	2002 171	-1.85
p29r29	2001 128	4.47	p31r27	2011 186	6.61	p37r26	1991 141	0.14
p29r29	1997 165	5.05	p31r27	1994 299	7.03	p37r26	2009 142	0.26
p29r29	1995 288	5.71	p31r27	2011 266	8.92	p37r26	1995 168	1.35
p29r29	2011 284	5.88	p31r29	2006 172	-3.49	p37r26	1995 264	1.68
p29r29	2010 105	6.19	p31r29	1989 189	-3.38	p37r26	1987 162	2.15
p29r29	1993 266	6.86	p31r29	2004 135	-2.66	p37r26	1991 269	2.26
p29r29	2011 156	8.37	p31r29	1989 269	-2.31	p37r26	1994 101	2.76
p29r29	2010 281	9.63	p31r29	2003 100	-2.24	p37r26	2013 169	3.40
			p31r29	2003 132	-1.84	p37r26	2011 276	7.32
			p31r29	1990 96	-1.65	p37r26	2011 212	9.14

6. References

- Abatzoglou JT: Development of gridded surface meteorological data for ecological applications and modelling, *Int. J. Climatol*, 33, 121–131, 2011.
- Acharya G: Approaches to valuing the hidden hydrological services of wetland ecosystems, *Ecol. Econ*, 35, 63–74, 2000.
- Ahnert F: Introduction to Geomorphology, John Wiley & Sons, New York, 1996.
- Allen CR, Angeler DG, Cumming GS, Carl F, and Twidwell D: Quantifying spatial resilience, *J. Appl. Ecol*, 53, 625–635, 2016.
- Ameli AA, Creed IF: Quantifying hydrologic connectivity of wetlands to surface water systems, *Hyrol. Earth Syst. Sci*, 21, 1791–1808, 2017.
- Beeri O and Phillips RL: Tracking palustrine water seasonal and annual variability in agricultural wetland landscapes using Landsat from 1997 to 2005, *Glob. Change Biol*, 13, 897–912, 2007.
- Belsley DA: Conditioning Diagnostics, Collinearity and Weak Data in Regression, John Wiley & Sons, New York, 1991.
- Belsley DA, Kuh E, and Welsch RE: Regression Diagnostics: Identifying Influential Data and Sources of Collinearity, John Wiley & Sons, New York, 1980.
- Betts MG, Ganio LM, Huso MMP, Som NA, Huettmann F, Bowman J, and Wintle BA: Comment on “Methods to account for spatial autocorrelation in the analysis of species distributional data: A review.”, *Ecography*, 32, 374–378, 2009.
- Blann KL, Anderson JL, Sands GR, and Vondracek B: Effects of agricultural drainage on aquatic ecosystems: A review, *Crit. Rev. Environ. Sci. Technol*, 39, 909–1001, 2009.
- Boschilia SM, Oliveira EF, and Thomaz SM: Do aquatic macrophytes co-occur randomly? An analysis of null models in a tropical floodplain, *Oecologia*, 156, 203–214, 2008. [PubMed: 18274779]
- Breusch TS and Pagan AR: A simple test for heteroskedasticity and random coefficient variation, *Econometrica*, 47, 1287–1294, 1979.
- Bryson RA and Hare FK: Climates of North America, in: *World Survey of Climatology*, Vol. 11, Lansberg HE, (Ed.), Elsevier, New York, 47 pp., 1974.

- Calhoun AJK, Mushet DM, Bell KP, Boix D, Fitsimons JA, and Isselin-Nondedeu F: Temporary wetlands: challenges and solutions to conserving a “disappearing” ecosystem, *Biol. Conserv*, 211, 3–11, 2017.
- Carroll RW, Pohl GM, Tracy J, Winter T, and Smith R: Simulation of a semipermanent wetland basin in the Cottonwood Lake Area, East-Central North Dakota, *J. Hydrol. Eng*, 1, 70–84, 2005.
- Christensen JR, Nash MS, and Neale A: Identifying riparian buffer effects on stream nitrogen in southeastern coastal plain watersheds, *Environ. Manage.* 52, 1161–1176, 2013. [PubMed: 23989318]
- Clayton L and Moran SR: Chronology of late Wisconsinan glaciation in middle North America, *Quaternary Sci. Rev*, 1, 55–82, 1982.
- Cohen MJ, Creed IF, Alexander L, Basu N, Calhoun A, Craft CB, D’Amico E, DeKeyser S, Fowler L, Golden H, Jawitz JW, Kalla P, Kirkman LK, Lane CR, Lang M, Leibowitz S, Lewis DB, Marton JM, McLaughlin DL, Mushet D, Raanan-Kipperwas H, Rains MC, Smith L, and Walls S: Do geographically isolated wetlands influence landscape functions?, *P. Natl. Acad. Sci. U.S.A.*, 113, 1978–1986, 2016.
- Cowardin LM, Carter V, Golet FC, and LaRoe ET: Classification of wetlands and deepwater habitats of the United States, U.S. Fish and Wildlife Service, Washington, DC, FWS/OBS-79/31, 1979.
- Daly C, Halbleib M, Smith JI, Gibson WP, Doggett MK, Taylor GH, Curtis J, and Pasteris PA: Physiographically-sensitive mapping of temperature and precipitation across the conterminous United States, *Int. J. Climatol*, 28, 2031–2064, 2008.
- De Laney TA: Benefits to downstream flood attenuation and water quality as a result of constructed wetlands in agricultural landscapes, *J. Soil Water Conserv*, 50, 620–626, 1995.
- Dore MHI: Climate change and changes in global precipitation patterns: What do we know?, *Environ. Int*, 31, 1167–1181, 2005. [PubMed: 15922449]
- Dormann CF, McPherson JM, Araujo MB, Bivand R, Bolliger J, Carl G, Davies RG, Hirzel A, Jetz W, Kissling WD, Kühn I, Ohlemüller R, Peres-Neto PR, Reineking B, Schröder B, Schurr FM, and Wilson R: Methods to account for spatial autocorrelation in the analysis of species distributional data: A review, *Ecography*, 30, 609–628, 2007.
- Dunn OJ: Multiple comparison among means, *J. Am. Stat. Assoc*, 56, 52–64, 1961.
- Eamus D, Hatton T, Cook P, and Colvin C: *Ecohydrology: vegetation function, water and resource management*, CSIRO Publishing, Australia, 360 pp., 2006.
- Euliss NH Jr. and Mushet DM: Water-level fluctuation in wetlands as a function of landscape condition in the Prairie Pothole Region, *Wetlands*, 16, 587–59, 1996.
- Evenson GR, Golden HE, Lane CR, D’Amico E: An improved representation of geographically isolated wetlands in a watershed-scale hydrologic model, *Hydrol Processes*, 2016.
- Fausey NR, Doering EJ, and Palmer ML: Purposes and benefits of drainage, in: *Farm drainage in the United States: History, status, and prospects*, Pavelis GA (Ed.), USDA Economic Research Service, Washington, DC, Misc. Publ. 1455, 4 pp., 1987.
- Feddema JJ: A revised Thornthwaite-type global climate classification, *Phys. Geogr*, 26, 442–466, 2005.
- Fleiss JL: *Statistical methods for rates and proportions* (2nd ed.), John Wiley & Sons, New York City, NY, 1981.
- Flint RF: *Glacial and Quaternary Geology*, John Wiley & Sons, New York City, NY, 1971.
- Forbes AD: Classification-algorithm evaluation: Five performance measures based on confusion matrices. *J. Clinical Monitoring*, 11(3), 189–206, 1995.
- Gleason RA and Tangen BA: Ecosystem services derived from wetland conservation practices in the United States prairie pothole region with an emphasis on the U.S., in: *Department of Agriculture Conservation Reserve and Wetlands Reserve Programs Professional Paper 1745: Floodwater storage*, Gleason RA, Laubhan MK, Euliss MK Jr. (Eds.), U.S. Geological Survey, Reston, VA, 7 pp., 2008.
- Golden HE, Sander HA, Lane CR, Zhao C, Price K, D’Amico E, Christensen JR: Relative effects of geographically isolated wetlands on streamflow: A watershed-scale analysis, *Ecohydrology*, 9(1), 21–38, 2016.

- Golden HE, Creed IF, Ali G, Basu NB, Neff BP, Rains MC, McLaughlin DL, Alexander LC, Ameli AA, Christensen JR, Evenson GR, Jones CN, Lane CR, and Lang M: Integrating geographically isolated wetlands into land management decisions, *Front. Ecol. Environ*, 15, 319–327, 2017. [PubMed: 30505246]
- Gornall J, Betts R, Burke E, Clark R, Camp J, Willett K, and Wiltshire A: Implications of climate change for agricultural productivity in the early twenty-first century, *Philos. Trans. R. Soc. B. Biol. Sci*, 365, 2973–2989, 2010.
- Green AA, Berman M, Switzer P, and Craig MD: A transformation for ordering multispectral data in terms of image quality with implications for noise removal, *IEEE T. Geosci. Remote*, 26, 65–74, 1988.
- Hamilton SK, Sippel SJ, and Melack JM: Comparison of inundation patterns among major South American floodplains, *J. Geophys. Res*, 107, 1–14, 2002.
- Hamilton SK, Sippel SJ, and Melack JM: Seasonal inundation patterns in two large savanna floodplains of South America: the Llanos de Moxos (Bolivia) and the Llanos del Orinoco (Venezuela and Colombia), *Hydrol. Process*, 18, 2103–2116, 2004.
- Hamon WR: Estimating potential evapotranspiration, *J. Hydr. Eng. Div. ASCE*, 87, 107–120, 1961.
- Hanushek EA: Efficient estimators for regressing regression coefficients, *Am. Stat*, 28, 66–67, 1974.
- Heiler G, Hein T, Schiemer F, and Bornette G: Hydrological connectivity and flood pulses as the central aspects for the integrity of a river-floodplain system, *Regul. River*, 11, 351–361, 1995.
- Heine RA, Lant CL, Sengupta RR: Development and comparison of approaches for automated mapping of stream channel networks. *Annals of the Assoc of Am Geographers*, 94(3), 477–490, 2004.
- Hendrickx J: Perturb: Tools for evaluating collinearity, R package version 2.05, <http://CRAN.R-project.org/package=perturb>, 2012.
- Hoffmann CC, Kjaergaard C, Uusi-Kämpä J, Hansen HC, and Kronvang B: Phosphorus retention in riparian buffers: Review of their efficiency, *J. Environ. Qual*, 38, 1942–1955, 2009. [PubMed: 19704138]
- Homer C, Dewitx J, Yang L, Jin S, Danielson P, Xian G, Coulston J, Herold N, Wickham J, and Megown K: Completion of the 2011 National Land Cover Database for the conterminous United States – representing a decade of land cover change information, *Photogramm. Eng. Rem. S*, 81, 345–354, 2015.
- Huang S, Dahal D, Young C, Chander G, and Liu S: Integration of Palmer Drought Severity Index and remote sensing data to simulate wetland water surface from 1910 to 2009 in Cottonwood Lake area, North Dakota, *Remote Sens. Environ*, 115, 3377–3389, 2011.
- Johnson WC, Millett BV, Gilmanov T, Voldseth RA, Guntenspergen GR, and Naugle DE: Vulnerability of northern prairie wetlands to climate change, *Bioscience*, 55, 863–872, 2005.
- Jones JW: Efficient wetland surface water detection and monitoring via Landsat: comparison with in situ data from the Everglades depth estimation network, *Remote Sens*, 7, 12503–12538, 2015.
- Kahara SN, Mockler RM, Higgins KF, Chipps SR, and Johnson RR: Spatiotemporal patterns of wetland occurrence in the Prairie Pothole Region of eastern South Dakota, *Wetlands*, 29, 678–689, 2009.
- Kantrud HA, Krapu GL, and Swanson GA: Prairie basin wetlands of the Dakotas: a community profile, U.S. Fish and Wildlife Service Biological Report 85(7.28), Washington, DC, 1989.
- King KW, Fausey NR, and Williams MR: Effect of subsurface drainage on streamflow in an agricultural headwater watershed, *J. Hydrol*, 519, 438–445, 2014.
- Kuppel S, Houspanossian J, Noretto MD, and Jobbágy EG: What does it take to flood the Pampas? Lessons from a decade of strong hydrological fluctuations, *Water Resour. Res*, 51, 2937–2950, 2015.
- LaBaugh JW, Winter TC, and Rosenberry DO: Hydrologic functions of prairie wetlands, *Great Plains Res*, 4, 17–37, 1998.
- Lewis JB and Linzer DA: Estimating regression models in which the dependent variable is based on estimates, *Polit. Anal*, 13, 345–364, 2005.

- Liaw A and Wiener M: Breiman and Cutler's random forests for classification and regression, R package version 4.6–12, R Foundation for Statistical Computing, Vienna, Austria, <https://www.stat.berkeley.edu/~breiman/RandomForests/>, 2005.
- Masek JG, Vermote EF, Saleous N, Wolfe R, Hall EF, Huemmrich F, Gao F, Kutler J, and Teng-Kui L: A Landsat surface reflectance data set for North America, 1990–2000, *IEEE Geosci. Remote S*, 3, 68–72, 2006.
- McCauley LA, Anteau MJ, Van Der Burg MP, and Wiltermuth MT: Land use and wetland drainage affect water levels and dynamics of remaining wetlands, *Ecosphere*, 6, 1–20, 2015.
- Millett B, Johnson WC, and Guntenspergen G: Climate trends of the North American Prairie Pothole Region 1906–2000, *Climatic Change*, 93, 243–267, 2009.
- Mitchell KE, Lohmann D, Houser PR, Wood EF, Schaake JC, Robock A, Cosgrove BA, Sheffield J, Duan Q, Luo L, Higgins RW, Pinker RT, Tarpley JD, Lettenmaier DP, Marshall CH, Entin JK, Pan M, Shi W, Koren V, Meng J, Ramsay BH, and Bailey AA: The multi-institution North American Land Data Assimilation System (NLDAS): Utilizing multiple GCIP products and partners in a continental distributed hydrological modeling system, *J. Geophys. Res.*, 109, D07S90, 2004.
- Mokrech M, Nicholls RJ, Richards JA, Henriques C, Holman IP, and Shackley S: Regional impact assessment of flooding under future climate and socio-economic scenarios for East Anglia and North West England, *Climatic Change*, 90, 31–55, 2008.
- Murphy MA, Evans JS, and Storfer A: Quantifying *Bufo boreas* connectivity in Yellowstone National Park with landscape genetics, *Ecology*, 91, 252–261, 2010. [PubMed: 20380214]
- Niemuth ND, Wangler B, and Reynolds RE: Spatial and temporal variation in wet area of wetlands in the prairie pothole region of North Dakota and South Dakota, *Wetlands*, 30, 1053–1064, 2010.
- NOAA National Climatic Data Center: Data Tools: 1981–2010 Normals, <http://www.ncdc.noaa.gov/cdo-web/datatools/normals>, last access: 12, January, 2017, 2014.
- Poff NL, Allan JD, Bain MB, Karr JR, Prestegard KL, Richter BD, Sparks RE, and Stromberg JC: The natural flow regime, *BioScience*, 47, 769–784, 1997.
- Pringle CM: Hydrologic connectivity and the management of biological reserves: a global perspective, *Ecol. Appl.*, 11, 981–998, 2001.
- Rosenberry DO, Stannard DI, Winter TC, and Martinez ML: Comparison of 13 equations for determining evapotranspiration from a prairie wetland, Cottonwood Lake area, North Dakota, USA, *Wetlands*, 24, 483–497, 2004.
- Sass GZ and Creed IF: Characterizing hydrodynamics on boreal landscapes using archived synthetic aperture radar imagery, *Hydrol. Process*, 22, 1687–1699, 2008.
- Saylor KL, Acevedo W, Soulard CE, and Taylor JL: Land cover trends dataset, 2000–2011, U.S. Geological Survey, 2015.
- Schmidt F, Persson A: Comparison of DEM data capture and topographic wetness indices, *Precision Agriculture*, 4(2), 179–192, 2003.
- Shaw DA, Vanderkamp G, Conly FM, Pietroniro A, and Martz L: The fill-spill hydrology of prairie wetland complexes during drought and deluge, *Hydrol. Process*, 26, 3147–3156, 2012.
- Skaggs RW, Breve MA, and Gilliam JW: Hydrologic and water quality impacts of agricultural drainage, *Crit. Rev. Environ. Sci. Technol.*, 24, 1–32, 1994.
- Sloan CE: Ground-water hydrology of Prairie Potholes in North Dakota, U.S. Geological Survey Professional Paper, 585, 1–27, 1972.
- Smith LC, Sheng Y, and MacDonald GM: A first pan-arctic assessment of the influence of glaciation, permafrost, topography and peatlands on northern hemisphere lake distribution, *Permafrost Periglac.*, 18, 201–208, 2007.
- Stokes CR, Popovnin V, Aleynikov A, Gurney SD, and Shahgedanova M: Recent glacier retreat in the Caucasus Mountains, Russia, and associated increase in supraglacial debris cover and supra-/proglacial lake development, *Ann. Glaciol.*, 46, 195–203, 2007.
- Turin G: An introduction to matched filters, *IRE T. Inform. Theor.*, 6, 311–329, 1960.
- U.S. Department of Agriculture, National Resources Conservation Service: Watershed Boundary Dataset, http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/water/watersheds/dataset/?cid=nrcs143_021625, last access: 20 September 2016, 2015.

- U.S. Department of Agriculture, Natural Resources Conservation Service: Soil Survey Geographic (SSURGO) Database, <https://sdmdataaccess.sc.egov.usda.gov>, last access: 25 May 2017, 2017.
- U.S. Fish and Wildlife Service: National Wetlands Inventory, <http://www.fws.gov/wetlands/>, last access: 1 September 2016, 2010.
- U.S. Geological Survey: The National Hydrography Dataset (NHD) concepts and content, available at: http://nhd.usgs.gov/chapter1/chp1_data_users_guide.pdf, 2010.
- U.S. Geological Survey: The National Hydrography Dataset (NHD), <ftp://nhdftp.usgs.gov/DataSets/Staged/States/FileGDB/HighResolution>, last access: 2 September 2017, 2013.
- van der Kamp GW, Stolte J, and Clark RG: Drying out of small prairie wetlands after conversion of their catchments from cultivation to permanent brome grass, *Hydrolog. Sci. J.*, 44, 387–397, 1999.
- Van Meter KJ and Basu NB: Signatures of human impact: size distributions and spatial organization of wetlands in the Prairie Pothole landscape, *Ecol. Appl.*, 25, 451–465, 2015. [PubMed: 26263667]
- Vanderhoof MK, Alexander LC, and Todd MJ: Temporal and spatial patterns of wetland extent influence variability of surface water connectivity in the Prairie Pothole Region, United States, *Landscape Ecol.*, 31, 805–824, 2016.
- Vanderhoof MK and Alexander LC: The role of lake expansion in altering the wetland landscape of the Prairie Pothole Region, *Wetlands*, 36, 309–321, 2016. [PubMed: 32669752]
- Vanderhoof MK, Christensen JR, and Alexander LC: Patterns and drivers for wetland connections in the Prairie Pothole Region, United States, *Wetl. Ecol. Manag.*, 25, 275–297, 2017. [PubMed: 32025096]
- Venables WN and Ripley BD: *Modern Applied Statistics with S*, Fourth Edition, Springer, New York, 2002.
- Villalobos-Jimenez G and Hassall C: Effects of the urban heat island on the phenology of Odonata in London, UK, *Int. J. Biometeorol.*, 61, 1337–1346, 2017. [PubMed: 28190181]
- Wagener T, Sivapalan M, Troch P, and Woods R: Catchment classification and hydrologic similarity, *Geogr. Compass*, 1, 901–931, 2007.
- Wiche GG: Lake levels, streamflow, and surface-water quality in the Devil's Lake Area, North Dakota, U.S. Geological Survey Fact Sheet, Bismarck, ND, 8 pp., 1996.
- Willmott CJ and Feddema JJ: A more rational climatic moisture index, *Prof. Geogr.*, 44, 84–88, 1992.
- Winter TC: The concept of hydrologic landscapes, *J. Am. Water Res. Assoc.*, 37, 335–349, 2001.
- Winter TC and Rosenberry DO: The interaction of ground water with Prairie Pothole wetlands in the Cottonwood Lake area, east-central North Dakota, 1979–1990, *Wetlands*, 15, 193–211, 1995.
- Winter TC and Rosenberry DO: Hydrology of prairie pothole wetlands during drought and deluge: 7-year study of the Cottonwood Lake wetland complex in North Dakota in the perspective of longer term measured and proxy hydrological records, *Climatic Change*, 40, 189–209, 1998.
- Yao T, Pu J, Lu A, Wang Y, and Yu W: Recent glacial retreat and its impact on hydrological processes on the Tibetan Plateau, China, and surrounding regions, *Arct. Antarct. Alp. Res.*, 39, 642–650, 2007.
- Zhang B, Schwartz FW, and Liu G: Systematics in the size structure of prairie pothole lakes through drought and deluge, *Water Resour. Res.*, 45, W04421, 2009.
- Zhu Z and Woodcock CE: Automated cloud, cloud shadow, and snow detection in multitemporal Landsat data: An algorithm designed specifically for monitoring land cover change, *Remote Sens. Environ.*, 152, 217–234, 2014.




Figure 1. Distribution of Landsat path/rows used to map surface-water extent and corresponding 8-digit Hydrological Units (HUC8s) used for further analysis in relation to the boundary of the Prairie Pothole Region (PPR). The p37r26 behaved dissimilarly from the PPR and similarly to the adjacent Northern Prairie (NP) in all regards and was therefore included with the NP for analyses organized by PPR and NP.




Figure 2. Theoretical figure showing the derived dependent variable defined as the slope of the statistical relationship between accumulated water and surface-water extent. Some areas show a steep slope or substantial increase in surface-water extent as more water becomes available via precipitation minus potential evapotranspiration (PET), while other areas show little to no change in surface-water extent, presumably as excess water leaves the system through runoff or infiltration.




Figure 3. Standard errors of the slope tended to increase as slopes increased for both A) discontinuous surface water (DCW) or surface water disconnected from the stream network and B) continuously connected water (CCW) or surface water connected to the stream network.




Figure 4. Mean surface-water abundance and the amount of “wetting up” varied substantially between different Landsat path/rows. Portions of the Northern Prairie (e.g., p26r30) showed relatively less surface-water extent and expansion (A and B) while portions of the Prairie Pothole Region (e.g., p29r29) showed relatively more surface-water extent and expansion (C and D). Note: not all water is visible at this zoomed-out scale. PHDI: Palmer Hydrological Drought Index




Figure 5. Examples of minor and substantial expansion of surface-water extent between historically dry and historically wet points in time. PHDI: Palmer Hydrological Drought Index.




Figure 6. The spatial distribution of the Surface Water Climate Variable (SWCV) values from the statistical relationships between available water, defined as precipitation minus potential evapotranspiration accumulated over the previous 9 months, and surface-water extent. Greater SWCV values indicate greater change in surface-water extent with increased available water. Surface-water extent was divided between A) disconnected surface water (DCW), or surface-water extent disconnected from the stream network, and B) continuously connected water (CCW), or surface-water extent connected to the stream network.




Figure 7. Distribution of slope and standard error values organized by Landsat path/row and primary path/row location, i.e., the Northern Prairie or the Prairie Pothole Region (PPR) for A) surface water that is disconnected from the stream network (DCW), and B) surface water that is connected to the stream network (CCW). HUC8: 8-digit Hydrological Units

A summary of the Landsat images utilized within each selected path/row. Landsat TM images were used for dates 2011 and earlier. Landsat 8 OLI images were used for 2013 forward.

Table 1.

Path/Row	PPR/Northern Prairie (NP) (primary)	Number of Images	Spring (DOY 60–151)	Summer (DOY 152–243)	Fall (DOY 244–335)	Year Range	Min. PHDI (%)	Max. PHDI (%)	Mean PHDI (%)
p26r30	NP	12	6	4	2	1987–2010	4	99	45
p26r32	NP	17	10	3	4	1988–2010	2	99	51
p27r30	PPR	9	3	4	2	1988–2008	4	99	54
p29r29	PPR	17	9	2	6	1990–2011	7	100	69
p30r30	PPR	13	5	5	3	1988–2013	2	100	45
p30r31	NP	15	6	5	4	1986–2011	5	94	38
p31r27	PPR	15	2	6	7	1990–2011	3	100	67
p31r29	PPR	13	6	5	2	1989–2011	7	99	45
p33r28	NP	15	8	2	5	1988–2015	1	99	49
p36r28	NP	16	7	7	2	1985–2013	2	96	38
p37r26	NP*	15	4	6	5	1987–2013	1	99	52
Total		157	66	49	42				

DOY: day of year; PHDI: Palmer Hydrological Drought Index.

* p37r26 was considered NP because of its dissimilarity with the rest of the PPR.

Table 2.

Landsat images and corresponding National Agricultural Imaging Program (NAIP) images used to validate the Landsat surface-water extent maps. Accuracy is presented here by Landsat image.

Landsat Path/Row	Landsat date	NAIP date(s)	Gap (days)	PHDI	SP12	Number of points	OE (%)	CE (%)	OA (%)	DC (%)	RB (%)
p26r32	28-Jun-04	23-Jun-04 and 07-Jul-04	-5 to +9 days	0.57	0.14	947	6.3	5.9	97.4	93.9	-0.5
p27r30	14-Jul-13	10-Jul-13 and 12-Jul-13	-4 to -2 days	-0.34	0.05	707	11.8	9.3	92.5	89.5	-2.7
p29r29	13-Oct-06	25-Sep-06	-18 days	2.3	-0.08	814	11.1	2.5	93.6	93.0	-8.8
p29r29	8-Oct-10	17-Sep-10 and 20-Sep-10	+18 to +21 days	9.63	3.06	959	1.9	3.3	97.4	96.4	1.4
p31r29	17-Jul-04	10-Jul-04 and 14-Jul-04	-7 to -3 days	-0.4	-0.04	1302	7.4	1.5	97.2	95.4	-6.0
p33r28	13-Jul-03	11-Jul-03 and 15-Jul-03	-2 to +2 days	-2.74	-0.91	908	10.6	27.0	85.5	80.4	22.5
p37r26	31-Jul-11	16-Jul-11 and 19-Jul-11	-15 to -12 days	2.96	1.29	498	16.8	9.7	90.2	86.6	-7.9

PHDI: Palmer Hydrological Drought Index, SP12: 12 month Standardized Precipitation Index, OE: omission error for water, CE: commission error for water, OA: overall accuracy, DC: Dice coefficient, RB: relative bias

Table 3.

Summary of accuracy statistics across all of the Landsat images validated using National Agricultural Imaging Program (NAIP) imagery.

	NAIP - Inundated	NAIP - Non-Inundated	Total
Landsat - Inundated	2052	183	2235
Landsat - Non-Inundated	190	3710	3900
Total	2242	3893	6135
Omission error for water (%)	8.5		
Commission error for water (%)	8.2		
Overall Accuracy (%)	93.9		
Dice Coefficient	91.7		
Relative Bias	0.0		

Table 4.

Independent variables considered in the landscape analysis and the distribution of values for each variable across the 8-digit hydrological units (HUC8s). Mean values for the HUC8s within the Prairie Pothole Region (PPR) and Northern Prairie (NP) are also shown with significant differences ($p < 0.01$) between the two groups, as determined by the Wilcoxon rank sum test, indicated by different superscript letters.

Independent Variables	Units	Range	25th %	50th %	75th %	PPR (avg)	NP (avg)	Source
Wetland and Stream Characteristics								
Stream density	m ha ⁻¹	0.1 to 26.1	7.2	11.4	15.0	7.8 ^a	14.5 ^b	High-Resolution NHD (USGS 2013)
Total wetland density	no ha ⁻¹	0 to 0.2	0.02	0.03	0.06	0.06 ^a	0.03 ^b	NWI (USFWS 2010)
Total wetland areal abundance	ha ha ⁻¹	0 to 0.7	0.02	0.03	0.08	0.08 ^a	0.05 ^b	NWI (USFWS 2010)
Portion of total water area from large features	%	0.1 to 97.8	32.1	44.7	58.0	45.0 ^a	47.2 ^a	NWI (USFWS 2010, Zhang et al., 2009)
Portion DCW of total surface water	%	0 to 100	10.6	24.4	50.0	44.5 ^a	22.8 ^b	Landsat and NHD (USGS 2013)
Climate Averages								
Moisture Index Average	~	-0.4 to 0.7	-0.1	-0.04	0.2	0.04 ^a	-0.03 ^a	PRISM (Daly et al., 2008)
Precipitation Average	mm yr ⁻¹	312.3 to 1007.8	490.3	599.6	790.8	641.5 ^a	624.3 ^a	PRISM (Daly et al., 2008)
PET Average	mm yr ⁻¹	496.2 to 683.0	564.2	595.5	628.9	595.5 ^a	594.8 ^a	PRISM (Daly et al., 2008)
Soil and Topography								
Available water storage (0–150 cm), weighted	cm	7.6 to 29.5	18.0	22.8	24.7	24.0 ^a	19.1 ^b	SSURGO (Soil Survey Staff, 2017)
Annual minimum depth to water table	cm	0.1 to 69.0	11	24.8	43.3	40.5 ^a	17.9 ^b	SSURGO (Soil Survey Staff, 2017)
K _{sat}	µm sec ⁻¹	2.1 to 107.7	8.4	13.8	22.5	21.4 ^a	21.2 ^a	SSURGO (Soil Survey Staff, 2017)
Slope gradient, weighted average	%	1.5 to 19.2	3.0	4.3	7.1	3.3 ^a	7.1 ^b	SSURGO (Soil Survey Staff, 2017)
Human Influence								
Agricultural land cover	%	0.1 to 92.0	25.2	62.8	80.5	72.6 ^a	39.6 ^b	2011 NLCD (Homer et al., 2015)
Percent drained by anthropogenic means	%	0 to 93.0	5.9	50.1	77.8	61.7 ^a	32.5 ^b	2011 NLCD and SSURGO

NHD: National Hydrography Dataset, NWI: National Wetland Inventory, PRISM: Parameter-elevation Regressions on Independent Slopes Model, SSURGO: Soil Survey Geographic Database, NLCD: National Land Cover Database, DCW: disconnected surface water, PET: potential evapotranspiration, avg: average

Table 5.

Spearman rank correlation values between the independent variables considered in the analysis. Bonferroni correction was applied to the p-values and significant correlations ($p < 0.05$) are starred.

Variable	DCW auto-covariate	CCW auto-covariate	Portion dis-connected	Stream density	Wetland density	Wetland areal abund.	Dominance of lg. water bodies	MI	Precip	PET	Avail water storage (0-150 cm)	Annual min depth to water table	Ksat	Slope gradient	Ag land cover	Percent drained
DCW autocovariate	1	na	0.45*	-0.66*	0.48*	0.48*	-0.04	0.03	-0.05	0.03	0.29	0.54*	0.21	-0.58*	0.33*	0.22
CCW autocovariate		1	-0.11	-0.16	0.15	0.27	0.18	-0.29	-0.26	0.15	0.05	-0.04	0.01	-0.16	-0.03	-0.07
Portion DCW of total water			1	-0.38*	0.32*	-0.09	-0.63*	0.33*	0.20	-0.05	0.37*	0.54*	0.11	-0.34*	0.46*	0.26
Stream density				1	-0.33*	-0.37*	-0.05	-0.34*	-0.21	0.09	-0.34*	-0.62*	-0.47*	0.66*	-0.33*	-0.2
Wetland density					1	0.79*	-0.02	0.24	0.19	-0.1	0.26	0.29	-0.03	-0.19	0.11	0.25
Wetland areal abundance						1	0.44	0.18	0.06	-0.1	0.21	0.26	-0.01	-0.29	0.05	0.23
Dominance of lg water bodies							1	-0.22	-0.16	0	-0.11	-0.1	0.08	0.1	-0.24	-0.01
MI								1	0.86*	-0.2	0.60*	0.67*	0.14	-0.41*	0.80*	0.64*
Precipitation									1	0.25	0.48*	0.44*	0.03	-0.17	0.66*	0.50*
PET										1	-0.07	-0.21	-0.2	0.12	-0.08	-0.16
Avail water storage (0-150 cm)											1	0.49*	-0.05	-0.44*	0.66*	0.51*
Annual min depth to water table												1	0.19	-0.61*	0.69*	0.57*
Ksat													1	-0.25	0.02	-0.07
Slope gradient														1	-0.63*	-0.32*
Agricultural land cover															1	0.63*

EPA Author Manuscript

EPA Author Manuscript

EPA Author Manuscript

DCW: surface water disconnected from the stream network, CCW: continuously connected surface water, MI: Moisture Index, PET: potential evapotranspiration, precip: precipitation, lg: large, ag: agricultural, Ksat: saturated hydraulic conductivity, na: not applicable

Table 6.

The percent of HUC8s across the study area that showed a significant relationship ($p < 0.05$) between surface-water extent and (1) precipitation (Precip or P) or (2) precipitation minus potential evapotranspiration (PET) for different accumulation periods.

Accumulated Period	Precip DCW (%)	P - PET DCW (%)	Inclusion of PET change (DCW)	Precip CCW (%)	P - PET CCW (%)	Inclusion of PET change (CCW)
3 months	19.4	27.1	7.6	15.3	28.5	13.2
6 months	5.6	31.9	26.4	9.0	33.3	24.3
9 months	20.8	59.7	38.9	27.1	48.6	21.5
12 months	45.8	50.7	4.9	42.4	41.0	-1.4
18 months	24.3	58.3	34.0	25.7	39.6	13.9
24 months	52.1	50.7	-1.4	43.8	37.5	-6.3
30 months	28.5	55.6	27.1	27.1	43.1	16.0
36 months	54.9	54.9	0.0	47.2	44.4	-2.8
HUC8s with a sig relationship in at least 1 time period	65.3	75.7	10.4	59.0	67.4	8.3

DCW: disconnected surface water; CCW: continuously, connected surface water.

Table 7. Surface-water extent conditions summarized for the Prairie Pothole Region (PPR) and adjacent Northern Prairie (NP).

Region	Path/ rows (all or part)	Total area (km ²)	Min (ha km ⁻²)	Max (ha km ⁻²)	Median (ha km ⁻²)	Added min to max (ha km ⁻²)	Reduction from median to min (%)	Increase from median to max (%)	Min (% of all) (area)	Max (% of all) (area)	Median (% of all) (area)
PPR TSW	7	146,309	3.51	11.99	6.33	8.48	44.6	89.2	~	~	~
NP TSW	9	173,026	1.62	4.07	2.45	2.45	33.9	66.1	~	~	~
PPR CCW	7	146,309	2.82	7.56	4.44	4.74	36.5	70.4	80.3	63.1	70.1
NP CCW	9	173,026	1.44	3.11	2.06	1.66	30.0	50.5	89.1	76.3	84.2
PPR DCW	7	146,309	0.69	4.42	1.90	3.73	63.4	133.4	19.7	36.9	29.9
NP DCW	9	173,026	0.18	0.97	0.39	0.79	54.4	149.2	10.9	23.7	15.8

TSW: total surface-water extent, CCW: continuously connected surface water that intersects the stream network, DCW: disconnected surface water or surface water that does not directly intersect the stream network.

Table 8.

Spearman rank correlation values between the dependent variables and each of the independent variables considered in the analysis. Bonferroni correction was applied to the p-values and significant correlations ($p < 0.05$) are starred. Relative variable importance as determined by random forest models are also presented for each variable (i.e., increase in node purity).

Variable	Response (DCW, 9 months)		Response (CCW, 9 months)	
	Spearman rank correlation	Increase in node purity	Spearman rank correlation	Increase in node purity
Autocovariate	0.79*	0.081	0.53*	0.108
Portion DCW is of total surface water	0.42*	0.012	-0.11	0.334 [†]
Stream density	-0.64*	0.036 [†]	-0.15	0.060
Wetland density	0.52*	0.048 [†]	0.27	0.057 [†]
Wetland areal abundance	0.51*	0.017 [†]	0.48*	0.855 [†]
Portion of total water from large features	-0.01	0.004	0.30	0.556 [†]
Moisture Index (average)	-0.03	0.005	-0.28	0.053 [†]
Precipitation (average)	-0.10	0.008 [†]	-0.33*	0.039 [†]
PET (average)	-0.06	0.011 [†]	-0.13	0.034
Available water storage (0–150 cm)	0.27	0.007	-0.01	0.061
Annual minimum depth to water table	0.56*	0.027 [†]	0.09	0.046
Ksat	0.04	0.004	-0.08	0.070 [†]
Slope gradient, weighted average	-0.59*	0.025 [†]	-0.22	0.072
Agricultural land cover	0.31	0.005	-0.05	0.035
Percent drained by anthropogenic means	0.22	0.004	-0.04	0.020

PET: potential evapotranspiration, Ksat: saturated hydraulic conductivity, DCW: disconnected surface water, CCW: continuously, connected surface water

[†] Variables selected by the random forest model selection process, using the R package rUtilities, when the autocovariate was not included.

Feasible generalized least square models with residual weights applied relating the response (of surface-water extent to water availability) to landscape-related variables. All variables included in the models were significant. DCW: surface water disconnected from the stream network, CCW: continuously connected surface water, SE: standard error, D.F.: degrees of freedom

Table 9.

Response of DCW water to water availability		Variables	Coefficients	SE	t-value
D.F. = 145		Intercept	0.17	0.01	12.84
F-statistic = 73.6		Autocovariate	0.03	0.004	6.32
adjusted R ² = 0.66		Wetland density	0.90	0.23	3.96
		Minimum depth to groundwater	0.0021	0.0006	3.29
		Percent anthropogenically drained	-0.0004	0.0003	-1.25
Response of CCW water to water availability		Variables	Coefficients	SE	t-value
D.F. = 144		Intercept	0.018	0.01	1.43
F-statistic = 69.4		Wetland areal abundance	0.96	0.07	14.42
adjusted R ² = 0.58		Wetland density	-0.43	0.21	-2.09
		Autocovariate	-0.12	0.01	-0.89

PET: potential evapotranspiration, Ksat: saturated hydraulic conductivity, DCW: disconnected surface water, CCW: continuously, connected surface water