

Brief Report

Butyrate production in the acetogen *Eubacterium limosum* is dependent on the carbon and energy source

Dennis Litty and Volker Müller* 

Department of Molecular Microbiology & Bioenergetics,
Institute of Molecular Biosciences, Goethe-University
Frankfurt am Main, Hessen, Germany.

Summary

***Eubacterium limosum* KIST612 is one of the few acetogenic bacteria that has the genes encoding for butyrate synthesis from acetyl-CoA, and indeed, *E. limosum* KIST612 is known to produce butyrate from CO but not from H₂ + CO₂. Butyrate production from CO was only seen in bioreactors with cell recycling or in batch cultures with addition of acetate. Here, we present detailed study on growth of *E. limosum* KIST612 on different carbon and energy sources with the goal, to find other substrates that lead to butyrate formation. Batch fermentations in serum bottles revealed that acetate was the major product under all conditions investigated. Butyrate formation from the C1 compounds carbon dioxide and hydrogen, carbon monoxide or formate was not observed. However, growth on glucose led to butyrate formation, but only in the stationary growth phase. A maximum of 4.3 mM butyrate was observed, corresponding to a butyrate:glucose ratio of 0.21:1 and a butyrate:acetate ratio of 0.14:1. Interestingly, growth on the C1 substrate methanol also led to butyrate formation in the stationary growth phase with a butyrate:methanol ratio of 0.17:1 and a butyrate:acetate ratio of 0.33:1. Since methanol can be produced chemically from carbon dioxide, this offers the possibility for a combined chemical-biochemical production of butyrate from H₂ + CO₂ using this**

acetogenic biocatalyst. With the advent of genetic methods in acetogens, butanol production from methanol maybe possible as well.

Introduction

Acetogens are a physiological group of strictly anaerobic bacteria that are characterized by a special pathway for CO₂ fixation, the Wood–Ljungdahl pathway (WLP) (Müller, 2003; Drake *et al.*, 2008; Ragsdale, 2008). The WLP is a branched linear pathway in which two mol of CO₂ are reduced to one mol of acetyl-CoA which is further converted to acetate in all species under most conditions (Müller and Frerichs, 2013). Moreover, some species can convert acetyl-CoA (or acetate) to ethanol or even to C₄ compounds such as butyrate (Daniell *et al.*, 2012; Jeong *et al.*, 2015; Bengelsdorf *et al.*, 2016). Therefore, acetogens have come into focus as biocatalysts for a CO₂-based bioeconomy and ethanol is already produced on an industrial scale using *Clostridium autoethanogenum* (Bengelsdorf *et al.*, 2018; Heffernan *et al.*, 2020). The addition of one carbon to the chain length of the product increases the value of the product by a factor of 1.5–3, depending on the product (Kim *et al.*, 2019). Butyrate is not a prime product to be produced but it can be reduced to butanol in a two-step enzymatic process and butanol is a highly desired biofuel (Dürre, 2016). Butyrate is produced naturally by only a few acetogens such as *Clostridium carboxidivorans* (Liou *et al.*, 2005), *Clostridium drakei* (Küsel *et al.*, 2000; Liou *et al.*, 2005), *Oxobacter pfenigii* (Krumholz and Bryant, 1985) and *E. limosum* strains such as KIST612 (Pacaud *et al.*, 1985; Loubiere and Lindley, 1991; Chang *et al.*, 1997). The latter has gained much interest for it produces butyrate from synthesis gas (syngas), a mixture of H₂, CO₂ and CO in different concentrations, depending on the source (Chang *et al.*, 2001; Park *et al.*, 2017). Syngas is an industrial waste stream that is already been used as feedstock for acetogenic conversion to ethanol (Dürre and Eikmanns, 2015; Humphreys and Minton, 2018).

Received 21 January, 2021; revised 9 February, 2021; accepted 9 February, 2021.

Microbial Biotechnology (2021) 14(6), 2686–2692
doi:10.1111/1751-7915.13779

Funding Information

Financial support by the BMBF-Project 'Biometchem' is gratefully acknowledged.

However, butyrate formation in *E. limosum* KIST612 was only observed in bioreactors with cell recycling (Chang *et al.*, 2001), but not in batch cultures or only to minor amounts when acetate was added to the culture (Park *et al.*, 2017). Methanol and formate are two promising, alternative feedstocks for the industrial production of biofuels using acetogens as they can be produced from many sustainable feedstocks including biomass, municipal solid waste, biogas as well as CO₂. One major advantage using these two feedstocks

is that they are fully soluble and therefore can overcome the challenges gaseous C1 feedstocks are facing due to their low mass transfer. In addition, formate and methanol can also be easily transported and stored.

The methyl group of methanol is channelled into the WLP by a methyltransferase system (Kremp *et al.*, 2018; Kremp and Müller, 2020) whereas formate is an intermediate of the pathway (Fig. 1). Unfortunately, it is not known whether these C1 substrates maybe converted to butyrate as well. Here, we have investigated




Fig. 1. The Wood-Ljungdahl pathway of CO₂ reduction. In the WLP, two molecules of CO₂ are reduced to the central intermediate acetyl-CoA. Entry points for other C1 substrates are indicated. Acetyl-CoA is the precursor of biomass and a wide range of natural products (blue). The pathways leading from acetyl-CoA to products are not complete and miss intermediates, reducing equivalents and ATP input/output. Only the pathway leading to butyrate is complete. CoFeSP, corrinoid/iron sulfur protein; THF, tetrahydrofolate; CoA, coenzyme A; [H], reducing equivalent.

the physiology of growth of *E. limosum* KIST612 on different substrates with a focus on the production of butyrate.

Results and discussion

Growth with and product formation from glucose

When *E. limosum* KIST612 was transferred to carbonate buffered basal medium (CBBM) (Chang *et al.*, 1999) containing different amounts of glucose ranging from 20 to 200 mM, growth rates were identical ($\mu = 0.34 \text{ h}^{-1}$) as were the final yields ($\text{OD}_{600} = 4.5$). To describe the growth physiology in more detail, cells were cultured on 20 mM glucose. On transfer of a glucose-adapted preculture to fresh medium, growth started immediately with a rate of 0.34 h^{-1} and proceeded for about 13 h, before the stationary phase started (Fig. 2). Parallel to growth, the glucose concentration dropped continuously with a rate of $1.4 \text{ mmol l}^{-1} \cdot \text{h}^{-1}$ to a residual concentration of 1.2 mM.

In parallel, the pH dropped from 7.2 to 4.5. Glucose consumption was paralleled by a production of acetate that reached a final concentration of 32.1 mM, corresponding to an acetate:glucose ratio of 1.7:1. Interestingly, at the end of the exponential growth phase butyrate was produced and butyrate production reached a steady state at around 17 h. The final butyrate concentration was 4.3 mM, corresponding to a butyrate:glucose ratio of 0.23:1. Even later, at around 13 h, ethanol formation started, but ethanol formation was very small with values in the 0.4–0.7 mM range. In total, the recovery of glucose in all end products was $75 \pm 3.9 \%$, not accounting for CO_2 . Growth on fructose led to similar growth characteristics (Fig. 2).

Growth with and product formation from $\text{H}_2 + \text{CO}_2$, CO or formate

The WLP accepts different C1 substrates with different oxidation/reduction states. Growth of *E. limosum*




Fig. 2. Growth of *E. limosum* on glucose and fructose. *E. limosum* KIST612 was grown at 37°C in 500 ml of carbonate-buffered basal medium (CBBM) (Chang *et al.*, 1999) with glucose under a N_2/CO_2 (80/20% [v/v]) atmosphere. Growth was followed by measuring the optical density (OD) at 600 nm.

A. Growth of *E. limosum* on 20 (●), 40 (○), 80 (▼), 120 (△), 160 (■) or 200 (□) mM glucose.

B. Growth and product portfolio of *E. limosum* on 20 mM glucose. OD_{600} (●) was determined photometrically, pH (◇) was determined with an Orion Basic electrode (Thermo Electron Corp. Witchford, UK). The concentration of glucose (■) was determined by a D-glucose/D-fructose assay kit (R-Biopharm, Pfungstadt, Germany). Acetate (▲), butyrate (▼) and ethanol (◆) were measured by gas chromatography as described previously (Jeong *et al.*, 2015).

C. Growth of *E. limosum* on 20 (●), 40 (○), 80 (▼), 120 (△), 160 (■) or 200 (□) mM fructose. All data points are mean \pm SEM; $N = 3$ independent experiments.




Fig. 3. Growth of *E. limosum* on H₂ + CO₂, CO or formate.

A. *E. limosum* was grown at 37°C in 500 ml of CBBM with overpressure of 1 bar H₂ + CO₂ (80/20% [v/v]). H₂ + CO₂ (■) was determined by gas chromatography as described previously (Bertsch and Müller, 2015). OD₆₀₀ (●), pH (◆) as well as acetate (▲), butyrate (▼) and isobutyrate (●) were determined as described in the legend to Fig. 2.

B. *E. limosum* was grown at 37°C in 500 ml of phosphate buffered basal medium (Chang *et al.*, 1999) with overpressure of 1 bar CO (100%) (■). CO was determined by gas chromatography as described previously (Bertsch and Müller, 2015). OD₆₀₀ (●), pH (◆) as well as acetate (▲), butyrate (▼), isobutyrate (●) and ethanol (▼) were determined as described in the legend to Fig. 2 (Bertsch and Müller, 2015).

C. *E. limosum* KIST612 was grown at 37°C in 500 ml of CBBM with 20 mM Na⁺-Formate (■) under a N₂/CO₂ (80/20% [v/v]) atmosphere. The concentration of formate was determined by a formate assay kit (R-Biopharm, Pfungstadt, Germany). OD₆₀₀ (●), pH (◆) as well as acetate (▲), butyrate (▼) and isobutyrate (●) were determined as described in the legend to Fig. 2. All data points are mean ± SEM; N = 3 independent experiments.

KIST612 on H₂ + CO₂ and CO have been described (Chang *et al.*, 2001); the entry points are shown in Fig. 1. Formate is an intermediate of the WLP but has


not been described as substrate for *E. limosum* KIST612. Growth on H₂ + CO₂ or CO proceeded with growth rates of 0.04 and 0.05 h⁻¹; acetate was the major

product, butyrate was only observed in trace amounts (Fig. 3). Cells also grew on formate with a growth rate similar to $H_2 + CO_2$ or CO (0.03 h^{-1}) but the final yield was much lower ($OD_{600} = 0.25$). Growth was accompanied by acetate production, but due to the consumption of formate, in sum, the pH increased slightly. Butyrate was not produced.

Growth with and product formation from methanol

E. limosum did grow on methanol as sole carbon and energy source, as predicted from its genome sequence (Roh *et al.*, 2011). Whereas *A. woodii* reached the maximum growth rate at 60 mM methanol (Kremp *et al.*, 2018), growth of *E. limosum* was already maximal at 20 mM methanol. When a methanol-adapted culture (two transfers) of *E. limosum* KIST612 was

transferred to CBBM with 20 mM methanol, growth started immediately with a doubling time of 17.61 h and proceed for about 60 h, before the stationary phase started (Fig. 4). Parallel to growth, the methanol concentration dropped continuously with a rate of $0.29\text{ mmol l}^{-1}\cdot\text{h}^{-1}$ to a residual concentration of 3.4 mM. Methanol consumption was accompanied by a production of acetate that reached a final concentration of 12 mM, corresponding to an acetate:methanol ratio of 0.53:1. Since acetogenesis from methanol according to Equation (1):



removes the CO_2 from the solution leading to alkalization parallel to acidification by acid production, the pH did not drop but increased slightly. Most important,




Fig. 4. Growth of *E. limosum* on methanol. *E. limosum* KIST612 was grown at 37°C in 500 ml of CBBM with methanol under a N_2/CO_2 (80/20% [v/v]) atmosphere.

A. Growth of *E. limosum* on 20 (●), 40 (○), 80 (▼), 100 (△), 120 (▲), 160 (■) or 200 (□) mM methanol.

B. *E. limosum* KIST612 was grown at 37°C in 500 ml of CBBM with 20 mM methanol under a N_2/CO_2 (80/20% [v/v]) atmosphere. OD_{600} (●), pH (◆) as well as acetate (▲), butyrate (▼), isobutyrate (●), ethanol (▼) and methanol (■) were determined as described in the legend to Fig. 2. All data points are mean \pm SEM; $N = 3$ independent experiments

again at the end of the exponential growth phase butyrate was produced and butyrate production reached a steady state at around 67 h. The final butyrate concentration was 3.7 mM, corresponding to a butyrate:methanol ratio of 0.17:1. Ethanol was not observed but traces of isobutyrate (0.5 mM). In sum, almost all substrate carbon (methanol + CO₂) was recovered in the major end products acetate and butyrate (99 ± 10.3%).

Conclusion

Here we describe for the first time that the acetogen *E. limosum* KIST612 produces butyrate from methanol. Growth on methanol requires the action of a methyltransferase system that transfers the methyl group from methanol to tetrahydrofolate and *E. limosum* KIST612 has a gene cluster similar to a previously suggested methanol-specific methyltransferase system of *A. woodii* (Kremp and Müller, 2020). Butyrate production from acetyl-CoA follows the pathways described for example for *Clostridium acetobutylicum* (Dürre *et al.*, 2002) involving thiolase, hydroxybutyryl-CoA dehydrogenase, crotonase and butyryl-CoA dehydrogenase with the exception, that the latter is electron-bifurcating and reduces ferredoxin alongside with crotonyl-CoA (Jeong *et al.*, 2015). As in clostridia, butanol could be produced from butyryl-CoA by two subsequent reduction steps (Dürre, 2016). With the establishment of first genetic methods from *E. limosum* KIST612 (Jeong *et al.*, 2020) it should be possible in the future to express butyryl-CoA dehydrogenases in *E. limosum* KIST612. This would make methanol a promising feedstock for acetogenic production of butyrate.

Acknowledgements

Financial support by the European Research Area Cofund on BioTechnologies (ERA CoBioTech; project BIOMETCHEM) is gratefully acknowledged.

Conflict of interest

The authors declare no conflict of interest.

References

Bengelsdorf, F.R., Beck, M.H., Erz, C., Hoffmeister, S., Karl, M.M., Riegler, P., *et al.* (2018) Bacterial anaerobic synthesis gas (syngas) and CO₂+H₂ Fermentation. *Adv Appl Microbiol* **103**: 143–221.

Bengelsdorf, F.R., Poehlein, A., Linder, S., Erz, C., Hummel, T., Hoffmeister, S., *et al.* (2016) Industrial acetogenic biocatalysts: a comparative metabolic and genomic analysis. *Front Microbiol* **7**: 1036.

Bertsch, J., and Müller, V. (2015) CO metabolism in the acetogen *Acetobacterium woodii*. *Appl Environ Microbiol* **81**: 5949–5956.

Chang, I.S., Kim, B.H., Kim, D.H., Lovitt, R.W., and Sung, H.C. (1999) Formulation of defined media for carbon monoxide fermentation by *Eubacterium limosum* KIST612 and the growth characteristics of the bacterium. *J Biosci Bioeng* **88**: 682–685.

Chang, I.S., Kim, B.H., Lovitt, R.W., and Bang, J.S. (2001) Effect of CO partial pressure on cell-recycled continuous CO fermentation by *Eubacterium limosum* KIST612. *Process Biochem* **37**: 411–421.

Chang, I.S., Kim, D.H., Kim, B.H., Shin, P.K., Yoon, J.H., Lee, J.S., and Park, Y.H. (1997) Isolation and identification of carbon monoxide utilizing anaerobe, *Eubacterium limosum* KIST612. *Kor J Appl Microbiol Biotechnol* **25**: 1–8.

Daniell, J., Köpke, M., and Simpson, S.D. (2012) Commercial biomass syngas fermentation. *Energies* **5**: 5372–5417.

Drake, H.L., Gößner, A.S., and Daniel, S.L. (2008) Old acetogens, new light. *Ann NY Acad Sci* **1125**: 100–128.

Dürre, P. (2016) Butanol formation from gaseous substrates. *FEMS Microbiol Lett* **363**: fnw040.

Dürre, P., Böhringer, M., Nakotte, S., Schaffer, S., Thormann, K., and Zickner, B. (2002) Transcriptional regulation of solventogenesis in *Clostridium acetobutylicum*. *J Mol Microbiol Biotechnol* **4**: 295–300.

Dürre, P., and Eikmanns, B.J. (2015) C1-carbon sources for chemical and fuel production by microbial gas fermentation. *Curr Opin Biotechnol* **35**: 63–72.

Heffernan, J.K., Valgepea, K., de Souza Pinto Lemgruber, R., Casini, I., Plan, M., Tappel, R., *et al.* (2020) Enhancing CO₂-valorization using *Clostridium autoethanogenum* for sustainable fuel and chemicals production. *Front Bioeng Biotechnol* **8**: 204.

Humphreys, C.M., and Minton, N.P. (2018) Advances in metabolic engineering in the microbial production of fuels and chemicals from C1 gas. *Curr Opin Biotechnol* **50**: 174–181.

Jeong, J., Bertsch, J., Hess, V., Choi, S., Choi, I.G., Chang, I.S., and Müller, V. (2015) Energy conservation model based on genomic and experimental analyses of a carbon monoxide-utilizing, butyrate-forming acetogen, *Eubacterium limosum* KIST612. *Appl Environ Microbiol* **81**: 4782–4790.

Jeong, J., Kim, J.-Y., Park, B., Choi, I.-G., and Chang, I.S. (2020) Genetic engineering system for syngas-utilizing acetogen, *Eubacterium limosum* KIST612. *Bioresour Technol Rep* **11**: 100452.

Kim, H., Jeon, B.S., and Sang, B.-I. (2019) An efficient new process for the selective production of odd-chain carboxylic acids by simple carbon elongation using *Megasphaera hexanoica*. *Sci Rep* **9**: 11999.

Kremp, F., and Müller, V. (2020) Methanol and methyl group conversion in acetogenic bacteria: Biochemistry, physiology and application. *FEMS Microbiol Rev*, in press. <https://doi.org/10.1093/femsre/fuaa040>

Kremp, F., Poehlein, A., Daniel, R., and Müller, V. (2018) Methanol metabolism in the acetogenic bacterium *Acetobacterium woodii*. *Environ Microbiol* **20**: 4369–4384.

- Krumholz, L.R., and Bryant, M.P. (1985) *Clostridium pfenigii* sp. nov. uses methoxyl groups of monobenzenoids and produces butyrate. *Int J Syst Bacteriol* **35**: 454–456.
- Küsel, K., Dorsch, T., Acker, G., Stackebrandt, E., and Drake, H.L. (2000) *Clostridium scatologenes* strain SL1 isolated as an acetogenic bacterium from acidic sediments. *Int J Syst Evol Microbiol* **50**: 537–546.
- Liou, J.S., Balkwill, D.L., Drake, G.R., and Tanner, R.S. (2005) *Clostridium carboxidivorans* sp. nov., a solvent-producing *Clostridium* isolated from an agricultural settling lagoon, and reclassification of the acetogen *Clostridium scatologenes* strain SL1 as *Clostridium drakei* sp. nov. *Int J Syst Evol Microbiol* **55**: 2085–2091.
- Loubiere, P., and Lindley, N.D. (1991) The use of acetate as an additional co-substrate improves methylotrophic growth of the acetogenic anaerobe *Eubacterium limosum* when CO₂ fixation is rate-limiting. *J Gen Microbiol* **137**: 2247–2251.
- Müller, V. (2003) Energy conservation in acetogenic bacteria. *Appl Environ Microbiol* **69**: 6345–6353.
- Müller, V., and Frerichs, J. (2013) Acetogenic bacteria. In: *Encyclopedia of Life Sciences*. Chichester: John Wiley & Sons Ltd. <https://onlinelibrary.wiley.com/doi/10.1002/9780470015902.a0020086.pub2>.
- Pacaud, S., Loubiere, P., and Goma, G. (1985) Methanol metabolism by *Eubacterium limosum* B2: Effects of pH and carbon dioxide on growth and organic acid production. *Curr Microbiol* **12**: 245–250.
- Park, S., Yasin, M., Jeong, J., Cha, M., Kang, H., Jang, N., et al. (2017) Acetate-assisted increase of butyrate production by *Eubacterium limosum* KIST612 during carbon monoxide fermentation. *Bioresour Technol* **245**: 560–566.
- Ragsdale, S.W. (2008) Enzymology of the Wood-Ljungdahl pathway of acetogenesis. *Ann NY Acad Sci* **1125**: 129–136.
- Roh, H., Ko, H.J., Kim, D., Choi, D.G., Park, S., Kim, S., et al. (2011) Complete genome sequence of a carbon monoxide-utilizing acetogen, *Eubacterium limosum* KIST612. *J Bacteriol* **193**: 307–308.