

ARTICLE

Open Access

Transcriptional regulation of flavonoid biosynthesis in *Artemisia annua* by AaYABBY5

Sadaf-Ilyas Kayani¹, Qian Shen¹, Saeed-ur Rahman¹, Xueqing Fu¹, Yongpeng Li¹, Chen Wang¹, Danial Hassani¹ and Kexuan Tang¹✉

Abstract

Artemisia annua is a medicinal plant rich in terpenes and flavonoids with useful biological activities such as antioxidant, anticancer, and antimalarial activities. The transcriptional regulation of flavonoid biosynthesis in *A. annua* has not been well-studied. In this study, we identified a YABBY family transcription factor, AaYABBY5, as a positive regulator of anthocyanin and total flavonoid contents in *A. annua*. AaYABBY5 was selected based on its similar expression pattern to the phenylalanine ammonia lyase (*PAL*), chalcone synthase (*CHS*), chalcone isomerase (*CHI*), and flavonol synthase (*FLS*) genes. A transient dual-luciferase assay in *Nicotiana bethamiana* with the AaYABBY5 effector showed a significant increase in the activity of the downstream *LUC* gene, with reporters *AaPAL*, *AaCHS*, *AaCHI*, and *AaUFGT*. The yeast one-hybrid system further confirmed the direct activation of these promoters by AaYABBY5. Gene expression analysis of stably transformed *AaYABBY5* overexpression, *AaYABBY5* antisense, and control plants revealed a significant increase in the expression of *AaPAL*, *AaCHS*, *AaCHI*, *AaFLS*, *AaFSLI*, *AaLDOX*, and *AaUFGT* in *AaYABBY5* overexpression plants. Moreover, their total flavonoid content and anthocyanin content were also found to increase. *AaYABBY5* antisense plants showed a significant decrease in the expression of flavonoid biosynthetic genes, as well as a decrease in anthocyanin and total flavonoid contents. In addition, phenotypic analysis revealed deep purple-pigmented stems, an increase in the leaf lamina size, and higher trichome densities in *AaYABBY5* overexpression plants. Together, these data proved that AaYABBY5 is a positive regulator of flavonoid biosynthesis in *A. annua*. Our study provides candidate transcription factors for the improvement of flavonoid concentrations in *A. annua* and can be further extended to elucidate its mechanism of regulating trichome development.

Introduction

Artemisia annua is a renowned plant rich in terpenes and flavonoids. Flavonoids are polyphenolic plant secondary metabolites. Anthocyanins, flavonols, flavanols, and proanthocyanidins (PAs) or condensed tannins are the major classes of flavonoids. These compounds show differential expression patterns according to plant growth or developmental stages and in a species-specific manner¹. Flavonoids perform various functions in plants, including antioxidant activity, protection against UV light,

defensive responses against plant pathogens, activation of nodulation genes in legumes, fertility, and auxin transport². Flavonoids are also beneficial for human health in many aspects as a nutritional source, as well as for the treatment of various diseases.

Increasing evidence recommends using the dried leaves of *A. annua*, which contain both artemisinin and flavonoids, to cure malaria more effectively than using only artemisinin³. A recent study showed that the combination of kaempferol and artesunate exerts potent antimalarial activity synergistically in mice infected with malarial plasmodium⁴. In a recent study, the mechanism for increased bioavailability of artemisinin when using dried leaves that supply artemisinin and flavonoids synergistically was shown⁵. A comparative study indicated the

Correspondence: Kexuan Tang (kxtang@sjtu.edu.cn)

¹Joint International Research Laboratory of Metabolic and Developmental Sciences, Plant Biotechnology Research Center, Fudan-SJTU-Nottingham Plant Biotechnology R&D Center, School of Agriculture and Biology, Shanghai Jiao Tong University, 200240 Shanghai, China

© The Author(s) 2021




Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

parallel biosynthesis of flavonoids and artemisinin at three growth stages, except for a few flavonoids that showed altered behavior. Furthermore, flavonoids of *A. annua* also exhibit efficacious antioxidant effects⁶, strong anti-inflammatory properties⁷, and potent anticancer characteristics⁸. To cure cancer, various flavonoids of *A. annua* have been reported to synergize with many anticancer drugs. The reported flavonoids that aid anticancer drugs include apigenin, eupatin, luteolin, silyben, kaempferol, and quercetin^{9–14}. In a recent study, potent anti-inflammatory activities of the flavonoids of *A. annua* were reported⁷.

A. annua can produce ~40 different flavonoids, which may vary depending upon the cultivar or Line¹⁵. Despite the massive abundance of flavonoid varieties in *A. annua* and their significant therapeutic potential, the flavonoid biosynthetic pathway and its transcriptional regulation have not been well-elucidated. A few studies have been reported on the regulation of flavonoid biosynthesis in *A. annua*^{16,17}. In other plant species, including *A. thaliana*, the flavonoid biosynthetic pathway has been thoroughly studied¹⁸ (Fig. 1). The upstream shikimate pathway provides the precursor amino acid phenylalanine, which is catalyzed by phenylalanine ammonia lyase (PAL), to

generate cinnamate¹⁹. Cinnamate is then transformed into 4-coumaroyl-CoA in the presence of cinnamate 4-hydroxylase (C4H) and 4-coumarate-CoA ligase (4CL)²⁰. The first committed step toward flavonoid biosynthesis involves 3x malonyl-CoA and 1x4-coumaroyl-CoA to generate chalcones (the basic skeleton of all flavonoids) in a condensation reaction catalyzed by chalcone synthase (CHS)^{21,22}. In the next step, chalcone is isomerized to the flavanone naringenin in the presence of chalcone isomerase (CHI)²³. In the following step, which generates dihydroflavonones, naringenin is oxidized by flavanone-3-hydroxylase (F3H) to yield dihydrokaempferol, which is subsequently hydroxylated by flavonoid 3'-hydroxylase (F3'H) or flavonoid 3',5'-hydroxylase (F3'5'H), producing dihydroquercetin or dihydromyricetin, respectively²⁴.

F3'H or F3'5'H may also directly perform the hydroxylation of naringenin to yield flavonones called eriodictyol and pentahydroxy-flavone, which upon hydroxylation generate dihydroquercetin and dihydromyricetin, respectively. Next, three dihydroflavonols are converted to anthocyanidians in a two-step reaction catalyzed by dihydroflavonol reductase (DFR) and leucoanthocyanidin dioxygenase (LDOX), successively. In



the first step, DFR is involved, which converts dihydroquercetin to leucocyanidin, dihydrokaempferol to leucopelargonidin, and dihydromyricetin to leucodelphinidin. The second step is involved in the LDOX-mediated catalytic oxidation of leucocyanidin to cyanidin, leucopelargonidin to pelargonidin, and leucodelphinidin to delphinidin. In the final step, cyanidin, pelargonidin, and delphinidin are glycosylated by the enzyme UDP-glucose flavonoid 3-*O*-glucosyl transferase (UFGT) to produce colorful and stable compounds called anthocyanins²⁵. Anthocyanin: cyanidin-3-glucoside may be further converted to peonidin-3-glucoside, and delphinidin-3-glucoside is converted to petunidin or malvidin-3-glucoside after being methylated by methyltransferases (MTs). Molecules called tannins or PAs are generated by the enzymes leucoanthocyanidin reductase (LAR) and anthocyanidin reductase (ANR), which catalyze the reduction of leucocyanidin to catechin or cyanidin to epicatechin, respectively.

Recently, several genes from the *A. annua* flavonoid biosynthetic pathway, including chalcone isomerase (*AaCHI*) phenylalanine ammonia lyase (*AaPAL1*), flavanone-3-hydroxylase (*AaF3'H*), and *AaFLS1*, have been cloned and characterized^{16,17,26,27}. A recent study showed that the overexpression of *AaCHI* in *A. annua* provides not only increased levels of flavonoids but also artemisinin²⁷. However, the transcriptional regulation of flavonoid biosynthesis in *A. annua* has not been well-elucidated. The overexpression of *MYC2* in *A. annua* resulted in an increased level of both anthocyanins and artemisinin²⁸. Most of the R2R3-MYB genes have been shown to play regulatory roles in flavonoid biosynthesis in *A. annua*. Recently, the MYB family transcription factor *AaTAR2* from *A. annua* was reported to synergistically regulate both artemisinin and flavonoids²⁹; however, the mechanism of the regulation of flavonoid biosynthesis was not found. In plants, flavonoid biosynthesis is tightly regulated by MBW complexes that interact with late biosynthetic genes. In *Arabidopsis thaliana*, this ternary complex comprises specific R2R3-MYB (PAP1, PAP2, MYB113, or MYB114) and bHLH transcription factors (GLABRA 3, GL3 or Enhancer of GLABRA 3, ECL3) that interact with WDR proteins (TTG1; TRANSPARENT TESTA GLABRA 1)^{30–34}. However, these complexes have not yet been identified in *A. annua*. The identification of transcription factors that regulate flavonoid biosynthesis (by direct means or via indirect interactions), the discovery of MBW complexes and their transcriptional control are important to obtain a better understanding of flavonoid biosynthesis in *A. annua*.

The YABBY gene family is a small family of transcription factors that are specifically found in only seed plants. Due to their small group and their role in primary and secondary plant metabolism, they are of great interest to

researchers. Over the last decade, several studies have been conducted on YABBY proteins; however, the function of YABBY family transcription factors in flavonoid biosynthesis in *A. annua* was not previously characterized. The characteristic features of the YABBY family are the N-terminal C₂C₂ zinc finger-like domain and C-terminal HMG domain^{35,36}. The zinc finger domain mediates protein–protein interactions, whereas the YAB domain mediates DNA binding. YABBY protein complexes generate homo or heterodimers between the YABBY proteins and form complexes with other proteins³⁷. Angiosperms such as *Arabidopsis* contain six YABBY genes in their genomes^{35,36}, which are classified into six groups, FIL + YAB3, YAB2, YAB5, CRC, and INO, and are found to be transcriptional regulators^{38,39}.

The primary function of YABBYs in plants is the regulation of lamina outgrowth and leaf development⁴⁰. YABBY proteins interact with LEUNIG or its homologs, and YAB-LUC complexes regulate adaxial cell polarity in leaves and the initiation of the apical meristem in shoots⁴¹. It was first thought that these transcription factors were limited to the regulation of primary metabolic processes in plants, and massive data on YABBY-mediated leaf development in various seed plants are available. However, during the last decade, a few studies have reported the role of YABBY transcription factors in secondary metabolism, such as the regulation of anthocyanins and glucosinolate through *AtFIL* in *A. thaliana* and the control of monoterpene biosynthesis by *MsYABBY5* in *Mentha spicata*^{42–44}. YABBY transcription factors also work in response to various stresses and plant defense responses; for example, *Arabidopsis YABBY10* is involved in drought and salt stress responses⁴⁵, and *AtFIL* knockout lines provide good resistance to bacterial infections⁴².

In our previous study, we cloned and characterized a YABBY gene; *AaYABBY5* is a positive regulator of artemisinin biosynthesis that directly binds to the promoters of *AaCYP71AV1*, and *AaDBR2*, activates their gene expression and results in a significant increase in the concentration of artemisinin. In this study, we elaborated on the role of *AaYABBY5* in the biosynthesis of flavonoids, including anthocyanins, and their regulation. *AaYABBY5* was selected for this study based on its similar expression pattern to that of the *PAL*, *CHI*, and *FLS* genes. *YABBY5* was found to be a direct activator of promoters of the *PAL*, *CHI*, *CHS*, and *UFGT* genes in the transient *Nicotiana benthamiana* infiltration system and yeast transformation system, respectively. In addition, comparative analysis of transgenic *A. annua* plants that overexpressed the *AaYABBY5* ORF, overexpressed the antisense RNA of *AaYABBY5*, plants transformed with empty pHB vector and untransformed/wild-type plants revealed a significant increase in the transcript levels of *AaPAL*, *AaCHI*, *AaCHS*, *AaFLS*, *AaFSII*, *AaLDOX*, and

AaUFGT, as well as enhanced flavonoid and anthocyanin production in *AaYABBY5* overexpression plants but a significant decrease in gene expression, flavonoids, and anthocyanins in *AaYABBY5* antisense plants compared to the control plants.

To our knowledge, in *A. annua*, the mechanism of the transcription factor-mediated direct activation of flavonoid biosynthesis genes has not been studied before. Previous data do not explain how flavonoids are regulated at the molecular level. Our study provided evidence of direct regulation of the *PAL* gene, which is present upstream of the flavonoid biosynthetic pathway; *AaCHS* and *AaCHI*, which regulate the first committed step of flavonoid biosynthesis and the next preceding step, respectively; and *UFGT*, which is involved in the conversion of precursor anthocyanidin molecules to anthocyanins. The present study broadens the knowledge on the direct regulation of flavonoids by *AaYABBY5* in *A. annua*. To our knowledge, we provide the first molecular mechanism of flavonoid regulation in *A. annua*.

Results

Flavonoid biosynthetic genes contain YABBY-binding motifs

The cloned promoter sequences of flavonoid biosynthetic genes used in the present study, *AaPAL*, *AaCHS*, *AaCHI*, *AaFLS*, *AaFSII*, and genes regulating anthocyanins *AaDFR*, *AaLDOX*, and *AaUFGT*, were analyzed for putative YABBY-binding sequences³⁷ using PlantPAN 3.0 (PlantPAN; <http://PlantPAN.itps.ncku.edu.tw>). Recent work, including ChIP and RNA-seq studies on YABBY-binding motifs present in soya bean and protein-binding microarrays in *A. thaliana*, has shown that these sites vary greatly among different species. YABBY-binding motifs are represented by AT-rich sites in Arabidopsis with consensus binding sequences defined as AATNATAA and AATNATTA. The homologous YABBY-binding motifs found in the promoter sequences are shown in Fig. 2a, with the positions marked by numbers. Except for *DFR*, YABBY-binding motifs were identified in all promoters.




Fig. 2 Bioinformatic analysis of the promoters of flavonoid biosynthetic genes, as well as the expression profile of YABBY family genes and flavonoid-regulating genes in *A. annua*. **a** Putative YABBY-binding sites predicted by PLANTPAN3.0 are shown. Positions on plus and minus strands are represented by numbers above and below the promoter sequence, respectively. YABBY-binding sites were found in all promoters except the *DFR* promoter sequence, for which no predicted YABBY-binding site was predicted. **b** Heatmap showing the expression profile of YABBY family genes, as well as flavonoid biosynthetic genes in six tissues. The color scale at the top represents the RPKM (reads per kilobase per million mapped reads) values. *AaYABBY5* (marked with a black dot) was selected as a potential transcription factor that might regulate flavonoid biosynthesis because of its similar expression pattern to the important flavonoid biosynthetic genes

Global expression profile of YABBY family genes and selection of AaYABBY5 as a potential transcription factor regulating flavonoid biosynthesis

The transcriptome data of six different tissues of *A. annua* were previously generated by our lab⁴⁶. Plant secondary metabolites are usually synthesized in a species- or tissue-specific manner, and secondary metabolites, including flavonoids, are synthesized in trichomes⁴⁷. To identify potential YABBY family genes that might regulate flavonoid biosynthesis, a heatmap was constructed to compare the expression of YABBY genes and flavonoid biosynthetic genes across tissues; trichomes, buds, stems, roots, leaves, and seeds (Fig. 2b).

We found two YABBY genes clustered with flavonoid pathway genes, showing higher expression in trichomes and buds. Among the YABBY family genes, AaYABBY5 (marked with a black dot) was found to be a candidate transcription factor that showed a transcription profile parallel to that of *PAL*, *CHI*, *DFR*, and *FLS* and showed higher expression in trichome, bud, stem, and leaf tissues. In our previous findings, we found that AaYABBY5 regulates *DBR2* and *CYP71AV1* that are involved in the artemisinin biosynthetic pathway⁴⁸. Here, we found that *AaYABBY5*, *DBR2*, and *CYP71AV1* showed similar expression patterns, i.e., higher expression in trichomes and buds, with a progressive decline in their expression in the stem tissues.

As this study focused on regulating flavonoid regulation, we hypothesized that AaYABBY5 might regulate *PAL*, *CHS*, *CHI*, and *FLS*, which showed similar expression patterns in trichomes and/or buds. Interestingly, *PAL* and *DBR2* (the YABBY5 target gene) showed similar expression in trichomes, buds, stems, roots, and leaves. *DFR* and *FSII* expression was found more in buds than in trichomes. *F3'H* expression was very different from the other genes present in the flavonoid pathway. *LDOX* and *UFGT* are enzymes that are involved in anthocyanin biosynthesis. In *A. annua*, anthocyanins are limited to stem tissues²⁸, and the expression of *LDOX* and *UFGT* was found to be higher in stem tissues than in other tissues. In a previous study, real-time PCR analysis of different tissues revealed that *AaYABBY5* transcripts are also found in the stem tissues of *A. annua*⁴⁸; therefore, we speculated that it might also regulate the *LDOX* and/or *UFGT* genes. Overall, it was speculated that YABBY5 might regulate *PAL*, *CHI*, *CHS*, and *FLS*, which are involved in early flavonoid biosynthesis, and *LDOX* and *UFGT*, which are present in the late flavonoid (anthocyanin) pathway. Further experiments were carried out to test this hypothesis.

AaYABBY5 significantly activates the promoters of AaPAL, AaCHI, AaCHS, and AaUFGT in transiently transformed *N. benthamiana*


Knowing putative YABBY-binding sites in the *AaPAL*, *AaCHI*, *AaCHS*, *AaDFR*, *AaFLS*, *AaFSII*, *AaLDOX*, and *AaUFGT* promoter sequences, a dual-luciferase assay was

performed, where *AaYABBY5* inserted in pEarleyGate 104-YFP was used as an effector and promoter sequences inserted into pGreenII 0800-LUC were used as reporters (Fig. 3a). Equal-sized infiltrated leaf discs for each combination of the reporter with the effector AaYABBY5 were analyzed by commercially available dual-LUC reagents (Promega, USA). Values greater than a twofold increase were taken into consideration, and lower values were negated. A significant increase in relative LUC/REN values was found for the *AaPAL*, *AaCHS*, *AaCHI*, and *AaUFGT* promoters. AaYABBY5 exhibited a 7.4-fold increase in the activity of the *PAL* promoter, a 3.2-fold increase in the activity of the *CHS* promoter, a 3.4-fold increase in the activity of the *CHI* promoter and a three fold increase in the activity of the *UFGT* promoter (Fig. 3b–d, i). The fold change was calculated from comparative values of each effector/reporter, and an empty vector was used as a negative control/reporter. The increase in the LUC/REN values corresponds to the intensity of LUC signals driven by the respective promoters in the presence of the AaYABBY5 protein. Based on these results, it was hypothesized that AaYABBY5 could activate these promoter sequences in *A. annua* either directly, by binding to putative YABBY-binding motifs, or indirectly, through some protein–protein interactions.

AaYABBY5 directly binds to promoter regions of AaPAL, AaCHS, AaCHI, and AaUFGT in the EGY48 yeast strain

Transactivation assays using *N. benthamiana* revealed that AaYABBY5 mediated a significant increase in the activities of *PAL*, *CHS*, *CHI*, and *UFGT* promoters in vivo. Therefore, to determine the molecular basis of this regulation and whether AaYABBY5 directly activates them, a Y1H assay was performed. The experiment demonstrated the binding of the pB42AD-AaYABBY5 fusion protein (blue color appearance), but not pB42AD alone (no color), to the *PAL*, *CHS*, *CHI*, and *UFGT* promoter sequences, indicated by the activation of the *lacZ* reporter gene, which produces β -galactosidase and cleaves the X-gal present in growth medium to a compound with a blue-colored phenotype. The experiment was repeated three times to validate the results. No colored phenotype was found for the *DFR*, *FLS*, *FSII*, and *LDOX* promoters (Fig. 4b).

The results of the Y1H assay were consistent with the findings of the dual-luciferase reporter assay. In other plants, including *A. thaliana*, flavonoid genes, and their transcriptional regulation are well-studied. The *A. thaliana* *FIL* gene has been previously reported to be a positive regulator of anthocyanins by activating the *MYB75* gene⁴². To our knowledge, for the first time, we found the molecular basis of the transcriptional regulation of flavonoid biosynthetic pathway genes, *PAL*, *CHS*, *CHI*, and *UFGT* in *A. annua*. These results indicated that AaYABBY5 might have the potential to regulate both flavonoid and anthocyanin biosynthesis in *A. annua*.




Therefore, these contents were measured and compared in *AaYABBY5* OE and *AaYABBY5* AnT. *A. annua* plants.

AaYABBY5*-overexpressing *A. annua* plants showed a consistent significant increase in the expression of genes from *AaPAL* to *AaUFGT

After analyzing the binding of *AaYABBY5* to the promoters of early (flavanone) and late (anthocyanin)

flavonoid biosynthetic genes, it was important to study its functions in *A. annua*. As expected, the gene expression analysis revealed an increase in the expression of all genes under study. The comparative expression of these genes indicated that *AaCHS* showed higher expression than all other genes under study. Similarly, the expression of *DFR*, *FSII*, and *UFGT* was lower than that of *PAL*, *CHS*, *CHI*, *FLS*, and *LDOX* (Fig. 4c-j).



Although *DFR*, *FLS*, *FSII*, and *LDOX* were not activated by AaYABBY5, a higher expression of these genes in AaYABBY5-OE plants was found. It is proposed that the increased flux provided by *PAL*, *CHS*, and *CHI* activates downstream pathway enzymes by increasing the concentration of substrates for the enzymes acting downstream: *DFR*, *FLS*, *FSII*, and *LDOX*. AaYABBY5 overexpression not only increased the expression of its direct target genes but also the flavonoid pathway under study.

AaYABBY5 antisense *A. annua* plants showed a significant decrease in the expression of flavonoid biosynthetic genes

The AaYABBY5 protein activates flavonoid biosynthetic genes, and the increased expression of AaYABBY5 led to a dramatic increase in the expression of flavonoid-regulating genes. Transcript levels of *PAL*, *CHI*, *CHS*, *DFR*, *FLS*, *FSII*, *LDOX*, and *UGFT* were analyzed in AaYABBY5 antisense RNA-containing plants. As expected, a significant decrease in the expression of the genes under study was found in AaYABBY5 AnT. plants

(Fig. 4c–j). Overall, from these findings, a clear understanding of *AaYABBY5*-regulated flavonoid biosynthesis was obtained. To validate the above findings, flavonoid and anthocyanin concentrations were measured and compared among *AaYABBY5* OE, *AaYABBY5* AnT, wild-type/control plants, and vector-containing plants.

***AaYABBY5* positively regulates flavonoid biosynthesis**

Flavonoids are polyphenolic plant secondary metabolites that are classified into different types. In this study, we found that *AaYABBY5* positively regulates the *PAL*, *CHI*, *CHS*, and *UGT* genes. The results of real-time PCR also verified the increased expression of these genes in *AaYABBY5* overexpression plants. These findings revealed a positive behavior of *AaYABBY5* toward flavonoid biosynthesis in *A. annua*. To justify this, flavonoid

contents from *AaYABBY5* overexpression plants, *AaYABBY5* antisense plants, and control plants were measured using the aluminum chloride (AlCl₃) colorimetric method with quercetin as a standard. As expected, the results revealed an increased concentration of flavonoids in *AaYABBY5* overexpression plants compared to *AaYABBY5* antisense or control plants (Fig. 5a), proving the function of *AaYABBY5* as a positive regulator of flavonoid biosynthesis.

***AaYABBY5* promotes anthocyanin biosynthesis, displaying a purple phenotype in *A. annua* stems**

Anthocyanins are secondary metabolites widely present in plant species and are responsible for the purple, bluish, and pinkish pigmentation of different plant parts. In this study, we observed deep purple pigmentation in the stems




Fig. 5 Comparative analysis of flavonoid and anthocyanin contents in *AaYABBY5*-OE plants, *AaYABBY5* AnT. plants, and control plants, as well as phenotypic analysis of transgenic plants. a Comparative analysis of total flavonoid content in *AaYABBY5*-OE plants, *AaYABBY5* AnT. plants, vector-containing (35S), and wild-type plants, showing a significant increase in the total flavonoid content of *AaYABBY5* OE plants compared to *AaYABBY5* AnT. plants and control ones. Similarly, AnT. plants showed a decrease in the concentration of flavonoids. **b–d** Phenotypic analysis of transgenic plants showing deep purple pigmentation in the stems of *AaYABBY5*-OE plants, whereas no purple-color phenotype was found in wild-type and/or antisense *AaYABBY5* plants. Scale bars = 8 cm in (b) and (c) and 4 cm in (d). **e** Comparative analysis of anthocyanin concentrations in *AaYABBY5*-OE plants, *AaYABBY5* AnT. plants, and vector-containing (35S) and wild-type plants, showing a significant increase in the total anthocyanin content of *AaYABBY5*-OE plants compared to *AaYABBY5* AnT. plants and control ones. In contrast, *AaYABBY5* AnT. plants showed a decrease in the concentration of anthocyanins. **f** Coloric representation of anthocyanin enrichment in the anthocyanin extract of *AaYABBY5*-OE plants showing the red color of the anthocyanin extract, an indication of an increased amount of anthocyanins. Scale bars = 4 cm. **g, h** Phenotypic analysis showing broader lamina in *AaYABBY5*-OE plants (ii) compared to *AaYABBY5* AnT. plants (i). Scale bars = 3.5 cm and 10 cm in g and h, respectively. The graph shows the mean values ± SD of three experimental replicates. Error bars show standard deviation for sample, n = 3. Statistical significance was determined using Student’s t test. **P < 0.01, *P < 0.05

of *AaYABBY5* overexpression plants, whereas *AaYABBY5* antisense plants and control plants showed no purple phenotype (Fig. 5b, c).

To test whether *AaYABBY5* is involved in the regulation of anthocyanin biosynthesis in *A. annua*, we measured and compared the total anthocyanin content of stem extracts of control plants and pHB/wild-type, *AaYABBY5*-overexpressing, and *AaYABBY5* antisense plants. It is known that anthocyanins produce a red color when treated with acids. Consistent with this, the anthocyanin extracts from purple-colored *AaYABBY5*-OE plants were red-colored in acidic medium (Fig. 5f). As expected, a significant increase in the concentration of anthocyanins was found in the stems of plants overexpressing *AaYABBY5* compared to the control and *AaYABBY5* antisense plants (Fig. 5e). Previous investigations have demonstrated that the YABBY family TF *AtFIL* is a positive regulator of anthocyanin biosynthesis in the model plant *A. thaliana* through activating the *MYB75* promoter; however, no direct link to the genes regulating anthocyanins or flavonoids was reported in *A. annua*⁴². *MYC2* has been reported to be responsible for an increase in the anthocyanin content, giving a purple phenotype to *AaMYC2*-overexpressing stems²⁸; however, the mechanism underlying this regulation was not studied. To our knowledge, we provide the first molecular basis of flavonoids, including anthocyanin regulation by the YABBY family transcription factor *AaYABBY5*. It was supposed that *AaYABBY5* activates anthocyanin biosynthesis through the direct activation of *UFGT* and upstream pathway genes.

***AaYABBY5* overexpression results in broader leaf lamina and increased trichome density**

The primary function of YABBY family transcription factors found in seed plants is the control of leaf development, increasing the size of the leaf lamina, and maintaining organ polarity⁵⁰; therefore, it was important to determine whether *AaYABBY5* overexpression and/or its downregulation in *A. annua* affected leaf morphology. Phenotypic analysis of *AaYABBY5*-OE plants, *AaYABBY5* AnT. plants, and control plants showed that the leaves of *AaYABBY5*-OE plants have broader leaf lamina, whereas in *AaYABBY5* AnT. plants, the leaf lamina was reduced. Leaves of *AaYABBY5* AnT. plants were radialized compared to that of OE plants and/control plants (Fig. 5g, h). Trichomes are the sites of secondary metabolite synthesis, and studies have reported that flavonoids are synthesized in trichomes⁴⁷.

Trichome development is a part of leaf development. Knowing that *AaYABBY5* regulates leaf lamina, the trichome densities on the leaf surfaces were compared among transgenic plants. The trichome densities on leaf surfaces from *AaYABBY5* overexpression plants, *AaYABBY5* antisense plants, and control plants with an

empty vector were calculated. A significant increase in the trichome densities of *AaYABBY5* overexpression plants was found compared to control plants, whereas for *AaYABBY5* antisense plants, trichome density was found to significantly decrease. These results show that *AaYABBY5* is also a positive regulator of trichome development (Fig. 6a, b).

AaHD1 (homeodomain protein 1) is involved in the initiation of both glandular and nonglandular trichomes in *A. annua*⁴⁹; however, no protein interactions between *AaYABBY5* and *AaHD1* were found. It is proposed that *AaYABBY5* might activate the promoter of *AaHD1*, which regulates trichome development. This research opened paths for future research where the molecular mechanism of the regulation of trichome development by *AaYABBY5* can be found.

Discussion

Artemisia annua is a renowned traditional Chinese medicinal plant that has been used in China for a long time as a treatment of fever, inflammation, and malaria, and its use now extends to Europe and North America⁵⁰. *A. annua* is rich in hydroxylated flavonoids and polymethoxylated flavonoids with useful biological activities, such as antioxidant, anticancer, and antimalarial activities⁵¹. The most interesting feature of *A. annua* flavonoids found recently is to synergize antimalarial and anticancer compounds, most importantly artemisinin^{3,50}. Despite the worthwhile effects of flavonoids, research on the molecular and transcriptional regulation of flavonoid biosynthesis in *A. annua* is limited. Recently, chalcone isomerase (*AaCHI*) phenylalanine ammonia lyase (*AaPAL1*), flavanone-3-hydroxylase (*AaF3'H*), and *AaFLS1* have been cloned and characterized as enzymes that take part in flavonoid biosynthesis^{16,17,26,27}. A few transcription factors from the bHLH and MYB families have been recently reported as regulators of flavonoids in *A. annua*. Examples include the overexpression of *MYC2* (which binds to G boxes present in the *AaCYP71AV1* and *AaDBR2* promoters) in *A. annua*, resulting in an increased level of both anthocyanins and artemisinin²⁸. Mostly, R2R3-MYB has been shown to play regulatory roles in flavonoid biosynthesis in *A. annua*. Recently, the MYB family transcription factor *AaTAR2* from *A. annua* was reported to synergistically regulate both artemisinin and flavonoids²⁹; however, the molecular mechanism regulating flavonoid production was not elucidated.

Arabidopsis *FIL* activates *MYB28*, which activates aliphatic glucosinolate biosynthetic genes⁴³. *AtYAB3* is a positive regulator of *MYB75*, an activator of anthocyanin biosynthesis⁴². *MsYABBY5* is a direct activator of *WRKY75*, which possibly represses terpene biosynthesis in *M. spicata*⁴⁴. *OsYABBY4* is a negative regulator of the gibberellin (GA) 20-oxidase 2 gene (*GA20ox2*), and *SLR1*




Fig. 6 Measurement of glandular trichome density and working model of regulation of flavonoid biosynthesis by AaYABBY5. **a** Trichome densities on the leaf surface of transformed and control plants were calculated, and images were captured using fluorescence microscopy (Olympus, Japan). The trichome densities of *AaYABBY5*-OE plants were higher than those of the control and *AaYABBY5* AnT plants. Scale bars represent 200 μ m. **b** Graphical representation of trichome densities of control, *AaYABBY5* AnT, and *AaYABBY5*-OE plants calculated as numbers per millimeter square (no./mm²). Data = mean values \pm standard deviation for $n = 4$. Error bars represent standard deviation for $n = 4$. Student's *t* test with paired and two-tailed distribution methods. * represents $P < 0.05$. **c** The *AaYABBY5* protein directly targets the promoters of *PAL*, lying upstream of the flavonoid biosynthetic pathway. The *PAL* gene, once activated, provides increased metabolic flux toward the first committed step of flavonoid biosynthesis. *AaYABBY5* also directly binds to and activates the *CHS* and *CHI* promoters, providing increasing concentrations of flavonones (total flavonoid content). Going through multistep reactions, in the final step, anthocyanidins, which are the precursors of anthocyanins, are converted into anthocyanins by *AaYABBY5* through the direct activation of the *AaUFGT* promoter. *AaYABBY5* also regulates trichome number and leaf lamina growth through an unknown mechanism

is involved in gibberellic acid responses in rice⁵². Because of the functions performed by YABBY genes in other plants and their role in anthocyanin biosynthesis in *A. thaliana*, it was important to determine the role of YABBY TFs in flavonoid biosynthesis. The role of YABBY family transcription factors in flavonoid biosynthesis in *A. annua* was not previously characterized. This study provides *AaYABBY5* as a novel regulator of flavonoid biosynthesis in *A. annua*.

We screened the promoters of *PAL* (from upstream precursors of the flavonoids pathway), *CHS* (involved in the first committed step toward artemisinin biosynthesis),

CHI, *DFR* (acting downstream of *CHS*) *FLS* and *FSII* as flavonols and flavones regulators, respectively, and *LDOX* and *UFGT* (which are responsible for anthocyanin and pigment formation), using dual-luciferase reporter assay and Y1H assay, and found that *PAL*, *CHS*, *CHI*, and *UFGT* are directly regulated by *AaYABBY5* (Figs. 3 and 4b). The direct binding of *AaYABBY5* to these promoter regions provided the molecular basis for the regulation of flavonoids in *A. annua*. The results of the Y1H assay were consistent with the results of the dual-Luc assay.

In *A. thaliana*, AtFIL was reported to be a target of JAZ3 and a regulator of anthocyanin biosynthesis by

activating the promoter of *MYB75* (*PAP1*); however, no direct interaction with flavonoid pathway genes was reported⁴². In plants, including *A. thaliana*, MYB family proteins have been proven to be candidate genes that regulate flavonoid biosynthesis. AtMYB12 has been found to be a regulator of *CHS* and *FLS* genes through Myb-recognition elements (MREs) present in the promoters of these genes⁵³. The R2R3-MYB transcription factor PAP1 displayed a purple phenotype in transgenic Arabidopsis by upregulating various genes involved in anthocyanin biosynthesis⁵⁴. Recently, the MYB family transcription factor AaTAR2 was found to synergistically regulate both artemisinin (indirectly) and flavonoids in *A. annua*; however, the molecular mechanism regulating flavonoid production was not clearly elucidated, and no direct interaction with the artemisinin biosynthetic pathway was given²⁹. Based on these findings, it was important to study the regulation of flavonoid biosynthetic pathway genes in *AaYABBY5* overexpression plants, *AaYABBY5* antisense plants, and control plants.

To confirm the role of *AaYABBY5* in the regulation of flavonoid biosynthesis, comparative expression and biochemical analysis of transgenic *A. annua* containing pHB-35S: *AaYABBY5*, pHB-35S: anti-*AaYABBY5*, pHB, and wild-type plants revealed a significant upregulation of the *AaPAL*, *AaCHI*, *AaCHS*, *AaFLS*, *AaFSII*, *AaLDOX*, and *AaUFGT* genes under study (Fig. 4c–j). We propose that *AaYABBY5* binds to the *PAL* promoter and activates *PAL* gene expression, which provides metabolic flux to the flavonoid pathway. The overexpression of *AaYABBY5* in *A. annua* provides elevated levels of the *AaYABBY5* protein, which binds to and activates the *PAL* gene. The increased activity of PAL converts more phenylalanine molecules to cinnamate, which provides precursor molecules for entry into the flavonoid pathway. This provides an increase in the concentration of the substrate molecules for the CHS enzyme. Knowing that *AaCHS* catalyzes the first committed step of the flavonoid pathway (Fig. 1) and considering our findings that *AaYABBY5* has a strong binding capability to the *AaCHS* promoter, it is supposed that increased *AaCHS* expression accelerates the biosynthesis of chalcones, which provides flux to downstream reactions, providing an increase in the expression of genes lying downstream of *AaCHS* in the flavonoid biosynthesis in *AaYABBY5*-OE *A. annua* plants. When *PAL* and *CHS* are activated, they perform their function and trigger the activation of downstream reactions, providing accelerated substrate recovery, as a result of which more enzyme molecules are activated, as found by increases in the transcript levels of *CHI*, *FLS*, *FSII*, *LDOX*, and *UFGT*. Among these, *CHI* and *UFGT* are also activated directly by *AaYABBY5*. Therefore, it is concluded that the increase in the concentration of flavonoids in OE *AaYABBY5* transgenic *A. annua* is governed by

AaYABBY5, which directly activates the *PAL*, *CHS*, and *CHI* genes. The results were further validated by analyzing *A. annua* plants with suppressed *AaYABBY5*, wherein the expression of genes regulating flavonoid biosynthesis and flavonoid concentration were found to be significantly decreased (Figs. 4c–j and 5a).

In *A. annua*, chalcone isomerase (*AaCHI*), phenylalanine ammonia lyase (*AaPAL1*), flavanone-3-hydroxylase (*AaF3'H*), and *AaFLS1* have been characterized as enzymes that take part in the flavonoid biosynthesis pathway^{16,17,26,27}. The results obtained in this study are in accordance with these previous findings that, when the expression of *PAL* or *CHI* is increased, it thereby enhances the flavonoid concentration. The term flavonoids used in this study indicates the concentration of flavones and flavonols according to the principle of the AlCl₃ method. Kaempferol and quercetin, are important flavonoids of *A. annua* that belong to the flavonol group, and a significant increase in the flavonoid concentration in *AaYABBY5*-OE plants might correspond to the enhanced accumulation of kaempferol and quercetin, which favors the synergistic role of *AaYABBY5* toward the regulation of flavonoids (present study) and artemisinin biosynthesis⁴⁸.

Anthocyanins are a group of flavonoids that are synthesized downstream of the *AaDFR* gene, governed by *AaLDOX* and *AaUFGT* in two-step reactions. Physiological analysis of transgenic plants and control plants revealed the presence of purple pigmentation in the stems and petioles of *AaYABBY5*-OE plants, whereas no colored phenotype was observed in the control and *AaYABBY5* AnT. plants (Fig. 5b, c). Gene expression analysis also revealed a significant increase in the expression of *AaDFR*, *AaLDOX*, and *AaUFGT*, as well as an increase in the concentration of anthocyanins in *AaYABBY5*-OE plants. As expected, the transcript levels of *AaDFR*, *AaLDOX*, and *AaUFGT* and the resulting anthocyanin concentrations were significantly decreased in *AaYABBY5* AnT. plants (Fig. 4f, i, j). *AaYABBY5* cannot directly bind to *DFR* and *LDOX* promoters. It is supposed that the activation of early flavonoid genes, *CHI* and *CHS*, results in the activation of later steps, which enables the rapid accumulation of anthocyanin precursor molecules; pelargonidin, delphinidin, and cyanidin are collectively named anthocyanidins. Precursor anthocyanins called anthocyanidins are glycosylated by the enzyme UDP-glucose flavonoid 3-*O*-glucosyl transferase (*UFGT*) to produce colorful and stable compounds called anthocyanins²⁵. We hypothesized that the increase in the concentration of anthocyanidins, which are substrate molecules for the *UFGT* enzyme, and direct activation of *AaUFGT* by *AaYABBY5* increases the rate of forward reaction toward anthocyanin biosynthesis governed by glucosylation. These findings prove that *AaYABBY5* is a positive regulator of anthocyanins.

Regarding the basic role of YABBY genes is in the regulation of leaf development, in this study, comparative phenotypic analysis of *AaYABBY5*-OE, *AaYABBY5*-AnT., and control plants showed that as expected, the leaf lamina was broader in overexpression plants (Fig. 5g–h (ii)), whereas radialized lamina was observed in antisense plants (Fig. 5g–h (i)). The differences can also be seen in Fig. 5d (i–iii). These results showed that *AaYABBY5*, in addition to regulating secondary metabolite biosynthesis, also regulates primary metabolism associated with leaf growth. Considering these results, trichomes, which are the sites of secondary metabolite synthesis in plants, were studied. Comparative trichome densities of *AaYABBY5*-OE, *AaYABBY5*-AnT., and control plants revealed higher trichome densities in *AaYABBY5*-OE plants than in the control plants and *AaYABBY5*-AnT. plants; however, the molecular mechanism of its regulation is the target of future research (Fig. 6a, b). A model representing the functions of *AaYABBY5* found in this study is given in Fig. 6c.

This study, for the first time, provided transcriptional regulation of flavonoids using YABBY family transcription factors, and this is the first report on the direct transcriptional regulation of flavonoid pathway genes in *A. annua*. Our previous data that *AaYABBY5* directly activates artemisinin biosynthetic pathway genes⁴⁸ and our present findings regarding *AaYABBY5*-mediated direct regulation of flavonoids, including anthocyanins, provide evidence of the parallel transcriptional regulation of artemisinin and flavonoid biosynthesis by *AaYABBY5* in *A. annua*, thus proving that *AaYABBY5* can be a good candidate gene to provide increasing concentrations of the two biologicals at the same time in *A. annua*.

Materials and methods

Cloning of promoters

The *A. annua* genomic assembly data present in NCBI were screened to find putative promoter sequences upstream of the initiation codon ATG using ORF sequences of *AaPAL*, *AaCHS*, *AaCHI*, *AaFLS*, *AaFSII*, *AaDFR*, *AaLDOX*, and *AaUFGT* as queries.

Approximately 2 kb *PAL* promoter, 1.6 kb *AaCHS* promoter, 1.4 kb *AaCHI* promoter, 1.4 kb *AaFLS* promoter, 1.2 kb *AaFSII* promoter, 1.2 kb *AaDFR* promoter, 1.9 kb *AaLDOX* promoter, and 1.7 kb promoter sequence of *AaUFGT* were cloned. Genomic DNA extracted from young leaves of *A. annua* was used as a template, and amplification was performed using a KOD Plus PCR kit (Toyobo). DNA bands purified using the DNA Gel Extraction Kit (AxyPrep) were ligated to the PLB simple vector (Tiangen Biotech, China) containing carbenicillin-resistance genes according to the manufacturer's instructions provided with the Lethal Based Fast Cloning Kit (Tiangen), followed by transformation into DH5 α -competent cells (Invitrogen).

Bioinformatic analysis of promoters of flavonoid biosynthetic genes

The cloned promoter sequences were confirmed by Sanger sequencing and analyzed for the prediction of putative YABBY-binding sites³⁷ using PLANTPAN3.0.

Identification of *AaYABBY5* as a potential flavonoid-regulating transcription factor

The transcriptomic data of six different tissues of *A. annua* generated by our research center were screened, and reads of each YABBY gene, flavonoid biosynthetic genes, and artemisinin biosynthetic genes in six different tissues (trichome, bud, stem, root, leaf, and seed) were obtained using a BLASTN search against each database with an *E* value $<1 \times 10^{-6}$, and read counts were normalized by calculating the value of reads per kilobase per million (RPKM). To predict the YABBY transcription factor, which may regulate flavonoid-regulating genes, coexpression analysis was performed on the basis of RNA sequencing data using Multi-Experiment Viewer (MeV4.9.0) software⁵⁵.

Plant material and growth conditions

The seeds of the *A. annua* cultivar Huhao #1 used in this study were collected from our previous stably transformed transgenic *A. annua* plants with overexpressed *AaYABBY5* (35S—*AaYABBY5*), silenced *AaYABBY5* (35S-anti-*AaYABBY5*), empty vector-containing plants (35S), and wild-type or control plants (untransformed). Details about construct preparation and *A. annua* transformation are given in ref. ⁴⁸. *A. annua* plants (transgenic and control) were also propagated from stem cuttings. For such cutting propagations, 8–10-cm long stems or side shoots were cut just below a leaf and propagated in plant trays supplemented with soil matrix. After extensive rooting development and proper growth as a complete plant, the cuttings were propagated to pots. In the transient agroinfiltration assay, *N. benthamiana* plants with young leaves and expanded lamina were used. Both plant types were grown under the same conditions of temperature: 25 °C \pm 2 °C and photoperiod: 16 h light:8 h dark.

Transient *N. benthamiana* infiltration system

To perform a transient infiltration assay in the *N. benthamiana* system using a dual-LUC kit, effector and reporter strains were prepared as follows. For the effector strain, the open-reading frame of *AaYABBY5* without its terminal codon was amplified using primer sequences specifically designed for cloning into the pENTR-TOPO vector, followed by subsequent recombination into the Gateway destination vector pEarleyGate 104-YFP (N-terminal YFP) (Invitrogen, USA). For the reporter construct, promoter sequences of *PAL*, *CHS*, *CHI*, *FLS*, *FSII*, *LDOX*, and *UFGT* were amplified with primer sequences

containing adapter sequences specific to the pGreenII 0800-LUC plasmid. The purified fragment was subsequently ligated into the pGreenII 0800-LUC plasmid through HindIII and PstI sites to generate pro:*LUC* constructs, according to a previous protocol⁴⁸. The plasmids were transformed into *A. tumefaciens* GV3101 competent cells with pSoup-p19 to help suppress gene silencing. The process of agrobacterium culture preparation, infiltration into the *N. benthamiana* leaf tissues, sample collection, and preparation for the measurement of LUC activity was followed according to a previous protocol⁴⁸.

Yeast one-hybrid system

For the Y1H assay, bait and prey constructs were prepared as follows. The promoter regions of *AaPAL*, *AaCHS*, *AaCHI*, *AaFLS*, *AaFSL*, *AaDFR*, *AaLDOX*, and *AaUFGT* were amplified using primer sequences with 5' adapter sequences specific to the placZ vector and subsequently ligated into placZ using one-step cloning, following the protocol given by ClonExpress II (Vazyme). Similarly, the open-reading frame of *AaYABBY5* with its terminal codon was amplified using primer sequences with 5' adapter sequences specific to the pB42AD vector (Addgene) and subsequently ligated into it. Both constructs were confirmed by sequencing. Promoter sequences cloned in placZ were used as bait against pB42AD- *AaYABBY5* prey.

The assay was performed according to the manual of the Matchmaker Gold Y1H system given in the *Yeast Protocols Handbook*, Clontech, (Japan). The combination of pB42AD-*AaYABBY5* with the respective promoter sequence in placZ was cotransformed into EGY48 competent cells. The combination of negative controls pB42AD- *AaYABBY5* with empty lacZ (lacZ-0), pB42AD-0 with the placZ promoter, and empty pB42AD (pB42AD-0) with empty placZ (placZ-0) was also transformed into EGY48 cells. Transformed yeast cells were streaked on SD-Tryptophan-Uracil (SD/-Trp-Ura). After 3 days, colonies were shifted to SD/-Trp-Ura supplemented with X-Gal (5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside). The experiment was repeated three times.

RNA extraction and gene expression analysis

To check the expression pattern using real-time PCR, RNA extraction and cDNA synthesis were performed using an RNAPrep Pure Plant Kit (Tiangen) and PrimeScript RT Reagents (Takara), respectively. Pure and good quality 500 ng RNA was used as a template for cDNA synthesis using reverse transcriptase enzyme. The cDNA was diluted in RNase-free ddH₂O at a 1:40 ratio. Six microliters of diluted cDNA was used as a template for real-time PCR analysis using the protocol of ref. ²⁸ with minor modifications.

SYBR Green Super real premix was used to prepare a master mix, and a Roche Light Cycler 96-well Real-Time PCR Machine (Roche, Switzerland) was used to quantify the expression. The PCR program was set up at 40 cycles, each containing 2 min at 95 °C for initial denaturation, 20 s at 95 °C for denaturation, 20 s at 54 °C for annealing, and 20 s at 72 °C for the extension. β -Actin was used as a standard. For each experimental and control sample, three repeats were used.

Preparation of flavonoid extracts and measurement of total flavonoid content

For quantitative determination of the total flavonoid content, leaf samples from each plant were collected and ground to powder form using liquid nitrogen (−70). Approximately 1 g of each leaf sample was placed in 50-ml glass flasks and mixed with 5 ml methanolic solution (70%). The extraction was performed twice using an ultrasonic processor (DL-720B) at a frequency of 55 Hz and a temperature of 30 °C for 35 min. The supernatants containing extracts were collected following centrifugation at 6000 rpm for 10 min. Approximately 3 ml of each extract was filtered through a 0.22- μ m nitrocellulose filter and transferred to glass vials.

The total flavonoid content of *A. annua* exudates was calculated using the aluminum chloride (AlCl₃) colorimetric method according to protocol⁵⁶. The working principle of the AlCl₃ colorimetric method is that AlCl₃ forms acid-stable complexes with the C-4 keto group and the C-3 or C-5 hydroxyl group of flavones and flavonols. It also forms complexes with the orthodihydroxyl groups in the A- or B-ring of flavonoids. For sample preparation, 10 μ L of each extract was diluted in 150 ml of deionized water, followed by the addition of 20 μ l potassium acetate (1 M) and 20 μ l of aluminum chloride (10%). The final volume of the 200 μ l reaction mixture was incubated at 37 °C for 30 min. The absorbance was measured at 415 nm. The total flavonoid content was calculated as quercetin equivalents. The absorbance of three independent biological repeats from each plant type: *AaYABBY5* OE, *AaYABBY5* AnT., as well as control plants, was measured in triplicate. Error bars indicate the SD of the average flavonoid concentration.

Preparation of anthocyanin extracts and measurement of anthocyanins

To extract anthocyanins/pigments from the stem tissues of *A. annua* plants, the following protocol⁵⁷ was used with minor modifications. Samples from each plant were collected and ground in liquid nitrogen to powder form. A 1 g sample was mixed with 10 ml of acidic methanol (70% methanol with 0.1% HCl) and processed by an ultrasonic processor (DL-720B) at 50 Hz for 40 min. The processed

samples were kept at 4°C overnight mixing and incubation under dark conditions.

The extracts were purified by centrifugation at 12000 rpm for 10 min by filtration. The supernatants were separated and filtered through 0.22- μ m nitrocellulose filters. Absorbance at 530 nm and 657 nm was measured using a microplate reader (BioTek, ELx 800). Relative anthocyanin content was found by $(A_{530}-0.25 \times A_{657})/FW$ (FW is the fresh weight of sample in grams). The absorbance of each sample was measured in triplicate. Three independent biological replicates from each plant type were used. Error bars indicate the standard deviation of the average anthocyanin concentration.

Measuring trichome density

To calculate the trichome densities, mature and healthy leaves from three-month-old *A. annua* plants were used. To ensure comparison at the same growth level, the 8th leaf below the meristem was selected from transformed and control plants. Images were captured using fluorescence microscopy (Olympus, Japan) under the following conditions: 5x magnification and 450–480 nm excitation wavelength. The images were analyzed by using ImageJ 1.51k software to measure the leaf area and calculate trichome numbers. Finally, trichome densities were calculated from the ratio of trichome number to the leaf area. The experiment was repeated three times, using the 8th leaf from three different plants of each line, to obtain statistically significant results.

Primer sequences

The primer sequences used in this study were prepared using Primer 3 software and are given in Supplementary Table S1.

Statistical analysis

Statistical analysis was performed using Student's *t* test with paired and two-tailed distribution methods. ** and * represent statistically significant group differences for $P < 0.01$ and < 0.05 , respectively.

Acknowledgements

This work was supported by the National Key R&D Program of China (2018YFA0900600), the Bill & Melinda Gates Foundation (OPP1199872), the SJTU Global Strategic Partnership Fund (2020SJTU-CORNELL), and the SJTU Trans-med Awards Research Program (20190104).

Author contributions

S.K. and K.T. planned and designed the research. S.K. performed the experiments. Y.L., D.H., and Q.S. helped with promoter cloning and vector construction. X.F. helped with the measurement of flavonoid contents. C.W. helped with capturing plant images. S.K. wrote the manuscript. S.R., Q.S., and K.T. revised the manuscript. All authors approved the manuscript for submission.

Data availability

All data supporting this study are included in the article and its supplementary files.

Conflict of interest

The authors declare no competing interests.

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41438-021-00693-x>.

Received: 16 April 2021 Revised: 24 July 2021 Accepted: 15 August 2021
Published online: 01 December 2021

References

- Debeaujon, I., Peeters, A. J., Léon-Kloosterziel, K. M. & Koornneef, M. The TRANSPARENT TESTA12 gene of *Arabidopsis* encodes a multidrug secondary transporter-like protein required for flavonoid sequestration in vacuoles of the seed coat endothelium. *Plant Cell* **13**, 853–871 (2001).
- Kitamura, S. Transport of flavonoids: from cytosolic synthesis to vacuolar accumulation. in *The Science of Flavonoids* (ed. Grotenwold, E) 123–146 (Springer New York, 2006).
- Weathers, P. J., Towler, M., Hassanali, A., Lutgen, P. & Engeu, P. O. Dried-leaf *Artemisia annua*: a practical malaria therapeutic for developing countries. *World J. Pharmacol.* **3**, 39–55 (2014).
- Ounjajean, S., Benjasak, N., Sae-lao, S. & Somsak, V. Kaempferol addition increases the antimalarial activity of artesunate in experimental mice. *J. Tropical Med.* **2020**, 6165928 (2020).
- Desrosiers, M. R. & Weathers, P. J. Artemisinin permeability via Caco-2 cells increases after simulated digestion of *Artemisia annua* leaves. *J. Ethnopharmacol.* **210**, 254–259 (2018).
- Song, Y. et al. Polyphenolic profile and antioxidant effects of various parts of *Artemisia annua* L. *Biomed. Chromatogr.* **30**, 588–595 (2016).
- Li, Y. J. et al. Flavonoids casticin and chrysosplenol D from *Artemisia annua* L. inhibit inflammation in vitro and in vivo. *Toxicol. Appl. Pharmacol.* **286**, 151–158 (2015).
- Ko, Y. S. et al. Polyphenols from *Artemisia annua* L. inhibit adhesion and EMT of highly metastatic breast cancer cells MDA-MB-231. *Phytother. Res.* **30**, 1180–1188 (2016).
- Henrich, C. J. et al. A high-throughput cell-based assay for inhibitors of ABCG2 activity. *J. Biomolecular Screen.* **11**, 176–183 (2006).
- Du, G. J. et al. Luteolin as a glycolysis inhibitor offers superior efficacy and lesser toxicity of doxorubicin in breast cancer cells. *Biochem. Biophys. Res. Commun.* **372**, 497–502 (2008).
- Kim, J. Y. et al. Quercetin sensitizes human hepatoma cells to TRAIL-induced apoptosis via Sp1-mediated DR5 up-regulation and proteasome-mediated c-FLIPs down-regulation. *J. Cell. Biochem.* **105**, 1386–1398 (2008).
- Siegelin, M. D., Reuss, D. E., Habel, A., Herold-Mende, C. & von Deimling, A. The flavonoid kaempferol sensitizes human glioma cells to TRAIL-mediated apoptosis by proteasomal degradation of survivin. *Mol. Cancer Therapeutics* **7**, 3566–3574 (2008).
- Szliszka, E., Czuba, Z. P., Jernas, K. & Król, W. Dietary flavonoids sensitize HeLa cells to tumor necrosis factor-related apoptosis-inducing ligand (TRAIL). *Int. J. Mol. Sci.* **9**, 56–64 (2008).
- Zhou, L. et al. Silibinin restores paclitaxel sensitivity to paclitaxel-resistant human ovarian carcinoma cells. *Anticancer Res.* **28**, 1119–1127 (2008).
- Ferreira, J. F., Simon, J. E. & Janick, J. N. Developmental studies of *Artemisia annua*: flowering and artemisinin production under greenhouse and field conditions. *Planta Med.* **61**, 167–170 (1995).
- Liu, S. et al. Comparative transcriptomic analysis of key genes involved in flavonoid biosynthetic pathway and identification of a flavonol synthase from *Artemisia annua* L. *Plant Biol.* **19**, 618–629 (2017).
- Xiong, S. et al. Molecular cloning and characterization of a flavanone 3-hydroxylase gene from *Artemisia annua* L. *Plant Physiol. Biochem.* **105**, 29–36 (2016).
- Quattrocchio, F. et al. PH4 of *Petunia* is an R2R3 MYB protein that activates vacuolar acidification through interactions with basic-helix-loop-helix transcription factors of the anthocyanin pathway. *Plant Cell* **18**, 1274–1291 (2006).
- Feng, F., Li, M., Ma, F. & Cheng, L. Phenylpropanoid metabolites and expression of key genes involved in anthocyanin biosynthesis in the shaded peel of apple fruit in response to sun exposure. *Plant Physiol. Biochem.* **69**, 54–61 (2013).
- Winkel-Shirley, B. Flavonoid biosynthesis. A colorful model for genetics, biochemistry, cell biology, and biotechnology. *Plant Physiol.* **126**, 485–493 (2001).

21. Dao, T. T., Linthorst, H. J. & Verpoorte, R. Chalcone synthase and its functions in plant resistance. *Phytochemistry Rev.: Proc. Phytochem. Soc. Eur.* **10**, 397–412 (2011).
22. Holton, T. A. & Cornish, E. C. Genetics and biochemistry of anthocyanin biosynthesis. *Plant Cell* **7**, 1071–1083 (1995).
23. Jez, J. M., Bowman, M. E., Dixon, R. A. & Noel, J. P. Structure and mechanism of the evolutionarily unique plant enzyme chalcone isomerase. *Nat. Struct. Biol.* **7**, 786–791 (2000).
24. Grotewold, E. The genetics and biochemistry of floral pigments. *Annu. Rev. Plant Biol.* **57**, 761–780 (2006).
25. Sui, X. et al. cDNA cloning and characterization of UDP-glucose:anthocyanidin 3-O glucosyltransferase in *Freesia hybrida*. *Plant Cell Rep.* **30**, 1209–1218 (2011).
26. Zhang, Y. et al. Molecular cloning and promoter analysis of the specific salicylic acid biosynthetic pathway gene phenylalanine ammonia-lyase (*AaPAL1*) from *Artemisia annua*. *Biotechnol. Appl. Biochem.* **63**, 514–524 (2016).
27. Ma, J., Fu, X., Zhang, T., Qian, H. & Zhao, J. Cloning and analyzing of chalcone isomerase gene (*AaCHI*) from *Artemisia annua*. *Plant Cell, Tissue Organ Cult.* **137**, 45–54 (2019).
28. Shen, Q. et al. The jasmonate-responsive *AaMYC2* transcription factor positively regulates artemisinin biosynthesis in *Artemisia annua*. *N. Phytologist* **210**, 1269–1281 (2016).
29. Zhou, Z. et al. *TRICHOME AND ARTEMISININ REGULATOR 2* positively regulates trichome development and artemisinin biosynthesis in *Artemisia annua*. *N. Phytologist* **228**, 932–945 (2020).
30. Zhang, F., Gonzalez, A., Zhao, M., Payne, C. T. & Lloyd, A. A network of redundant bHLH proteins functions in all TTG1-dependent pathways of *Arabidopsis*. *Development* **130**, 4859–4869 (2003).
31. Zimmermann, I. M., Heim, M. A., Weisshaar, B. & Uhrig, J. F. Comprehensive identification of *Arabidopsis thaliana* MYB transcription factors interacting with R/B-like BHLH proteins. *Plant J.* **40**, 22–34 (2004).
32. Bernhardt, C., Zhao, M., Gonzalez, A., Lloyd, A. & Schiefelbein, J. The bHLH genes *GL3* and *EGL3* participate in an intercellular regulatory circuit that controls cell patterning in the *Arabidopsis* root epidermis. *Development* **132**, 291–298 (2005).
33. Gonzalez, A., Zhao, M., Leavitt, J. M. & Lloyd, A. M. Regulation of the anthocyanin biosynthetic pathway by the TTG1/bHLH/Myb transcriptional complex in *Arabidopsis* seedlings. *Plant J.* **53**, 814–827 (2008).
34. Li, S. F. et al. The *Arabidopsis* MYB5 transcription factor regulates mucilage synthesis, seed coat development, and trichome morphogenesis. *Plant Cell* **21**, 72–89 (2009).
35. Bowman, J. L. The YABBY gene family and abaxial cell fate. *Curr. Opin. Plant Biol.* **3**, 17–22 (2000).
36. Bowman, J. L. & Smyth, D. R. *CRABS CLAW*, a gene that regulates carpel and nectary development in *Arabidopsis*, encodes a novel protein with zinc finger and helix-loop-helix domains. *Development* **126**, 2387–2396 (1999).
37. Franco-Zorrilla, J. M. et al. DNA-binding specificities of plant transcription factors and their potential to define target genes. *Proc. Natl Acad. Sci. USA* **111**, 2367–2372 (2014).
38. Yamada, T., Ito, M. & Kato, M. YABBY2-homologue expression in lateral organs of *Amborella trichopoda* (Amborellaceae). *Int. J. Plant Sci.* **165**, 917–924 (2004).
39. Lee, J. Y. H. et al. Recruitment of *CRABS CLAW* to promote nectary development within the eudicot clade. *Development* **132**, 5021–5032 (2005).
40. Siegfried, K. R. et al. Members of the YABBY gene family specify abaxial cell fate in *Arabidopsis*. *Development* **126**, 4117–4128 (1999).
41. Stähle, M. I., Kuehlich, J., Staron, L., Von Arnim, A. G. & Golz, J. F. YABBYs and the transcriptional corepressors *LEUNIG* and *LEUNIG_HOMOLOG* maintain leaf polarity and meristem activity in *Arabidopsis*. *Plant Cell* **21**, 3105–3118 (2009).
42. Boter, M. et al. *FILAMENTOUS FLOWER* is a direct target of *JAZ3* and modulates responses to jasmonate. *Plant Cell* **27**, 3160–3174 (2015).
43. Douglas, S. J., Li, B., Kliebenstein, D. J., Nambara, E. & Riggs, C. D. A novel filamentous flower mutant suppresses brevipedicellus developmental defects and modulates glucosinolate and auxin levels. *PLoS ONE* **12**, e0177045–e0177045 (2017).
44. Wang, Q. et al. Metabolic engineering of terpene biosynthesis in plants using a trichome-specific transcription factor *Ms YABBY5* from spearmint (*Mentha spicata*). *Plant Biotechnol. J.* **14**, 1619–1632 (2016).
45. Zhao, S. et al. Genome-wide analysis of the YABBY family in soybean and functional identification of *GmYABBY10* involvement in high salt and drought stresses. *Plant Physiol. Biochem.* **119**, 132–146 (2017).
46. Shen, Q. et al. The genome of *Artemisia annua* provides insight into the evolution of Asteraceae family and artemisinin biosynthesis. *Mol. Plant* **11**, 776–788 (2018).
47. Tattini, M., Gravano, E., Pinelli, P., Mulinacci, N. & Romani, A. J. Flavonoids accumulate in leaves and glandular trichomes of *Phillyrea latifolia* exposed to excess solar radiation. *N. Phytologist* **148**, 69–77 (2000).
48. Kayani, S. et al. The YABBY family transcription factor *AaYABBY5* directly targets cytochrome P450 monooxygenase (*CYP71AV1*) and double-bond reductase 2 (*DBR2*) involved in artemisinin biosynthesis in *Artemisia Annua*. *Front. Plant Sci.* **10**, 1084 (2019).
49. Yan, T. et al. *HOMEODOMAIN PROTEIN 1* is required for jasmonate-mediated glandular trichome initiation in *Artemisia annua*. *N. Phytologist* **213**, 1145–1155 (2017).
50. Dhingra, V., Vishweshwar Rao, K. & Lakshmi Narasu, M. Current status of artemisinin and its derivatives as antimalarial drugs. *Life Sci.* **66**, 279–300 (2000).
51. Kim, W. S. et al. Anti-inflammatory, antioxidant and antimicrobial effects of artemisinin extracts from *Artemisia annua* L. *The Korean. J. Physiol. Pharmacol.* **19**, 21–27 (2015).
52. Yang, C., Ma, Y. & Li, J. The rice YABBY4 gene regulates plant growth and development through modulating the gibberellin pathway. *J. Exp. Bot.* **67**, 5545–5556 (2016).
53. Mehrtens, F., Kranz, H., Bednarek, P. & Weisshaar, B. The *Arabidopsis* transcription factor *MYB12* is a flavonol-specific regulator of phenylpropanoid biosynthesis. *Plant Physiol.* **138**, 1083–1096 (2005).
54. Tohge, T. et al. Functional genomics by integrated analysis of metabolome and transcriptome of *Arabidopsis* plants over-expressing an MYB transcription factor. *Plant J.* **42**, 218–35 (2005).
55. Saeed, A. I. et al. *TM4*: a free, open-source system for microarray data management and analysis. *BioTechniques* **34**, 374–378 (2003).
56. Kumari, A. & Pandey-Rai, S. Enhanced arsenic tolerance and secondary metabolism by modulation of gene expression and proteome profile in *Artemisia annua* L. after application of exogenous salicylic acid. *Plant Physiol. Biochem.* **132**, 590–602 (2018).
57. Kubasek, W. L. et al. Regulation of flavonoid biosynthetic genes in germinating *Arabidopsis* seedlings. *Plant Cell* **4**, 1229–1236 (1992).