

Construction of Spirooxindole Analogues Engrafted with Indole and Pyrazole Scaffolds as Acetylcholinesterase Inhibitors

Mohammad Shahidul Islam, Abdullah Mohammed Al-Majid, Mohammad Azam, Ved Prakash Verma,* Assem Barakat,* Matti Haukka, Abdullah A. Elgazar, Amira Mira, and Farid A. Badria



Cite This: *ACS Omega* 2021, 6, 31539–31556



Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: Twenty-five new hits of spirooxindole analogs **8a–y** grafted with indole and pyrazole scaffolds were designed and constructed *via* a [3+2]-cycloaddition (32CA) reaction starting from three components: new chalcone-based indole and pyrazole scaffolds **5a–d**, substituted isatins **6a–c**, and secondary amines **7a–d**. The potency of the compounds were assessed in modulating cholinesterase (AChE) activity using Ellman's method. Compounds **8i** and **8y** showed the strongest acetylcholine esterase inhibition (AChEI) with IC₅₀ values of 24.1 and 27.8 μM, respectively. Molecular docking was used to study their interaction with the active site of hAChE.



INTRODUCTION

Neurodegeneration is a key aspect of a large number of diseases that come under the umbrella of neurodegenerative disease. Of these different disorders, the most notable are Parkinson's disease, Huntington's disease, and Alzheimer's disease (AD). Alzheimer's disease (AD) symptoms are memory loss, impairment of cognitive functions, and dementia. AD involves two major neuropathological hallmarks causing neuronal dysfunctions and cell death: the presence of extracellular amyloid β-peptide (Aβ) deposits (senile plaques) and aggregates of the hyperphosphorylated tau protein (neurofibrillary tangles)¹ along with the tau hyperphosphorylation are the most proposed pathogenetic mechanisms,² and mitochondrial cascade hypothesis has attracted much interest recently.³ Other much debatable AD hypotheses are the tau hypothesis,⁴ cholesterol hypothesis,⁵ inflammatory hypothesis,⁶ oxidative stress hypothesis,⁷ metal hypothesis,⁸ vascular hypothesis,⁹ and cell cycle hypothesis.¹⁰ Up to date, rivastigmine, galantamine, and donepezil represent the only ChE inhibitors approved for AD treatment, differing in chemical structures and pharmacologic and pharmacokinetic profiles. To design and discover a new agent that might work as ChE inhibitors is a challenge.

Heterocycles having azoles as a core structure have been discovered for several applications.¹¹ Pyrazoles are one of the important heterocycles, which exhibited significant properties in material sciences,¹² agriculture development,¹³ medicine,¹⁴ and pharmacological applications.¹⁵ Among pharmacological applications are antibiotic,¹⁶ sensors,¹⁷ pesticide,¹⁸ antibacterial,¹⁹ and antifungal activities.²⁰ On the other hand, several

molecules grafted with the pyrazole scaffold have shown high efficacy toward antiviral,²¹ antitumor,²² anti-inflammatory,²³ antioxidant,²⁴ and antidepressant activities.²⁵ Indeed, many drugs incorporated the pyrazole moiety employed for the treatment of metabolic disorder diseases such as Alzheimer's,²⁶ Parkinson's,²⁷ and neuroprotective,²⁸ which makes this pharmacophore very attractive for drug discovery. One representative example of advanced glycation inhibitors reported by Han et al. is that this agent is based on the pyrazole-5-carboxamide as a core structure.²⁹ However, Turkan et al. have reported substituted pyrazole derivatives, which have been discovered as potent cholinesterase inhibitors.³⁰

Spirooxindoles exhibit a broad range of biological effects and are well-tolerated in biomedical applications.³¹ Their applications use AChEs.³² Kia et al. reported representative examples, including spirooxindoles grafted with piperidine and pyrrolizine scaffolds, which are found to be beneficial for AChE (compound III; IC₅₀ = 3.36 μM or 2.28 ± 0.07 μg/mL) (Figure 1),^{32d} and another representative example based on mono- and *bis*-spiro-pyrrolidines, where the hit IV shows high efficacy against AChE with an IC₅₀ value of 2.35 μM (Figure 1).^{32f} Chigurupati et al. reported indolopyrazolines with the

Received: July 26, 2021

Accepted: November 1, 2021

Published: November 12, 2021



ACS Publications

© 2021 The Authors. Published by
American Chemical Society

31539

<https://doi.org/10.1021/acsomeg.1c03978>
ACS Omega 2021, 6, 31539–31556







Figure 1. Significant acetylcholinesterase (AChE) inhibitory activity of representative spirooxindole analogues.

Scheme 1. Synthesis of the Chalcone Engrafted with Indole and Pyrazole Scaffolds 5a–d



high biochemical application against AChE inhibition (**V**; $\text{IC}_{50} = 13.77 \pm 0.25 \mu\text{M}$), respectively.³³ Extending our recent efforts on the development of cholinesterase inhibition, Barakat et al. reported a new series of spirooxindoles grafted with the benzo[*b*]thiophene scaffold, which were found to exhibit moderate potential against AD (Figure 1).^{32b,34} The above reports inspired the investigation of several pharmacophores inside the rigid spirooxindole privileged structure, such as indole and pyrazole scaffolds, which might act as better AChE inhibitors.


Several progresses have been devoted toward the synthesis of spirooxindoles recently.³⁵ In between these approaches, the [3 + 2] cycloaddition reaction protocol was efficient and promising to afford the spirooxindole privileged structures with several stereogenic centers.³⁶

In this paper, we described the [3 + 2] cycloaddition reaction approach for the synthesis of new spirooxindoles based on a new chalcone grafted with indole and pyrazole motifs. Many substituted isatins and amino acids were also investigated. The biochemical potential of AChE was also studied.


RESULTS AND DISCUSSION

Synthesis of Spirooxindole Analogs 8a–y Engrafted with Indole and Pyrazole Scaffolds. The indole and pyrazole scaffolds as interesting pharmacophores are combined into the spirooxindole analogs for exploring the acetylcholinesterase (AChE) inhibitory activity. Initially, we synthesized the new chalcones **5a–d** by aldol condensation of acetylpyrazole and substituted indole-3-carbaldehyde in basic

Scheme 2. Synthesis of the Spirooxindole Analogues Engrafted with Indole and Pyrazole Scaffolds 8a–y



Scheme 3. Proposed Approach for the [3 + 2] Cycloaddition Reaction, Explaining the Regio- and Diastereoselective Synthesis



condition under reflux for 72 h. The new chalcone-based indole and pyrazole scaffolds, which act as a synthon for the [3 + 2] cycloaddition (32CA) reaction, are depicted in Scheme 1. Subsequently, the new series of spirooxindole analogues tethered indole and pyrazole scaffolds were constructed *via* a one-pot multicomponent reaction approach³⁷ in MeOH under reflux conditions for 2–24 h (Scheme 2). Twenty-five

examples were achieved by variation of many substituted isatins **6a–c** with different electronic effects (isatin **6a**; 6-chloroisatin **6b**; 5-nitroisatin **6c**) with four different amino acids **7a–d** (thioproline **7a**, octahydro-1*H*-indole-2-carboxylic acid **7b**; L-proline **7c**; sacrosine **7d**). The spirooxindole analogues engrafted with indole and pyrazole scaffolds were isolated in a single regio- and diastereoselective isomer in

acceptable to excellent chemical yield up to 86%. The proposed mechanism, as shown in **Scheme 3**, proceeded *via* an *ortho/endo* 32CA approach.³⁸ The optical rotation of the synthesized compounds was measured, and the regio- and diastereoselectivity of the cycloadducts were confirmed. Single-crystal X-ray diffraction analysis of compound **8c** confirmed that our hypothesis belongs to the final stereoselectivity of the cycloadducts of the final compounds. The absolute configurations of products **8a–y** were assigned based on the obtained x-ray diffraction analysis as follows for the 4 stereogenic centers as R, S, R, S.

Crystal Structure Description. The X-ray structure of **8c** showing atom numbering and thermal ellipsoids drawn at a 30% probability level is shown in **Figure 2**. The structure




Figure 2. Atom numbering and thermal ellipsoids at a 30% probability level for **8c**.

agreed very well with the spectral analyses and revealed the presence of asymmetric centers at C9, C21, C10, and C13. The crystal data and structure refinement details are depicted in **Table 1**. The compound crystallized in the triclinic system and the *P*1 space group with unit cell parameters of $a = 12.71740(10)$ Å, $b = 15.72830(10)$ Å, $c = 19.3853(2)$ Å, and $\beta = 105.9640(10)$ °. The unit cell volume is $3727.97(6)$ Å³ with $Z = 4$. The asymmetric unit comprised one molecule and two chloroform molecules as the crystal solvent. The selected bond distances and angles are listed in **Table S1** (Supporting Information).

The molecular packing in **8c** is controlled by strong N–H...O and N–H...N hydrogen bonds as well as weak C—H...X interactions (X=Cl, Br, N, or O). The corresponding hydrogen bond parameters are listed in **Table S2** (Supporting Information) and are shown in the left part of **Figure 3**. The hydrogen bond network is shown in the right part of the same figure.

Acetylcholine Esterase Inhibitory Activity. The ability of the synthesized compounds to inhibit acetylcholine esterase (AChE) was evaluated using Ellman's method.³⁹ Compounds **8i** and **8y** showed the strongest acetylcholine esterase inhibition (AChEI) with IC₅₀ values of 24.1 and 27.8 μM, respectively.

Four Compounds; **8c**, **8d**, **8f**, **8h**, **8j**, **8w**, and **8x** showed moderate activity inhibitory activity (with IC₅₀ ≤ 50 μM). Compounds **8m**, **8o**, **8p**, **8q**, and **8s** had weak activity (with

Table 1. Crystal Data

	8c
CCDC	2105913
empirical formula	$C_{34}H_{28}BrCl_6N_5O_2S$
fw	863.28
temp (K)	120(2)
λ (Å)	1.54184
cryst syst	monoclinic
space group	<i>P</i> 21/n
a (Å)	12.71740(10)
b (Å)	15.72830(10)
c (Å)	19.3853(2)
β (deg)	105.9640(10)
V (Å ³)	3727.97(6)
Z	4
ρ_{calc} (Mg/m ³)	1.538
μ (Mo K α) (mm ⁻¹)	6.305
no. reflns.	38891
unique reflns.	7824
completeness to $\theta = 67.684^\circ$	100.0%
GOOF (F^2)	1.024
R_{int}	0.0352
R_1^a ($I \geq 2\sigma$)	0.0354
wR ₂ ^b ($I \geq 2\sigma$)	0.0861

$$^a R_1 = \sum ||F_o|| - |F_c|| / \sum |F_o|. \quad ^b wR_2 = [\sum [w(F_o^2 - F_c^2)^2] / \sum [w(F_o^2)]^{1/2}]^{1/2}.$$

IC₅₀ values 65–90 μM), while compounds **8b**, **8e**, **8q**, **8k**, **8e**, **9r**, **8r**, **8t**, **8u**, and **8v** were in active with IC₅₀ > 100 μM (**Table 2**).

Molecular Docking Study. Molecular docking has been used extensively to identify and explain the molecular mechanism of several lead compounds in drug discovery.⁴⁰ The software validation revealed that it was able to reproduce the experimental pose with root-mean-square deviation (RMSD) equal to 0.39 (**Figure 4**). Since *in vitro* enzyme inhibition showed that compounds **8i**, **8h**, and **8y** are the most active among the synthesized compounds with moderate inhibition activity, we used molecular docking to gain insights into the molecular interaction of the compound with the active site of hACHe.

In the context of the total energy required for binding, the cocrystallized ligand achieved much better binding affinity than the 3 compounds, yet their binding energy was found to be reasonable with respect to their moderate activity in the enzyme inhibition assay, as presented in **Table 3**. Post docking analysis showed that compound **8i** has different binding modes rather than compounds **8h** and **8y**, as the first one tends to bind in the middle of the gorge of hACHe. On the other hand, compounds **8h** and **8y** showed most of their interactions with amino acids located at the entrance of the gorge. For example, compound **8i** was able to interact with amino acids in diverse sites in the enzyme such as TRP-86 and TYR-337 in the anionic site, Ser-203 and HIS-447 in the catalytic site, PHE-295 and PHE-297 in the acyl binding site, and formed 2 hydrogen bonds with SER-125 and GLU-202, as depicted in **Figure 5**, these interactions are well reported to be important for achieving good inhibitory activity.⁴¹

On the other hand, compound **8y** showed interactions only with TYR-337 and PHE-295 from the anionic and acyl binding sites, respectively, but with less hydrophobic interaction than **8i**. Nevertheless, they were able to access the peripheral anionic site by interacting with amino acids, such as TYR-124,




Figure 3. Hydrogen bond contacts (left) and packing of molecular units *via* hydrogen bonding interactions (right).

TRY-341, and TRP-286, as shown in Figures 6 and 7, which might explain their ability to achieve the moderate inhibitory effect as observed in the enzyme inhibition assay.

CONCLUSIONS

We summarized in this paper that we had synthesized 25 hits based on a spirooxindole scaffold grafted with two other pharmacophores, including indole and pyrazole moieties. The results of the AChE assay show that compounds **8i** and **8y** show the strongest acetylcholine esterase inhibition (AChEI) with IC₅₀ values of 24.1 and 27.8 μM, respectively. The AChE activity exhibited promising results, which make them candidates for further research.

EXPERIMENTAL SECTION

General information. Phenylhydrazine, acetylacetone, N,N-dimethylformamide-dimethyl acetal (DMF-DMA), Piperidine, and NaOH were purchased from Aldrich and used as received. All of the indole derivatives were purchased from Aldrich and used as is. All bases were used as received (in air) or dried under vacuum at 100 °C (under an inert atmosphere). All solvents were used as received when experiments were conducted in air. Flash chromatography was performed on 100–200 mesh silica gel. ¹H- and ¹³C-nuclear magnetic resonance (NMR) spectra were recorded on JEOL-700 MHz spectrometers at ambient temperature in CDCl₃ & DMSO-d₆, which were purchased from Sigma-Aldrich. Chemical shifts (ppm) are referenced to the residual solvent peak. Coupling constants, J, are given in hertz. Abbreviations used in the designation of the signals: s = singlet, d = doublet, dd = doublet of doublets, ddd = doublet of doublet of doublets, dt = doublet of triplets, t = triplet, td = triplet of doublets, and m = multiplet. All melting points were measured on a Gallenkamp melting point apparatus in open glass capillaries and are uncorrected. IR Spectra were measured as KBr pellets on a Nicolet 6700 FT-IR spectrophotometer. Specific rotations were recorded in 'A KRÜSS Optronic GmbH P8000 polarimeter.

Synthesis of 1-(5-Methyl-1-phenyl-1H-pyrazol-4-yl)-ethanone (3). N,N-Dimethylformamide-dimethyl acetal (DMF-DMA) (17.85 g, 0.15 mol) was added to acetylacetone **1** (0.1 mol) and stirred for 10 min at ambient temperature, followed by the addition of phenylhydrazine derivative **2** (0.1 mol), and the reaction mixture was heated at 70–80 °C for 24

h. The completion of the reaction was monitored by thin-layer chromatography (TLC) (20% EA/n-hexane). The reaction mixture was then allowed to cool and kept in a fridge for 24 h and a white solid precipitated out, which was isolated by simple filtration and washed with diethyl ether to afford pure white product **3** (5 g, 25 mmol, 25% yield). The MLs part was concentrated and purified using a column to afford another 6 g of pure white product **3** (30 mmol, 30%). The overall yield of acetyl-pyrazole-3 (11.0 g, 55%).⁴²

m.p.: 88–90 °C; ¹H NMR (400 MHz, CDCl₃): δ (ppm) = 7.99 (s, 1H, pyrazole-H), 7.52–7.46 (m, 2H, Ar-H), 7.44 (d, J = 6.8 Hz, 1H, Ar-H), 7.41–7.36 (m, 2H, Ar-H), 2.56 (s, 3H, COCH₃), 2.47 (s, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃): δ (ppm) = 193.56 (CO), 143.05, 141.97, 138.58, 129.36, 128.89, 125.61, 121.16, 28.75 (COCH₃), 12.44 (CH₃); IR (KBr, cm⁻¹): 3060, 3001, 1660, 1597, 1545, 1502, 1457, 1399, 1382, 1363, 1276, 1238, 1196, 1171, 1011, 937, 884, 866, 769, 718, 690, 662, 637, 558; [anal. calcd. for C₁₂H₁₂N₂O: C, 71.98; H, 6.04; N, 13.99; found: C, 72.07; H, 6.01; N, 13.94]; LC/MS (ESI, m/z): found 201 [M + H]⁺; exact mass 200.09 for C₁₂H₁₂N₂O. All of the analytical data are in accordance with the reported literature.⁴²













Synthesis of Chalcones (5a–d) from Acetyl-pyrazole (3) and Substituted Indole-3-Carboxyaldehyde (4a–d) (GP1). Compound **3** (1 g, 5 mmol) and substituted indole-3-carboxyaldehyde **4a–d** (5 mmol) were dissolved in ethanol (20 mL) in a 100 mL round bottom flask. Piperidine (950 mg, 2 equiv) was added to the reaction mixture and refluxed at 80 °C 48–72 h. The completion of the reaction was monitored by TLC (30% EA/n-hexane). Then, the solid precipitated out, which was isolated by simple filtration and washed with ethanol to afford pure white/yellow product **5a–d** (70–80% yield).

Synthesis of (E)-3-(1H-Indol-3-yl)-1-(5-methyl-1-phenyl-1H-pyrazol-4-yl)prop-2-en-1-one (5a). Following the general procedure (GP1), acetyl-pyrazole **3** (1.0 g, 5.0 mmol) and indole-3-carboxyaldehyde **4a** (0.87 g, 6.0 mmol) produce pyrazolenone-**5a** (yield 0.9 g, 55%); m.p.: 222–224; ¹H NMR (400 MHz, DMSO-d₆): δ (ppm) = δ 11.85 (s, 1H, NH), 8.59 (s, 1H, Ar-H), 8.17–8.11 (m, 1H, Ar-H), 8.08 (s, 1H, Ar-H), 8.00 (d, J = 15.4 Hz, 1H, CH=CH), 7.58 (d, J = 4.3 Hz, 4H, Ar-H), 7.54–7.41 (m, 3H, Ar-H & CH=CH), 7.29–7.17 (m, 2H, Ar-H), 2.62 (s, 3H, CH₃); ¹³C NMR (100 MHz, DMSO-d₆): δ (ppm) = δ 184.19 (CO), 142.69, 141.53, 138.53, 137.47,

Table 2. AChE Inhibitory Activity of the Synthesized Spirooxindole Analogues Engrafted with Indole and Pyrazole Scaffolds 8a–y

#	Chemical Structure	IC ₅₀ (μM)	#	Chemical Structure	IC ₅₀ (μM)
8a		57.1	8n		>100
8b		>100	8o		75.9
8c		41.2	8p		78.4
8d		36.5	8q		91.2
8e		>100	8r		>100
8f		33.4	8s		65.5
8g		>100	8t		>100

Table 2. continued

8h		31.3	8u		>100
8i		24.1	8v		>100
8j		36.8	8w		47.7
8k		>100	8x		39.5
8l		>100	8y		27.8
8m		80.3	STD		1.2

136.61, 132.48, 129.30, 128.56, 125.41, 125.20, 122.58, 121.61, 121.00, 120.51, 118.15, 112.55, 112.33, 12.20 ($\underline{\text{CH}_3}$); IR (KBr, cm^{-1}) $\nu_{\text{max}} = 3158, 3106, 3047, 2877, 1649, 1638, 1581, 1521, 1502, 1453, 1384, 1347, 1279, 1255, 1229, 1217, 1190, 1140, 1115, 1072, 1064, 1036, 1034, 1004, 974, 945, 893, 879, 852, 834, 762, 734, 714, 696, 656, 641, 603, 593, 558, 507; LC/MS (ESI, m/z): 328.2 [$\text{M} + \text{H}]^+$, exact mass 327.14 for $\text{C}_{21}\text{H}_{17}\text{N}_3\text{O}$.$

Synthesis of (E)-1-(5-Methyl-1-phenyl-1*H*-pyrazol-4-yl)-3-(1-methyl-1*H*-indol-3-yl)prop-2-en-1-one (5b). Following the general procedure (GP1), acetyl-pyrazole 3 (1.0 g, 5.0 mmol) and indole-1-methyl-3-carboxyaldehyde 4b (0.95 g, 6.0 mmol) produce pyrazolenone-5b (yield 1.0 g, 56%); m.p.: 182–183; ^1H NMR (400 MHz, DMSO- d_6): δ (ppm) = δ 8.58 (s, 1H, Ar-H), 8.15 (d, $J = 7.8$ Hz, 1H, Ar-H), 8.04 (s, 1H, Ar-H), 7.95

(d, $J = 15.5$ Hz, 1H, CH=CH), 7.58 (d, $J = 4.4$ Hz, 3H, Ar-H), 7.57–7.49 (m, 3H, Ar-H), 7.46 (d, $J = 15.5$ Hz, 1H, CH=CH), 7.33–7.25 (m, 2H, Ar-H), 3.85 (s, 3H, $\underline{\text{CH}_3}$), 2.62 (s, 3H, $\underline{\text{CH}_3}$); ^{13}C NMR (100 MHz, DMSO- d_6): δ (ppm) = δ 184.09 ($\underline{\text{CO}}$), 142.71, 141.53, 141.49, 138.51, 137.93, 135.98, 135.89, 129.30, 128.56, 125.63, 125.40, 122.66, 121.59, 121.26, 120.67, 120.61, 118.17, 118.10, 111.56, 110.75, 110.66, 32.97 ($\underline{\text{CH}_3}$), 12.15 ($\underline{\text{CH}_3}$); IR (KBr, cm^{-1}) $\nu_{\text{max}} = 3108, 3045, 1643, 1569, 1524, 1501, 1472, 1464, 1386, 1373, 1341, 1281, 1259, 1219, 1187, 1178, 1157, 1132, 1075, 1039, 1003, 937, 855, 841, 821, 771, 751, 699, 681, 654, 540; LC/MS (ESI, m/z): found 342.2 [$\text{M} + \text{H}]^+$, exact mass 341.15 for $\text{C}_{22}\text{H}_{19}\text{N}_3\text{O}$.$

Synthesis of (E)-3-(5-Bromo-1*H*-indol-3-yl)-1-(5-methyl-1-phenyl-1*H*-pyrazol-4-yl)prop-2-en-1-one (5c). Following the general procedure (GP1), acetyl-pyrazole 3 (1.0 g, 5.0 mmol)




Figure 4. Donepezil (Red) docked in the active site of AChE (4EY7) and overlaid with cocrystallized ligand (Green) RMSD = 0.39.

Table 3. Binding Energy of Compound Docked in the Binding Site of the hAChE Active Site^a

	compound	total energy	VDW	H-bond
1.	8i	-127.335	-121.335	-6
2.	8y	-126.589	-118.253	-8.336
3.	8h	-123.962	-118.983	-4.979
4.	cocrystallized ligand	-149.55	-139.4	-10.15

^aVDW = Van der Waals force and H-Bond = Hydrogen bond.

and 5-bromoindole-3-carboxyaldehyde **4c** (1.34 g, 6.0 mmol) produce pyrazolenone-**5c** (yield 0.85 g, 42%); m.p.: 214–215; ¹H NMR (400 MHz, DMSO-*d*₆): δ (ppm) = δ 12.02 (s, 1H, NH), 8.63 (s, 1H, Ar-H), 8.26 (d, *J* = 1.9 Hz, 1H, Ar-H), 8.14

(s, 1H, Ar-H), 7.95 (d, *J* = 15.5 Hz, 1H, CH=CH), 7.60–7.55 (m, 4H, Ar-H + CH=CH), 7.53–7.49 (m, 1H, Ar-H), 7.46 (d, *J* = 3.9 Hz, 1H, Ar-H), 7.45 (d, *J* = 3.1 Hz, 1H, Ar-H), 7.35 (dd, *J* = 8.6, 1.9 Hz, 1H, Ar-H), 2.61 (s, 3H, CH₃); ¹³C NMR (100 MHz, DMSO-*d*₆): δ (ppm) = δ 184.22 (C=O), 142.77, 141.76, 138.50, 136.07, 135.67, 132.99, 129.30, 128.56, 127.02, 125.39, 125.18, 122.39, 121.52, 119.03, 114.25, 113.74, 112.17, 12.19 (CH₃); IR (KBr, cm⁻¹) ν_{max} = 3153, 3077, 3030, 2934, 2900, 1643, 1559, 1500, 1454, 1433, 1394, 1370, 1299, 1273, 1243, 1223, 1182, 1136, 1095, 1038, 1022, 1007, 956, 938, 881, 851, 824, 789, 758, 689, 660, 636, 609, 554; LC/MS (ESI, *m/z*): found 406.6 [M-(₇₉Br) + H]⁺, 408.1 [M-(₈₁Br) + H]⁺; exact mass 405.05 for C₂₁H₁₆BrN₃O.




Figure 5. **8i** docked in the active site of AChE (4EY7) and corresponding two-dimensional (2D) presentation. The H-bond is represented by green dotted lines, hydrophobic interactions are represented by magenta dotted lines, and Pi-sulfur interaction is represented by orange dotted lines.




Figure 6. **8y** docked in the active site of AChE (4EY7) and corresponding 2D presentation. The H-bond is represented by green dotted lines, hydrophobic interactions are represented by magenta dotted lines, and Pi-sulfur interaction is represented by orange dotted lines.




Figure 7. **8h** docked in the active site of AChE (4EY7) and corresponding 2D presentation. The H-bond is represented by green dotted lines, hydrophobic interactions are represented by magenta dotted lines, and Pi-sulfur interaction is represented by orange dotted lines.

Synthesis of (*E*)-3-(5-Chloro-1*H*-indol-3-yl)-1-(5-methyl-1-phenyl-1*H*-pyrazol-4-yl) prop-2-en-1-one (5d**).** Following the general procedure (GP1), acetyl-pyrazole **3** (1.0 g, 5.0 mmol) and 5-chloroindole-3-carboxyaldehyde **4d** (1.07 g, 6.0 mmol) produce pyrazolenone-**5d** (yield 0.9 g, 50%); m.p.: 237–238; ¹H NMR (400 MHz, DMSO-*d*₆): δ (ppm) = δ 11.95 (bs, 1H, NH), 8.65 (s, 1H, Ar-H), 8.20–8.05 (m, 2H, Ar-H), 7.95 (d, *J* = 15.4 Hz, 1H, CH=CH), 7.57 (d, *J* = 4.4 Hz, 4H, Ar-H), 7.50 (dd, *J* = 8.7, 4.6 Hz, 2H, Ar-H), 7.46 (d, *J* = 15.7 Hz, 1H, CH=CH), 7.23 (dd, *J* = 8.2, 2.1 Hz, 1H, Ar-H), 2.61 (s, 3H, CH₃); ¹³C NMR (100 MHz, DMSO-*d*₆): δ (ppm) = δ 184.24 (C=O), 142.78, 141.80, 138.52, 135.86, 135.76, 133.34, 129.30, 128.56, 126.35, 125.73, 125.40, 122.60, 121.53, 119.55, 118.92, 113.81, 112.30, 12.19 (CH₃); IR (KBr, cm⁻¹) ν_{max} = 3107, 2896, 1648, 1637, 1576, 1518, 1500, 1453, 1395, 1383, 1373, 1314, 1251, 1212, 1142, 1123, 1072, 1047, 1005, 973, 944, 897, 863, 837, 794, 765, 740, 697, 651, 643, 609, 594, 575;

LC/MS (ESI, *m/z*): found 362.1 [M₍₃₅Cl) + H]⁺, 364.1 [M₍₃₇Cl) + H]⁺; exact mass 361.09 for C₂₁H₁₆ClN₃O.

Synthesis of Spirooxindole Derivatives **8a–y.** General Procedure (GP2). Chalcones **5a–d** (0.25 mmol), isatin derivatives **6a–c** (0.25 mmol), and amino acids **7a–d** (1.5 equiv, 0.37 mmol) were dissolved in methanol (20 mL), and the reaction mixture was refluxed for 2–4 h. Finally, the products were isolated by flash column chromatography, using 1–3% MeOH/DCM to afford pyrazole spirooxindole **8a–y**.

(*3R,6'S,7'R,7a'S*)-7'-(1*H*-Indol-3-yl)-6'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-1',6',7',7a'-tetrahydro-3'*H*-spiro[indoline-3,5'-pyrrolo[1,2-*c*]thiazol]-2-one (**8a**). Following the general procedure (GP2), chalcone **5a** (82 mg, 0.25 mmol), isatin **6a** (37 mg, 0.25 mmol), and thiopropine **7a** (50 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield light yellow solid compound **8a**; yield (85 mg, 62%); m.p.: 138–139 °C; [α]_D²⁵ =

-16.19° (c 0.13, MeOH); ^1H NMR (700 MHz, CDCl_3) δ (ppm) = 8.59 (s, 1H, NH), 8.33 (s, 1H, NH), 8.10 (d, J = 7.1 Hz, 1H, Ar-H), 7.92 (s, 1H, Ar-H), 7.72 (d, J = 7.6 Hz, 1H, Ar-H), 7.40–7.33 (m, 4H, Ar-H), 7.28 (d, J = 2.5 Hz, 1H, Ar-H), 7.20–7.17 (m, 2H, Ar-H), 7.17–7.14 (m, 3H, Ar-H), 7.04–7.01 (m, 1H, Ar-H), 6.64 (d, J = 7.7 Hz, 1H, Ar-H), 4.79 (d, J = 11.8 Hz, 1H, CH_{CO}), 4.66–4.61 (m, 1H, NCH), 4.25 (dd, J = 11.8, 9.7 Hz, 1H, NCHCH), 3.94 (d, J = 10.5 Hz, 1H, NCH₂), 3.61 (d, J = 10.5 Hz, 1H, NCH₂), 3.11–3.03 (m, 2H, SCH₂), 1.92 (s, 3H, CH₃); ^{13}C NMR (176 MHz, CDCl_3) δ (ppm) = 191.09 (CO), 180.75 (CO), 143.50, 141.28, 141.03, 138.31, 136.74, 129.76, 129.29, 129.15, 128.88, 126.46, 125.47, 124.08, 123.02, 122.28, 122.27, 121.00, 119.93, 119.84, 113.65, 111.58, 109.65, 75.12, 73.75, 63.86, 55.16, 43.58, 37.41, 11.75 (CH₃); IR (KBr, cm^{-1}) ν_{max} = 3288, 3058, 2921, 1733, 1722, 1717, 1699, 1694, 1682, 1674, 1668, 1661, 1652, 1619, 1615, 1597, 1538, 1532, 1504, 1470, 1456, 1393, 1337, 1222, 1119, 935, 875, 807, 749, 695; [anal. calcd. for $\text{C}_{32}\text{H}_{27}\text{N}_5\text{O}_2\text{S}$: C, 70.44; H, 4.99; N, 12.83; found: C, 70.31; H, 5.05; N, 12.91]; LC/MS (ESI, m/z): found 546.4 [M + H]⁺, exact mass 545.19 for $\text{C}_{32}\text{H}_{27}\text{N}_5\text{O}_2\text{S}$.

(3*R*,6'*S*,7'*R*,7*a'**S*)-6'-(5-Methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-7'-(1-methyl-1*H*-indol-3-yl)-1',6',7',7*a'*-tetrahydro-3'H-spiro[indoline-3,5'-pyrrolo[1,2-c]thiazol]-2-one (**8b**). Following the general procedure (GP2), chalcone **5b** (85 mg, 0.25 mmol), isatin **6a** (37 mg, 0.25 mmol), and thioproline **7a** (50 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield light yellow solid compound **8b**; yield (101 mg, 72%); m.p.: 186–187 °C; $[\alpha]_D^{25} = -12.53^\circ$ (c 0.13, MeOH); ^1H NMR (700 MHz, CDCl_3) δ (ppm) = 8.80 (s, 1H, NH), 8.09 (d, J = 7.24 Hz, 1H, Ar-H), 7.92 (s, 1H, Ar-H), 7.72 (d, J = 7.63 Hz, 1H, Ar-H), 7.39–7.35 (m, 3H, Ar-H), 7.29 (ddd, J = 8.1, 1.4, 0.7 Hz, 1H, Ar-H), 7.25 (dd, J = 6.9, 1.3 Hz, 1H, Ar-H), 7.21 (dd, J = 5.0, 1.4 Hz, 1H, Ar-H), 7.19 (s, 1H, Ar-H), 7.19–7.16 (m, 2H, Ar-H), 7.16–7.12 (m, 2H, Ar-H), 7.03 (td, J = 7.6, 1.1 Hz, 1H, Ar-H), 6.65 (d, J = 7.73 Hz, 1H, Ar-H), 4.77 (d, J = 11.8 Hz, 1H, CH_{CO}), 4.67–4.61 (m, 1H, NCH), 4.22 (dd, J = 11.8, 9.8 Hz, 1H, NCHCH), 3.94 (d, J = 10.6 Hz, 1H, NCH₂), 3.74 (s, 3H, CH₃), 3.61 (d, J = 10.5 Hz, 1H, NCH₂), 3.07 (d, J = 4.3 Hz, 2H, SCH₂), 1.92 (s, 3H, CH₃); ^{13}C NMR (176 MHz, CDCl_3) δ (ppm) = 191.09 (CO), 180.96 (CO), 143.43, 141.25, 141.21, 138.27, 137.40, 129.70, 129.23, 129.03, 128.81, 127.60, 126.88, 125.45, 124.02, 122.13, 121.80, 120.97, 119.90, 119.38, 112.03, 109.75, 109.58, 75.20, 73.86, 63.89, 55.20, 43.45, 37.35, 32.80 (CH₃), 11.67 (CH₃); IR (KBr, cm^{-1}) ν_{max} = 3237, 2923, 1738, 1733, 1722, 1699, 1694, 1682, 1674, 1668, 1661, 1652, 1645, 1634, 1622, 1615, 1597, 1557, 1538, 1532, 1505, 1470, 1456, 1398, 1329, 1229, 1180, 1156, 1115, 1069, 1013, 934, 801, 741, 695; [anal. calcd. for $\text{C}_{33}\text{H}_{29}\text{N}_5\text{O}_2\text{S}$: C, 70.82; H, 5.22; N, 12.51; found: C, 71.01; H, 5.15; N, 12.39]; LC/MS (ESI, m/z): found 560.4 [M + H]⁺, exact mass 559.14 for $\text{C}_{33}\text{H}_{29}\text{N}_5\text{O}_2\text{S}$.

(3*R*,6'*S*,7'*R*,7*a'**S*)-7'-(5-Bromo-1*H*-indol-3-yl)-6'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-1',6',7',7*a'*-tetrahydro-3'H-spiro[indoline-3,5'-pyrrolo[1,2-c]thiazol]-2-one (**8c**). Following the general procedure (GP2), chalcone **5c** (102 mg, 0.25 mmol), isatin **6a** (37 mg, 0.25 mmol), and thioproline **7a** (50 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield light yellow solid compound **8c**; yield (59 mg, 38%); m.p.: 180–181 °C; $[\alpha]_D^{25} =$

-12.53° (c 0.11, MeOH); ^1H NMR (700 MHz, CDCl_3) δ (ppm) = 9.05 (s, 1H, NH), 8.66 (s, 1H, NH), 8.18 (s, 1H, Ar-H), 7.98 (s, 1H, Ar-H), 7.69 (d, J = 7.5 Hz, 1H, Ar-H), 7.38–7.35 (m, 4H, Ar-H), 7.20–7.18 (m, 1H, Ar-H), 7.15–7.13 (m, 2H, Ar-H), 7.03 (d, J = 7.6 Hz, 1H, Ar-H), 6.62 (d, J = 7.7 Hz, 1H, Ar-H), 4.63 (d, J = 11.6 Hz, 1H, CH_{CO}), 4.50–4.49 (ddd, J = 9.4, 6.2, 2.5 Hz, 1H, NCH), 4.21 (m, 1H, NCHCH), 3.92 (d, J = 10.4 Hz, 1H, NCH₂), 3.60 (d, J = 10.4 Hz, 1H, NCH₂), 3.08–3.04 (m, 1H, SCH₂), 3.03–3.98 (m, 1H, SCH₂), 1.91 (s, 3H, CH₃); ^{13}C NMR (176 MHz, CDCl_3) δ (ppm) = 191.02 (CO), 180.91 (CO), 143.59, 141.14, 138.23, 135.18, 129.84, 129.33, 128.96, 128.58, 127.75, 127.12, 125.47, 125.19, 124.00, 123.78, 122.33, 122.04, 120.91, 113.74, 113.18, 113.03, 109.80, 75.06, 74.25, 65.41, 54.90, 42.63, 37.28, 11.70 (CH₃); IR (KBr, cm^{-1}) ν_{max} = 3273, 2924, 1721, 1668, 1617, 1597, 1537, 1503, 1470, 1455, 1396, 1331, 1283, 1223, 1179, 1099, 934, 884, 807, 795, 762, 752, 694, 656, 604; [anal. calcd. for $\text{C}_{32}\text{H}_{26}\text{BrN}_5\text{O}_2\text{S}$: C, 61.54; H, 4.20; N, 11.21; found: C, 61.43; H, 4.35; N, 11.27]; LC/MS (ESI, m/z): found 624.7 [M(₇₉Br) + H]⁺, 626.1 [M(₈₁Br) + H]⁺; exact mass 623.10 for $\text{C}_{32}\text{H}_{26}\text{BrN}_5\text{O}_2\text{S}$.

(3*R*,6'*S*,7'*R*,7*a'**S*)-7'-(5-Chloro-1*H*-indol-3-yl)-6'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-1',6',7',7*a'*-tetrahydro-3'H-spiro[indoline-3,5'-pyrrolo[1,2-c]thiazol]-2-one (**8d**). Following the general procedure (GP2), chalcone **5d** (91 mg, 0.25 mmol), isatin **6a** (37 mg, 0.25 mmol), and thioproline **7a** (50 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield light yellow solid compound **8d**; yield (76 mg, 52%); m.p.: 210–212 °C; $[\alpha]_D^{25} = -12.03^\circ$ (c 0.11, MeOH); ^1H NMR (700 MHz, CDCl_3) δ (ppm) = 8.61 (s, 1H, NH), 8.38 (s, 1H, NH), 8.00 (s, 1H, Ar-H), 7.86 (s, 1H, Ar-H), 7.64 (d, J = 7.6 Hz, 1H, Ar-H), 7.33 (dd, J = 10.7, 7.0 Hz, 3H, Ar-H), 7.23–7.18 (m, 2H, Ar-H), 7.09 (dd, J = 15.4, 8.3 Hz, 4H, Ar-H), 7.00–6.96 (m, 1H, Ar-H), 6.60 (d, J = 7.8 Hz, 1H, Ar-H), 4.58 (d, J = 11.6 Hz, 1H, CH_{CO}), 4.50–4.42 (m, 1H, NCH), 4.16 (t, J = 10.6 Hz, 1H, NCHCH), 3.88 (d, J = 10.4 Hz, 1H, NCH₂), 3.55 (d, J = 10.4 Hz, 1H, NCH₂), 3.06–2.94 (m, 2H, SCH₂), 1.86 (s, 3H, CH₃); ^{13}C NMR (176 MHz, CDCl_3) δ (ppm) = 190.95 (CO), 180.78 (CO), 143.60, 141.17, 141.01, 138.27, 134.95, 129.88, 129.35, 129.06, 128.97, 127.87, 125.71, 125.50, 124.00, 122.75, 122.40, 120.91, 119.13, 113.90, 112.54, 109.76, 74.12, 64.53, 54.95, 42.77, 37.30, 11.72 (CH₃); IR (KBr, cm^{-1}) ν_{max} = 3284, 2926, 1720, 1668, 1652, 1616, 1597, 1538, 1504, 1470, 1397, 1329, 1283, 1268, 1223, 1180, 1103, 934, 893, 806, 796, 763, 752, 694, 683, 658, 605; [anal. calcd. for $\text{C}_{32}\text{H}_{26}\text{ClN}_5\text{O}_2\text{S}$: C, 66.26; H, 4.52; N, 12.07; found: C, 66.14; H, 4.63; N, 12.15]; LC/MS (ESI, m/z): found 580.6 [M(₃₅Cl) + H]⁺, 582.3 [M(₃₇Cl) + H]⁺; exact mass 579.15 for $\text{C}_{32}\text{H}_{26}\text{ClN}_5\text{O}_2\text{S}$.

(3*R*,6'*S*,7'*R*,7*a'**S*)-6-Chloro-7'-(1*H*-indol-3-yl)-6'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-3',6',7',7*a'*-tetrahydro-1*H*-spiro[indoline-3,5'-pyrrolo[1,2-c]thiazol]-2-one (**8e**). Following the general procedure (GP2), chalcone **5a** (82 mg, 0.25 mmol), 6-chloroisatin **6b** (46 mg, 0.25 mmol), and thioproline **7a** (50 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield light yellow solid compound **8e**; yield (44 mg, 30%); m.p.: 188–190 °C; $[\alpha]_D^{25} = -18.78^\circ$ (c 0.11, MeOH); ^1H NMR (700 MHz, DMSO-*d*₆): δ (ppm) = 11.00 (s, 1H, NH), 10.68 (s, 1H, NH), 7.87 (d, J = 7.8 Hz, 1H, Ar-H), 7.84 (d, J = 2.5 Hz, 1H, Ar-H), 7.54 (d, J = 8.2 Hz, 1H, Ar-H), 7.51–7.45 (m, 4H, Ar-

H), 7.38–7.32 (m, 3H, Ar-H), 7.11–7.06 (m, 2H, Ar-H), 7.04 (d, J = 8.2 Hz, 1H, Ar-H), 6.67 (s, 1H, Ar-H), 4.73 (d, J = 11.8 Hz, 1H, CHCO), 4.33–4.29 (m, 1H, NCH), 4.13–4.09 (m, 1H, NCHCH), 3.78 (d, J = 10.3 Hz, 1H, NCH_{2(a)}), 3.43 (d, J = 8.2 Hz, 1H, NCH_{2(b)}), 3.07–2.99 (m, 2H, SCH₂), 1.92 (s, 3H, CH₃); ¹³C NMR (176 MHz, DMSO-*d*₆) δ (ppm) = 190.34 (CO), 178.87 (CO), 143.68, 142.64, 140.80, 137.91, 136.59, 133.90, 129.93, 129.31, 128.87, 128.07, 126.65, 126.45, 126.21, 125.26, 123.43, 122.45, 121.19, 120.54, 118.77, 112.02, 109.44, 73.87, 73.77, 62.86, 54.93, 53.92, 43.19, 36.70, 11.22 (CH₃); IR (KBr, cm⁻¹) ν _{max} = 3295, 2928, 1723, 1717, 1699, 1668, 1645, 1615, 1598, 1538, 1533, 1504, 1483, 1456, 1398, 1379, 1338, 1325, 1282, 1244, 1224, 1183, 1125, 1096, 1072, 926, 852, 811, 764, 743, 694, 658, 594, 529; [anal. calcd. for C₃₂H₂₆ClN₅O₂S: C, 66.26; H, 4.52; N, 12.07; found: C, 66.12; H, 4.67; N, 12.22]; LC/MS (ESI, *m/z*): found 580.5 [M(₃₅Cl) + H]⁺, 582.3 [M(₃₅Cl) + H]⁺; exact mass 579.15 for C₃₂H₂₆ClN₅O₂S.

(3*R*,6'*S*,7'R,7a'S)-7'-(1H-Indol-3-yl)-6'-(5-methyl-1-phenyl-1H-pyrazole-4-carbonyl)-5-nitro-1',6',7',7a'-tetrahydro-3'H-spiro[indoline-3,5'-pyrrolo[1,2-c]thiazol]-2-one (**8f**). Following the general procedure (GP2), chalcone **5a** (82 mg, 0.25 mmol), 5-nitroisatin **6c** (48 mg, 0.25 mmol), and thioproline **7a** (50 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield yellow solid compound **8f**; yield (49 mg, 33%); m.p.: 213–214 °C; $[\alpha]_D^{25} = -36.24^\circ$ (c 0.15, MeOH); ¹H NMR (700 MHz, DMSO-*d*₆) δ (ppm) = 11.26 (s, 1H, NH), 11.04 (s, 1H, NH), 8.46 (s, 1H, Ar-H), 8.17 (d, J = 8.2 Hz, 1H, Ar-H), 7.89–7.84 (m, 2H, Ar-H), 7.55 (s, 1H, Ar-H), 7.50–7.47 (m, 2H, Ar-H), 7.46 (d, J = 7.5 Hz, 1H, Ar-H), 7.36 (d, J = 6.9 Hz, 1H, Ar-H), 7.32–7.30 (m, 2H, Ar-H), 7.09–7.10 (m, 2H, Ar-H), 6.88 (d, J = 8.6 Hz, 1H, Ar-H), 4.79 (d, J = 12.0 Hz, 1H, CHCO), 4.41–4.36 (m, 1H, NCH), 4.14–4.09 (m, 1H, NCHCH), 3.80 (d, J = 10.8 Hz, 1H, NCH_{2(a)}), 3.50 (d, J = 10.8 Hz, 1H, NCH_{2(b)}), 3.08 (t, J = 3.6 Hz, 2H, SCH₂), 1.86 (s, 3H, CH₃); ¹³C NMR (176 MHz, DMSO-*d*₆) δ (ppm) = 190.24 (CO), 179.44 (CO), 148.63, 142.79, 141.34, 140.76, 137.81, 136.64, 129.34, 128.93, 128.08, 126.92, 126.46, 126.07, 125.26, 124.34, 123.83, 121.25, 120.47, 118.89, 118.54, 111.42, 109.90, 74.15, 73.58, 62.93, 54.48, 43.59, 36.77, 11.25 (CH₃); IR (KBr, cm⁻¹) ν _{max} = 3252, 2924, 2853, 1736, 1729, 1665, 1652, 1626, 1598, 1526, 1504, 1478, 1455, 1398, 1338, 1300, 1248, 1223, 1198, 1180, 1124, 1098, 1069, 932, 907, 828, 807, 763, 743, 694, 557; [anal. calcd. for C₃₂H₂₆N₆O₄S: C, 65.07; H, 4.44; N, 14.23; found: C, 64.91; H, 4.56; N, 14.04]; LC/MS (ESI, *m/z*): found 591.7 [M + H]⁺, exact mass 590.17 for C₃₂H₂₆N₆O₄S.

(3*R*,6'*S*,7'R,7a'S)-6-Chloro-6'-(5-methyl-1-phenyl-1H-pyrazole-4-carbonyl)-7'-(1-methyl-1H-indol-3-yl)-1',6',7',7a'-tetrahydro-3'H-spiro[indoline-3,5'-pyrrolo[1,2-c]thiazol]-2-one (**8g**). Following the general procedure (GP2), chalcone **5b** (85 mg, 0.25 mmol), 6-chloroisatin **6b** (46 mg, 0.25 mmol), and thioproline **7a** (50 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield light yellow solid compound **8g**; yield (55 mg, 37%); m.p.: 164–165 °C; $[\alpha]_D^{25} = -41.94^\circ$ (c 0.12, MeOH); ¹H NMR (700 MHz, CDCl₃) δ (ppm) = 9.23 (s, 1H, NH), 8.07 (d, J = 7.9 Hz, 1H, Ar-H), 7.94 (s, 1H, Ar-H), 7.64 (d, J = 8.1 Hz, 1H, Ar-H), 7.40 (d, J = 6.2 Hz, 2H, Ar-H), 7.36 (s, 1H, Ar-H), 7.29 (d, J = 8.2 Hz, 1H, Ar-H), 7.25–7.22 (m, 1H, Ar-H), 7.21–7.18 (m, 2H, Ar-H), 7.16 (d, J = 8.1 Hz, 2H, Ar-H), 7.00 (d, J

= 8.1 Hz, 1H, Ar-H), 6.51 (s, 1H, Ar-H), 4.75 (d, J = 11.8 Hz, 1H, CHCO), 4.63–4.60 (m, 1H, NCH), 4.21–4.15 (m, 1H, NCHCH), 3.93 (d, J = 10.6 Hz, 1H, NCH_{2(a)}), 3.74 (s, 3H, NCH₃), 3.56 (d, J = 10.7 Hz, 1H, NCH_{2(b)}), 3.09–3.02 (m, 2H, SCH₂), 1.99 (s, 3H, CH₃); ¹³C NMR (176 MHz, CDCl₃) δ (ppm) = 190.90 (CO), 180.77 (CO), 143.66, 141.19, 138.14, 137.42, 135.36, 130.05, 129.39, 129.11, 128.65, 127.59, 127.10, 126.86, 125.53, 122.49, 121.90, 120.90, 119.84, 119.50, 111.84, 110.17, 109.63, 74.78, 73.80, 63.95, 55.17, 43.58, 37.44, 32.85 (NCH₃), 11.71 (CH₃); IR (KBr, cm⁻¹) ν _{max} = 3234, 2922, 2849, 1733, 1717, 1699, 1667, 1652, 1612, 1598, 1544, 1537, 1532, 1503, 1483, 1454, 1401, 1377, 1327, 1261, 1223, 1182, 157, 1123, 1071, 1012, 925, 808, 765, 738, 694, 612, 529; [anal. calcd. for C₃₃H₂₈ClN₅O₂S: C, 66.71; H, 4.75; N, 11.79; found: C, 66.85; H, 4.62; N, 12.03]; LC/MS (ESI, *m/z*): found 594.5 [M(₃₅Cl) + H]⁺, 596.0 [M(₃₇Cl) + H]⁺; exact mass 593.17 for C₃₃H₂₈ClN₅O₂S.

(3*R*,6'*S*,7'R,7a'S)-6'-5-Methyl-1-phenyl-1H-pyrazole-4-carbonyl)-7'-(1-methyl-1H-indol-3-yl)-5-nitro-3',6',7',7a'-tetrahydro-1'H-spiro[indoline-3,5'-pyrrolo[1,2-c]thiazol]-2-one (**8h**). Following the general procedure (GP2), chalcone **5b** (85 mg, 0.25 mmol), 5-nitroisatin **6c** (48 mg, 0.25 mmol), and thioproline **7a** (50 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 16 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield yellow solid compound **8h**; yield (41 mg, 27%); m.p.: 172–173 °C; $[\alpha]_D^{25} = -22.87^\circ$ (c 0.13, MeOH); ¹H NMR (700 MHz, CDCl₃) δ (ppm) = 9.78 (s, 1H, NH), 8.73 (s, 1H, Ar-H), 8.14 (dd, J = 27.8, 8.4 Hz, 2H, Ar-H), 7.99 (s, 1H, Ar-H), 7.47–7.41 (m, 3H, Ar-H), 7.40 (d, J = 4.3 Hz, 1H, Ar-H), 7.35 (d, J = 8.2 Hz, 1H, Ar-H), 7.30 (d, J = 8.2 Hz, 1H, Ar-H), 7.23–7.19 (m, 1H, Ar-H), 7.16 (dd, J = 6.7, 3.1 Hz, 2H, Ar-H), 6.69 (d, J = 9.2 Hz, 1H, Ar-H), 4.89 (d, J = 11.5 Hz, 1H, CHCO), 4.73 (m, 1H NCH), 4.30 (t, J = 10.7 Hz, 1H, NCHCH), 3.99 (d, J = 10.4 Hz, 1H, NCH_{2(a)}), 3.79 (s, 3H, CH₃), 3.51 (d, J = 10.5 Hz, 1H, NCH_{2(b)}), 3.16 (d, J = 4.4 Hz, 2H, SCH₂), 2.01 (s, 3H, CH₃); ¹³C NMR (176 MHz, CDCl₃) δ (ppm) = 190.04 (CO), 180.99 (CO), 146.95, 143.91, 142.89, 141.14, 137.92, 137.52, 129.47, 129.26, 128.66, 128.04, 127.74, 127.12, 126.55, 125.38, 125.06, 124.92, 121.96, 120.90, 119.75, 111.02, 109.77, 73.37, 63.71, 55.11, 44.29, 37.47, 32.86 (CH₃), 11.86 (CH₃); IR (KBr, cm⁻¹) ν _{max} = 3208, 2926, 2859, 1736, 1716, 1699, 1682, 1678, 1668, 1652, 1622, 1615, 1598, 1524, 1504, 1475, 1455, 1404, 1337, 1221, 1177, 1123, 1103, 932, 833, 805, 765, 741, 694, 556; [anal. calcd. for C₃₃H₂₈N₆O₄S: C, 65.55; H, 4.67; N, 13.90; found: C, 65.67; H, 4.81; N, 14.04]; LC/MS (ESI, *m/z*): found 605.6 [M + H]⁺, exact mass 604.19 for C₃₃H₂₈N₆O₄S.

(3*R*,6'*S*,7'R,7a'S)-7'-(5-Bromo-1H-indol-3-yl)-6'-(5-methyl-1-phenyl-1H-pyrazole-4-carbonyl)-5-nitro-3',6',7',7a'-tetrahydro-1'H-spiro[indoline-3,5'-pyrrolo[1,2-c]thiazol]-2-one (**8i**). Following the general procedure (GP2), chalcone **5c** (102 mg, 0.25 mmol), 5-nitroisatin **6c** (48 mg, 0.25 mmol), and thioproline **7a** (50 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 16 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield yellow solid compound **8i**; yield (82 mg, 49%); m.p.: 250–251 °C; $[\alpha]_D^{25} = -126.29^\circ$ (c 0.19, MeOH); ¹H NMR (700 MHz, DMSO-*d*₆) δ (ppm) = 11.30 (d, J = 2.6 Hz, 1H, NH), 11.25 (s, 1H, NH), 8.45 (d, J = 2.4 Hz, 1H, Ar-H), 8.17 (dd, J = 8.6, 2.4 Hz, 1H, Ar-H), 7.98 (d, J = 1.9 Hz, 1H, Ar-H), 7.84 (s, 1H, Ar-H), 7.67 (d, J = 2.6 Hz, 1H, Ar-H), 7.51–7.48 (m, 2H, Ar-H), 7.48–7.45 (m, 1H, Ar-H), 7.35–7.31 (m, 3H, Ar-H), 7.21

(dd, $J = 8.5, 1.9$ Hz, 1H, Ar-H), 6.87 (d, $J = 8.6$ Hz, 1H, Ar-H), 4.67 (d, $J = 11.9$ Hz, 1H, CHCO), 4.37–4.33 (m, 1H, NCH), 4.10 (dd, $J = 11.8, 9.5$ Hz, 1H, NCHCH), 3.79 (d, $J = 10.7$ Hz, 1H, NCH_{2(a)}), 3.50 (d, $J = 10.6$ Hz, 1H, NCH_{2(b)}), 3.09–3.04 (m, 2H, SCH₂), 1.86 (s, 3H, CH₃); ¹³C NMR (176 MHz, DMSO-*d*₆) δ (ppm) = 190.22 (CO), 179.25 (CO), 148.59, 142.75, 141.34, 140.74, 137.77, 135.16, 129.31, 128.90, 128.07, 126.90, 125.42, 125.21, 124.29, 123.85, 123.69, 120.73, 120.38, 113.84, 111.50, 111.42, 109.86, 73.95, 73.48, 63.46, 54.14, 42.83, 36.56, 11.21 (CH₃); IR (KBr, cm⁻¹) ν _{max} = 3383, 3101, 2855, 1747, 1729, 1649, 1622, 1598, 1530, 1504, 1463, 1454, 1412, 1337, 1290, 1253, 1222, 1199, 1173, 1097, 933, 880, 830, 799, 753, 693, 607; [anal. calcd. for C₃₂H₂₅BrN₆O₄S: C, 57.40; H, 3.76; N, 12.55; found: C, 57.36; H, 3.84; N, 12.59]; LC/MS (ESI, *m/z*): found 669.6 [M-(₇₉Br) + H]⁺, 671.5 [M-(₈₁Br) + H]⁺; exact mass 668.08 for C₃₂H₂₅BrN₆O₄S.

(*R*,*S*,*S*,*R*,*S*,*R*)-7'-(5-Chloro-1*H*-indol-3-yl)-6'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-5-nitro-3',6',7',7*a*'-tetrahydro-1*H*-spiro[indoline-3,5'-pyrrolo[1,2-*c*]thiazol]-2-one (**8j**). Following the general procedure (GP2), chalcone **5d** (91 mg, 0.25 mmol), 5-nitroisatin **6c** (48 mg, 0.25 mmol), and thioproline **7a** (50 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield yellow solid compound **8j**; yield (53 mg, 40%); m.p.: 196–197 °C; $[\alpha]_D^{25} = -53.73^\circ$ (c 0.10, MeOH); ¹H NMR (700 MHz, DMSO-*d*₆) δ (ppm) = 11.30 (s, 1H, NH), 11.27 (s, 1H, NH), 8.46 (s, 1H, Ar-H), 8.18 (d, $J = 8.6$ Hz, 1H, Ar-H), 7.85 (s, 1H, Ar-H), 7.69 (s, 1H, Ar-H), 7.53–7.46 (m, 2H, Ar-H), 7.48 (d, $J = 5.5$ Hz, 1H, Ar-H), 7.39 (d, $J = 8.6$ Hz, 1H, Ar-H), 7.34–7.31 (m, 3H, Ar-H), 7.12 (d, $J = 8.6$ Hz, 1H, Ar-H), 6.88 (d, $J = 8.7$ Hz, 1H, Ar-H), 4.68 (d, $J = 11.7$ Hz, 1H, CHCO), 4.39–4.35 (m, 1H, NCH), 4.12 (t, $J = 12.1$ Hz, 1H, NCHCH), 3.80 (d, $J = 10.9$ Hz, 1H, NCH_{2(a)}), 3.51 (d, $J = 10.9$ Hz, 1H, NCH_{2(b)}), 3.11–3.05 (m, 2H, SCH₂), 1.87 (s, 3H, CH₃); ¹³C NMR (176 MHz, DMSO-*d*₆) δ (ppm) = 190.27 (CO), 179.33 (CO), 148.63, 142.79, 141.39, 140.77, 137.80, 134.98, 129.35, 128.94, 127.38, 126.94, 126.66, 126.45, 125.62, 124.33, 123.50, 121.22, 120.41, 117.74, 113.42, 111.62, 109.90, 74.00, 73.49, 62.93, 54.94, 42.87, 36.62, 11.23 (CH₃); IR (KBr, cm⁻¹) ν _{max} = 3344, 3270, 2932, 2861, 1725, 1657, 1623, 1598, 1530, 1503, 1477, 1454, 1339, 1296, 1224, 1180, 1101, 931, 892, 797, 764, 693, 614, 553; [Anal. Calcd. for C₃₂H₂₅ClN₆O₄S: C, 61.49; H, 4.03; N, 13.44; found: C, 61.35; H, 3.91; N, 13.62]; LC/MS (ESI, *m/z*): found 625.8 [M-(₃₅Cl) + H]⁺, 627.2 [M-(₃₇Cl) + H]⁺; exact mass 624.13 for C₃₂H₂₅ClN₆O₄S.

(*R*,*S*,*S*,*R*,*S*,*R*)-1'-(1*H*-Indol-3-yl)-2'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-1',2',4*a*',5',6',7',8',8*a*',9',9*a*'-decahydrospiro[indoline-3,3'-pyrrolo[1,2-*a*]indol]-2-one (**8k**). Following the general procedure (GP2), chalcone **5a** (82 mg, 0.25 mmol), isatin **6a** (37 mg, 0.25 mmol), and octahydro-1*H*-indole-2-carboxylic acid **7b** (64 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield light yellow solid compound **8k**; yield (125 mg, 86%); m.p.: 200–201 °C; $[\alpha]_D^{25} = -18.76^\circ$ (c 0.12, MeOH); ¹H NMR (700 MHz, CDCl₃) δ (ppm) = 8.33 (s, 1H, NH), 8.04 (d, $J = 1.2$ Hz, 1H, Ar-H), 8.04–8.02 (m, 1H, Ar-H), 7.95 (s, 1H, NH), 7.42 (d, $J = 7.4$ Hz, 1H, Ar-H), 7.40–7.34 (m, 3H, Ar-H), 7.33–7.29 (m, 1H, Ar-H), 7.20–7.14 (m, 5H, Ar-H), 7.13–7.09 (m, 1H, Ar-H), 7.06–7.02 (m, 1H, Ar-H), 6.57 (d, $J = 7.7$ Hz, 1H, Ar-H), 4.96 (d, $J = 11.9$ Hz, 1H, CHCO), 4.57–4.52 (m, 1H, NCH), 4.20 (dd, $J = 12.0, 10.0$

Hz, 1H, NCHCH), 3.27 (q, $J = 3.8$ Hz, 1H, NCH), 2.19–2.14 (m, 1H, NCHCH), 1.93 (s, 3H, CH₃), 1.85–1.78 (m, 2H, CH₂), 1.59–1.44 (m, 4H, CH₂), 1.20–1.15 (m, 1H, CH₂), 1.07–1.01 (m, 1H, CH₂), 1.01–0.95 (m, 2H, CH₂); ¹³C NMR (176 MHz, CDCl₃) δ (ppm) = 191.46 (CO), 181.76 (CO), 143.42, 141.62, 140.65, 138.43, 136.60, 129.25, 128.94, 128.74, 128.53, 127.06, 125.41, 125.10, 121.99, 121.78, 121.74, 121.26, 120.04, 119.48, 114.40, 111.35, 109.61, 72.74, 70.23, 66.94, 57.75, 45.57, 41.98, 38.40, 28.59, 27.78, 24.86, 20.13, 11.88 (CH₃); IR (KBr, cm⁻¹) ν _{max} = 3291, 2926, 2854, 1716, 1699, 1694, 1683, 1668, 1660, 1652, 1617, 1597, 1557, 1538, 1505, 1475, 1456, 1398, 1373, 1330, 1221, 1179, 1099, 1011, 934, 870, 790, 741, 694; [Anal. Calcd. for C₃₇H₃₅N₅O₂: C, 76.40; H, 6.06; N, 12.04; found: C, 76.28; H, 6.14; N, 12.11]; LC/MS (ESI, *m/z*): found 582.5 [M + H]⁺, exact mass 581.28 for C₃₇H₃₅N₅O₂.

(*R*,*R*,*S*,*R*,*S*,*R*)-6-Chloro-1'-(1*H*-indol-3-yl)-2'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-5-nitro-3',6',7',7*a*'-tetrahydro-1*H*-spiro[indoline-3,5'-pyrrolo[1,2-*a*]indol]-2-one (**8l**). Following the general procedure (GP2), chalcone **5a** (82 mg, 0.25 mmol), 6-chloroisatin **6b** (46 mg, 0.25 mmol), and octahydro-1*H*-indole-2-carboxylic acid **7b** (64 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield light yellow solid compound **8l**; yield (131 mg, 85%); m.p.: 205–206 °C; $[\alpha]_D^{25} = -20.56^\circ$ (c 0.11, MeOH); ¹H NMR (700 MHz, CDCl₃) δ (ppm) = 8.55 (s, 1H, NH), 8.32 (s, 1H, NH), 8.10 (d, $J = 2.8$ Hz, 1H, Ar-H), 8.00 (d, $J = 7.6$ Hz, 1H, Ar-H), 7.42–7.38 (m, 3H, Ar-H), 7.32 (d, $J = 8.4$ Hz, 2H, Ar-H), 7.19–7.11 (m, 5H, Ar-H), 7.01 (d, $J = 8.0$ Hz, 1H, Ar-H), 6.41 (d, $J = 2.3$ Hz, 1H, Ar-H), 4.96 (d, $J = 12.0$ Hz, 1H, CHCO), 4.56–4.52 (m, 1H, NCH), 4.15 (dd, $J = 12.0, 10.0$ Hz, 1H, NCHCH), 3.22 (q, $J = 3.8$ Hz, 1H, NCH), 2.19–2.14 (m, 1H, NCHCH), 2.00 (s, 3H, CH₃), 1.82–1.76 (m, 2H, CH₂), 1.59–1.55 (m, 1H, CH₂), 1.54–1.46 (m, 3H, CH₂), 1.20–1.16 (m, 1H, CH₂), 1.07–0.96 (m, 3H, CH₂); ¹³C NMR (176 MHz, CDCl₃): δ (ppm) = 191.36 (CO), 181.77 (CO), 143.61, 142.07, 141.60, 138.28, 136.59, 134.55, 129.39, 129.03, 128.68, 126.99, 125.46, 123.59, 122.06, 121.81, 121.56, 121.16, 119.95, 119.55, 114.15, 111.40, 110.13, 72.39, 70.19, 67.06, 57.81, 45.69, 41.98, 38.35, 28.65, 27.77, 24.81, 20.14, 11.86 (CH₃); IR (KBr, cm⁻¹) ν _{max} = 3298, 2926, 2851, 1725, 1668, 1652, 1614, 1598, 1538, 1532, 1504, 1484, 1455, 1447, 1398, 1373, 1323, 1284, 1243, 1222, 1179, 1130, 1072, 1011, 937, 924, 869, 795, 764, 741, 694, 660, 597, 584, 524; [Anal. Calcd. for C₃₇H₃₄ClN₅O₂: C, 72.12; H, 5.56; N, 11.37; found: C, 71.96; H, 5.63; N, 11.28]; LC/MS (ESI, *m/z*): found 616.6 [M-(₃₅Cl) + H]⁺, 618.3 [M-(₃₇Cl) + H]⁺; exact mass 615.24 for C₃₇H₃₄ClN₅O₂.

(*R*,*R*,*S*,*R*,*S*,*R*)-1'-(1*H*-Indol-3-yl)-2'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-5-nitro-3',6',7',7*a*'-tetrahydro-1*H*-spiro[indoline-3,5'-pyrrolo[1,2-*a*]indol]-2-one (**8m**). Following the general procedure (GP2), chalcone **5a** (82 mg, 0.25 mmol), 5-nitroisatin **6c** (48 mg, 0.25 mmol), and octahydro-1*H*-indole-2-carboxylic acid **7b** (64 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 4 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield yellow solid compound **8m**; yield (103 mg, 66%); m.p.: 210–211 °C; $[\alpha]_D^{25} = -15.63^\circ$ (c 0.13, MeOH); ¹H NMR (700 MHz, CDCl₃) δ (ppm) = 9.19 (s, 1H, NH), 8.27–8.22 (m, 2H, Ar-H), 8.11 (s, 1H, NH), 8.06 (d, $J = 8.6$ Hz, 1H, Ar-H),

8.02 (d, $J = 7.9$ Hz, 1H, Ar-H), 7.42–7.37 (m, 3H, Ar-H), 7.34 (d, $J = 7.0$ Hz, 1H, Ar-H), 7.20 (s, 1H, Ar-H), 7.18–7.14 (m, 2H, Ar-H), 7.10 (d, $J = 7.5$ Hz, 2H, Ar-H), 6.42 (d, $J = 8.7$ Hz, 1H, Ar-H), 5.03 (d, $J = 12.0$ Hz, 1H, CHCO), 4.63–4.58 (m, 1H, NCH), 4.21 (t, $J = 11.0$ Hz, 1H, NCHCH), 3.24 (q, $J = 3.7$ Hz, 1H, NCH), 2.23–2.18 (m, 1H, NCHCH), 1.95 (s, 3H, CH₃), 1.90–1.86 (m, 1H, CH₂), 1.84–1.81 (m, 1H, CH₂), 1.61–1.56 (m, 1H, CH₂), 1.53–1.46 (m, 3H, CH₂), 1.22–1.17 (m, 1H, CH₂), 1.08–0.98 (m, 2H, CH₂), 0.93–0.89 (m, 1H, CH₂); ¹³C NMR (176 MHz, CDCl₃): δ (ppm) = 190.53 (CO), 182.11 (CO), 146.67, 143.82, 142.71, 141.51, 138.12, 136.64, 129.48, 129.20, 127.14, 126.80, 126.02, 125.38, 123.96, 122.21, 121.96, 121.13, 120.02, 119.71, 113.58, 111.43, 109.26, 72.52, 70.05, 67.02, 57.95, 46.02, 42.11, 38.27, 28.56, 27.72, 24.77, 20.05, 11.95 (CH₃); IR (KBr, cm⁻¹) ν _{max} = 3341, 2926, 2854, 1733, 1668, 1653, 1623, 1598, 1533, 1503, 1477, 1455, 1397, 1338, 1221, 1178, 1125, 1096, 1072, 1011, 930, 833, 764, 742, 695, 660, 555; [anal. calcd. for C₃₇H₃₄N₆O₄: C, 70.91; H, 5.47; N, 13.41; found: C, 71.06; H, 5.39; N, 13.49]; LC/MS (ESI, m/z): found 627.5 [M + H]⁺; exact mass 626.26 for C₃₇H₃₄N₆O₄.

(1'R,2'S,3R,9a'R)-6-Chloro-2'-(5-methyl-1-phenyl-1H-pyrazole-4-carbonyl)-1'-(1-methyl-1H-indol-3-yl)-1',2',4a',5',6',7',8',8a',9',9a'-decahydrospiro[indoline-3,3'-pyrrolo[1,2-a]indol]-2-one (**8n**). Following the general procedure (GP2), chalcone **5b** (85 mg, 0.25 mmol), 6-chloroisatin **6b** (46 mg, 0.25 mmol), and octahydro-1H-indole-2-carboxylic acid **7b** (64 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield light yellow solid compound **8n**; yield (134 mg, 85%); m.p.: 172–173 °C; $[\alpha]_D^{25} = -19.29^\circ$ (c 0.10, MeOH); ¹H NMR (700 MHz, CDCl₃) δ (ppm) = 8.06 (s, 1H, NH), 8.01 (d, $J = 7.9$ Hz, 1H, Ar-H), 7.44–7.40 (m, 3H, Ar-H), 7.31 (d, $J = 8.0$ Hz, 1H, Ar-H), 7.27 (s, 1H, Ar-H), 7.23–7.20 (m, 1H, Ar-H), 7.17 (d, $J = 7.7$ Hz, 3H, Ar-H), 7.10 (s, 1H, Ar-H), 7.02 (d, $J = 8.0$ Hz, 1H, Ar-H), 6.43 (s, 1H, Ar-H), 4.93 (d, $J = 12.0$ Hz, 1H, CHCO), 4.56–4.52 (m, 1H, NCH), 4.16–4.11 (m, 1H, NCHCH), 3.73 (s, 3H, NCH₃), 3.23 (q, $J = 3.8$ Hz, 1H, NCH), 2.20–2.15 (m, 1H, NCHCH), 2.02 (s, 3H, CH₃), 1.84–1.79 (m, 2H, CH₂), 1.60–1.56 (m, 1H, CH₂), 1.55–1.48 (m, 3H, CH₂), 1.21–1.17 (m, 1H, CH₂), 1.08–0.96 (m, 3H, CH₂); ¹³C NMR (176 MHz, CDCl₃): δ (ppm) = 191.30 (CO), 181.64 (CO), 143.59, 142.02, 141.57, 138.33, 137.27, 134.52, 129.38, 128.99, 127.44, 127.12, 126.40, 125.46, 123.59, 121.66, 121.56, 121.16, 120.13, 119.09, 112.69, 110.05, 109.36, 72.29, 70.32, 67.20, 57.79, 45.60, 42.00, 38.38, 32.83 (NCH₃), 28.64, 27.78, 24.84, 20.11, 11.88 (CH₃); IR (KBr, cm⁻¹) ν _{max} = 3057, 2926, 2849, 1733, 1668, 1652, 1615, 1538, 1532, 1504, 1484, 1455, 1404, 1373, 1328, 1282, 1224, 1179, 1130, 1071, 1012, 935, 924, 869, 796, 764, 740, 694; [anal. calcd. for C₃₈H₃₆ClN₅O₂: C, 72.43; H, 5.76; N, 11.11; found: C, 72.52; H, 5.71; N, 11.03]; LC/MS (ESI, m/z): found 630.6 [M(₃₅Cl) + H]⁺, 632.3 [M(₃₇Cl) + H]⁺; exact mass 629.26 for C₃₈H₃₆ClN₅O₂.

(1'R,2'S,3R,9a'R)-2'-(5-Methyl-1-phenyl-1H-pyrazole-4-carbonyl)-1'-(1-methyl-1H-indol-3-yl)-5-nitro-1',2',4a',5',6',7',8',8a',9',9a'-decahydrospiro[indoline-3,3'-pyrrolo[1,2-a]indol]-2-one (**8o**). Following the general procedure (GP2), chalcone **5b** (85 mg, 0.25 mmol), 5-nitroisatin **6c** (48 mg, 0.25 mmol), and octahydro-1H-indole-2-carboxylic acid **7b** (64 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 4 h and purified by column chromatography

100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield yellow solid compound **8o**; yield (119 mg, 74%); m.p.: 185–186 °C; $[\alpha]_D^{25} = -32.74^\circ$ (c 0.14, MeOH); ¹H NMR (700 MHz, CDCl₃) δ (ppm) = 11.03 (s, 1H, NH), 8.16 (d, $J = 7.6$ Hz, 2H, Ar-H), 7.98 (s, 1H, Ar-H), 7.86 (d, $J = 7.9$ Hz, 1H, Ar-H), 7.52–7.50 (m, 2H, Ar-H), 7.49–7.45 (m, 2H, Ar-H), 7.38–7.33 (m, 3H, Ar-H), 7.16–7.13 (m, 1H, Ar-H), 7.10–7.07 (m, 1H, Ar-H), 6.85 (d, $J = 9.1$ Hz, 1H, Ar-H), 4.92 (d, $J = 12.1$ Hz, 1H, CHCO), 4.25–4.12 (m, 1H, NCH), 4.13 (dd, $J = 12.1$, 9.9 Hz, 1H, NCHCH), 3.71 (s, 3H, NCH₃), 3.22 (q, $J = 3.5$ Hz, 1H, NCH), 2.19–2.13 (m, 1H, NCHCH), 2.05–1.99 (m, 1H, CH₂), 1.81 (s, 3H, CH₃), 1.65–1.61 (m, 1H, CH₂), 1.55–1.50 (m, 1H, CH₂), 1.48–1.43 (m, 1H, CH₂), 1.39–1.27 (m, 2H, CH₂), 1.15–1.10 (m, 1H, CH₂), 1.04–0.97 (m, 1H, CH₂), 0.96–0.90 (m, 1H, CH₂), 0.80–0.74 (m, 1H, CH₂); ¹³C NMR (176 MHz, CDCl₃): δ (ppm) = 190.37 (CO), 181.03 (CO), 148.21, 142.53, 141.42, 141.08, 137.89, 136.71, 129.31, 128.82, 128.03, 127.10, 126.54, 126.41, 125.20, 125.07, 123.11, 121.17, 120.49, 119.19, 118.56, 111.66, 109.68, 71.51, 70.53, 66.16, 56.87, 44.37, 41.52, 37.53, 32.33 (NCH₃), 27.69, 27.27, 24.37, 19.27, 11.08 (CH₃); IR (KBr, cm⁻¹) ν _{max} = 3056, 2927, 2854, 1738, 1733, 1716, 1699, 1694, 1682, 1674, 1668, 1661, 1652, 1645, 1634, 1622, 1615, 1598, 1557, 1538, 1524, 1519, 1505, 1475, 1464, 1456, 1436, 1427, 1398, 1373, 1338, 1223, 1178, 1127, 1072, 1011, 930, 833, 795; [anal. calcd. for C₃₈H₃₆N₆O₄: C, 71.23; H, 5.66; N, 13.12; found: C, 71.31; H, 5.74; N, 13.27]; LC/MS (ESI, m/z): found 641.5 [M + H]⁺; exact mass 640.26 for C₃₈H₃₆N₆O₄.

(1'R,2'S,3R,9a'R)-1'-(5-Bromo-1H-indol-3-yl)-6-chloro-2'-(5-methyl-1-phenyl-1H-pyrazole-4-carbonyl)-1',2',4a',5',6',7',8',8a',9',9a'-decahydrospiro[indoline-3,3'-pyrrolo[1,2-a]indol]-2-one (**8p**). Following the general procedure (GP2), chalcone **5c** (102 mg, 0.25 mmol), 6-chloroisatin **6b** (46 mg, 0.25 mmol), and octahydro-1H-indole-2-carboxylic acid **7b** (64 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield light yellow solid compound **8p**; yield (155 mg, 89%); m.p.: 199–200 °C; $[\alpha]_D^{25} = -26.47^\circ$ (c 0.12, MeOH); ¹H NMR (700 MHz, CDCl₃) δ (ppm) = 8.72 (s, 1H, NH), 8.63 (s, 1H, NH), 8.12 (s, 1H, Ar-H), 8.05 (s, 1H, Ar-H), 7.43–7.39 (m, 3H, Ar-H), 7.31 (d, $J = 8.0$ Hz, 1H, Ar-H), 7.23 (d, $J = 1.5$ Hz, 2H, Ar-H), 7.15–7.12 (m, 2H, Ar-H), 7.11 (d, $J = 2.5$ Hz, 1H, Ar-H), 7.02 (dd, $J = 8.0$, 1.9 Hz, 1H, Ar-H), 6.35 (s, 1H, Ar-H), 4.79 (d, $J = 12.0$ Hz, 1H, CHCO), 4.38–4.33 (m, 1H, NCH), 4.09 (dd, $J = 12.1$, 10.1 Hz, 1H, NCHCH), 3.21 (q, $J = 3.7$ Hz, 1H, NCH), 2.22–2.17 (m, 1H, NCHCH), 1.98 (s, 3H, CH₃), 1.85–1.78 (m, 2H, CH₂), 1.60–1.56 (m, 1H, CH₂), 1.54–1.45 (m, 3H, CH₂), 1.22–1.17 (m, 1H, CH₂), 1.09–0.97 (m, 3H, CH₂); ¹³C NMR (176 MHz, CDCl₃): δ (ppm) = 191.05 (CO), 181.83 (CO), 143.75, 142.03, 141.48, 138.22, 135.08, 134.65, 129.46, 129.36, 129.17, 129.15, 128.70, 127.14, 125.45, 125.03, 123.42, 122.38, 122.12, 121.67, 121.05, 114.32, 112.87, 110.15, 72.37, 70.86, 67.65, 57.77, 44.84, 41.96, 38.35, 28.57, 27.77, 24.77, 20.11, 11.80 (CH₃); IR (KBr, cm⁻¹) ν _{max} = 3273, 2926, 2853, 1733, 1699, 1674, 1652, 1615, 1557, 1538, 1532, 1504, 1483, 1456, 1398, 1373, 1322, 1278, 1220, 1179, 1130, 1102, 1072, 936, 932, 884, 869, 797, 765, 693, 670, 595, 525; [anal. calcd. for C₃₇H₃₃BrClN₅O₂: C, 63.94; H, 4.79; N, 10.08; found: C, 64.09; H, 4.86; N, 10.13]; LC/MS (ESI, m/z): found 695.0 [M(₃₅Cl/₇₉Br) + H]⁺, 696.3 [M(₃₇Cl/₈₁Br) + H]⁺, 698.1 [M(₃₇Cl + ₈₁Br) + H]⁺; exact mass 693.15 for C₃₇H₃₃BrClN₅O₂.

(*1R,2'S,3R,9a'R*)-1'-(5-Bromo-1*H*-indol-3-yl)-6-chloro-2'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-1',2',4a',5',6',7',8',8a',9',9a'-decahydrospiro[indoline-3,3'-pyrrolo[1,2-a]indol]-2-one (**8q**). Following the general procedure (GP2), chalcone **5c** (102 mg, 0.25 mmol), 5-nitroisatin **6c** (48 mg, 0.25 mmol), and octahydro-1*H*-indole-2-carboxylic acid **7b** (64 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 4 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield yellow solid compound **8q**; yield (120 mg, 68%); m.p.: 190–199 °C; $[\alpha]_D^{25} = -167.95^\circ$ (c 0.10, MeOH); ¹H NMR (700 MHz, DMSO-*d*₆): δ (ppm) = 11.17 (d, *J* = 2.7 Hz, 1H, NH), 11.01 (s, 1H, NH), 8.26 (d, *J* = 2.3 Hz, 1H, Ar-H), 8.15 (dd, *J* = 8.6, 2.4 Hz, 1H, Ar-H), 8.04 (s, 1H, Ar-H), 8.03 (d, *J* = 1.9 Hz, 1H, Ar-H), 7.57 (d, *J* = 2.6 Hz, 1H, Ar-H), 7.53–7.50 (m, 2H, Ar-H), 7.48–7.45 (m, 1H, Ar-H), 7.36 (d, *J* = 7.1 Hz, 2H, Ar-H), 7.29 (d, *J* = 8.6 Hz, 1H, Ar-H), 7.18 (dd, *J* = 8.5, 1.9 Hz, 1H, Ar-H), 6.84 (d, *J* = 8.6 Hz, 1H, Ar-H), 4.88 (d, *J* = 12.0 Hz, 1H, CHCO), 4.17–4.11 (m, 2H, NCH + NCHCH), 3.23 (q, *J* = 3.4 Hz, 1H, NCH), 2.18–2.10 (m, 2H NCHCH₂), 1.81 (s, 3H, CH₃), 1.63–1.58 (m, 1H, CH₂), 1.54–1.50 (m, 1H, CH₂), 1.49–1.45 (m, 1H, CH₂), 1.38–1.28 (m, 2H, CH₂), 1.15–1.10 (m, 1H, CH₂), 1.03–0.97 (m, 1H, CH₂), 0.96–0.90 (m, 1H, CH₂), 0.77–0.72 (m, 1H, CH₂); ¹³C NMR (176 MHz, DMSO-*d*₆): δ (ppm) = 190.50 (CO), 180.99 (CO), 148.17, 142.49, 141.54, 141.20, 137.91, 134.91, 129.31, 128.92, 128.81, 128.03, 126.51, 126.40, 125.20, 125.06, 123.62, 123.52, 123.33, 121.21, 120.55, 113.43, 112.57, 111.14, 109.61, 71.50, 70.72, 66.35, 56.77, 43.80, 41.56, 37.06, 27.65, 27.29, 24.42, 19.25, 11.08 (CH₃); IR (KBr, cm⁻¹) ν_{\max} = 3363, 2930, 2854, 1734, 1762, 1621, 1599, 1528, 1503, 1478, 1454, 1398, 1337, 1221, 1178, 1126, 1098, 1081, 928, 884, 866, 831, 795, 765, 753, 693, 659, 598, 553; [anal. calcd. for C₃₇H₃₃BrN₆O₄: C, 63.94; H, 4.79; N, 10.08; found: C, 64.09; H, 4.86; N, 10.13]; LC/MS (ESI, *m/z*): found 705.8 [M(₇₉Br) + H]⁺, 707.5 [M(₈₁Br) + H]⁺, exact mass 704.17 for C₃₇H₃₃BrN₆O₄.

(*1R,2'S,3R,9a'R*)-6-Chloro-1'-(5-chloro-1*H*-indol-3-yl)-2'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-1',2',4a',5',6',7',8',8a',9',9a'-decahydrospiro[indoline-3,3'-pyrrolo[1,2-a]indol]-2-one (**8r**). Following the general procedure (GP2), chalcone **5d** (91 mg, 0.25 mmol), 6-chloroisatin **6b** (46 mg, 0.25 mmol), and octahydro-1*H*-indole-2-carboxylic acid **7b** (64 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield light yellow solid compound **8r**; yield (148 mg, 91%); m.p.: 168–169 °C; $[\alpha]_D^{25} = -121.51^\circ$ (c 0.12, MeOH); ¹H NMR (700 MHz, CDCl₃) δ (ppm) = 8.77 (d, *J* = 11.7 Hz, 1H, NH), 8.63 (s, 1H, NH), 8.13 (d, *J* = 2.0 Hz, 1H, Ar-H), 7.91 (d, *J* = 2.0 Hz, 1H, Ar-H), 7.45–7.39 (m, 3H, Ar-H), 7.31 (d, *J* = 8.0 Hz, 1H, Ar-H), 7.28–7.26 (m, 1H, Ar-H), 7.15–7.12 (m, 3H, Ar-H), 7.11 (dd, *J* = 8.6, 2.0 Hz, 1H, Ar-H), 7.02 (dd, *J* = 8.0, 1.9 Hz, 1H, Ar-H), 6.35 (s, 1H, Ar-H), 4.81 (d, *J* = 12.0 Hz, 1H, CHCO), 4.39–4.36 (m, 1H, NCH), 4.10 (dd, *J* = 11.9, 10.0 Hz, 1H, NCHCH), 3.22 (q, *J* = 3.7 Hz, 1H, NCH), 2.19–2.14 (m, 1H, NCHCH), 1.99 (s, 3H), 1.83–1.76 (m, 2H, CH₂), 1.58–1.53 (m, 1H, CH₂), 1.52–1.44 (m, 3H, CH₂), 1.19–1.15 (m, 1H, CH₂), 1.07–0.95 (m, 3H, CH₂); ¹³C NMR (176 MHz, CDCl₃): δ (ppm) = 191.10 (CO), 181.85 (CO), 143.75, 142.06, 141.49, 138.22, 134.65, 129.46, 129.16, 128.69, 128.47, 127.14, 125.45, 123.43, 122.60, 122.47, 121.65, 121.05, 119.07, 114.34, 112.48, 110.16, 72.39, 70.79, 67.65, 57.78, 44.90, 41.96, 38.35, 28.58, 27.77, 24.77, 20.12, 11.79 (CH₃);

IR (KBr, cm⁻¹) ν_{\max} = 3287, 2925, 2854, 1723, 1661, 1652, 1615, 1538, 1533, 1504, 1484, 1398, 1373, 1320, 1277, 1221, 1178, 1131, 1104, 1072, 736, 926, 893, 869, 796, 763, 694, 606, 525; [anal. calcd. for C₃₇H₃₃Cl₂N₅O₂: C, 68.31; H, 5.11; N, 10.76; found: C, 68.25; H, 5.07; N, 10.84]; LC/MS (ESI, *m/z*): found 650.8 [M(₃₅Cl) + H]⁺, 652.0 [M(₃₇Cl) + H]⁺, 654.0 [M(₃₇Cl) + ₃₇Cl] + H]⁺; exact mass 649.2 for C₃₇H₃₃Cl₂N₅O₂.

(*1'R,2'S,3R,9a'R*)-1'-(5-Chloro-1*H*-indol-3-yl)-2'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-5-nitro-1',2',4a',5',6',7',8',8a',9',9a'-decahydrospiro[indoline-3,3'-pyrrolo[1,2-a]indol]-2-one (**8s**). Following the general procedure (GP2), chalcone **5d** (91 mg, 0.25 mmol), 5-nitroisatin **6c** (48 mg, 0.25 mmol), and octahydro-1*H*-indole-2-carboxylic acid **7b** (64 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 4 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield yellow solid compound **8s**; yield (132 mg, 80%); m.p.: 172–173 °C; $[\alpha]_D^{25} = -11.15^\circ$ (c 0.10, MeOH); ¹H NMR (700 MHz, DMSO-*d*₆): δ (ppm) = 11.15 (d, *J* = 2.6 Hz, 1H, NH), 11.01 (s, 1H, NH), 8.25 (d, *J* = 2.3 Hz, 1H, Ar-H), 8.15 (dd, *J* = 8.6, 2.3 Hz, 1H, Ar-H), 8.03 (s, 1H, Ar-H), 7.89 (d, *J* = 2.1 Hz, 1H, Ar-H), 7.58 (d, *J* = 2.5 Hz, 1H, Ar-H), 7.53–7.49 (m, 2H, Ar-H), 7.49–7.45 (m, 1H, Ar-H), 7.38–7.35 (m, 2H, Ar-H), 7.33 (d, *J* = 8.6 Hz, 1H, Ar-H), 7.07 (dd, *J* = 8.6, 2.0 Hz, 1H, Ar-H), 6.84 (d, *J* = 8.6 Hz, 1H, Ar-H), 4.88 (d, *J* = 12.0 Hz, 1H, CHCO), 4.15 (m, 2H, NCH + NCHCH), 3.23 (q, *J* = 3.4 Hz, 1H, NCH), 2.19–2.09 (m, 2H, CH₂), 1.81 (s, 3H, CH₃), 1.62–1.59 (m, 1H, CH₂), 1.55–1.50 (m, 1H, CH₂), 1.49–1.44 (m, 1H, CH₂), 1.39–1.27 (m, 2H, CH₂), 1.14–1.10 (m, 1H, CH₂), 1.03–0.97 (m, 1H, CH₂), 0.97–0.91 (m, 1H, CH₂), 0.77–0.72 (m, 1H, CH₂); ¹³C NMR (176 MHz, DMSO-*d*₆): δ (ppm) = 190.50 (CO), 180.99 (CO), 148.17, 142.49, 141.53, 141.20, 137.91, 134.69, 129.31, 128.81, 128.18, 128.03, 126.51, 126.40, 125.20, 123.81, 123.15, 121.00, 120.54, 118.20, 112.96, 112.64, 109.61, 71.51, 70.67, 66.34, 56.78, 43.83, 41.56, 37.09, 27.65, 27.29, 24.42, 19.25, 11.07 (CH₃); IR (KBr, cm⁻¹) ν_{\max} = 3349, 2928, 2853, 1732, 1661, 1623, 1598, 1537, 1525, 1504, 1455, 1398, 1337, 1220, 1178, 1128, 1099, 1078, 931, 892, 867, 833, 798, 766, 755, 694, 552; [anal. calcd. for C₃₇H₃₃ClN₆O₄: C, 67.22; H, 5.03; N, 12.71; found: C, 67.34; H, 12.63; N, 12.56]; LC/MS (ESI, *m/z*): found 661.6 [M(₃₅Cl) + H]⁺, 663.3 [M(₃₇Cl) + H]⁺; exact mass 660.23 for C₃₇H₃₃ClN₆O₄.

(*1'R,2'S,3R,7a'R*)-6-Chloro-1'-(1*H*-indol-3-yl)-2'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-1',2',5',6',7',7a'-hexahydrospiro[indoline-3,3'-pyrrolizin]-2-one (**8t**). Following the general procedure (GP2), chalcone **5a** (82 mg, 0.25 mmol), 6-chloroisatin **6b** (46 mg, 0.25 mmol), and L-proline **7c** (43 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and MeOH/CH₂Cl₂ (3:97) to yield light yellow solid compound **8t**; yield (103 mg, 73%); m.p.: 165–166 °C; $[\alpha]_D^{25} = -32.67^\circ$ (c 0.12, MeOH); ¹H NMR (700 MHz, CDCl₃): δ (ppm) = 9.13 (s, 1H, NH), 8.33 (s, 1H, NH), 8.09–8.04 (m, 1H, Ar-H), 7.93 (s, 1H, Ar-H), 7.41–7.35 (m, 3H, Ar-H), 7.32–7.30 (m, 1H, Ar-H), 7.29 (d, *J* = 8.1 Hz, 1H, Ar-H), 7.23 (d, *J* = 2.4 Hz, 1H, Ar-H), 7.19–7.13 (m, 4H, Ar-H), 6.99 (dd, *J* = 8.0, 1.9 Hz, 1H, Ar-H), 6.63 (d, *J* = 1.9 Hz, 1H, Ar-H), 4.89 (d, *J* = 11.4 Hz, 1H, CHCO), 4.49–4.46 (m, 1H, NCH), 4.19 (dd, *J* = 11.5, 9.9 Hz, 1H, NCHCH), 2.80–2.75 (m, 1H, NCH_{2(a)}), 2.70–2.66 (m, 1H, NCH_{2(b)}), 2.09 (s, 3H, CH₃), 2.06–2.02 (m, 1H, CH₂), 1.97–1.91 (m, 1H CH₂), 1.90–1.85

(m, 1H CH_2), 1.79–1.73 (m, 1H CH_2); ^{13}C NMR (176 MHz, CDCl_3): δ (ppm) = 191.91 ($\underline{\text{CO}}$), 181.47 ($\underline{\text{CO}}$), 143.56, 142.53, 141.27, 138.24, 136.68, 134.93, 129.33, 128.97, 128.86, 126.77, 125.58, 124.44, 122.32, 122.10, 121.88, 121.27, 119.90, 119.67, 114.21, 111.44, 110.70, 73.77, 70.74, 65.90, 48.38, 44.74, 31.06, 27.28, 12.00 ($\underline{\text{CH}_3}$); IR (KBr, cm^{-1}) $\nu_{\text{max}} = 3273$, 3056, 2962, 2867, 1733, 1661, 1653, 1613, 1536, 1531, 1502, 1486, 1453, 1396, 1324, 1285, 1244, 1222, 1185, 1131, 1072, 936, 922, 813, 794, 763, 741, 694, 526; [anal. calcd. for $\text{C}_{33}\text{H}_{28}\text{ClN}_5\text{O}_2$: C, 70.52; H, 5.02; N, 12.46; found: C, 70.66; H, 4.91; N, 12.53]; LC/MS (ESI, m/z): found 562.4 [$\text{M}(\text{Cl}) + \text{H}]^+$, 564.1 [$\text{M}(\text{Cl}) + \text{H}]^+$; exact mass 561.19 for $\text{C}_{33}\text{H}_{28}\text{ClN}_5\text{O}_2$.

(*1'R,2'S,3R,7a'R*)-6-Chloro-2'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-1'-(1-methyl-1*H*-indol-3-yl)-1',2',5',6',7',7a'-hexahydrospiro[indoline-3,3'-pyrrolizin]-2-one (**8u**). Following the general procedure (GP2), chalcone **5b** (85 mg, 0.25 mmol), 6-chloroisatin **6b** (46 mg, 0.25 mmol), and L-proline **7c** (43 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and $\text{MeOH}/\text{CH}_2\text{Cl}_2$ (3:97) to yield light yellow solid compound **8u**; yield (124 mg, 87%); m.p.: 150–151 °C; $[\alpha]_D^{25} = -23.35^\circ$ (c 0.11, MeOH); ^1H NMR (700 MHz, CDCl_3): δ (ppm) = 9.02 (s, 1H, NH), 8.07 (d, $J = 7.9$ Hz, 1H, Ar- $\underline{\text{H}}$), 7.89 (s, 1H, Ar- $\underline{\text{H}}$), 7.41–7.35 (m, 3H, Ar- $\underline{\text{H}}$), 7.30–7.26 (m, 2H, Ar- $\underline{\text{H}}$), 7.23–7.20 (m, 1H, Ar- $\underline{\text{H}}$), 7.17 (dd, $J = 16.0, 7.6$ Hz, 3H, Ar- $\underline{\text{H}}$), 7.13 (s, 1H, Ar- $\underline{\text{H}}$), 6.99 (dd, $J = 8.0, 1.9$ Hz, 1H, Ar- $\underline{\text{H}}$), 6.65 (d, $J = 1.9$ Hz, 1H, Ar- $\underline{\text{H}}$), 4.85 (d, $J = 11.5$ Hz, 1H, CHCO), 4.50–4.45 (m, 1H, NCH), 4.16 (dd, $J = 11.5, 10.0$ Hz, 1H, NCHCH), 3.73 (s, 3H, NCH_3), 2.82–2.76 (m, 1H, $\text{NCH}_{2(a)}$), 2.70–2.65 (m, 1H, $\text{NCH}_{2(b)}$), 2.09 (s, 3H, CH_3), 2.07–2.01 (m, 1H, CH_2), 1.98–1.92 (m, 1H, CH_2), 1.90–1.86 (m, 1H, CH_2), 1.78–1.73 (m, 1H, CH_2); ^{13}C NMR (176 MHz, CDCl_3): δ (ppm) = 191.89 ($\underline{\text{CO}}$), 181.46 ($\underline{\text{CO}}$), 143.51, 142.51, 141.24, 138.29, 137.38, 134.91, 129.32, 128.93, 128.86, 127.21, 126.92, 125.58, 124.47, 121.86, 121.71, 121.29, 120.07, 119.21, 112.71, 110.64, 109.44, 73.67, 70.80, 66.02, 48.43, 44.66, 32.82 ($\underline{\text{NCH}_3}$), 30.96, 27.19, 12.01 ($\underline{\text{CH}_3}$); IR (KBr, cm^{-1}) $\nu_{\text{max}} = 3216$, 3058, 2958, 2872, 1733, 1662, 1614, 1536, 1503, 1484, 1455, 1398, 1326, 1285, 1227, 1184, 1130, 1172, 1012, 935, 922, 815, 795, 765, 740, 694; [anal. calcd. for $\text{C}_{34}\text{H}_{30}\text{ClN}_5\text{O}_2$: C, 70.89; H, 5.25; N, 12.16; found: C, 71.11; H, 5.16; N, 12.01]; LC/MS (ESI, m/z): found 576.5 [$\text{M}(\text{Cl}) + \text{H}]^+$, 578.2 [$\text{M}(\text{Cl}) + \text{H}]^+$; exact mass 575.21 for $\text{C}_{34}\text{H}_{30}\text{ClN}_5\text{O}_2$.

(*1'R,2'S,3R,7a'R*)-1'-(5-Bromo-1*H*-indol-3-yl)-2'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-1',2',5',6',7',7a'-hexahydrospiro[indoline-3,3'-pyrrolizin]-2-one (**8v**). Following the general procedure (GP2), chalcone **5c** (102 mg, 0.25 mmol), isatin **6a** (37 mg, 0.25 mmol), and L-proline **7c** (43 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and $\text{MeOH}/\text{CH}_2\text{Cl}_2$ (3:97) to yield light yellow solid compound **8v**; yield (136 mg, 90%); m.p.: 178–179 °C; $[\alpha]_D^{25} = -37.86^\circ$ (c 0.10, MeOH); ^1H NMR (700 MHz, DMSO- d_6): δ (ppm) = 11.16 (s, 1H, NH), 10.35 (s, 1H, NH), 8.10 (s, 1H, Ar- $\underline{\text{H}}$), 7.85 (d, $J = 3.0$ Hz, 1H, Ar- $\underline{\text{H}}$), 7.51–7.46 (m, 3H, Ar- $\underline{\text{H}}$), 7.44 (d, $J = 9.1$ Hz, 1H, Ar- $\underline{\text{H}}$), 7.39 (d, $J = 7.9$ Hz, 1H, Ar- $\underline{\text{H}}$), 7.34–7.29 (m, 3H, Ar- $\underline{\text{H}}$), 7.19 (d, $J = 10.5$ Hz, 1H, Ar- $\underline{\text{H}}$), 7.14–7.10 (m, 1H, Ar- $\underline{\text{H}}$), 6.98–6.94 (m, 1H, Ar- $\underline{\text{H}}$), 6.65 (d, $J = 10.3$ Hz, 1H, Ar- $\underline{\text{H}}$), 4.68 (d, $J = 11.7$ Hz, 1H, CHCO), 4.09 (t, $J = 12.0$ Hz, 1H, NCHCH), 4.05–4.01 (m, 1H, NCH), 2.62–2.57 (m, 1H, $\text{NCH}_{2(a)}$), 2.39–2.35 (m, 1H,

$\text{NCH}_{2(b)}$), 1.94–1.90 (m, 1H, CH_2), 1.89 (s, 3H, CH_3), 1.87–1.84 (m, 1H, CH_2), 1.80–1.75 (m, 1H, CH_2), 1.75–1.69 (m, 1H, CH_2); ^{13}C NMR (176 MHz, DMSO- d_6): δ (ppm) = 191.33 ($\underline{\text{CO}}$), 179.87 ($\underline{\text{CO}}$), 142.37, 141.96, 140.86, 138.03, 135.10, 129.29, 129.08, 128.74, 128.70, 127.85, 125.27, 125.23, 123.96, 123.53, 121.32, 120.99, 120.78, 113.57, 113.33, 111.21, 109.43, 73.20, 71.04, 65.39, 47.42, 43.25, 30.49, 27.08, 11.21 ($\underline{\text{CH}_3}$); IR (KBr, cm^{-1}) $\nu_{\text{max}} = 3270$, 2958, 2928, 2869, 1717, 1662, 1653, 1616, 1597, 1538, 1504, 1470, 1457, 1397, 1330, 1268, 1221, 1190, 1104, 936, 884, 793, 754, 694, 678, 659, 610; [anal. calcd. for $\text{C}_{33}\text{H}_{28}\text{BrN}_5\text{O}_2$: C, 65.35; H, 4.65; N, 11.55; found: C, 65.24; H, 4.77; N, 11.49]; LC/MS (ESI, m/z): found 606.7 [$\text{M}(\text{Br}) + \text{H}]^+$, 608.4 [$\text{M}(\text{Br}) + \text{H}]^+$; exact mass 605.14 for $\text{C}_{33}\text{H}_{28}\text{BrN}_5\text{O}_2$.

(*1'R,2'S,3R,7a'R*)-1'-(5-Chloro-1*H*-indol-3-yl)-2'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-5-nitro-1',2',5',6',7',7a'-hexahydrospiro[indoline-3,3'-pyrrolizin]-2-one (**8w**). Following the general procedure (GP2), chalcone **5d** (91 mg, 0.25 mmol), 5-nitroisatin **6c** (48 mg, 0.25 mmol), and L-proline **7c** (43 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 6 h and purified by column chromatography 100–200 mesh silica gel and $\text{MeOH}/\text{CH}_2\text{Cl}_2$ (3:97) to yield yellow solid compound **8w**; yield (74 mg, 49%); m.p.: 225–226 °C; $[\alpha]_D^{25} = -67.91^\circ$ (c 0.13, MeOH); ^1H NMR (700 MHz, DMSO- d_6): δ (ppm) = 11.18 (s, 1H, NH), 11.16 (s, 1H, NH), 8.14 (dd, $J = 8.6, 2.3$ Hz, 1H, Ar- $\underline{\text{H}}$), 8.12 (d, $J = 2.3$ Hz, 1H, Ar- $\underline{\text{H}}$), 7.93 (d, $J = 2.1$ Hz, 1H, Ar- $\underline{\text{H}}$), 7.84 (s, 1H), 7.55 (d, $J = 2.5$ Hz, 1H, Ar- $\underline{\text{H}}$), 7.51–7.48 (m, 2H, Ar- $\underline{\text{H}}$), 7.47–7.44 (m, 1H, Ar- $\underline{\text{H}}$), 7.36 (d, $J = 8.6$ Hz, 1H, Ar- $\underline{\text{H}}$), 7.31 (d, $J = 8.6$ Hz, 2H, Ar- $\underline{\text{H}}$), 7.08 (dd, $J = 8.6, 2.0$ Hz, 1H, Ar- $\underline{\text{H}}$), 6.88 (d, $J = 8.6$ Hz, 1H, Ar- $\underline{\text{H}}$), 4.75 (d, $J = 11.4$ Hz, 1H, CHCO), 4.15–4.09 (m, 2H, $\text{NCH} + \text{NCHCH}$), 2.70–2.65 (m, 1H, $\text{NCH}_{2(a)}$), 2.46–2.42 (m, 1H, $\text{NCH}_{2(b)}$), 1.99–1.94 (m, 1H, CH_2), 1.92 (s, 3H, CH_3), 1.90–1.88 (m, 1H, CH_2), 1.84–1.76 (m, 2H, CH_2); ^{13}C NMR (176 MHz, DMSO- d_6): δ (ppm) = 191.15 ($\underline{\text{CO}}$), 180.27 ($\underline{\text{CO}}$), 148.48, 142.63, 141.67, 140.69, 137.82, 134.92, 129.30, 128.86, 127.70, 126.56, 126.22, 125.19, 124.53, 123.23, 122.78, 121.03, 120.59, 118.14, 113.15, 112.52, 109.86, 72.44, 70.67, 65.39, 47.41, 43.52, 30.02, 27.09, 11.21 ($\underline{\text{CH}_3}$); IR (KBr, cm^{-1}) $\nu_{\text{max}} = 3408$, 3378, 3111, 3058, 2973, 2861, 2836, 1748, 1645, 1622, 1598, 1525, 1503, 1479, 1455, 1417, 1392, 1373, 1338, 1293, 1253, 1221, 1188, 1174, 1137, 1100, 1076, 986, 938, 930, 887, 860, 850, 832, 798, 763, 755, 694, 655, 610, 554; [anal. calcd. for $\text{C}_{33}\text{H}_{27}\text{ClN}_6\text{O}_4$: C, 65.29; H, 4.48; N, 13.84; found: C, 65.39; H, 4.56; N, 13.87]; LC/MS (ESI, m/z): found 607.5 [$\text{M}(\text{Cl}) + \text{H}]^+$, 609.2 [$\text{M}(\text{Cl}) + \text{H}]^+$; exact mass 606.18 for $\text{C}_{33}\text{H}_{27}\text{ClN}_6\text{O}_4$.

(*2'R,3'S,4'R*)-6-Chloro-4'-(1*H*-indol-3-yl)-1'-methyl-3'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)spiro[indoline-3,2'-pyrrolidin]-2-one (**8x**). Following the general procedure (GP2), chalcone **5a** (82 mg, 0.25 mmol), 6-chloroisatin **6b** (46 mg, 0.25 mmol), and sacrosine **7d** (34 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and $\text{MeOH}/\text{CH}_2\text{Cl}_2$ (3:97) to yield light yellow solid compound **8x**; yield (48 mg, 35%); m.p.: 140–141 °C; $[\alpha]_D^{25} = -19.37^\circ$ (c 0.10, MeOH); ^1H NMR (700 MHz, CDCl_3): δ (ppm) = 8.51 (s, 1H, NH), 8.19 (s, 1H, NH), 8.08 (d, $J = 9.5$ Hz, 1H, Ar- $\underline{\text{H}}$), 7.73 (s, 1H, Ar- $\underline{\text{H}}$), 7.45–7.38 (m, 3H, Ar- $\underline{\text{H}}$), 7.33–7.30 (m, 1H, Ar- $\underline{\text{H}}$), 7.26 (d, $J = 1.2$ Hz, 1H, Ar- $\underline{\text{H}}$), 7.20–7.14 (m, 5H, Ar- $\underline{\text{H}}$), 6.92 (dd, $J = 8.0, 1.9$ Hz, 1H, Ar- $\underline{\text{H}}$), 6.49 (d, $J = 1.8$ Hz, 1H, Ar- $\underline{\text{H}}$), 4.80–4.73 (m, 1H, NCH_2CH), 4.45 (d, $J = 9.6$ Hz, 1H, CHCO), 3.86–3.81 (m, 1H, $\text{NCH}_{2(a)}$), 3.49–3.45

(m, 1H, $\text{NCH}_{2(b)}$), 2.29 (s, 3H, NCH_3), 2.17 (s, 3H, CH_3); ^{13}C NMR (176 MHz, CDCl_3): δ (ppm) = 192.45 (CO), 180.34 (CO), 143.46, 142.23, 140.55, 138.20, 136.60, 134.48, 129.44, 129.18, 127.71, 126.87, 126.77, 125.71, 122.53, 122.21, 122.16, 121.44, 120.04, 119.75, 115.62, 111.30, 109.77, 73.69, 64.47, 59.54, 35.99, 35.30 (NCH_3), 11.91 (CH_3); IR (KBr, cm^{-1}) $\nu_{\text{max}} = 3278, 3072, 2934, 2826, 2797, 1726, 1666, 1654, 1612, 1597, 1534, 1503, 1485, 1552, 1398, 1322, 1284, 1240, 1216, 1179, 1119, 1094, 1069, 936, 910, 882, 814, 798, 766, 718, 693, 592; [anal. calcd. for $\text{C}_{31}\text{H}_{26}\text{ClN}_5\text{O}_2$: C, 69.46; H, 4.89; N, 13.07; found: C, 69.31; H, 4.97; N, 12.93]; LC/MS (ESI, m/z): found 536.4 [$\text{M}^{+}(\text{Cl}) + \text{H}^{+}$], 538.2 [$\text{M}^{+}(\text{Cl}) + \text{H}^{+}$]; exact mass 535.18 for $\text{C}_{31}\text{H}_{26}\text{ClN}_5\text{O}_2$.$

(*2'R,3'S,4'R*)-4'-(5-Bromo-1*H*-indol-3-yl)-6-chloro-1'-methyl-3'-(5-methyl-1-phenyl-1*H*-pyrazole-4-carbonyl)-spiro[indoline-3,2'-pyrrolidin]-2-one (**8y**). Following the general procedure (GP2), chalcone **5c** (102 mg, 0.25 mmol), 6-chloroisatin **6b** (46 mg, 0.25 mmol), and sacrosine **7d** (34 mg, 0.37 mmol) in methanol (20 mL) were refluxed for 3 h and purified by column chromatography 100–200 mesh silica gel and $\text{MeOH}/\text{CH}_2\text{Cl}_2$ (3:97) to yield light yellow solid compound **8y**; yield (73 mg, 47%); m.p.: 158–159 °C; $[\alpha]_D^{25} = -35.63^\circ$ (c 0.12, MeOH); ^1H NMR (700 MHz, CDCl_3): δ (ppm) = 8.99 (s, 1H, NH), 8.42 (s, 1H, NH), 8.18 (s, 1H, Ar-H), 7.77 (s, 1H, Ar-H), 7.44–7.38 (m, 3H, Ar-H), 7.22 (dd, $J = 8.7, 2.0$ Hz, 2H, Ar-H), 7.18–7.11 (m, 4H, Ar-H), 6.92 (dd, $J = 8.0, 1.8$ Hz, 1H, Ar-H), 6.54 (d, $J = 1.8$ Hz, 1H, Ar-H), 4.70 (q, $J = 9.8$ Hz, 1H, NCH_2CH_2), 4.36 (d, $J = 9.4$ Hz, 1H, CH_2CO), 3.75 (t, $J = 9.5$ Hz, 1H, $\text{NCH}_{2(a)}$), 3.46 (t, $J = 9.5$ Hz, 1H, $\text{NCH}_{2(b)}$), 2.27 (s, 3H, NCH_3), 2.19 (s, 3H, CH_3); ^{13}C NMR (176 MHz, CDCl_3): δ (ppm) = 192.35 (CO), 180.67 (CO), 143.63, 142.34, 140.51, 138.14, 135.16, 134.58, 129.47, 129.24, 128.70, 127.64, 126.53, 125.71, 125.00, 123.27, 122.55, 122.51, 121.38, 115.62, 113.00, 112.75, 110.02, 73.75, 64.57, 59.54, 35.58, 35.26 (NCH_3), 11.93 (CH_3); IR (KBr, cm^{-1}) $\nu_{\text{max}} = 3276, 3071, 2936, 2843, 2795, 1722, 1668, 1652, 1613, 1598, 1537, 1504, 1484, 1554, 1397, 1321, 1280, 1244, 1219, 1181, 1115, 1096, 1068, 938, 916, 883, 813, 793, 765, 714, 694, 593; [anal. calcd. for $\text{C}_{31}\text{H}_{25}\text{BrClN}_5\text{O}_2$: C, 60.55; H, 4.10; N, 11.39; found: C, 60.41; H, 4.15; N, 11.47]; LC/MS (ESI, m/z): found 614.8 [$\text{M}^{+}(\text{Cl}/^{79}\text{Br}) + \text{H}^{+}$], 616.1 [$\text{M}^{+}(\text{Cl}/^{81}\text{Br}) + \text{H}^{+}$], 618.0 [$\text{M}^{+}(\text{Cl}) + \text{Br}^{+}$ + H^{+}]; exact mass 613.09 for $\text{C}_{31}\text{H}_{25}\text{BrClN}_5\text{O}_2$.$

Acetylcholine Esterase (AChE) Inhibitory Assay (AChEI). AChEI activity was measured using Ellman's method as previously described and provided in the Supporting Information.^{39b,c}

Molecular Docking Study. The protocol for the molecular docking study to investigate the binding mode of the most active compounds are provided in the Supporting Information.⁴³

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsomegaf.1c03978>.

AChE assay; molecular docking study protocol; copies of NMR and MS spectrum; and also X-ray single-crystal analysis of compound **8c** (PDF)

AUTHOR INFORMATION

Corresponding Authors

Ved Prakash Verma – Department of Chemistry, Banasthali Vidyapith, Banasthali 304022 Rajasthan, India;
✉ orcid.org/0000-0003-2142-2459;
Email: vedprakash079@gmail.com

Assem Barakat – Department of Chemistry, College of Science, King Saud University, Riyadh 11451, Saudi Arabia;
Department of Chemistry, Faculty of Science, Alexandria University, Ibrahimia, Alexandria 21321, Egypt;
✉ orcid.org/0000-0002-7885-3201; Email: ambarakat@ksu.edu.sa

Authors

Mohammad Shahidul Islam – Department of Chemistry, College of Science, King Saud University, Riyadh 11451, Saudi Arabia; Department of Chemistry, Banasthali Vidyapith, Banasthali 304022 Rajasthan, India

Abdullah Mohammed Al-Majid – Department of Chemistry, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

Mohammad Azam – Department of Chemistry, College of Science, King Saud University, Riyadh 11451, Saudi Arabia;
✉ orcid.org/0000-0002-4274-2796

Matti Haukka – Department of Chemistry, University of Jyväskylä, FI-40014 Jyväskylä, Finland; ✉ orcid.org/0000-0002-6744-7208

Abdullah A. Elgazar – Department of Pharmacognosy, Faculty of Pharmacy, Kafrelsheikh University, Kafrelsheikh 33516, Egypt

Amira Mira – Department of Pharmacognosy, Faculty of Pharmacy, Mansoura University, Mansoura 35516, Egypt

Farid A. Badria – Department of Pharmacognosy, Faculty of Pharmacy, Mansoura University, Mansoura 35516, Egypt

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acsomegaf.1c03978>

Notes

The authors declare no competing financial interest.

▼ The current research was carried out in the current affiliation.

ACKNOWLEDGMENTS

The authors would like to extend their sincere appreciation to Researchers Supporting Project Number (RSP-2021/64), King Saud University, Riyadh, Saudi Arabia.

REFERENCES

- Hardy, J. A.; Higgins, G. A. Alzheimer's disease: the amyloid cascade hypothesis. *Science* **1992**, *256*, 184–186.
- Mohandas, E.; Rajmohan, V.; Raghunath, B. Neurobiology of Alzheimer's disease. *Indian J. Psychiatry* **2009**, *51*, 55.
- Swerdlow, R. H.; Burns, J. M.; Khan, S. M. The Alzheimer's disease mitochondrial cascade hypothesis. *J. Alzheimer's Dis.* **2010**, *20*, S265–S279.
- Maccioni, R. B.; Fariñas, G.; Morales, I.; Navarrete, L. The revitalized tau hypothesis on Alzheimer's disease. *Arch. Med. Res.* **2010**, *41*, 226–231.
- Wood, W. G.; Li, L.; Müller, W. E.; Eckert, G. P. Cholesterol as a causative factor in Alzheimer's disease: a debatable hypothesis. *J. Neurochem.* **2014**, *129*, 559–572.
- Zotova, E.; Nicoll, J. A.; Kalaria, R.; Holmes, C.; Boche, D. Inflammation in Alzheimer's disease: relevance to pathogenesis and therapy. *Alzheimer's Res. Ther.* **2010**, *2*, 1–9.

- (7) Markesberry, W. R. Oxidative stress hypothesis in Alzheimer's disease. *Free Radical Biol. Med.* **1997**, *23*, 134–147.
- (8) Bush, A. I.; Tanzi, R. E. Therapeutics for Alzheimer's disease based on the metal hypothesis. *Neurotherapeutics* **2008**, *5*, 421–432.
- (9) de la Torre, J. C. The vascular hypothesis of Alzheimer's disease: bench to bedside and beyond. *Neurodegener. Dis.* **2010**, *7*, 116–121.
- (10) Neve, R. L.; McPhie, D. L. The cell cycle as a therapeutic target for Alzheimer's disease. *Pharmacol. Ther.* **2006**, *111*, 99–113.
- (11) (a) de Oliveira Viana, J.; Monteiro, A. F. M.; Filho, J. M. B.; Scotti, L.; Scotti, M. T. The Azoles in Pharmacochemistry: Perspectives on the Synthesis of New Compounds and Chemoinformatic Contributions. *Curr. Pharm. Des.* **2020**, *25*, 4702–4716. (b) Teleb, M.; Rizk, O. H.; Zhang, F.-X.; Fronczeck, F. R.; Zamponi, G. W.; Fahmy, H. Design, synthesis and pharmacological evaluation of some substituted dihydropyrimidines with L-/T-type calcium channel blocking activities. *Bioorg. Chem.* **2019**, *83*, 354–366.
- (12) Radi, S.; El Massaoudi, M.; Bacquet, M.; Degoutin, S.; Adarsh, N.; Robeyns, K.; Garcia, Y. A novel environment-friendly hybrid material based on a modified silica gel with a bispyrazole derivative for the removal of Zn II, Pb II, Cd II and Cu II traces from aqueous solutions. *Inorg. Chem. Front.* **2017**, *4*, 1821–1831.
- (13) Ouyang, G.; Cai, X.-J.; Chen, Z.; Song, B.-A.; Bhadury, P. S.; Yang, S.; Jin, L.-H.; Xue, W.; Hu, D.-Y.; Zeng, S. Synthesis and antiviral activities of pyrazole derivatives containing an oxime moiety. *J. Agric. Food Chem.* **2008**, *56*, 10160–10167.
- (14) Keter, F. K.; Darkwa, J. Perspective: the potential of pyrazole-based compounds in medicine. *BioMetals* **2012**, *25*, 9–21.
- (15) Ramesh, B.; Bhalgat, C. M. Novel dihydropyrimidines and its pyrazole derivatives: Synthesis and pharmacological screening. *Eur. J. Med. Chem.* **2011**, *46*, 1882–1891.
- (16) Kumar, V.; Kaur, K.; Gupta, G. K.; Sharma, A. K. Pyrazole containing natural products: synthetic preview and biological significance. *Eur. J. Med. Chem.* **2013**, *69*, 735–753.
- (17) Diculescu, V. C.; Chioreea-Paquin, A.-M.; Oliveira-Brett, A. M. Applications of a DNA-electrochemical biosensor. *TrAC, Trends Anal. Chem.* **2016**, *79*, 23–36.
- (18) Graillot, V.; Tomasetig, F.; Cravedi, J.-P.; Audebert, M. Evidence of the in vitro genotoxicity of methyl-pyrazole pesticides in human cells. *Mutat. Res., Genet. Toxicol. Environ. Mutagen.* **2012**, *748*, 8–16.
- (19) Çetin, A.; Bildirici, I. A study on synthesis and antimicrobial activity of 4-acyl-pyrazoles. *J. Saudi Chem. Soc.* **2018**, *22*, 279–296.
- (20) Ramírez-Prada, J.; Robledo, S. M.; Vélez, I. D.; del Pilar Crespo, M.; Quiroga, J.; Abonia, R.; Montoya, A.; Svetaz, L.; Zacchino, S.; Insuasty, B. Synthesis of novel quinoline-based 4, 5-dihydro-1H-pyrazoles as potential anticancer, antifungal, antibacterial and antiprotozoal agents. *Eur. J. Med. Chem.* **2017**, *131*, 237–254.
- (21) El-Sabbagh, O. I.; Baraka, M. M.; Ibrahim, S. M.; Pannecouque, C.; Andrei, G.; Snoeck, R.; Balzarini, J.; Rashad, A. A. Synthesis and antiviral activity of new pyrazole and thiazole derivatives. *Eur. J. Med. Chem.* **2009**, *44*, 3746–3753.
- (22) Insuasty, B.; Tigreros, A.; Orozco, F.; Quiroga, J.; Abonia, R.; Nogueras, M.; Sanchez, A.; Cobo, J. Synthesis of novel pyrazolic analogues of chalcones and their 3-aryl-4-(3-aryl-4, 5-dihydro-1H-pyrazol-5-yl)-1-phenyl-1H-pyrazole derivatives as potential antitumor agents. *Bioorg. Med. Chem.* **2010**, *18*, 4965–4974.
- (23) Bandgar, B. P.; Gawande, S. S.; Bodade, R. G.; Gawande, N. M.; Khobragade, C. N. Synthesis and biological evaluation of a novel series of pyrazole chalcones as anti-inflammatory, antioxidant and antimicrobial agents. *Bioorg. Med. Chem.* **2009**, *17*, 8168–8173.
- (24) Pérez-Fernández, R.; Goya, P.; Elguero, J. A review of recent progress (2002–2012) on the biological activities of pyrazoles. *Arkivoc* **2014**, *2014*, 233–293.
- (25) Özdemir, A.; Altintop, M. D.; Kaplancıklı, Z. A.; Can, Ö. D.; Demir Özkan, Ü.; Turan-Zitouni, G. Synthesis and Evaluation of New 1, 5-Diaryl-3-[4-(methyl-sulfonyl) phenyl]-4, 5-dihydro-1H-pyrazole Derivatives as Potential Antidepressant Agents. *Molecules* **2015**, *20*, 2668–2684.
- (26) Kling, A.; Jantos, K.; Mack, H.; Hornberger, W.; Drescher, K.; Nimmrich, V.; Relo, A.; Wicke, K.; Hutchins, C. W.; Lao, Y.; et al. Discovery of Novel and Highly Selective Inhibitors of Calpain for the Treatment of Alzheimer's Disease: 2-(3-Phenyl-1 H-pyrazol-1-yl)-nicotinamides. *J. Med. Chem.* **2017**, *60*, 7123–7138.
- (27) (a) Ahsan, N.; Mishra, S.; Jain, M. K.; Surolia, A.; Gupta, S. Curcumin Pyrazole and its derivative (N-(3-Nitrophenylpyrazole) Curcumin inhibit aggregation, disrupt fibrils and modulate toxicity of Wild type and Mutant α -Synuclein. *Sci. Rep.* **2015**, *5*, No. 9862. (b) Estrada, A. A.; Chan, B. K.; Baker-Glenn, C.; Beresford, A.; Burdick, D. J.; Chambers, M.; Chen, H.; Dominguez, S. L.; Dotson, J.; Drummond, J.; et al. Discovery of highly potent, selective, and brain-penetrant aminopyrazole leucine-rich repeat kinase 2 (LRRK2) small molecule inhibitors. *J. Med. Chem.* **2014**, *57*, 921–936.
- (28) Neelarapu, R.; Holzle, D. L.; Velaparthi, S.; Bai, H.; Brunsteiner, M.; Blond, S. Y.; Petukhov, P. A. Design, synthesis, docking, and biological evaluation of novel diazide-containing isoxazole-and pyrazole-based histone deacetylase probes. *J. Med. Chem.* **2011**, *54*, 4350–4364.
- (29) Han, Y. T.; Kim, K.; Choi, G.-I.; An, H.; Son, D.; Kim, H.; Ha, H.-J.; Son, J.-H.; Chung, S.-J.; Park, H.-J.; et al. Pyrazole-S-carboxamides, novel inhibitors of receptor for advanced glycation end products (RAGE). *Eur. J. Med. Chem.* **2014**, *79*, 128–142.
- (30) (a) Turkan, F.; Çetin, A.; Taslimi, P.; Gülcin, İ. Some pyrazoles derivatives: Potent carbonic anhydrase, α -glycosidase, and cholinesterase enzymes inhibitors. *Arch. Pharm.* **2018**, *351*, No. 1800200. (b) Turkan, F.; Çetin, A.; Taslimi, P.; Karaman, M.; Gülcin, İ. Synthesis, biological evaluation and molecular docking of novel pyrazole derivatives as potent carbonic anhydrase and acetylcholinesterase inhibitors. *Bioorg. Chem.* **2019**, *86*, 420–427.
- (31) (a) Brandão, P.; Marques, C.; Burke, A. J.; Pineiro, M. The application of isatin-based multicomponent-reactions in the quest for new bioactive and druglike molecules. *Eur. J. Med. Chem.* **2021**, *211*, 113102. (b) Lotfy, G.; Aziz, Y. M. A.; Said, M. M.; El Sayed, H.; El Sayed, H.; Abu-Serie, M. M.; Teleb, M.; Dömling, A.; Barakat, A. Molecular hybridization design and synthesis of novel spirooxindole-based MDM2 inhibitors endowed with BCL2 signaling attenuation; a step towards the next generation p53 activators. *Bioorg. Chem.* **2021**, No. 105427. (c) Aziz, Y. M. A.; Lotfy, G.; Said, M. M.; El Ashry, E. S. H.; El Tamany, E. S. H.; Soliman, S.; Abu-Serie, M. M.; Teleb, M.; Yousuf, S.; Dömling, A.; Domingo, L. R.; Barakat, A. Design, Synthesis, Chemical and Biochemical Insights on to Novel Hybrid Spirooxindoles-Based p53-MDM2 Inhibitors with Potential Bcl2 Signaling Attenuation. *Front. Chem.* **2021**, 915.
- (32) (a) Choi, S. S.; Lee, S.-R.; Kim, S. U.; Lee, H. J. Alzheimer's disease and stem cell therapy. *Exp. Neurobiol.* **2014**, *23*, 45. (b) Giacobini, E. Selective inhibitors of butyrylcholinesterase. *Drugs Aging* **2001**, *18*, 891–898. (c) Kia, Y.; Osman, H.; Kumar, R. S.; Murugaiyah, V.; Basiri, A.; Perumal, S.; Razak, I. A. A facile chemo-, regio- and stereoselective synthesis and cholinesterase inhibitory activity of spirooxindole–pyrrolizine–piperidine hybrids. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 2979–2983. (d) Kia, Y.; Osman, H.; Kumar, R. S.; Basiri, A.; Murugaiyah, V. Synthesis and discovery of highly functionalized mono- and bis-spiro-pyrrolidines as potent cholinesterase enzyme inhibitors. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 1815–1819. (e) Kia, Y.; Osman, H.; Kumar, R. S.; Murugaiyah, V.; Basiri, A.; Perumal, S.; Wahab, H. A.; Bing, C. S. Synthesis and discovery of novel piperidone-grafted mono- and bis-spirooxindole-hexahydropyrrolizines as potent cholinesterase inhibitors. *Bioorg. Med. Chem.* **2013**, *21*, 1696–1707. (f) Kia, Y.; Osman, H.; Kumar, R. S.; Basiri, A.; Murugaiyah, V. Ionic liquid mediated synthesis of mono- and bis-spirooxindole-hexahydropyrrolidines as cholinesterase inhibitors and their molecular docking studies. *Bioorg. Med. Chem.* **2014**, *22*, 1318–1328. (g) Arumugam, N.; Almansour, A. I.; Kumar, R. S.; Kotresha, D.; Saiswaroop, R.; Venketesh, S. Dispiropyrrolidinyl-piperidone embedded indeno [1, 2-b] quinoxaline heterocyclic hybrids: Synthesis, cholinesterase inhibitory activity and their molecular docking simulation. *Bioorg. Med. Chem.* **2019**, *27*, 2621–2628. (h) Barakat, A.; Soliman, S. M.; Alshahrani, S.; Islam, M. S.; Ali,

- M.; Al-Majid, A. M.; Yousuf, S. Synthesis, X-ray Single Crystal, Conformational Analysis and Cholinesterase Inhibitory Activity of a New Spiropyrrolidine Scaffold Tethered Benzo [b] Thiophene Analogue. *Crystals* **2020**, *10*, 120. (i) Goyal, D.; Kaur, A.; Goyal, B. Benzofuran and indole: promising scaffolds for drug development in Alzheimer's disease. *ChemMedChem* **2018**, *13*, 1275–1299. (j) Kumar, R. S.; Ali, M. A.; Osman, H.; Ismail, R.; Choon, T. S.; Yoon, Y. K.; Wei, A. C.; Pandian, S.; Manogaran, E. Synthesis and discovery of novel hexacyclic cage compounds as inhibitors of acetylcholinesterase. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 3997–4000.
- (33) Chigurupati, S.; Selvaraj, M.; Mani, V.; Selvarajan, K. K.; Mohammad, J. I.; Kaveti, B.; Bera, H.; Palanimuthu, V. R.; Teh, L. K.; Salleh, M. Z. Identification of novel acetylcholinesterase inhibitors: Indolopyrazoline derivatives and molecular docking studies. *Bioorg. Chem.* **2016**, *67*, 9–17.
- (34) Barakat, A.; Alshahrani, S.; Al-Majid, A. M.; Ali, M.; Altowyan, M. S.; Islam, M. S.; Almary, A. S.; Ashraf, S.; Ul-Haq, Z. Synthesis of a New Class of Spirooxindole–Benzo [b] Thiophene-Based Molecules as Acetylcholinesterase Inhibitors. *Molecules* **2020**, *25*, 4671.
- (35) (a) Mei, G.-J.; Shi, F. Catalytic asymmetric synthesis of spirooxindoles: recent developments. *Chem. Commun.* **2018**, *54*, 6607–6621. (b) Saranya, P.; Neetha, M.; Aneela, T.; Anilkumar, G. Transition metal-catalyzed synthesis of spirooxindoles. *RSC Adv.* **2021**, *11*, 7146–7179.
- (36) (a) Wang, C.-S.; Zhu, R.-Y.; Zheng, J.; Shi, F.; Tu, S.-J. Enantioselective construction of spiro [indoline-3, 2'-pyrrole] framework via catalytic asymmetric 1, 3-dipolar cycloadditions using allenes as equivalents of alkynes. *J. Org. Chem.* **2015**, *80*, 512–520. (b) Dai, W.; Jiang, X.-L.; Wu, Q.; Shi, F.; Tu, S.-J. Diastereo-and enantioselective construction of 3, 3'-pyrrolidinylidispriooxindole framework via catalytic asymmetric 1, 3-dipolar cycloadditions. *J. Org. Chem.* **2015**, *80*, 5737–5744. (c) Wang, Y.-M.; Zhang, H.-H.; Li, C.; Fan, T.; Shi, F. Catalytic asymmetric chemoselective 1, 3-dipolar cycloadditions of an azomethine ylide with isatin-derived imines: diastereo-and enantioselective construction of a spiro [imidazolidine-2, 3'-oxindole] framework. *Chem. Commun.* **2016**, *52*, 1804–1807.
- (37) (a) Altowyan, M. S.; Barakat, A.; Al-Majid, A. M.; Al-Ghulikah, H. Spiroindolone Analogues as Potential Hypoglycemic with Dual Inhibitory Activity on α -Amylase and α -Glucosidase. *Molecules* **2019**, *24*, 2342. (b) Altowyan, M. S.; Barakat, A.; Al-Majid, A. M.; Al-Ghulikah, H. Spiroindolone analogues bearing benzofuran moiety as a selective cyclooxygenase COX-1 with TNF- α and IL-6 inhibitors. *Saudi J. Biol. Sci.* **2020**, *27*, 1208–1216. (c) Barakat, A.; Islam, M. S.; Ghawas, H. M.; Al-Majid, A. M.; El-Senduny, F. F.; Badria, F. A.; Elshaier, Y. A. M.; Ghabbour, H. A. Substituted spirooxindole derivatives as potent anticancer agents through inhibition of phosphodiesterase 1. *RSC Adv.* **2018**, *8*, 14335–14346. (d) Islam, M. S.; Ghawas, H. M.; El-Senduny, F. F.; Al-Majid, A. M.; Elshaier, Y. A.; Badria, F. A.; Barakat, A. Synthesis of new thiazolo-pyrrolidine-(spirooxindole) tethered to 3-acylindole as anticancer agents. *Bioorg. Chem.* **2019**, *82*, 423–430. (e) Al-Majid, A. M.; Soliman, S. M.; Haukka, M.; Ali, M.; Islam, M. S.; Shaik, M. R.; Barakat, A. Design, Construction, and Characterization of a New Regioisomer and Diastereomer Material Based on the Spirooxindole Scaffold Incorporating a Sulphone Function. *Symmetry* **2020**, *12*, 1337. (f) Aziz, y. a.; Lotfy, G.; Said, M. M.; El Sayed, H.; El Ashry, H.; El Tamany, H.; Soliman, S. M.; Haukka, M.; Barakat, A. A.; et al. Synthesis and characterization of a new spirooxindole grafted pyrrolidino/piperidine moiety. *Record. Pharm. Biomed. Sci.* **2021**, *5*, 1–7. (g) Islam, M. S.; Haukka, M.; Soliman, S. M.; Al-Majid, A. M.; Rahman, A. M.; Bari, A.; Barakat, A. Regio-and Stereoselective Synthesis of Spiro-heterocycles Bearing the Pyrazole Scaffold via [3+2] Cycloaddition Reaction. *J. Mol. Struct.* **2021**, *1250*, No. 131711.
- (38) (a) Ríos-Gutiérrez, M.; Domingo, L. R. Unravelling the mysteries of the [3+ 2] cycloaddition reactions. *Eur. J. Org. Chem.* **2019**, *2019*, 267–282. (b) R Domingo, L.; Chamorro, E.; Perez, P. Understanding the high reactivity of the azomethine ylides in [3 + 2] cycloaddition reactions. *Lett. Org. Chem.* **2010**, *7*, 432–439.
- (c) Domingo, L. R.; Kula, K.; Ríos-Gutiérrez, M. Unveiling the reactivity of cyclic azomethine ylides in [3+ 2] cycloaddition reactions within the molecular electron density theory. *Eur. J. Org. Chem.* **2020**, *2020*, 5938–5948. (d) Domingo, L. R.; Ríos-Gutiérrez, M.; Pérez, P. A molecular electron density theory study of the role of the copper metalation of azomethine ylides in [3+ 2] cycloaddition reactions. *J. Org. Chem.* **2018**, *83*, 10959–10973.
- (39) (a) Ellman, G. L.; Courtney, K. D.; Andres, V.; Featherstone, R. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.* **1961**, *7*, 88–95. (b) Mira, A.; Yamashita, S.; Kataoka, Y.; Shimizu, K. In vitro neuroprotective activities of compounds from Angelica shikokiana Makino. *Molecules* **2015**, *20*, 4813–4832. (c) Abdel Bar, F. M.; Elimam, D. M.; Mira, A. S.; El-Senduny, F. F.; Badria, F. A. Derivatization, molecular docking and in vitro acetylcholinesterase inhibitory activity of glycyrrhizin as a selective anti-Alzheimer agent. *Nat. Prod. Res.* **2019**, *33*, 2591–2599.
- (40) Elgazar, A. A.; Knany, H. R.; Ali, M. S. Insights on the molecular mechanism of anti-inflammatory effect of formula from Islamic traditional medicine: An in-silico study. *J. Tradit. Complementary Med.* **2019**, *9*, 353–363.
- (41) (a) Cheung, J.; Rudolph, M. J.; Burshteyn, F.; Cassidy, M. S.; Gary, E. N.; Love, J.; Franklin, M. C.; Height, J. J. Structures of human acetylcholinesterase in complex with pharmacologically important ligands. *J. Med. Chem.* **2012**, *55*, 10282–10286. (b) Bacalhau, P.; San Juan, A. A.; Marques, C. S.; Peixoto, D.; Goth, A.; Guarda, C.; Silva, M.; Arantes, S.; Caldeira, A. T.; Martins, R.; Burke, A. J. New cholinesterase inhibitors for Alzheimer's disease: Structure Activity Studies (SARs) and molecular docking of isoquinolone and azepanone derivatives. *Bioorg. Chem.* **2016**, *67*, 1–8.
- (42) Alinezhad, H.; Tajbakhsh, M.; Zare, M. Catalyst-free one-pot synthesis of 1, 4, 5-trisubstituted pyrazoles in 2, 2, 2-trifluoroethanol. *J. Fluorine Chem.* **2011**, *132*, 995–1000.
- (43) (a) Inc., C. C. G. C. M. O. E. M. S. S. W., Suite #910, Montreal, QC, Canada, H3A 2R7. (b) Hsu, K.-C.; Chen, Y.-F.; Lin, S.-R.; Yang, J.-M. iGEMDOCK: a graphical environment of enhancing GEMDOCK using pharmacological interactions and post-screening analysis. *BMC Bioinf.* **2011**, *12*, 1–11. (c) Studio, D. J. A. I. S. D., CA, USA (2009). “version 2.5”.