



Erratum: Behavioral Abnormalities in Knockout and Humanized Tau Mice

OPEN ACCESS

Approved by:

Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*Correspondence:

Frontiers Production Office
production.office@frontiersin.org

Specialty section:

This article was submitted to
Neuroendocrine Science,
a section of the journal
Frontiers in Endocrinology

Received: 29 October 2021

Accepted: 29 October 2021

Published: 18 November 2021

Citation:

Frontiers Production Office (2021)
Erratum: Behavioral Abnormalities in
Knockout and Humanized Tau Mice.
Front. Endocrinol. 12:804873.
doi: 10.3389/fendo.2021.804873

Frontiers Production Office*

Frontiers Media SA, Lausanne, Switzerland

Keywords: Alzheimer's disease, MAPT, Tau protein, insulin, anxiety, metabolism, memory

An Erratum on:

Behavioral Abnormalities in Knockout and Humanized Tau Mice

By Gonçalves RA, Wijesekara N, Fraser PE and De Felice FG (2020). *Front. Endocrinol.* 11:124.
doi: 10.3389/fendo.2020.00124

Due to a production error, there was a mistake in **Figure 4A** as published. Image 4A has a blank box, containing no data. The corrected **Figure 4A** appears below.

The publisher apologizes for this mistake.

The original version of this article has been updated.

Copyright © 2021 Frontiers Production Office. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 4 | Phospho-tau and tau oligomers are increased in multiple brain regions of hTau mice. Immunoblot analysis of tau oligomers in **(A)** cortical, **(B)** hippocampal, and **(C)** hypothalamic lysates from 20-week-old WT, TauKO, and hTau mice. Immunoblot analysis of pTauSer199Ser202/TotalTau ratio in **(D)** cortical (n= 8 WT; 6 hTau), **(E)** hippocampal (n = 7 WT; 9 hTau), and **(F)** hypothalamic (n = 6 WT; 6 hTau) lysates from 20-week-old WT, TauKO, and hTau mice.