

Concise report

IL11 is elevated in systemic sclerosis and IL11-dependent ERK signalling underlies TGF β -mediated activation of dermal fibroblasts

Eleonora Adami ^{1,*}, Sivakumar Viswanathan ^{1,*}, Anissa A. Widjaja¹, Benjamin Ng², Sonia Chothani¹, Nevin Zhihao², Jessie Tan², Pei Min Lio², Benjamin L. George¹, Umut Altunoglu³, Kakaly Ghosh⁴, Bhairav S. Paleja⁵, Sebastian Schafer^{1,2}, Bruno Reversade^{3,4,5,6}, Salvatore Albani⁷, Andrea Low Hsiu Ling^{8,9}, Steven O'Reilly ^{10,#} and Stuart A. Cook ^{1,2,11,#}

Abstract

Objectives. Interleukin 11 (IL11) is highly upregulated in skin and lung fibroblasts from patients with systemic sclerosis (SSc). Here we tested whether IL11 is mechanistically linked with activation of human dermal fibroblasts (HDFs) from patients with SSc or controls.

Methods. We measured serum IL11 levels in volunteers and patients with early diffuse SSc and manipulated IL11 signalling in HDFs using gain- and loss-of-function approaches that we combined with molecular and cellular phenotyping.

Results. In patients with SSc, serum IL11 levels are elevated as compared with healthy controls. All transforming growth factor beta (TGF β) isoforms induced IL11 secretion from HDFs, which highly express IL11 receptor α -subunit and the glycoprotein 130 (gp130) co-receptor, suggestive of an autocrine loop of IL11 activity in HDFs. IL11 stimulated ERK activation in HDFs and resulted in HDF-to-myofibroblast transformation and extracellular matrix secretion. The pro-fibrotic action of IL11 in HDFs appeared unrelated to STAT3 activity, independent of TGF β upregulation and was not associated with phosphorylation of SMAD2/3. Inhibition of IL11 signalling using either a neutralizing antibody against IL11 or siRNA against *IL11RA* reduced TGF β -induced HDF proliferation, matrix production and cell migration, which was phenocopied by pharmacological inhibition of ERK.

Conclusions. These data reveal that autocrine IL11-dependent ERK activity alone or downstream of TGF β stimulation promotes fibrosis phenotypes in dermal fibroblasts and suggest IL11 as a potential therapeutic target in SSc.

Key words: systemic sclerosis, IL11, IL11RA, TGF β 2, antibody therapy, fibrosis, neutralizing antibody

Rheumatology key messages

- Serum IL11 levels are elevated in patients with diffuse systemic sclerosis compared with healthy controls.
- IL11-dependent ERK signalling is required for TGF β to elicit a profibrotic response in dermal fibroblasts.
- Neutralizing IL11 antibodies block TGF β 2-driven myofibroblast activation in systemic sclerosis fibroblasts.

¹Cardiovascular and Metabolic Disorders Program, Duke-National University of Singapore Medical School, ²National Heart Research Institute Singapore, National Heart Centre Singapore, Singapore, ³Department of Medical Genetics, Koç University, School of Medicine, Istanbul, Turkey, ⁴Genome Institute of Singapore, Human Genetics and Therapeutics Laboratory, ⁵Institute of Molecular and Cellular Biology, A*STAR, ⁶Department of Paediatrics, National University of Singapore, ⁷Translational Immunology Institute (TII), SingHealth-DukeNUS Academic Medical Centre, ⁸Department of Rheumatology and Immunology, Singapore General Hospital, ⁹Duke-National University of Singapore Medical School, Singapore, ¹⁰Department of Biosciences, Durham University, Durham and

¹¹MRC-London Institute of Medical Sciences, Hammersmith Hospital Campus, London, UK

Submitted 9 October 2020; accepted 5 February 2021

Correspondence to: Stuart A. Cook, 8 College Road 169857, Duke-NUS Medical School, Singapore. E-mail: stuart.cook@duke-nus.edu.sg

*These authors contributed equally to this work.

#Co-senior authors.

Introduction

Dermal fibrosis is a pathological hallmark of chronic inflammatory skin conditions such as SSc, atopic dermatitis and psoriasis. In SSc, fibrosis also occurs in visceral organs, notably the lung, and is accompanied by proliferative vasculopathy [1, 2] and inflammation [3].

TGF β family proteins are the quintessential mediators of fibroblast activation across organs and diseases, including SSc [4]. The TGF β 2 isoform is notably upregulated in fibroblasts and skin biopsies from patients with SSc as compared with healthy individuals, at both the RNA and the protein level [5, 6]. Thus, anti-fibrotic strategies for patients with SSc need to be carefully selected based on the ability to inhibit TGF β 2 signalling and those targeting integrin-mediated TGF β activation may not be effective. Unfortunately, targeting TGF β in general is associated with serious side effects, which have precluded the use of anti-TGF β reagents in the clinic, to date [7, 8].

Interleukin (IL) 11 has recently been identified as a TGF β -responsive profibrotic cytokine [9, 10], and its gene has been shown to be one of the most highly upregulated in fibroblasts from SSc lesional skin and in pulmonary fibroblasts of patients with interstitial lung disease due to SSc [6, 11]. Myofibroblasts are the major cell type that underlie fibrosis and secrete copious amounts of extracellular matrix while also being pro-inflammatory. Given that TGF β -driven myofibroblast activation is IL11-dependent in fibroblasts from some tissues and stromal cells more generally [12], we set out to characterize IL11 as a potential fibrogenic factor in dermal fibroblasts from controls and patients with SSc.

Methods

Detailed methods are provided in the [supplementary data](#) file that accompanies this paper, which includes detailed methods on the following: patient cohort, cell culture, recombinant proteins and chemicals, sequencing of the *IL11RA* mutation, RNA sequencing, immunofluorescence staining, ELISA, immunoblotting, siRNA knockdown, RT-qPCR, high-content imaging assays, half-maximal inhibitory concentration measurement, migration assay and statistical analysis.

Results

IL11 levels are elevated in the serum of patients with early diffuse SSc

We measured circulating IL11 levels in serum from a cohort of 21 patients with early diffuse SSc (<2 years from onset of non-Raynaud's symptoms) by ELISA (Human Quantikine IL11, R&D Systems, Minneapolis, MN, USA) and found that IL11 levels were significantly elevated in the serum of SSc patients compared with sex-matched healthy controls (Fig. 1A, mean SSc: 27.2(10.5) pg/ml; mean controls: 9.6(5.5) pg/ml; $P < 0.001$; further details

in [Supplementary Table S1](#) and [Supplementary Fig. S1](#), available at *Rheumatology* online). Thus IL11 is not only highly expressed in skin and lung fibroblasts from patients with SSc [6, 11] but is also systematically elevated.

IL11 signalling drives HDF-to-myofibroblast transformation

To begin to examine whether IL11 plays a role in dermal fibroblast biology, we performed RNA-sequencing (RNA-seq) of primary human dermal fibroblasts (HDFs). Analyses revealed that IL11 receptor α -subunit (IL11RA) is one of the most highly expressed members of the IL6 family of receptors (Fig. 1B), which signal via gp130. TGF β receptor (TGF β R) 2 transcripts were also highly expressed, whereas TGF β R1 and TGF β R3 RNA levels were less abundant (Fig. 1B). Immunofluorescence staining of HDFs revealed strong expression of IL11RA at the protein level (Fig. 1C).

We stimulated HDFs with TGF β 1, TGF β 2 or TGF β 3 and found that all isoforms induced IL11 secretion (Fig. 1D). These data provide evidence that HDFs are both a source and a target of IL11 and that IL11 signalling, perhaps in an autocrine mode, is related to TGF β -stimulated effects in HDFs.


To assess whether IL11 secretion from HDFs following TGF β stimulation was related to profibrotic phenotypes, we stimulated HDFs with IL11, TGF β 1, TGF β 2 or TGF β 3 in the presence of either non-targeting control siRNA or *IL11RA* siRNA (siIL11RA) (Fig. 1E-F; [Supplementary Fig. S2A-E](#), available at *Rheumatology* online). All stimuli induced HDF proliferation and secretion of the extracellular matrix proteins collagen I and periostin. Inhibition of IL11 signalling by siIL11RA inhibited the profibrotic effects of not only IL11 but also all TGF β isoforms. These data show that IL11 is a profibrotic factor in HDFs and that TGF β -driven HDF activation is dependent, at least in part, on autocrine IL11 activity.

Dissection of signalling events downstream of IL11 activation in dermal fibroblasts

To investigate the signalling mechanisms underlying the profibrotic effects of IL11 in skin fibroblasts, we employed the use of HDFs obtained from an individual with Kreiborg–Pakistani syndrome (MIM: 614188). This individual lacks IL11 signalling due to a homozygous p.Arg296Trp loss-of-function mutation in *IL11RA*, which we confirmed by sequencing ([Supplementary Fig. S2F](#), available at *Rheumatology* online) [13]. We compared these *IL11RA* mutant fibroblasts to a healthy age- and ethnically matched control and treated both control and mutant HDFs with either low dose/physiological (10 ng/ml) or high dose/supra-physiological (1000 ng/ml) recombinant human IL11 (Fig. 1G).

Consistent with data from fibroblasts from other organs, physiological concentrations (10 ng/ml) of IL11 maximally activated extracellular signal-regulated kinase (ERK) in control human HDFs at an early time point (15 min) and sustained this activation throughout the

Fig. 1 IL11 levels are increased in the serum of patients with systemic sclerosis and IL11 has SMAD-independent profibrotic activity in human dermal fibroblasts



(A) Serum IL11 levels in patients with diffuse SSc and sex-matched healthy controls. Data: $n=21$ per group (Supplementary Table S1, available at *Rheumatology* online), whiskers from min. to max. value, line at median. Statistics: two-tailed t -test. (B) RNA-seq of primary human dermal fibroblasts ($n=4$). RNA expression values are shown as transcripts per million (TPM) for receptors of the TGF β and IL6 cytokine family. (C) IL11RA staining in unstimulated HDFs (scale bar, 100 μ m). (D) IL11 secretion by HDFs stimulated with three TGF β isoforms (5 ng/ml) for 24 h. Data ($n=3$): mean (s.d.). Statistics: one-way ANOVA, Dunnett's multiple testing correction. (E) Quantification of signal intensity/area (collagen I, periostin) and percentage of EdU-staining positive HDFs subjected to siRNA knock-down for *IL11RA* and stimulation with TGF β 1, TGF β 2, TGF β 3 or IL11 (5 ng/ml) for 24 h. NT: non targeting siRNA control. (F) Collagen I staining representative images (periostin and EdU staining images are shown in Supplementary Fig. S2E, available at *Rheumatology* online) of HDFs that had been subjected to siRNA knockdown for *IL11RA* and stimulation with the indicated factors for 24 h. (G) Western blots assessing protein levels of phospho(p)ERK, ERK, pSTAT3, STAT3, α SMA, GAPDH in primary dermal fibroblasts of control and from an individual carrying a homozygous germline loss-of-function p.Arg296Trp mutation in *IL11RA*. (H and I) Western blot of SMAD2 and pSMAD2/3 expression in HDFs following stimulation with TGF β 2 or IL11 (5 ng/ml; 15 min or 24 h). EdU: 5-ethynyl-2'-deoxyuridine; ERK: extracellular signal-regulated kinase; GAPDH: glyceraldehyde-3-phosphate dehydrogenase; HDF: human dermal fibroblast; STAT: signal transducer and activator of transcription.

stimulation period (24 h). While IL11 has been shown to stimulate signal transducer and activator of transcription (STAT) in some contexts, we observed negligible phosphorylation of STAT3 with physiological IL11 levels and limited phosphorylation with a supraphysiological dose in HDFs (Fig. 1G). IL11-induced ERK and STAT3 phosphorylation were absent in IL11-stimulated HDFs from the IL11RA p.Arg296Trp mutant individual. IL11 stimulation of HDFs resulted in increased α -smooth muscle actin (α SMA) expression that was absent in IL11RA p.Arg296Trp HDFs (Fig. 1G).

We next assessed the potential dependency of the pro-fibrotic effects of IL11 on canonical TGF β 2 and/or SMAD2/3 activity. We observed that following stimulation of HDFs with IL11 there was no increase in TGF β levels (Supplementary Fig. S2G and H, available at *Rheumatology* online) and no detectable SMAD2/3 phosphorylation at either early or late time points (Fig. 1H and I). These data are consistent with findings in cardiac fibroblasts [9] and rule out a substantive role for a TGF β -dependent feed-forward loop downstream of IL11 activity in HDFs.

Inhibition of IL11 signalling blocks TGF β 2-stimulated dermal fibroblast- to-myofibroblast transformation

Next, we sought to understand whether the use of a neutralizing IL11 antibody (X203) [14] could block activation of HDFs. We first tested the efficacy of X203 in inhibiting the secretion of MMP2 from TGF β 2-stimulated HDFs [14]. X203 dose-dependently inhibited TGF β 2-induced MMP2 secretion by both primary HDFs (orange line) and SSc fibroblasts (purple line) with half-maximal inhibitory concentration (IC₅₀) of 0.34 nM and 0.31 nM, respectively (Fig. 2A).

In migration assays, we found that IL11 or TGF β 2 promoted HDF migration and that X203 reduced the migratory capacity of TGF β 2-stimulated HDFs as compared with control (Fig. 2B and C and Supplementary Fig. S3A and B, available at *Rheumatology* online). Furthermore, inhibition of IL11 signalling with X203 in TGF β -stimulated HDFs markedly reduced cell proliferation and the expression of collagen I and periostin (Fig. 2D and E and Supplementary Fig. S3C–E, available at *Rheumatology* online).

ERK signalling is important for IL11 effects in cardiac fibroblasts [9] and we found that TGF β 2 or IL11 stimulated ERK phosphorylation to a similar degree in HDFs (Supplementary Fig. S3F, available at *Rheumatology* online). We tested the mechanistic importance of ERK signalling and found that pharmacological inhibition of ERK activity blocked the profibrotic effects of all TGF β isoforms with the same effect as neutralizing IL11 antibody (Fig. 2D and E and Supplementary Fig. S3C–E, G, available at *Rheumatology* online).

Finally and importantly, the use of IL11 neutralizing antibodies in SSc fibroblasts was shown to recapitulate the results seen in non-diseased HDFs. At baseline, SSc fibroblasts highly express IL11RA (Supplementary Fig. S3H, available at *Rheumatology* online) and stimulation

of SSc fibroblasts with IL11 or TGF β 2 induced α SMA expression (Fig. 2F and G), collagen I and periostin, and also increased cell proliferation (Supplementary Fig. S3I, available at *Rheumatology* online). Following TGF β 2 stimulation, inhibition of IL11 signalling with either X203 or ERK inhibition (U0126, 10 μ M) resulted in a reduction in all profibrotic phenotypes tested (Supplementary Fig. S3I, available at *Rheumatology* online).


Discussion

While many factors stimulate HDF activation, members of the TGF β superfamily are regarded as the prototypical activators. Using fibroblasts explanted from healthy controls and patients with diffuse SSc treated with TGF β , Sargent *et al.* [15] identified IL6 and IL11, among others, as TGF β -responsive genes *in vitro*. More recently, IL11 has been found to be highly upregulated in SSc skin fibroblasts [6] and in pulmonary fibroblasts of patients suffering from SSc-associated interstitial lung disease [11]. Of all the IL6 family cytokines, IL11 is the only member whose expression correlates with lung fibrosis severity in patients with idiopathic pulmonary fibrosis [10]. While targeting IL6 in SSc has proved unsuccessful for the primary end point of improvement in skin sclerosis (focuSSced trial, NCT02453256) [16], perhaps reflecting the low levels of IL6 receptor (IL6R) on fibroblasts [9], the role of IL11 has remained unexplored.

In this study, we show that patients with early diffuse SSc have elevated serum levels of IL11 and that HDFs highly express IL11RA but not IL6R. In keeping with data on fibroblasts from other organs [12], we show that IL11 is required for the profibrotic effects of TGF β in HDFs and that this is primarily driven via non-canonical ERK signalling. This is in agreement with previous work describing the central role for the ERK1/2 cascade in relation to TGF β -stimulated SSc fibroblast activation *in vitro* [17–19] and skin fibrosis *in vivo* [19]. We conclude that non-canonical TGF β and IL11 signalling act through ERK to cause fibrosis and that the effect downstream of TGF β is IL11-dependent.

Some studies have suggested a role for STAT3 in the profibrotic effects of TGF β [20–22]. We found that IL11 consistently induced ERK phosphorylation concomitant with α SMA upregulation, while STAT3 phosphorylation was limited, not sustained and was only apparent at supraphysiological IL11 levels. Thus, we think it is unlikely that STAT3 activity is directly related to IL11 effects on fibrosis phenotypes although indirect mechanisms could play a role. Our data also rule out a dependency of IL11 on canonical TGF β signalling as IL11-stimulated HDFs, while exhibiting pro-fibrotic phenotypes, do not upregulate TGF β and show no evidence of SMAD2/3 activation. We point out that it is possible that IL11 induces reactive oxidative species in HDFs, given that IL11 promotes NADPH oxidase 4 (NOX4) upregulation in hepatocytes and NOX4 is strongly upregulated in the skin of patients with SSc [6, 23, 24], and this requires further study.

Fig. 2 IL11 antibodies block TGF β 2-driven activation of skin fibroblasts from SSc patients and controls



(A) Dose-response curve and X203 IC₅₀ value in inhibiting MMP2 secretion by HDFs and SSc dermal fibroblasts stimulated with TGF β 2. (X203 serial 4-fold dilution from 4 μ g/ml to 61 pg/ml; line at $x = 2 \mu$ g/ml; total $n = 2$ per group.) (B) Transwell migration indices of HDFs following 24h stimulation with TGF β 2 (5 ng/ml) and in the presence of IgG or X203 (2 μ g/ml). Data: $n = 4$, mean with s.d. Statistics: one-way ANOVA, Tukey's multiple testing correction. (C) Migration assay representative images; scale bar, 150 μ m. (D) Representative images of collagen I, periostin and EdU stainings of HDFs, following a 24 h stimulation with IL11 (5 ng/ml), or a combination of TGF β 2 (5 ng/ml) and IgG (2 μ g/ml), IL11 neutralizing antibody (X203, 2 μ g/ml) or ERK inhibitor (U0126, 10 μ M); scale bar, 100 μ m. (E) Quantification of signal intensity/area (collagen I, periostin) and percentage of EdU-staining positive cells relative to experiment in (D). Statistics: one-way ANOVA, Tukey's multiple testing correction. (F and G) Representative images and quantification of percentage of α SMA-staining positive cells following a 24 h stimulation of SSc fibroblasts with the indicated factors. Data: whiskers from min. to max. value, line at median. Statistics: one-way ANOVA, Tukey's multiple testing correction. α SMA: α -smooth muscle actin; EdU: 5-ethynyl-2'-deoxyuridine; ERK: extracellular signal-regulated kinase; HDF: human dermal fibroblast.

SSc remains a disease for which no treatment has proven uniformly effective. Unfortunately, anti-TGF β interventions have generic and recurrent concerns over efficacy and safety [7, 8]. More recently, molecules targeting integrin-mediated TGF β activation have been developed to attempt to bypass toxicities. However, TGF β 2 is the most dysregulated TGF β isoform in SSc patients and this isoform is refractory to integrin-mediated activation. Thus the utility of integrin targeting therapies in SSc is open to question and other approaches may be needed.

In conclusion, we present evidence for the involvement of IL11, a TGF β responsive and pro-fibrotic cytokine that is highly upregulated in SSc fibroblasts, in HDF activation. If IL11 is shown to promote skin fibrosis in *in vivo* disease models, IL11 may represent a new and accessible therapeutic target for dermal fibrosis in SSc and perhaps other conditions.

Acknowledgements

We would also like to thank the NHCS sequencing team for their technical expertise. S.A.C., E.A. and S.V. designed the study and planned experiments. S.V., E.A., A.A.W., B.N., J.T. and B.L.G. carried out molecular biology experiments; S.V., E.A., A.A.W. and B.N. analysed the data. L.P.M. prepared RNA-seq libraries and S.C. and N.Z. analysed sequencing data. Human samples acquisition and preparation: S.O.R., A.L.H.L., S.A., B.R., U.A., K.G. and B.S.P. E.A., A.A.W., S.S. and S.A.C. prepared the manuscript with input from co-authors.

Funding: This research is supported by the National Medical Research Council (NMRC), Singapore STaR awards (NMRC/STaR/0029/2017), NMRC Centre Grant to the NHCS, MOH-CIRG18nov-0002, Goh Foundation, Tanoto Foundation, NMRC CIRG14nov021. S.A. is supported by the NMRC (NMRC/STaR/020/2013, NMRC/MOHAFCAT2/2/08, TCR15Jun006, NMRC/CIRG/1460/2016, and NMRC/CG/M003/2017), by Duke-NUS, BMRC (SPF2014/005), and by Duke-NUS and SingHealth core funding. A.A.W. is supported by NMRC/OFYIRG/0053/2017.

Disclosure statement: S.A.C., S.S., A.A.W. and B.N. are co-inventors on a number of patent applications relating to the role of IL11 in human diseases that include the published patents: WO2017103108, WO2017103108 A2, WO 2018/109174 A2, WO 2018/109170 A2. S.A.C. and S.S. are co-founders and shareholders of Enleofen Bio PTE Ltd, a company (which S.A.C. is a director of) that developed anti-IL11 therapeutics, which was acquired for further development by Boehringer Ingelheim.

Data availability statement

Data are available under GEO accession number (GSE162966).

Supplementary data

Supplementary data are available at *Rheumatology* online.

References

- Allanore Y, Simms R, Distler O *et al.* Systemic sclerosis. *Nat Rev Dis Primers* 2015;1:15002.
- Matucci-Cerinic M, Kahaleh B, Wigley FM. Review: evidence that systemic sclerosis is a vascular disease. *Arthritis Rheum* 2013;65:1953–62.
- Worrell JC, O'Reilly S. Bi-directional communication: conversations between fibroblasts and immune cells in systemic sclerosis. *J Autoimmun* 2020;113: 102526.
- Zeisberg M, Kalluri R. Cellular mechanisms of tissue fibrosis. 1. Common and organ-specific mechanisms associated with tissue fibrosis. *Am J Physiol Cell Physiol* 2013;304:C216–25.
- Shin JY, Beckett JD, Bagirzadeh R *et al.* Epigenetic activation and memory at a *TGFB2* enhancer in systemic sclerosis. *Sci Transl Med* 2019;11: eaaw0790. doi: 10.1126/scitranslmed.aaw0790.
- Denton CP, Ong VH, Xu S *et al.* Therapeutic interleukin-6 blockade reverses transforming growth factor-beta pathway activation in dermal fibroblasts: insights from the faSScinat clinical trial in systemic sclerosis. *Ann Rheum Dis* 2018;77:1362–71.
- Denton CP, Merkel PA, Furst DE *et al.*; Cat-192 Study Group; Scleroderma Clinical Trials Consortium. Recombinant human anti-transforming growth factor β 1 antibody therapy in systemic sclerosis: a multicenter, randomized, placebo-controlled phase I/II trial of CAT-192. *Arthritis Rheum* 2007;56:323–33.
- Varga J, Pasche B. Transforming growth factor β as a therapeutic target in systemic sclerosis. *Nat Rev Rheumatol* 2009;5:200–6.
- Schafer S, Viswanathan S, Widjaja AA *et al.* IL-11 is a crucial determinant of cardiovascular fibrosis. *Nature* 2017;552:110–5.
- Ng B, Dong J, D'Agostino G *et al.* Interleukin-11 is a therapeutic target in idiopathic pulmonary fibrosis. *Sci Transl Med* 2019;11:eaaw1237.
- Lindahl GE, Stock CJ, Shi-Wen X *et al.* Microarray profiling reveals suppressed interferon stimulated gene program in fibroblasts from scleroderma-associated interstitial lung disease. *Respir Res* 2013;14:80.
- Cook SA, Schafer S. Hiding in plain sight: interleukin-11 emerges as a master regulator of fibrosis, tissue integrity, and stromal inflammation. *Annu Rev Med* 2020;71: 263–76.
- Nieminen P, Morgan NV, Fenwick AL *et al.* Inactivation of IL11 signaling causes craniosynostosis, delayed tooth eruption, and supernumerary teeth. *Am J Hum Genet* 2011;89:67–81.
- Widjaja AA, Singh BK, Adami E *et al.* Inhibiting interleukin 11 signaling reduces hepatocyte death and liver fibrosis, inflammation, and steatosis in mouse models of

- nonalcoholic steatohepatitis. *Gastroenterology* 2019;157:777–92.e14.
- 15 Sargent JL, Milano A, Bhattacharyya S *et al.* A TGF β -responsive gene signature is associated with a subset of diffuse scleroderma with increased disease severity. *J Invest Dermatol* 2009;130:694–705.
- 16 Khanna D, Lin CJF, Furst DE *et al.* Tocilizumab in systemic sclerosis: a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Respir Med* 2020;8:963–74.
- 17 Akhmetshina A, Dees C, Pilecky M *et al.* Rho-associated kinases are crucial for myofibroblast differentiation and production of extracellular matrix in scleroderma fibroblasts. *Arthritis Rheum* 2008;58:2553–64.
- 18 Reich N, Maurer B, Akhmetshina A *et al.* The transcription factor Fra-2 regulates the production of extracellular matrix in systemic sclerosis. *Arthritis Rheum* 2010;62:280–90.
- 19 Gerber EE, Gallo EM, Fontana SC *et al.* Integrin-modulating therapy prevents fibrosis and autoimmunity in mouse models of scleroderma. *Nature* 2013;503:126–30.
- 20 Chakraborty D, Šumová B, Mallano T *et al.* Activation of STAT3 integrates common profibrotic pathways to promote fibroblast activation and tissue fibrosis. *Nat Commun* 2017;8:1130.
- 21 Dees C, Tomcik M, Palumbo-Zerr K *et al.* JAK-2 as a novel mediator of the profibrotic effects of transforming growth factor β in systemic sclerosis. *Arthritis Rheum* 2012;64:3006–15.
- 22 O'Reilly S, Ciechomska M, Cant R, van Laar JM. Interleukin-6 (IL-6) trans signaling drives a STAT3-dependent pathway that leads to hyperactive transforming growth factor- β (TGF- β) signaling promoting SMAD3 activation and fibrosis via Gremlin protein. *J Biol Chem* 2014;289:9952–60.
- 23 Piera-Velazquez S, Makul A, Jiménez SA. Increased expression of NADPH oxidase 4 in systemic sclerosis dermal fibroblasts: regulation by transforming growth factor β . *Arthritis Rheumatol* 2015;67:2749–58.
- 24 Widjaja AA, Dong J, Adami E, Viswanathan S, Ng B, Singh BK. Redefining Interleukin 11 as a regeneration-limiting hepatotoxin. *bioRxiv*, doi: 10.1101/830018, 4 November 2019, preprint: not peer reviewed.