

2. M'barek L, Sakka S, Meghdiche F, et al. MTHFR (C677T, A1298C), FV Leiden polymorphisms, and the prothrombin G20210A mutation in arterial ischemic stroke among young tunisian adults. *Metab Brain Dis.* 2021;36(3):421–428.
3. Schneider S, Kornejeva A, Vibo R, Kõrv J. Risk factors and etiology of young ischemic stroke patients in Estonia. *Stroke Res Treat.* 2017;2017:8075697.
4. Li F, Yang L, Yang R, et al. Ischemic stroke in young adults of Northern China: characteristics and risk factors for recurrence. *Eur Neurol.* 2017;77:115–122.
5. Pezzini A, Grassi M, Lodigiani C, et al. Italian Project on Stroke in Young Adults (IPSYS) Investigators. Predictors of long-term recurrent vascular events after ischemic stroke at young age. *Circulation.* 2014;129(16):1668–1676. doi:10.1161/circulationaha.113.005663
6. M'barek L, Sakka S, Meghdiche F, et al. Traditional risk factors and combined genetic markers of recurrent ischemic stroke in adults. *J Thromb Haemost.* 2021;19(10):2596–2604. doi:10.1111/jth.15448
7. Zheng S, Yao B. Impact of risk factors for recurrence after the first ischemic stroke in adults: a systematic review and meta-analysis. *J Clin Neurosci.* 2019;60:24–30.

Received: 14 October 2021 | Accepted: 18 October 2021

DOI: 10.1111/jth.15580

Persistent endotheliopathy in the pathogenesis of long COVID syndrome: Comment from von Meijenfeldt et al.

To the Editor

With great interest we read the recent brief report by Dr. Fogarty and colleagues providing evidence for persistent endothelial cell activation in convalescent COVID-19 patients.¹ The authors showed that plasma levels of von Willebrand factor (VWF), factor VIII (FVIII), and soluble thrombomodulin were elevated at a median of 68 days after initial COVID-19 symptom resolution or discharge from the hospital. Interestingly, not all these patients experienced severe disease. Levels of endothelial injury markers were associated with long-term symptoms, including dyspnoea, fatigue, and concentration impairment, often referred to as long COVID. The authors intriguingly suggest that endothelial injury contributes to the pathogenesis of this syndrome. In accordance with Fogarty et al. we have previously shown sustained hemostatic alterations in plasma of patients 4 months after hospital admission with COVID-19 infection.² Here, we would like to share follow-up data from this cohort at 8 and 12 months after initial COVID-19, and associate the hemostatic status at these time points with functional impairment post-COVID-19 to assess whether hemostatic changes may indeed be linked to long COVID. We included 44 patients from the COMMUNITY (COVID-19 Immunity) study who were admitted to Danderyd Hospital, Stockholm, Sweden, with COVID-19 between April and June 2020, as described previously.³ Blood samples were drawn on admission and at 4, 8, and 12 months after hospital discharge. We performed various assays to assess the hemostatic status of these patients, including plasma levels of hemostatic proteins, a thrombin generation assay, and a functional assay of fibrinolysis. These methods and other specifics of this cohort have been described elsewhere.³ We compared the hemostatic status of patients at 8- and 12-month follow-up with that on admission, and at

the 4-month follow-up (the latter have been published previously²). We included previously described reference values of the various tests performed established in 29 healthy controls. To assess functional impairment post-COVID-19, the Sheehan Disability Scale was used, and determined by questionnaires at the 8- and 12-month follow-ups. Participants were asked to score the impact of their symptoms on work life, social life, and family life. Functional impairment was scored as follows; 0: none, 1–3: mild, 4–6: moderate, 7–10: marked. All participants gave informed consent and the study was approved by the Stockholm Ethical Review Board.

Table 1 displays the hemostatic status of COVID-19 patients on admission and 4, 8, and 12 months after hospital discharge. D-dimer levels that are known to be markedly elevated during COVID-19 infection declined to comparably low levels at 4-, 8-, and 12-month follow-up, although levels remained numerically higher than in healthy controls. VWF and FVIII also decreased over time and were comparable to levels in healthy controls 12 months after hospital discharge. Plasminogen activator inhibitor type 1 (PAI-1) levels were substantially elevated on admission and at 4 months after discharge,³ and, despite a gradual decline, remained significantly elevated both 8 and 12 months after hospital discharge compared to levels found in healthy controls, which may be related to obesity, which is known to be associated with elevated PAI-1 levels.⁴ In addition, clot lysis time, which is determined in part by PAI-1 plasma levels, remained significantly prolonged at 8- and 12-month follow-ups compared to healthy controls. Thrombin generation potential was still elevated at 8 months post-hospital discharge, but was comparable to healthy controls at 12 months. Interestingly, a recent study using rotational thromboelastometry to assess clot formation and fibrinolysis showed normalization of these parameters at 6 months after discharge of the intensive care for COVID-19.⁵

TABLE 1 Hemostasis tests in COVID-19 patients on admission and at 4-month, 8-month, and 12-month follow-up compared to healthy controls

	Healthy controls (n = 29)	COVID-19 patients on admission (n = 44)	COVID-19 patients 4-month follow-up (n = 44)	COVID-19 patients 8-month follow-up (n = 44)	COVID-19 patients 12-month follow-up (n = 44)	P-value Healthy controls vs. 8-month/1-year follow-up
Standard hemostasis tests						
D-dimer (ng/mL)	290 [205–445]	1110 [640–1910] ^{a,b,c}	390 [313–510]	420 [258–555]	360 [295–605]	.03/.04
Additional hemostasis tests						
Factor VIII (%)	136 [114–157]	218 [162–282] ^{a,b,c}	159 [132–196] ^e	151 [123–187]	149 [125–181]	.03/.06
VWF (%)	108 [83–128]	348 [248–413] ^{a,b,c}	120 [86–153]	118 [88–152]	100 [82–142]	.27/.97
PAI-1 (ng/mL)	0.60 [0.10–0.75]	2.60 [1.83–4.78] ^b	2.80 [0.70–4.90] ^d	1.08 [0.52–2.61]	1.54 [0.77–2.63]	<.01/<.001
Thrombin generation assay						
ETP (nM IIa*min)	606 [422–773]	796 [586–932] ^f	855 [699–980] ^e	760 [625–852] ^f	623 [469–718]	<.01/.69
Peak (nM IIa)	167 [123–215]	215 [169–267]	243 [196–279] ^e	242 [201–269] ^f	192 [153–230]	<.001/.11
Lag time (min)	2.00 [1.67–2.00]	2.65 [2.00–3.00] ^{a,b,c}	2.00 [1.67–2.33] ^{d,e}	2.00 [1.67–2.00]	2.00 [1.67–2.00]	.59/.48
Velocity index (nM IIa/min)	77 [62–112]	97 [67–123] ^{a,b}	134 [94–164] ^e	139 [117–154] ^f	100 [80–131]	<.001/.03
Clot lysis time (min)	66 [62–70]	82 [70–91]	77 [67–85]	77 [69–89]	74 [67–91]	<.001/<.01

Note: The results are presented as median [interquartile range]. Comparisons between patients on admission and follow-up were made using repeated measures analysis of variance. Comparisons with healthy controls were made with the use of the Mann-Whitney U test. A P-value <.05 was considered statistically significant.

Abbreviations: ETP, endogenous thrombin potential; PAI-1, plasminogen activator inhibitor type 1; VWF, von Willebrand factor.

^aSignificant difference between admission vs. 4-month follow-up.

^bSignificant difference between admission vs. 8-month follow-up.

^cSignificant difference between admission vs. 12-month follow-up.

^dSignificant difference between 4-month vs. 8-month follow-up.

^eSignificant difference between 4-month vs. 12-month follow-up.

^fSignificant difference between 8-month vs. 12-month follow-up.

Next, we assessed functional impairment of convalescent COVID-19 patients 8 and 12 months after hospital discharge. Notably, approximately 15% of patients reported marked impairment of their functioning 1 year after initial COVID-19. Overall, the impact of complaints on the three aspects of life decreased over time (Wilcoxon signed ranked test, median [interquartile range (IQR)] Sheehan score 8 vs. 12 months; work life 5 [0–5] vs. 2 [0–5] $P = .007$, social life 3 [0–5] vs. 3 [0–6] $P = .071$, family life 2 [0–6] vs. 2 [0–5] $P = .011$). Hemostatic markers at 8 and 12 months were not associated with severity of post-COVID complaints in any of the domains of life. In addition, we compared hemostatic markers between patients with none or mild complaints with patients with moderate to marked complaints at 8 and 12 months post-hospital discharge, and found no difference in hemostatic markers between these two groups in any of the domains of life (Table S1 in supporting information).

In conclusion, our results suggest that endothelial cell activation and plasma hypercoagulability assessed with thrombin generation assays persist up to 8 months after hospital discharge, and normalize after 1 year. A hyperfibrinolytic state persists at 12 months, but this may be related to obesity. In this study, hemostatic markers were not associated with functional impairment scored by patients, which contrasts with the data from Fogarty and colleagues, which were scored at a median of 68 days after symptom resolution or hospital discharge. We determined functional impairment scores at 8 and 12 months after initial disease. Functional impairment scores at 4 months were not available in our cohort, nor did we perform a 6-min walking test as described by Fogarty et al. We concur that ongoing endothelial cell activation and hypercoagulability may contribute to long COVID, although our data indicate that larger and more detailed investigations into factors that contribute to long COVID are warranted.

FUNDING INFORMATION

Region Stockholm; Knut & Alice Wallenberg foundation

CONFLICTS OF INTEREST

The authors have no conflicts of interest to disclose.

AUTHOR CONTRIBUTIONS

Conception and design: FM, CT, TL; patient inclusion: SH, CT; data acquisition: SH, CT; laboratory analyses: JA; analysis: FM, TL; interpretation: FM, SH, JA, CT, TL; supervision: CT, TL; drafting of manuscript: FM, TL; revision of the manuscript: SH, CT.

Fien A. von Meijenfeldt¹ 

Sebastian Havervall²

Jelle Adelmeijer¹

Charlotte Thalin²

Ton Lisman¹ 

¹*Surgical Research Laboratory, Department of Surgery, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands*

²*Division of Internal Medicine, Department of Clinical Sciences, Danderyd Hospital, Karolinska Institutet, Stockholm, Sweden*

Correspondence

Ton Lisman, Department of Surgery, University Medical Center Groningen, BA33, Hanzeplein 1, 9713 GZ Groningen, The Netherlands.
Email: j.a.lisman@umcg.nl

ORCID

Fien A. von Meijenfeldt  <https://orcid.org/0000-0003-0274-2414>

Ton Lisman  <https://orcid.org/0000-0002-3503-7140>

REFERENCES

1. Fogarty H, Townsend L, Morrin H, et al. Persistent endotheliopathy in the pathogenesis of long COVID syndrome. *J Thromb Haemost.* 2021;19(10):2546-2553.
2. von Meijenfeldt FA, Havervall S, Adelmeijer J, et al. Sustained prothrombotic changes in COVID-19 patients 4 months after hospital discharge. *Blood Adv.* 2021;5(3):756-759.
3. von Meijenfeldt FA, Havervall S, Adelmeijer J, et al. Prothrombotic changes in patients with COVID-19 are associated with disease severity and mortality. *Res Pract Thromb Haemost.* 2020;5(1):132-141.
4. Dietrich K, Ball GD, Mitchell LG. Increased plasminogen activator inhibitor results in a hypofibrinolytic state in adolescents with obesity: in vivo and ex vivo evidence. *Br J Haematol.* 2016;175(2):300-307.
5. Hulshof AM, Braeken DCW, Ghossein-Doha C, et al. Hemostasis and fibrinolysis in COVID-19 survivors 6 months after intensive care unit discharge. *Res Pract Thromb Haemost.* 2021;5(6):e12579.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.