


BMJ Open Effectiveness and safety of manual acupuncture therapy in patients with post-stroke depression: a protocol for a systematic review and meta-analysis

Wei Liu ,^{1,2} Chang Rao,^{1,2} Qi Zhao,^{1,2} Yuzheng Du,^{1,2} Xi Nan,^{1,2} Zefang Li,^{1,2} Chunsheng Yin^{1,2}

To cite: Liu W, Rao C, Zhao Q, *et al*. Effectiveness and safety of manual acupuncture therapy in patients with post-stroke depression: a protocol for a systematic review and meta-analysis. *BMJ Open* 2021;**11**:e048686. doi:10.1136/bmjopen-2021-048686

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-048686>).

WL, CR and QZ contributed equally.

Received 04 January 2021
Accepted 29 October 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹First Teaching Hospital of Tianjin University of Traditional Chinese Medicine, Acupuncture Department, Tianjin, China

²National Clinical Research Center for Chinese Medicine Acupuncture and Moxibustion, Tianjin, China

Correspondence to

Dr Yuzheng Du;
druyuzheng@163.com

ABSTRACT

Introduction Acupuncture is widely used on the rehabilitation of stroke survivors, including hemiplegia, constipation, emotional disorders and so on. Although the effectiveness of manual acupuncture therapy on post-stroke depression (PSD) has been confirmed by multiple randomised controlled trials, there were few meta-analyses focused on the connection between different techniques, durations or other detailed operations of manual acupuncture and their effectiveness of improving the depression severity and quality of life for patients with PSD.

Methods and analysis A systematic search will be performed on English databases (PubMed, the Cochrane Library, Medline and Embase), Chinese databases (China National Knowledge Infrastructure, WanFang Data, VIP and Chinese biomedical databases) and Japanese databases (Japan Science and Technology Agency and CiNii). The retrieval time limit will be from the establishment of the database to November 2020. Two researchers will independently screen the literature works, extract data and evaluate the quality of the included studies. Meta-analysis will be conducted by using STATA V.14.0 and Review Manager V.5.3.

Ethics and dissemination The results of this meta-analysis will be disseminated through publication in peer-reviewed journals or conference presentations. The data used in this meta-analysis will not contain individual patient data; therefore, ethical approval is not required.

PROSPERO registration number CRD42020222825.

INTRODUCTION

Stroke is currently the second leading cause of death worldwide, the burden of which has increased substantially over the past few decades due to expanding population numbers and aging as well as the increased prevalence of modifiable stroke risk factors.^{1,2} Depression is a common and recurrent psychiatric disorder that starts shortly after stroke and affects patients in the long term. A meta-analysis of the frequency of depression after stroke shows that approximately one-third of stroke survivors experience depression at any time point in the first year.³ Depression

Strengths and limitations of this study

- To the best of our knowledge, this study is the first meta-analysis especially focused on the effectiveness of manual acupuncture therapy for patients with post-stroke depression.
- Compared with previous studies, we will extract more detailed information on the treatment schedule of acupuncture (acupoints selection, twist technique, retention time, frequency, etc) in order to provide more analytical basis for subgroup analysis and sensitivity analysis.
- The electronic search will only include randomised controlled trials published in English, Chinese and Japanese that could limit the inclusion of studies.

after stroke is independently associated with poor health outcomes, including increasing mortality, disability, anxiety and lowering quality of life (QoL).⁴ In addition, there is a two-way relationship between depression and stroke: stroke could increase the risk of post-stroke depression (PSD); meanwhile, depression is an independent risk factor for stroke and stroke mortality.^{5,6}

This bidirectional relationship makes it more difficult to develop the treatment of PSD. Currently, few guidelines mentioned the assessment, treatment or prevention for it.⁷ For depressive disorder, Canadian Network for Mood and Anxiety Treatments, the American Psychiatric Association and the World Federation of Societies of Biological Psychiatry guidelines supported that selective serotonin uptake inhibitors (SSRIs) could be used as first-line treatment.^{8–10} But the pharmacotherapy of PSD needs to be more cautious, as some studies^{11–13} showed that the use of SSRI may relate to the potential risk of haemorrhagic stroke.

Acupuncture, a historic complementary therapy from China, has potential beneficial effects on improving dependency, global

neurological deficiency, and some specific neurological impairments for people with stroke in the convalescent stage.¹⁴ In the treatment of depression, a recent meta-analysis¹⁵ suggests that acupuncture combined with antidepressant medication is effective for the treatment of depression and has an early onset of action, safe and well tolerated over the first 6-week treatment period. However, few systematic reviews or meta-analysis focused on the effectiveness of acupuncture in treating PSD, although the number of papers related to this area has an upward trend recently.^{16 17}

Besides, there are no meta-analysis focusing on the effectiveness of manual acupuncture on improving the depression severity and QoL of patients post stroke, so far. What's more, there is a general problem of high heterogeneity in existing meta-analysis. One recent meta-analysis¹⁸ showed that the curative effect of acupuncture for post-stroke cognitive impairment may be related to manipulation and retention time; however, most of the existing meta-analysis on PSD did not conduct subgroup analysis for such content due to the lack of attention to the details of acupuncture treatment. Therefore, we considered that the higher heterogeneity may be relevant with the difference in the type of acupuncture (manual acupuncture, electroacupuncture, dry needle, etc) and the treatment schedule (acupoints selection, twist technique, retention time, frequency, etc). Hence, we would like to extract the detailed description of manual acupuncture treatment in the included articles and conduct subgroup analysis according to them.

Objectives

The primary purpose of this meta-analysis is to examine the efficacy of manual acupuncture in improving depression severity in individuals with PSD. Secondary aims are to evaluate its role in enhancing QoL and to assess the safety of this treatment.

METHODS AND ANALYSIS

This systematic review protocol has registered in PROSPERO. It will follow the new edition of the Cochrane handbook for systematic reviews of interventions¹⁹ and be reported according to the Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols.²⁰

Criteria for considering studies for this review

Types of studies

Randomised controlled trials (RCTs) in English, Chinese and Japanese will be included. Animal studies or studies with incomplete data will be excluded.

Participants

We will include patients who suffered PSD. The diagnosis of stroke should be based on CT, MRI or clinical criteria. Meanwhile, depression should be diagnosed according to the International Classification of Diseases Tenth Edition, the Diagnostic and Statistical Manual of Mental Disorders,

Chinese Classification of Mental Disorders or Hamilton Depression Rating Scale (HAMD).^{21–23}

Types of interventions

The relevant RCTs will be included if the following criteria were met: (1) using manual acupuncture alone, or in combination with another rehabilitation therapy, or in combination with pharmacotherapy in experiment group (EG); (2) using rehabilitation therapy other than manual acupuncture, pharmacotherapy, sham acupuncture or no treatment in control group (CG). In addition, other kinds of acupuncture therapies, such as electroacupuncture, dry needle, laser needle or acupoint injection, could not be used as interventions in EG or CG.

Types of outcomes measures

Primary outcomes

Depression severity

It is evaluated mainly by HAMD, Montgomery-Asberg Depression Rating Scale (MADRS), Beck Depression Inventory (BDI), Zung Self-Rating Depression Scale (SDS), etc.

If the included studies used two or more of above scales, we will give preference to clinician-rated scales. Following hierarchy will be applied¹: HAMD,² MADRS,³ BDI,⁴ SDS and⁵ all other depression scales.

Secondary outcomes

1. QoL: evaluated mainly by the Medical Outcomes Study Short Form 36, the Stroke Specific Quality of Life Scale or the WHO Quality of Life Scale.
2. Safety: evaluated mainly by the total numbers and severity of adverse events.

Search methods for identification of studies

The following 10 databases will be searched from establishment to November 2020: PubMed, the Cochrane Library, Medline, Embase, Japan Science and Technology Agency, CiNii (National Institute of Informatics), China National Knowledge Infrastructure, WanFang Data, VIP and Chinese Biomedical Databases. The combination of free words and medical subject headings, including 'depression, depressive disorder, acupuncture therapy, acupuncture, needle, needling, stroke, etc', will be used as the retrieval mode. The search strategy for the Cochrane Library is presented in [table 1](#).

Study selection and data extraction

EndNote V.X8.2 will be used to manage studies. First, duplicate literature will be excluded by electronic-based and manual-based steps in EndNote. Second, two reviewers will independently screen the titles and abstracts and select the studies which meet the eligibility criteria. If there are disagreements, the third reviewer will be consulted. The evaluators will read the full text of the included literature, and then preliminarily extracted relevant data, mainly including the following information: (1) inclusion and exclusion criteria; (2) the number of included samples (total number of cases, the number

Table 1 Search strategy for the Cochrane Library

ID	Search
#1	MeSH descriptor: [Stroke] explode all trees
#2	poststroke or post-stroke. ti,ab,kw
#3	MeSH descriptor: [Cerebral Infarction] explode all trees
#4	MeSH descriptor: [Cerebral Hemorrhage] explode all trees
#5	#1 or #2 or #3 or #4
#6	MeSH descriptor: [Depression] explode all trees
#7	depress* or affective disorder or affective symptoms. ti,ab,kw
#8	#6 or #7
#9	MeSH descriptor: [Acupuncture] explode all trees
#10	acupunctur* or acupoint* or needl*. ti,ab,kw
#11	#9 or #10
#12	#5 and #8 and #11

of cases in the treatment group and the number of cases in the CG); (3) grouping method and process; (4) basic data of the included research samples (mainly including gender, age and disease); (5) the intervention of the treatment group and the CG: (a) the treatment method, drug dose, treatment frequency, course of treatment, etc and (b) a detailed description of manual acupuncture treatment, including acupoints selection, twist technique, retention time, frequency, etc(6) and evaluation of the final research results (including the treatment efficiency of different treatment measures, the scale score at the beginning and end point, etc).

Quality assessment

The quality of evidence for main outcomes will be assessed by the Grades of Recommendation, Assessment, Development, and Evaluation (GRADE) approach. Two reviewers will do this independently through GRADEpro Guideline Development Tool. GRADE approach provides guidance for rating quality of evidence and grading strength of recommendations for healthcare. It has important implications for those summarising evidence for systematic reviews.²⁴ It assessed a body of evidence by referring to the concepts of the GRADE system, and determined and recorded the quality of a body of evidence for each clinical question; there are four quality levels: high, moderate, low and very low.²⁵

Assessment of heterogeneity, sensitivity analysis and subgroup analysis

We will use the I^2 statistic to assess the heterogeneity. If the I^2 value is below 50%, the fixed effect model will be used. Otherwise, sensitivity analysis will be conducted to explore the main sources of heterogeneity, after which, the random effect model will be used if the I^2 is still equal or greater than 50%. Both types of effect sizes will be presented with 95% CIs, and p values <0.05 will be regarded as statistically significant.

Meanwhile, subgroup analysis will also be conducted to explore the main sources of heterogeneity. Compared with previous studies, we will extract more detailed information on the treatment schedule of acupuncture, which could provide us more analytical basis for subgroup analysis. If the necessary information is available, subgroup analyses will be carried out according to certain factors (acupoints selection, twist technique, retention time, frequency, period of treatment and different types of CG). After grouping, two or more groups of studies will be analysed and compared in order to explore the causes of high heterogeneity.

Data synthesis

Continuous outcomes will be calculated as mean differences (MDs) or standardised mean differences (SMDs). If different scales are used to measure continuous outcomes, like depression severity and QOL, SMD will be used as a measure of effect size in efficacy outcome. It is calculated as the difference in mean outcome between groups divided by the SD of outcome among participants. If the same scale is used in the included literature, MD will be used. In addition, safety outcome will be the number of participants who dropped out due to adverse effects and the number of participants who reported at least one adverse event or effect. For these dichotomous outcomes, the OR will be calculated as the effect estimate.

Assessment of the risk of bias in individual studies

According to Cochrane Handbook for Systematic Reviews of Interventions, version 6 (<https://training.cochrane.org/handbook/current/chapter-08>), the risk of bias 2.0 (ROB 2.0) tool will be used to mean the methodological quality and the risk of bias of the included studies. One researcher assessed the ROB of included studies by using ROB 2.0 and another researcher confirmed the judgement. If there are any differences, the third researcher will be asked to solve the problem.

Publication bias

STATA V.14.0 will be used to evaluate publication bias. Begg's test and Egger's test will be used to assess the publication bias of the included trials and form the publication bias plot.

Patient and public involvement

It was not appropriate or possible to involve patients or the public in the design, or conduct, or reporting, or dissemination plans of our research.

DISCUSSION

This meta-analysis will focus on the different techniques, durations or other detailed operations of manual acupuncture applied in patients with PSD to explore their influence on depression severity and QoL. PSD are generally more disabled.²⁶ As an important part of traditional Chinese medicine, acupuncture plays an important role in clinical treatment.²⁷ In terms of clinical efficacy, acupuncture can assist in eliminating negative emotions

by significantly improving the functional communication and language function,²⁸ cognitive²⁹ and limb movement function³⁰ of stroke patients. At the mechanism level, acupuncture can modulate glutamate receptor and excitatory amino acid transporter expression,³¹ down-regulated the levels of unclear factor kappa light chain enhancer of activated B cells protein, inducible nitric oxide synthase and nitric oxide,³² so as to achieve the purpose of relieving PSD.

Ethics and dissemination

The results of this meta-analysis will be disseminated through publication in peer-reviewed journals or conference presentations. The data used in this meta-analysis will not contain individual patient data; therefore, ethical approval is not required.

Contributors WL and CR conceived, designed and wrote this protocol. WL provided a clinical perspective, especially to the manual acupuncture. QZ provided the writing and modification of part of the article. YD is the guarantor of this review and approved the final manuscript. XN, ZL and CY provided a preliminary data retrieval.

Funding This work was supported by National Key Research and Development Project (grant number: 2019YFC0840709).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Wei Liu <http://orcid.org/0000-0001-7974-849X>

REFERENCES

- GBD 2015 Neurological Disorders Collaborator Group. Global, regional, and national burden of neurological disorders during 1990–2015: a systematic analysis for the global burden of disease study 2015. *Lancet Neurol* 2017;16:877–97.
- Katan M, Luft A. Global burden of stroke. *Semin Neurol* 2018;38:208–11.
- Hackett ML, Pickles K. Part I: frequency of depression after stroke: an updated systematic review and meta-analysis of observational studies. *International Journal of Stroke* 2014;9:1017–25.
- Ayerbe L, Ayis S, Crichton S, et al. The long-term outcomes of depression up to 10 years after stroke; the South London stroke register. *Journal of Neurology, Neurosurgery & Psychiatry* 2014;85:514–21.
- Villa RF, Ferrari F, Moretti A. Post-Stroke depression: mechanisms and pharmacological treatment. *Pharmacol Ther* 2018;184:131–44.
- Bartoli F, Lillia N, Lax A, et al. Depression after stroke and risk of mortality: a systematic review and meta-analysis. *Stroke Res Treat* 2013;2013:1–11.
- Towfighi A, Ovbiagele B, El Husseini N, et al. Poststroke depression: a scientific statement for healthcare professionals from the American heart Association/American stroke association. *Stroke* 2017;48:e30–43.
- Kennedy SH, Lam RW, McIntyre RS, et al. Canadian network for mood and anxiety treatments (CANMAT) 2016 clinical guidelines for the management of adults with major depressive disorder: section 3. pharmacological treatments. *Can J Psychiatry* 2016;61:540–60.
- Practice guideline for the treatment of patients with major depressive disorder (revision). American psychiatric association. *Am J Psychiatry* 2000;157:1–45.
- Bauer M, Pfennig A, Severus E, et al. World Federation of societies of biological psychiatry (WFSBP) guidelines for biological treatment of unipolar depressive disorders, part 1: update 2013 on the acute and continuation treatment of unipolar depressive disorders. *World J Biol Psychiatry* 2013;14:334–85.
- Hackam DG, Mrkobrada M. Selective serotonin reuptake inhibitors and brain hemorrhage: a meta-analysis. *Neurology* 2012;79:1862–5.
- Shin D, Oh YH, Eom C-S, et al. Use of selective serotonin reuptake inhibitors and risk of stroke: a systematic review and meta-analysis. *J Neurol* 2014;261:686–95.
- Biffi A, Scotti L, Corrao G. Use of antidepressants and the risk of cardiovascular and cerebrovascular disease: a meta-analysis of observational studies. *Eur J Clin Pharmacol* 2017;73:487–97.
- Yang A, HM W, Tang JL. Acupuncture for stroke rehabilitation. *Cochrane Database Syst Rev* 2016;2016:CD004131.
- Chan Y-Y, Lo W-Y, Yang S-N, et al. The benefit of combined acupuncture and antidepressant medication for depression: a systematic review and meta-analysis. *J Affect Disord* 2015;176:106–17.
- Li X-B, Wang J, Xu A-D, et al. Clinical effects and safety of electroacupuncture for the treatment of post-stroke depression: a systematic review and meta-analysis of randomised controlled trials. *Acupunct Med* 2018;36:284–93.
- Chan Y-Y, Lo W-Y, Yang S-N, et al. The benefit of combined acupuncture and antidepressant medication for depression: a systematic review and meta-analysis. *J Affect Disord* 2015;176:106–17.
- Liu W, Rao C, Du Y. The effectiveness and safety of manual acupuncture therapy in patients with poststroke cognitive impairment: a meta-analysis. *Neural Plasticity* 2020;2020:1–15.
- Cumpston M, Li T, Page MJ, et al. Updated guidance for trusted systematic reviews: a new edition of the Cochrane Handbook for systematic reviews of interventions. *Cochrane Database Syst Rev* 2019;10:ED000142.
- Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1.
- Sheehan DV, Lecrubier Y, Sheehan KH, et al. The Mini-International neuropsychiatric interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry* 1998;59 Suppl 20:22–33.
- Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry* 1960;23:56–62.
- Zou Y-Z, Cui J-F, Han B, et al. Chinese psychiatrists views on global features of CCMD-III, ICD-10 and DSM-IV. *Asian J Psychiatry* 2008;1:56–9.
- Guyatt GH, Oxman AD, Schünemann HJ, et al. Grade guidelines: a new series of articles in the Journal of clinical epidemiology. *J Clin Epidemiol* 2011;64:380–2.
- Yoshida M, Kinoshita Y, Watanabe M, et al. JSGE clinical practice guidelines 2014: Standards, methods, and process of developing the guidelines. *J Gastroenterol* 2015;50:4–10.
- Blöchl M, Meissner S, Nestler S. Does depression after stroke negatively influence physical disability? A systematic review and meta-analysis of longitudinal studies. *J Affect Disord* 2019;247:45–56.
- Robinson N, Lorenc A, Ding W, et al. Exploring practice characteristics and research priorities of practitioners of traditional acupuncture in China and the EU-A survey. *J Ethnopharmacol* 2012;140:604–13.
- Zhang B, Han Y, Huang X, et al. Acupuncture is effective in improving functional communication in post-stroke aphasia: A systematic review and meta-analysis of randomized controlled trials. *Wien Klin Wochenschr* 2019;131:221–32.
- Wang S, Yang H, Zhang J, et al. Efficacy and safety assessment of acupuncture and nimodipine to treat mild cognitive impairment after cerebral infarction: a randomized controlled trial. *BMC Complement Altern Med* 2016;16:361.
- Yang A, HM W, Tang JL. Acupuncture for stroke rehabilitation. *Cochrane Database Syst Rev* 2016;2016:CD004131 <https://pubmed.ncbi.nlm.nih.gov/27562656/>
- Tu C-H, MacDonald I, Chen Y-H. The effects of acupuncture on glutamatergic neurotransmission in depression, anxiety, schizophrenia, and Alzheimer's disease: a review of the literature. *Front Psychiatry* 2019;10:14.
- Pu R, Shao R-H, Lu J, et al. [Acupuncture alleviated depressive-like behavior by down-regulating nf-kb, inos and no levels in prefrontal cortex of depression rats]. *Zhen Ci Yan Jiu* 2018;43:226–30.