


Priorities and Preferences of People Living with Dementia or Cognitive Impairment – A Systematic Review

Hannah Wehrmann¹
Bernhard Michalowsky¹ 
Simon Lepper¹
Wiebke Mohr¹
Anika Raedke¹
Wolfgang Hoffmann^{1,2}

¹Translational Health Care Research, German Center for Neurodegenerative Diseases (DZNE) Site Rostock/Greifswald, Greifswald, Germany; ²Epidemiology of Health Care and Community Health, Institute for Community Medicine & University Medicine Greifswald (UMG), Greifswald, Germany

Background: Knowledge about the priorities and preferences of people living with dementia (PwD) might help to individualize treatment, care, and support, which could improve patient-related outcomes. This study aimed to summarize preferences of PwD or people with mild cognitive impairment (MCI), considering all relevant aspects of health care and everyday life.

Methods: We conducted a systematic literature review and included studies about patient preferences published in English between January 1, 1990 and October 28, 2019. The inclusion criteria were that preferences were elicited directly by PwD or patients with MCI. We used the International Consortium for Health Outcomes Management value set for dementia to categorize the preferences into the following topics: a) clinical status, b) symptoms, functioning, and quality of life, and c) sustainability of care.

Results: Of 578 initially identified studies, 45 met the inclusion criteria. Patients preferred to be informed about the diagnosis as early as possible, especially for anticipatory care planning. They ranked caregiver quality of life as their highest priority. They preferred not to be a burden to others more than their caregivers' mood, their own functional status, or their own distressing behaviors. Furthermore, PwD are eager to participate in medical decisions, especially in those about creating an everyday life routine. PwD preferred their own quality of life, self-efficacy, and emotional well-being. Institutionalized PwD preferred individualized and person-centered care. According to the sustainability of care, PwD preferred to maintain close bonds with their family at the end of their life and wanted to be treated with empathy.

Conclusion: This systematic review provides essential insights into cognitively impaired patients' preferences, which are rarely considered in treatment, care, and support services. Further studies should evaluate whether considering preferences in treatment and care or daily living can improve patient-reported outcomes.

Keywords: dementia, patient preference, patient outcome assessment, decision-making

Introduction

Dementia represents one of the most significant public health challenges. Worldwide, approximately 50 million people live with dementia.^{1,2} A diagnosis as early as possible is essential to initiate evidence-based treatment and care and to better cope with the disease. The progressive nature of the disease leads to declining self-responsibility levels, self-determination, and autonomy, which are associated with an increasing need for care. Decisional capacity steadily decreases, but persons with dementia (PwD) wish to be acknowledged in all disease stages and aspects of care, make individual decisions, and be involved in decisions about treatment and care, daily living, and support.^{3–7}

Correspondence: Bernhard Michalowsky
Tel +49 3834 86 75 07
Email bernhard.michalowsky@dzne.de

Dementia care should promote patient autonomy and be person-centered and, thus, be preference-based, respecting the patient's values and the need to improve the patient's sense of self-efficiency.^{8,9} It is also important to offer patients a purpose in life, open up perspectives for them, support them, and transfer responsibility to them.¹⁰ Furthermore, informal caregivers need to be considered.¹¹ In the progression of the diseases, many PwD experience emotional changes and difficulties with how to express and manage their feelings, which can lead to more intense emotional reactions, which can be a great challenge for caregivers.¹² Additionally, it can be a challenge to meet the expectations of PwD when the share of responsibilities changes between the person with dementia and his or her caregiver.¹³

It is often assumed that informal caregivers know the preferences of PwD best. However, some studies found that more than one-third of caregivers could not predict PwD preferences correctly.¹⁴ Most caregivers try to compensate for the decreasing abilities to support the patient's well-being but might sometimes project their own preferences instead of putting themselves into the position of the PwD.^{14,15} Furthermore, caregivers' preference prediction for PwD can subconsciously be influenced by how the caregivers' own life is affected by certain decisions.⁷ Such preference divergences have also been elicited between nursing care staff and patients. Dementia patients in long-term care appreciate autonomy, whereas staff assumed safety and a homey ambience were the most important, demonstrating the existence of discrepancies between professionals and patients.¹⁶

Knowing preferences for PwD could inform recommendations for dementia care practices, increase adherence to treatment, therapies, and care, and improve and individualize interventions and patient-related outcomes, including health-related quality of life (QoL).¹⁷⁻¹⁹ A systematic review that summarizes the existing qualitative and quantitative evidence about PwD preferences is currently lacking. Therefore, this systematic review aimed to summarize the preferences of PwD, including all relevant domains of treatment, care, support, and everyday life.

Methods

Search Strategy

Studies published in English between January 1, 1990 and October 28, 2019 were identified through searches of the electronic databases PubMed, Science Direct, and Cochrane. Key phrases included "dementia", "Alzheimer", or

"Alzheimer's disease" in combination with "preferences" or "priority". Titles and abstracts were reviewed for dementia terms, whereas only the title was considered for preference terms. In parallel, a search of MeSH terms was conducted in the PubMed database to find all articles that used the terms "mild cognitive impairment" or "dementia" in combination with "patient preference". The systematic literature search process is presented in [Figure 1](#).

Data Extraction

Studies were included if i) patients were diagnosed with dementia, Alzheimer's disease, or mild cognitive impairment or ii) priorities and preferences were directly elicited from PwD or patients with MCI. All studies were included, irrespective of the quantitative or qualitative methods used to elicit preferences. Studies were excluded if they reported animal experiments or if no abstract was available. Additionally, systematic reviews and studies eliciting preferences with hypothetical patients from the general population were excluded. A total of 578 studies were found. After removing 34 duplicates, 544 studies remained to be screened for titles and abstracts by two independent reviewers. A third reviewer was involved in discussing disagreements. Overall, 446 studies were excluded after title and abstract screening, and 98 studies remained for full text screening. After full text screening, an additional 54 studies were rejected, resulting in the inclusion of 44 studies. The study flow chart of the screening procedure is demonstrated in [Figure 1](#).

Data Description

To categorize the heterogeneous studies with respect to their evaluated domains, we used the Standard Set of Dementia Criteria of the International Consortium for Health Outcomes Management (ICHOM).²⁰ Experts and patient representatives developed this standardized set of outcomes to focus on what is most important for PwD. As this systematic review was focused on preferences of PwD and not the disease in general, the Standard Set was modified and adapted. Specifically, the category "disease and progression" was replaced by the categories "treatment and care" and "diagnosis". Given that some studies used modern assistive technology, we added the category "technology". Caregiver QoL was removed, as preferences of PwD and not of their caregivers were the focus of this study. In cases in which study outcomes overlapped across two or more categories, three reviewers discussed the categorization. The final set categorized the elicited preferences of PwD into the following three main domains and subcategories: (A)




Figure 1 Study flow chart (CONSORT). Data from ICHOM standard set for Dementia. Available from: <https://connect.ichom.org/standard-sets/dementia/>.²⁰

clinical status including i) treatment and care and ii) diagnosis, (B) symptoms, functioning, and QoL, including i) cognitive, ii) social, iii) daily living, and iv) overall QoL, and (C) sustainability of care, including i) time to full-time care, ii) end-of-life care, and iii) assistive technology.

Results

Characteristics of the Included Studies

Most of the studies were published in Europe (n = 21) and the USA (n = 14). The ages of the study participants ranged from

57 to 87 years. Sample sizes differed substantially, ranging from two to 2028. An overview, including summaries of the included studies, is presented in Table 1.

Diagnosis

PwD and caregivers emphasized the importance of receiving a formal dementia diagnosis. Several studies found that more than 86% of PwD preferred to be informed about the dementia diagnosis.^{37–39} Those PwD who receive a formal diagnosis show a higher QoL related to

Table 1 Study Characteristics and Results Preference Studies of People with Dementia (PwD)

Area	Author/Year	Population/Country	Aim of Study	Mean Age	Study Description	n	Preferences
Symptoms, Functioning and Quality of Life							
Cognitive	Raez et al 2013 ³⁷	PwD under a 24-hour care by FM or staff, USA	To identify stimuli to ↑ engagement and to ↓ negative effects and problem behavior	82 years	Paired-stimulus preference assessments	8 PwD	<ul style="list-style-type: none"> • The use of a single-array presentation is as effective as the use of a three-array presentation to identify preferences • For some PwD, preferences remain stable
	Wijk et al 1997 ¹¹	PwD recruited from a psychiatric clinic, Sweden	To examine the ability to detect color naming/discrimination	80 years	Descriptive preference assessments	50 PwD	<ul style="list-style-type: none"> • Cognitive decline influenced the naming of mixed colors • preferences for colors are stable despite the disease
	McMurtry et al 2009 ⁷⁴	Japanese PwD who presented at a memory disorder clinic, Hawaii	To determine the influence of dementia on the native speech of bilingual PwD	76 years	Language assessments	2 PwD	<ul style="list-style-type: none"> • A regression to the primary language of bilingual PwD may help clinicians to identify dementia
	Lombrozo et al 2007 ⁷⁴	HA from Boston, PwD from a general hospital, USA	To examine explanatory judgments in PwD	84 years	Explanation choice task, causal-belief task	24 HA, 17 PwD	<ul style="list-style-type: none"> • Unlike HA, PwD prefer teleological explanations
	Halpern et al 2007 ²³	PwD living at home, HA from a senior citizen center, USA	To explore the stability of art preferences	78 years	Comprehensive evaluation and neuropsychological tests	16 PwD, 27 HA	<ul style="list-style-type: none"> • Aesthetic preferences can be preserved in PwD and are not dependent on explicit memory
	Sanzani Maserati et al 2019 ²²	PwD were recruited from to a cognitive disorder center of Bologna, Italy	To explore the emotional state of PwD through color choice preference	77 years	Lüscher colour test	121 PwD, 68 HA	<ul style="list-style-type: none"> • With a higher severity of the disease, auxiliary colors like violet and brown are preferred, whereas black and gray are rejected
	Morzek et al 2017 ⁶³	PwD at a general acute care hospital in Dresden, Germany	To identify picture categories that are liked and easily remembered by PwD	78 years	PwD votes of preferences and the ability to recall pictures were tested	37 PwD	<ul style="list-style-type: none"> • A main effect of familiarity on positive votes and recall pictures was seen
	Feliciano et al 2009 ²⁵	PwD living in 6 different long-term care settings, USA	To examine the use of stimuli in behavioral management	86 years	Preference assessment	11 PwD	<ul style="list-style-type: none"> • Using preferred items in behavioral management ↓ agitated behaviors and partially ↓ depressive symptoms
	Van de Ven et al 2017 ²⁸	PwD living in 23 different care networks in the Netherlands	To describe the decision-making process of PwD, shared decision-making	80 years	Qualitative study, a secondary analysis of 117 interviews	23 PwD, 44 CG, 45 Prof.	<ul style="list-style-type: none"> • Collaborative deliberation model is extended by engaging the network, recognizing the need for a decision, defining/ evaluating decisions, trying out alternatives ...
	Lopis et al 2017 ⁵⁴	PwD were recruited from a local memory center, France	To determine if eye contact influences the likeability/memory relating to faces in HA/PwD	81 years	Identifying faces they had previously seen during a surprise recognition test	19 PwD, 40 HA	<ul style="list-style-type: none"> • PwD show a positive correlation between ratings of likeability and recognition scores
Social							

Daily living	Cohen-Mansfield et al 2010 ³⁰	PwD out of 7 nursing homes in Maryland, USA	To study whether activities/stimuli based on PwD interests result in more engagement than other activities	86 years	The expanded version of the self-identify question-naire ⁷⁵ was used to identify PwD interests	193 PwD	<ul style="list-style-type: none"> • PwD with current interests in art, pets, and music were more engaged with stimuli that were related to these interests than PwD without these interests
	Cohen-Mansfield et al 2009 ⁵⁵	PwD at four nursing homes in Maryland, USA	To examine the influence of stimulus attributes on the engagement of PwD	86 years	I-on-I interview, systematic observations via OME	69 PwD	<ul style="list-style-type: none"> • Preference for the work-related rather than the manipulative stimuli, preference for small rather than large blocks
	Chong et al 2012 ²⁹	Australian patients	To show PwD attitudes toward physical activity	76 years	Analysis of focus groups, individual interviews	36 PwD, 14 HA	<ul style="list-style-type: none"> • Preferred activity program: simple/light/safe activities, accessible group setting, adjusted to individual income/interest
	Mungas et al 1990 ⁶	PwD at the UC Davis Medical Center, Sacramento, USA	To compare the sweet food cravings of PwD to HA	76 years	A food preference survey	45 PwD, 43 HA	<ul style="list-style-type: none"> • Compared to normal controls, PwD show an increased preference for sugar and sweet foods
	Milte et al 2017 ³²	PwD recruited from rest homes, Australia	To describe the food experience of PwD in rest homes	78 years	I-on-I interview, focus groups with PwD and FM	19 PwD and FM	<ul style="list-style-type: none"> • Maintaining choice and individual preferences around food is important to PwD
	Ortega et al 2017 ⁷	PwD living at home, USA	To find the most preferred item in preference assessments	84 years	Paired-stimulus procedure to assess preferences of 4 edible and 4 leisure items	14 PwD	<ul style="list-style-type: none"> • A leisure activity was the most preferred in the preference assessment
	Van't Leven et al 2019 ³¹	Dutch patients	To test how activating dyadic interventions fit the needs and preferences of PwD and CG	78 years	Qualitative design with semistructured interviews	27 PwD, 34 CG, 19 Prof.	<ul style="list-style-type: none"> • Five factors influenced dyad "fit": timing, need for an activity, lifestyle, apart-on-together, and meaning of (lost) activity

(Continued)

Table 1 (Continued).

Overall quality of life	Bohn et al 2018 ³⁹	PwD living in long-term care facilities, Edmonton area, Canada	To study if PwD prefer emotional gratification, despite their cognitive decline	64 years	Social activity preference card-sort task	75 HA, 23 PwD	<ul style="list-style-type: none"> PwD prefer emotionally meaningful ends and retain high levels of well-being
	Barrios et al 2016 ¹⁸	Couples of the summer 2014 HABIT session at Mayo Clinic, Rochester, USA	Ranking of behavioral outcomes of PwD, pre and post ranking after participation	77 years	Ranking task	16 dyads (PwD +HA)	<ul style="list-style-type: none"> Preferences of PwD in descending order: PwDs QoL, self-efficacy, CGs QoL, depression of CG, CG burden, daily functioning, anxiety of CG
	Smith et al 2018 ³³	HABIT program completers (2008–2014), USA	Outcome and treatment preferences of PwD	73 years	Ranking task	29 PwD, 54 spouses	<ul style="list-style-type: none"> Preferences of PwD in descending order: QoL, self-efficacy, depression, CG QoL, PwDs anxiety/, CG burden
	Shelton et al 2016 ³⁴	American patients recruited from the northern Ohio region, USA	To understand how actual and perceived incongruence of care preferences affects the social well-being of PwD/CG	73 years	In-depth interviews	128 dyads (PwD +CG)	<ul style="list-style-type: none"> Actual incongruence for socioemotional care preferences was a predictor of greater relationship strain and worse mood for PwD, whereas perceived incongruence for socioemotional care preferences was related to lower QoL
	Reamy et al 2011 ³⁵	Dyads from research/service organizations, USA	To find discrepancies in care preferences of PwD and CG	76 years	Interviews	266 dyads (PwD +CG)	<ul style="list-style-type: none"> CG underestimated the PwDs values for autonomy, burden, control, family, and safety
Clinical status							
Treatment	Harrison Denning et al 2016 ²⁶	Dyads were identified by psychiatrists in the UK	To explore if CG and PwD have the same end-of-life care preference	79 years	Cross-sectional study, LSPQ	60 dyads (PwD +CG)	<ul style="list-style-type: none"> CG had low to moderate agreement with PwD on preferences for end-of-life care treatment
	Dickinson et al 2013 ⁴³	PwD identified by health-care prof. in the UK	To investigate views of PwD on ACP		Qualitative study with semistructured interviews	17 PwD, 29 CG	<ul style="list-style-type: none"> 5 barriers to ACP: lack of knowledge, finding the right timing, lack of support, preference for informal plans vs written documentation, lack of choice around future care
	Cohen et al 2019 ⁴⁵	PwD recruited from 64 Boston-area facilities, USA	To examine concordance between advance directives and proxy care preferences among PwD	87 years	A cluster randomized clinical trial	328 dyads (PwD +HA)	<ul style="list-style-type: none"> The most prevalent directive: DNR, least common: antibiotics Concordance between directives and each level of care preference: comfort, 7%, basic, 49%, and intensive, 58%

Area	Author/Year	Population/Country	Aim of Study	Mean Age	Study Description	N	Main Outcome
Diagnosis	Jung et al 2017 ³⁷	PwD visiting the psychiatric outpatient clinic of Seoul University Hospital, Korea	To identify preferences regarding the disclosure of dementia and the awareness of ACP	75 years	Structured questionnaire	98 PwD, 62 FM	<ul style="list-style-type: none"> A disclosure of the dementia diagnosis was favored The majority of PwD also agreed on the necessity of ACP
	Mahieux et al 2018 ³⁹	PwD attending a memory clinic in Paris, France	To report disclosure preferences of PwD	78 years	Prospective single-center study	737 PwD, 268 HA	<ul style="list-style-type: none"> 85% wished to be informed of a dementia diagnosis, 7% did not want to know, and 7% were not sure
	Hellström et al 2012 ⁴²	PwD recruited from two memory clinics in Sweden	To report the understanding and experiences of PwD	71 years	Qualitative interviews	28 PwD, 20 HA, 9 CG	<ul style="list-style-type: none"> Regarding couples, the PwD are the ones deciding about the disclosure of their disease to others
	Elson et al 2005 ³⁸	Patients over 65, UK	To report the disclosure preference of PwD	76 years	Qualitative interviews	32 PwD, 4 HA	<ul style="list-style-type: none"> 2/3 did not know the cause of their memory complaint 70% of the PwD wanted to know the diagnosis
	Bamford et al 2016 ⁴⁰	PwD living in the UK	To assess preferences for diagnostic processes (PET and SPECT), comparison with HA	76 years	Card sorting exercise relating to importance (ranking)	68 PwD, 59 CG, 30 HA	<ul style="list-style-type: none"> PwD prioritized accuracy over other scan characteristics, followed by helpful staff, comfort, and companion
	Walker et al 2017 ⁷²	PwD recruited from the Alzheimer's association, Australia	To show PwDs experience with dementia assessment services	80 years	Qualitative, semistructured in-depth interviews	9 PwD, 7 CG	<ul style="list-style-type: none"> An important contact during a dementia assessment is the GP
	Mate et al 2012 ³⁶	PwD recruited by 169 GPs in Australia	To examine the predictors of QoL of PwD	84 years	Cross-sectional study	167 PwD, 1861 others	<ul style="list-style-type: none"> Satisfaction with GP communication is associated with higher QoL
	Sustainability of Care						
Technology	LaMonica et al 2017 ⁵¹	PwD recruited from Healthy Brain Aging Clinic, Sydney Australia	To show PwD internet use and interest in eHealth technology	68 years	HBA E-Health questionnaire ⁷⁸	160 PwD, 61 HA	<ul style="list-style-type: none"> PwD preferred e-health intervention: memory strategy training
	O'Philbin et al 2018 ²²	PwD were recruited through the Join Dementia research database, UK	To explore preferences of PwD in relation to digital life story work (DLSW)	62 years	CG responded to DCE and PwD completed an online survey	67 CGs, 17 PwD	<ul style="list-style-type: none"> Most preferred setting: individual one-to-one Most preferred use of the DLSW was to share memories
	De Sant'Anna et al 2010 ⁷⁹	PwD recruited from the Broca Geriatric Day Care Hospital in Paris, France	To compare the use of keyboard, mouse pad, and a computer screen of PwD with HA		Pilot study, qualitative and quantitative analyses	8 PwD, 10 HA	<ul style="list-style-type: none"> Basic computers can be accessible to PwD under certain conditions

(Continued)

Table 1 (Continued).

T-t-F-t Care	Dickins et al 2018 ¹⁰	PwD living in Australia	To understand how CG/FM conceptualize the issue of risk	66 years	83 semistructured interviews	53 prof., 20 HA, 7 PwD	<ul style="list-style-type: none"> There is no single approach to risks that can be applied to all PwD
	Groenewoud et al 2015 ⁵⁰	PwD living in the Netherlands	Influences on PwD health-care decisions	57 years	DCE, 11 attributes	421 PwD, 984 others	<ul style="list-style-type: none"> Expertise was the most valued attribute when choosing a health-care provider
	Snebye et al 2016 ⁷	PwD living at home, Norway	To explore ethical dilemmas concerning PwD autonomy	83 years	Qualitative, hermeneutic design	9 PwD, 9 FM, 9 Prof.	<ul style="list-style-type: none"> Three ethical dilemmas: PwD autonomy vs CG need to prevent harm, beneficence of FM/CG, autonomy of FM
EoL care	Ayalon et al 2012 ⁴⁷	Couples recruited at two psychogeriatric clinics in Israel	To evaluate concordance in end-of-life preferences between PwD and their spouses	76 years	Face-to-face interviews	53 couples (PwD +HA)	<ul style="list-style-type: none"> Moderate agreement regarding end-of-life care between PwD and their spouses
	Dening et al 2013 ⁴⁸	PwD and CG attending memory assessment service, UK	To explore PwD preferences for end-of-life care	83 years	Nominal group technique	12 PwD, 11 CG	<ul style="list-style-type: none"> Moderate agreement between PwD and their CG Preferences of PwD in descending order: maintain family links and independence, feel safe, not be a burden
	Hill et al 2017 ¹⁴	PwD were recruited through internet databases like Join Dementia research, UK	To identify the aspects of end-of-life care for PwD	73 years	Q-methodology qualitative and quantitative techniques	14 PwD, 43 CG	<ul style="list-style-type: none"> There is no universal opinion on what is important about end-of-life care for PwD
	Goodman et al 2013 ⁴⁴	PwD living in 6 care homes in the UK	To explore how PwD discuss their priorities and preferences for end-of-life care	85 years	Exploratory, qualitative study, interviews	18 PwD	<ul style="list-style-type: none"> For PwD, the experiences of living and dying in a care home are inextricably linked End-of-life care can be improved by documenting the preoccupations, key relationships, and wishes about everyday care of PwD
	Mulqueen et al 2017 ⁴⁹	PwD living in a residential care facility, Ireland	To explore preferences of PwD for end-of-life care	83 years	Nominal group technique	6 PwD, 6 CG	<ul style="list-style-type: none"> Preferences of PwD in descending order: comfort, family presence, communication, familiar staff and surroundings

Abbreviations: ↑, increase; ↓, decrease; CG, caregiver; PwD, people with dementia; HA, healthy adults; Prof., professionals; OME, observational measurement of engagement; LSPQ, Life Support Preference Questionnaire; T-t-F-t care, time-to-full-time care.

social life and the environment than PwD who have not received a formal diagnosis.³⁶ Another study highlighted that PwD would rather receive a false-positive diagnosis than staying undiagnosed.⁴⁰ Approximately 40% of PwD prefer that their family is present while receiving the diagnosis, and 30% of PwD prefer to be alone.³⁹ However, some PwD also express a feeling of “falling into a black hole” after they receive the diagnosis and explain that it takes time until they receive access to support services.⁴¹

In many relationships, PwD decide on the disclosure of the diagnosis to relatives and friends. Concerning this disclosure, preferences are divergent. Some people want no one to know, others only tell their spouses, some only want their family to know about the disease, and some feel fine with everybody knowing about their diagnosis. PwD express various reasons concerning their wish not to disclose: some never try to talk about it, whereas others feel regret after informing others about their medical condition. Others presume a change in others’ perceptions after disclosure and want to avoid being treated differently.⁴²

Treatment and Care

Studies have shown that it is difficult to identify the right moment to start advanced care planning (ACP). Most PwD have reported that right after the diagnosis is not good timing for ACP. It is often hard for both PwD and their relatives to cope with the new diagnosis. Sometimes this process of developing a coping strategy takes years.⁴³ Generally, willingness to plan is present, but most PwD prefer informal plans over written documentation and trust their family members to make the right decisions. PwD are not aware of all components in ACP. They described a lack of knowledge concerning all the opportunities, including the range of available services. Most PwD are not aware that the progression of dementia in the future might impede them from expressing preferences, eg, the preferred place of care. However, some PwD, for example, plan their funeral, ie, choosing songs and paying for a funeral plan because they do not want their relatives to spend too much money.^{43,44} When PwDs were asked about future scenarios, receiving invasive life-prolonging treatments, eg, cardiopulmonary resuscitation and tube feeding, was less preferred. Concordance between each level of care preference and directives that PwD want to withhold were as follows: comfort care 7%, primary care 49%, and intensive care 58%. Proxies preferred comfort care the most, followed by primary care and intensive care,

concluding that better alignment between preferences for comfort-focused care and advance directives is needed in advanced dementia.^{26,45}

Daily Living, Activities and Social Life

Studies have revealed that PwD have a positive attitude toward physical activity, as evidenced by their endorsing statements that described physical activity as being “important for health”, “good”, “enjoyable”, and “social”. The most preferred ones were “simple/light/safe” and “affordable” exercises with “accessible” settings, preferably in a group. In general, leisure activities were the most preferred. PwD agitation was found to be reduced by including their preferences for leisure activities in behavioral management.²⁵ The use of different kinds of stimuli that reflect past and present interests of PwD helps to increase engagement and responsiveness as well as to support positive behavior.³⁰ Most PwD prefer activities that are suggested by their general practitioner. Exercises should be adapted to individual needs and expectations. “A lack of company” and “memory” prevented PwD from doing exercises.²⁹ Joint-activating interventions to fit both PwD and their informal caregivers were influenced by factors, such as “timing”, “need for activity”, “lifestyle”, “apart or together (with their spouse)”, and “meaning of life”. Many dyads share similar values, such as “keeping active”, “getting out of the house everyday”, and “staying mobile”. Programs helping dyads organize activities and adapt lifestyles are highly appreciated.³¹

In addition to daily living activities, the food preferences of PwD are very heterogeneous and individual. However, PwD want to be involved in related daily life routines and decisions on mealtimes, meal sizes, and food options.³² Halpern et al²³ also found that PwD can state their aesthetic preferences and that these preferences did not change over time and with cognitive decline.²³ According to the design of the living environment, auxiliary colors, such as brown and violet, are most preferred, whereas black and gray are least preferred.^{21,22} Additionally, pictures that patients are able to relate to in terms of familiarity and urban and natural characteristics seem to be suitable for use as environmental cues. Pictures can further enhance the ambiance or serve as prompts for communication and interaction.⁵³

Social interaction has been found to slow down the progression of the disease in PwD.²⁵ It may be challenging to understand what PwD want to express because their verbal expression does not necessarily reflect what they mean and

act upon. Observing PwD behaviors and emotions can yield essential indications. In shared decision-making, PwD sometimes prefer to rely on their caregiver.²⁶ Some claim to be dependent on their caregiver because they cannot make decisions on their own. Additionally, it should be considered that PwD often need a great deal of time for their decisions, which may change over time.²⁷ Researchers have suggested adopting the Model of Collaborative Deliberation for the context of care for PwD by drawing particular attention to recognize and define decision-making.²⁸

According to the method of communication, PwD prefer teleological explanations over mechanical ones. Lombrozo et al²⁴ asked study participants questions, such as “Why are there eyes?”. Participants chose the (teleological) answer “so people and animals can see.”. The mechanistic explanation is “because bodies have special cells that combine to produce eyes”. McMurtray et al⁷⁴ additionally revealed that bilingual PwD relapse more and often return to their primary language during the progression of dementia.

Quality of Life

PwD and caregivers state that PwD QoL and self-efficacy are most important. Functional status, patient and caregiver mood, caregiver burden, and PwD memory performance have been ranked as the most essential aspects of QoL.^{18,33} Incongruence between socioemotional care preferences of both the caregiver and the PwD was found to be associated with lower QoL of PwD. Additionally, perceived incongruence of care preferences was found to negatively influence the mood of PwD and worsen the relationship between caregivers and PwD. Following this, a lack of correlation of preferences can predict a decrease in the QoL of PwD and adverse social and psychological outcomes.³⁴ Many caregivers underestimate preferences of PwD regarding autonomy, being a burden to others, control, family, and safety.³⁵ Generally, both PwD and their spouses have altruistic preferences (ie, they put the other’s needs before their own).^{18,33}

Technology

More than 80% of PwD regularly use a computer. Nearly everyone has a phone, and many of them know how to send text messages. More than 90% have access to the Internet at home, and every fourth person reported facing issues with using it. Only a few PwD use social media, but the use of e-mail is widespread. The majority expressed that a website designed for health issues in old age (eg, tracking physical

and cognitive health conditions) would be beneficial.⁵¹ A “digital life story book” to share memories with others is preferred by PwD.⁵²

Time to Full-Time Care

One of the major concerns for PwD is becoming less independent over time in areas, such as mobility (ie, driving a car), dealing with finances and work, and self-care. PwD try to upkeep as many activities for as long as possible. PwD understand that safety must be ensured either by someone helping them or by task modification. However, they want to have a purpose in life, and meaningful activity is crucial to them. PwD state that they can better cope with the disease when they are more active.¹⁰ Another study focused on the preferences for the selection of health-care services and providers, revealing that expertise is most important for PwD.⁵⁰

End-of-Life Care

More than 50% of PwD did not state their opinion on end-of-life decisions to their family caregivers. However, those PwD who stated their preferences usually had a different opinion than their relatives. The incongruence of preferences occurs, for example, in 48% of cases concerning CPR and tube feeding and in 60% of cases of artificial ventilation.^{46,47} At the end of life, PwD prefer to maintain family links, maintain independence, feel safe, not be a burden, be treated with respect and dignity, have a choice in their place of care, engage in pleasurable activities, experience person-centered care, be in touch with the world, and have comfortable care. However, most important for caregivers was to ensure good quality care at all times.⁴⁸ Hill et al¹⁴ identified four main preferences that are most important for PwD at the end of life: family involvement, living in the present, autonomy, and individuality. If PwD are unable to express their wishes, they prefer their family to make decisions. They highly value having their family and friends close during that time. PwD want to be cared for with compassion and want to be seen as individuals who can maintain hobbies and interests. Preserving their independence in self-care, such as eating and taking medications, as long as possible is of great importance for many PwD.¹⁴

PwD self-esteem and confidence related to expressing opinions are affected by their disease awareness and the related memory loss, which makes many PwD feel as if they have nothing to say that would be worth listening to. The feeling of being understood and well cared for is

important for PwD to entrust themselves to the staff. PwD in care homes at the end of life have stated that it is an excellent experience to live in a care home and values it as “his home” now. Furthermore, losses have a significant impact on whether PwD see a purpose in life or not. Frequently, PwD cannot think about their own care needs because their thoughts are overshadowed by past experiences (eg, a loss of a family member or spouse). Others prefer to return home to their family but know that it is not possible because they would be a burden.⁴⁴ Mulqueen et al⁴⁹ found that nurses do not always precisely predict what patients consider most important. PwD value comfort, family presence, familiar staff, and surroundings most, whereas nurses thought they most preferred good communication, pain management, ACP, and care by professional staff.⁴⁹

Cognitive Ability to State Preferences

Some studies have evaluated whether patients with cognitive decline are able to state their preferences, whether preferences are stable over time instead of associated with cognitive decline, and how cognitive decline affects the communication of preferences. Wijk et al²¹ evaluated the ability of color naming, color discrimination, and color preference in Alzheimer’s disease, revealing that the ability to discriminate and name colors was affected but that preferences for color are stable over time, despite cognitive decline.²¹

Discussion

This systematic literature review summarizes the preferences of PwD from 44 publications with very heterogeneous results, capturing preferences for the diagnosis and disclosure of the disease, aspects that have to be considered in the medical decision-making process physical and everyday life activities, QoL and self-efficiency as well as concordance on care preference and end-of-life care. The review revealed that PwD are able to state their preferences according to these domains and that these preferences are stable over time, even though the cognition of PwD declines and the disease progresses. Additionally, proxy preference ratings (ie, statement of preferences of caregivers for the PwD) have been found to provide useful aspects of PwD preferences, but they are not perfect substitutes for patients’ preferences. As a result, possible differences should be taken into account within decision-making processes.

The majority of PwD wants to be informed about the diagnosis as early as possible.^{37–39} In 2000, 40% of patients received a formal diagnosis.⁶⁰ A few years later, less than 50% received a formal dementia diagnosis.^{61,62} In the last 10 years, there has been increasing evidence that the prevalence of dementia is stagnating or even decreasing in some countries.^{63–65} Röhr et al⁶⁵ and Wolters et al⁶⁶ confirmed that there is evidence of decreasing age-specific incidence rates in industrialized nations. In Germany, the prevalence (incidence) of dementia decreased from 2.2% (0.4%) in 2015 to 2.0% (0.3%) in 2019, causing a decrease in the number of PwD, despite continued demographic changes.⁸⁰ Many physicians are somewhat skeptical of the ascribed effects of evidence-based and dementia-specific treatment opportunities, such as anti-dementia drugs.⁶⁷ For this reason and because of potential side effects, some experts have advised against the use of anti-dementia medication.⁶⁸ Some health authorities stopped covering the costs of such drugs. Therapeutic nihilism that the GPs are forming, is a disincentive for the state of that diagnostic workup. Therefore, practitioners’ current diagnostic behavior is not in line with patients’ preferences for an early and uncovered dementia diagnosis.

In previous times, it was believed that PwD do not need to know much about the disease due to missing curative treatments, but this perception has changed over the years.^{69,70} One reason was that PwD might not cope well with the diagnosis, which can deteriorate the relationship between the doctor and the patient. On the other hand, advantages were seen in reducing uncertainty and having sufficient time to organize social support services, appropriate treatments, and plans for the future when symptoms start to worsen.⁷¹ Furthermore, a dementia diagnosis can be a relief for older people who perceive memory loss without knowing the cause. It is assumed that coping with the disease is easier in early stages of dementia. PwD in these stages can still be meaningfully involved in conversations about ACP.^{36,61,62} Additionally, QoL was found to be positively associated with an early diagnosis, as it improves freedom and self-determination.^{19,36} Hence, clinicians and practitioners should not avoid discussing the disease with their patients.³⁶ Future research should evaluate the best ways to deliver the diagnosis and minimize negative emotional and psychological impacts, such as fear, as well as whether an early diagnosis is associated with better patient-reported outcomes later on.³⁷

Concerning postdiagnostic treatment and care, PwD prefer to focus on the present, and it is difficult to discuss and decide on treatment and care in the future, considering end-of-life care options.⁴⁸ Family members' and caregivers' support plays a significant role in current and future decisions around treatment and care, even though PwD should not ultimately leave important decisions regarding their own future in their relatives' hands. Uncertainty about the future can be reduced by including joint anticipatory planning between PwD and their families. Some PwD make informal flexible plans, but usually only in areas they perceive as necessary (eg, to decrease stress for their family). Additionally, families have to be prepared for potential changes in future care realities.^{26,43} This process should be initiated as early as possible to comply with PwD preferences as closely as possible for the current and future treatment and care situations.⁴⁵ Concerning the choice between behavioral treatment options, memory compensation training was found to be the most popular among PwD. Following this, it seems that many PwD are always aware of their memory problems and try to work against the disease progression.^{18,34}

Preferences could also play a crucial role in the daily life of PwD. Therefore, it would be beneficial to observe the emotions of PwD during daily life routines because they can agree or disagree with the current living situation, allowing PwD to be engaged in certain decisions without being a burden to others.²⁸ An introduction of behavior plans that used PwD-preferred items (such as personal photos, books, or music) or activities can reduce behavioral and neuropsychiatric symptoms and, thus, relieve the burden of caregivers and health professionals. However, not only is it essential for patients' behaviors to be considered in daily life, but specific behaviors of other people in the surroundings could also affect PwD. A majority of cognitive processes can be positively influenced by direct gaze because mere perception improves the likability of faces and helps with remembering them.⁵⁴

PwD want to be active and engaged and need a purpose to conclude a task. Identifying and carrying out activities that satisfy PwD might keep them physically and cognitively active for longer, which, in turn, could improve the living and caring situation.²⁵ The implementation of such activities in the daily routine could be beneficial. Such activity programs should use work-related stimuli to improve engagement duration and attention.⁵⁵ We identified various ways to retain high levels of well-being for PwD. Meaningful activities, including social interaction, provide satisfaction by

giving patients a purpose in life.⁵⁹ Dickins et al¹⁰ found that PwD in early stages look for activities that match their cognitive abilities, especially everyday activities.

Within their daily life routine, PwD want to be surrounded by a social environment. Being alone with no company and having memory problems are some of the main aspects that hindered them from being active. Simple, light, and safe exercises are preferred, such as walking.^{29,55} PwD are always more engaged in activities that suit their main interests.³⁰ Therefore, they prefer leisure activities over edible items. Furthermore, the method of communication should be considered because PwD prefer teleological explanations over mechanistic ones.²⁴ Such aspects should be considered within decision-making processes with PwD as well as within daily living and communication. Regarding preferences for food, PwD do not differ from nondemented controls. However, a systematic "one-size-fits-all" approach, as is often used in nursing homes, is not appropriate for either nonimpaired adults or PwD. As one aspect of person-centered care, PwD have to be able to keep control and be engaged in various aspects of food. A range of food choices and individual mealtimes should always be offered. Additionally, in cases where PwD are diagnosed with dysphagia, meals should still be taste-modified regularly and aligned with individual tastes.³² Concerning the living environment, different studies^{21,22} have revealed that pictures with familiar content or positive emotions are easier to recall for PwD and could be used as cues for the design and setup of the environment and ambience of PwD homes as well as within conversations and interactions with PwD.

Thus, individual preferences of PwD alongside their available resources have to be determined individually. As the disease progresses, isolation and the loss of commonplace occupations are some of the main challenges PwD face. Over time, autonomy declines, but most PwD still wish to remain in their homes. There is no general approach defining at which point in time PwD should leave their homes. To be dependent on others can be accepted by PwD to maintain activities that determine their daily life at home, which could still improve or maintain their well-being.^{10,38} Preferences for end-of-life care depend on the characteristics, individual ideas, and personal needs of PwD. Due to their cognitive decline, PwD sometimes feel unhelpful and not worth listening to. In such cases, caregivers need to show empathy and sensitivity to PwD to support their feelings of safety and trust.^{14,44} Home is considered a familiar place where

PwD feel safe, surrounded by people who care and know them very well.^{44,73} Communication is also a vital part of end-of-life care, as PwD want to be kept informed about what is happening around them and their health condition. As PwD cannot comprehend the future very well, it is necessary to make them aware of their influence on a variety of aspects in present and future care. Evidence-based guidelines need to be adapted individually by caregivers and clinicians to allow a person-centered approach and to reassure the excellent quality of care at all times.^{14,44}

This systematic review has shown a variety of heterogeneous preferences, that patient preferences are present in all aspects of care and daily living, and that PwD can state their preferences. Even though the disease progresses, preferences remain stable over time, irrespective of cognitive decline. Therefore, PwD preferences should always be taken into account. However, there is a lack of quantitative preference studies which identify the most and least preferred aspects of diagnosis, treatment, and care, and PwD daily life routines, as well as quantitative differences between these aspects. Until now, there has been a lack of studies evaluating whether a strict consideration of PwD preferences could improve patient-related outcomes, such as PwD QoL. Therefore, further research is needed. Such studies should be conducted to create a fundamental basis to extend existing evidence-based guidelines based on the clinical efficacy and effectiveness of interventions.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Deuschl G, Maier W. S3-Leitlinie Demenzen. Leitlinien für Diagnostik und Therapie in der Neurologie. (25.1. 2016). Deutsche Gesellschaft für Neurologie, Hrsg. Im Internet. [S3-guidelines Dementia. Guidelines for Diagnostic and Therapy in Neurology. (25.01.2016). German Society for Neurology]. Available from: www.dgn.org/leitlinien. Accessed November 7, 2020.
2. World Health Organisation. Dementia; 2020. Available from: www.who.int/news-room/fact-sheets/detail/dementia. Accessed November 7, 2020.
3. Samsi K, Manthorpe J. Everyday decision-making in dementia: findings from a longitudinal interview study of people with dementia and family carers. *Int Psychogeriatr*. 2013;25(6):949–961. doi:10.1017/S1041610213000306
4. Fetherstonhaugh D, Tanzia L, Nay R. Being central to decision making means I am still here!: the essence of decision making for people with dementia. *J Aging Stud*. 2013;27(2):143–150. doi:10.1016/j.jaging.2012.12.007
5. Hamann J, Bronner K, Margull J, et al. Patient participation in medical and social decisions in Alzheimer's disease. *J Am Geriatr Soc*. 2011;59(11):2045–2052. doi:10.1111/j.1532-5415.2011.03661.x
6. Hirschman KB, Joyce CM, James BD, et al. Do Alzheimer's disease patients want to participate in a treatment decision, and would their caregivers let them? *Gerontologist*. 2005;45(3):381–388. doi:10.1093/geront/45.3.381
7. Smebye KL, Kirkevold M, Engedal K. How do persons with dementia participate in decision making related to health and daily care? A multi-case study. *BMC Health Serv Res*. 2012;12(1):241. doi:10.1186/1472-6963-12-241
8. Edvardsson D, Winblad B, Sandman P-O. Person-centred care of people with severe Alzheimer's disease: current status and ways forward. *Lancet Neurol*. 2008;7(4):362–367. doi:10.1016/S1474-4422(08)70063-2
9. Dworkin R. *Autonomy and the Demented Self*. The Milbank Quarterly; 1986:4–16.
10. Dickins M, Goeman D, O'Keefe F, et al. Understanding the conceptualisation of risk in the context of community dementia care. *Soc Sci Med*. 2018;208:72–79. doi:10.1016/j.socscimed.2018.05.018
11. Aneshensel CS, Pearlin LI, Mullan JT, Zarit SH, Whitlatch CJ. *Profiles in Caregiving: The Unexpected Career*. Elsevier; 1995.
12. Alzheimer's Society. The psychological and emotional impact of dementia. 2020. Available from: <https://www.alzheimers.org.uk/get-support/help-dementia-care/understanding-supporting-person-dementia-psychological-emotional-impact>. Accessed November 7, 2020.
13. Eters L, Goodall D, Harrison BE. Caregiver burden among dementia patient caregivers: a review of the literature. *J Am Acad Nurse Pract*. 2008;20(8):423–428. doi:10.1111/j.1745-7599.2008.00342.x
14. Hill SR, Mason H, Poole M, et al. What is important at the end of life for people with dementia? The views of people with dementia and their carers. *Int J Geriatr Psychiatry*. 2017;32(9):1037–1045. doi:10.1002/gps.4564
15. Shalowitz DI, Garrett-Mayer E, Wendler D. The accuracy of surrogate decision makers: a systematic review. *Arch Intern Med*. 2006;166(5):493–497. doi:10.1001/archinte.166.5.493
16. Popham C, Orrell M. What matters for people with dementia in care homes? *Aging Ment Health*. 2012;16(2):181–188. doi:10.1080/13607863.2011.628972
17. Okonkwo O, Griffith HR, Belue K, et al. Medical decision-making capacity in patients with mild cognitive impairment. *Neurology*. 2007;69(15):1528–1535. doi:10.1212/01.wnl.0000277639.90611.d9
18. Barrios PG, González RP, Hanna SM, et al. Priority of treatment outcomes for caregivers and patients with mild cognitive impairment: preliminary analyses. *Neurol Ther*. 2016;5(2):183–192. doi:10.1007/s40120-016-0049-1
19. Dröes R-M, Boelens-van Der Knoop ECC, Bos J, et al. Quality of life in dementia in perspective: an explorative study of variations in opinions among people with dementia and their professional caregivers, and in literature. *Dementia*. 2006;5(4):533–558. doi:10.1177/1471301206069929
20. ICHOM. Standard Set for Dementia. Available from: <https://connect.ichom.org/standard-sets/dementia/>. Accessed November 8, 2020.
21. Wijk H, Berg S, Sivik L, et al. Colour discrimination, colour naming and colour preferences among individuals with Alzheimer's disease. *Int J Geriatr Psychiatry*. 1999;14(12):1000–1005. doi:10.1002/(SICI)1099-1166(199912)14:12<1000::AID-GPS46>3.0.CO;2-E
22. Stanzani Maserati M, Mitolo M, Medici F, et al. Color choice preference in cognitively impaired patients: a look inside Alzheimer's disease through the use of Lüscher Color Diagnostic. *Front Psychol*. 2019;10:1951. doi:10.3389/fpsyg.2019.01951
23. Halpern AR, Ly J, Elkin-Frankston S, et al. "I know what I like": stability of aesthetic preference in Alzheimer's patients. *Brain Cogn*. 2008;66(1):65–72. doi:10.1016/j.bandc.2007.05.008
24. Lombrozo T, Kelemen D, Zaitchik D. Inferring design: evidence of a preference for teleological explanations in patients with Alzheimer's disease. *Psychol Sci*. 2007;18(11):999–1006. doi:10.1111/j.1467-9280.2007.02015.x

25. Feliciano L, Steers ME, Elite-Marcandonatou A, et al. Applications of preference assessment procedures in depression and agitation management in elders with dementia. *Clin Gerontol.* 2009;32(3):239–259. doi:10.1080/07317110902895226
26. Harrison Denning K, King M, Jones L, et al. Advance care planning in dementia: do family carers know the treatment preferences of people with early dementia? *PLoS One.* 2016;11(7):e0159056. doi:10.1371/journal.pone.0159056
27. Raetz PB, LeBlanc LA, Baker JC, et al. Utility of the multiple-stimulus without replacement procedure and stability of preferences of older adults with dementia. *J Appl Behav Anal.* 2013;46(4):765–780. doi:10.1002/jaba.88
28. van de Ven LG, Smits C, Elwyn G, et al. Recognizing decision needs: first step for collaborative deliberation in dementia care networks. *Patient Educ Couns.* 2017;100(7):1329–1337. doi:10.1016/j.pec.2017.01.024
29. Chong TW, Doyle CJ, Cyarto EV, et al. Physical activity program preferences and perspectives of older adults with and without cognitive impairment. *Asia Pac Psychiatry.* 2014;6(2):179–190. doi:10.1111/appy.12015
30. Cohen-Mansfield J, Marx MS, Thein K, et al. The impact of past and present preferences on stimulus engagement in nursing home residents with dementia. *Aging Ment Health.* 2010;14(1):67–73. doi:10.1080/13607860902845574
31. Van't Leven N, de Lange J, Prick A-E, et al. How do activating interventions fit the personal needs, characteristics and preferences of people with dementia living in the community and their informal caregivers? *Dementia.* 2019;18(1):157–177. doi:10.1177/1471301216662378
32. Milte R, Shulver W, Killington M, et al. Struggling to maintain individuality—describing the experience of food in nursing homes for people with dementia. *Arch Gerontol Geriatr.* 2017;72:52–58. doi:10.1016/j.archger.2017.05.002
33. Smith GE, Chandler M, Fields JA, et al. A survey of patient and partner outcome and treatment preferences in mild cognitive impairment. *J Alzheimers Dis.* 2018;63(4):1459–1468. doi:10.3233/JAD-171161
34. Shelton EG, Orsulic-Jeras S, Whitlatch CJ, Szabo SM. Does it matter if we disagree? The impact of incongruent care preferences on persons with dementia and their care partners. *Gerontologist.* 2018;58(3):556–566.
35. Reamy AM, Kim K, Zarit SH, et al. Understanding discrepancy in perceptions of values: individuals with mild to moderate dementia and their family caregivers. *Gerontologist.* 2011;51(4):473–483. doi:10.1093/geront/gnr010
36. Mate KE, Pond CD, Magin PJ, et al. Diagnosis and disclosure of a memory problem is associated with quality of life in community based older Australians with dementia. *Int Psychogeriatr.* 2012;24(12):1962–1971. doi:10.1017/S1041610212001111
37. Jung JH, Kim MJ, Choi S-H, et al. Do patients want to listen to a diagnosis of Dementia in Korea? Preferences on disclosing a diagnosis of dementia and discussing advance care planning in elderly patients with memory concerns and their families. *Psychiatry Investig.* 2017;14(6):779–785. doi:10.4306/pi.2017.14.6.779
38. Elson P. Do older adults presenting with memory complaints wish to be told if later diagnosed with Alzheimer's disease? *Int J Geriatr Psychiatry.* 2006;21(5):419–425. doi:10.1002/gps.1485
39. Mahieux F, Herr M, Ankri J. What are the preferences of patients attending a memory clinic for disclosure of Alzheimer's disease? *Rev Neurol (Paris).* 2018;174(7–8):564–570. doi:10.1016/j.neurol.2017.10.016
40. Bamford C, Olsen K, Davison C, et al. Is there a preference for PET or SPECT brain imaging in diagnosing dementia? The views of people with dementia, carers, and healthy controls. *Int Psychogeriatr.* 2016;28(1):123. doi:10.1017/S1041610215001039
41. Byszewski AM, Molnar FJ, Aminzadeh F, et al. Dementia diagnosis disclosure: a study of patient and caregiver perspectives. *Alzheimer Dis Assoc Disord.* 2007;21(2):107–114. doi:10.1097/WAD.0b013e318065c481
42. Hellström I, Torres S. A wish to know but not always tell—couples living with dementia talk about disclosure preferences. *Aging Ment Health.* 2013;17(2):157–167. doi:10.1080/13607863.2012.742491
43. Dickinson C, Bamford C, Exley C, et al. Planning for tomorrow whilst living for today: the views of people with dementia and their families on advance care planning. *Int Psychogeriatr.* 2013;25(12):2011–2021. doi:10.1017/S1041610213001531
44. Goodman C, Amador S, Elmore N, et al. Preferences and priorities for ongoing and end-of-life care: a qualitative study of older people with dementia resident in care homes. *Int J Nurs Stud.* 2013;50(12):1639–1647. doi:10.1016/j.ijnurstu.2013.06.008
45. Cohen SM, Volandes AE, Shaffer ML, et al. Concordance between proxy level of care preference and advance directives among nursing home residents with advanced dementia: a cluster randomized clinical trial. *J Pain Symptom Manage.* 2019;57(1):37–46. e1. doi:10.1016/j.jpainsymman.2018.09.018
46. Tsai C-F, Lee Y-T, Lee W-J, et al. Depression of family caregivers is associated with disagreements on life-sustaining preferences for treating patients with dementia. *PLoS One.* 2015;10(7):e0133711. doi:10.1371/journal.pone.0133711
47. Ayalon L, Bachner YG, Dwolatzky T, et al. Preferences for end-of-life treatment: concordance between older adults with dementia or mild cognitive impairment and their spouses. *Int Psychogeriatr.* 2012;24(11):1798. doi:10.1017/S1041610212000877
48. Denning KH, Jones L, Sampson EL. Preferences for end-of-life care: a nominal group study of people with dementia and their family carers. *Palliat Med.* 2013;27(5):409–417. doi:10.1177/0269216312464094
49. Mulqueen K, Coffey A. Preferences of residents with dementia for end of life care. *Nurs Older People.* 2017;29(2):26–30. doi:10.7748/nop.2017.e862
50. Groenewoud S, Van Exel NJA, Bobinac A, et al. What influences patients' decisions when choosing a health care provider? Measuring preferences of patients with knee arthrosis, chronic depression, or Alzheimer's disease, using discrete choice experiments. *Health Serv Res.* 2015;50(6):1941–1972. doi:10.1111/1475-6773.12306
51. LaMonica HM, English A, Hickie IB, et al. Examining internet and eHealth practices and preferences: survey study of Australian older adults with subjective memory complaints, mild cognitive impairment, or dementia. *J Med Internet Res.* 2017;19(10):e358. doi:10.2196/jmir.7981
52. O'Philbin L, Woods B, Holmes E. People with dementia and caregiver preferences for digital life story work service interventions. A discrete choice experiment and digital survey. *Aging Ment Health.* 2020;24(2):353–361. doi:10.1080/13607863.2018.1525606
53. Motzek T, Bueter K, Marquardt G. Investigation of eligible picture categories for use as environmental cues in dementia-sensitive environments. *HERD.* 2017;10(4):64–73. doi:10.1177/1937586716679403
54. Lopis D, Baltazar M, Geronikola N, et al. Eye contact effects on social preference and face recognition in normal ageing and in Alzheimer's disease. *Psychol Res.* 2019;83(6):1292–1303. doi:10.1007/s00426-017-0955-6
55. Cohen-Mansfield J, Dakheel-Ali M, Thein K, et al. The impact of stimulus attributes on engagement of nursing home residents with dementia. *Arch Gerontol Geriatr.* 2009;49(1):1–6. doi:10.1016/j.archger.2008.04.002
56. DeLeon IG, Iwata BA. Evaluation of a multiple-stimulus presentation format for assessing reinforcer preferences. *J Appl Behav Anal.* 1996;29(4):519–533. doi:10.1901/jaba.1996.29-519
57. DeLeon IG, Iwata BA, Roscoe EM. Displacement of leisure reinforcers by food during preference assessments. *J Appl Behav Anal.* 1997;30(3):475–484. doi:10.1901/jaba.1997.30-475

58. Bojak SL, Carr JE. On the displacement of leisure items by food during multiple-stimulus preference assessments. *J Appl Behav Anal.* 1999;32(4):515–518. doi:10.1901/jaba.1999.32-515
59. Bohn L, Kwong See ST, Fung HH. Preference for emotionally meaningful activity in Alzheimer's disease. *Aging Ment Health.* 2019;23(11):1578–1585. doi:10.1080/13607863.2018.1506750
60. Johnson H, Bouman WP, Pinner G. On telling the truth in Alzheimer's disease: a pilot study of current practice and attitudes. *Int Psychogeriatr.* 2000;12(2):221–229. doi:10.1017/S1041610200006347
61. Alzheimer's Association. 2015 Alzheimer's disease facts and figures. *Alzheimers Dement.* 2015;11(3):332–384. doi:10.1016/j.jalz.2015.02.003
62. Van Hout H, Vernooij-Dassen MJ, Jansen DA, et al. Do general practitioners disclose correct information to their patients suspected of dementia and their caregivers? A prospective observational study. *Aging Ment Health.* 2006;10(2):151–155. doi:10.1080/13607860500310468
63. Prince M, Ali G-C, Guerchet M, et al. Recent global trends in the prevalence and incidence of dementia, and survival with dementia. *Alzheimers Res Ther.* 2016;8(1):23. doi:10.1186/s13195-016-0188-8
64. Stephan BCM, Birdi R, Tang EYH, et al. Secular trends in dementia prevalence and incidence worldwide: a systematic review. *J Alzheimers Dis.* 2018;66(2):653–680. doi:10.3233/JAD-180375
65. Roehr S, Pabst A, Luck T, et al. Is dementia incidence declining in high-income countries? A systematic review and meta-analysis. *Clin Epidemiol.* 2018;10:1233–1247. doi:10.2147/CLEP.S163649
66. Wolters FJ, Chibnik LB, Waziry R, et al. Twenty-seven-year time trends in dementia incidence in Europe and the United States: the Alzheimer cohorts consortium. *Neurology.* 2020;95(5):e519–e531. doi:10.1212/WNL.00000000000010022
67. Glinz D, Gloy VL, Monsch AU, et al. Acetylcholinesterase inhibitors combined with memantine for moderate to severe Alzheimer's disease: a meta-analysis. *Swiss Med Wkly.* 2019;149:w20093.
68. Fink HA, Linskens EJ, MacDonald R, et al. Benefits and harms of prescription drugs and supplements for treatment of clinical Alzheimer-type dementia. *Ann Intern Med.* 2020;172(10):656–668. doi:10.7326/M19-3887
69. Woods R. Discovering the person with Alzheimer's disease: cognitive, emotional and behavioural aspects. *Aging Ment Health.* 2001;5(sup1):7–16. doi:10.1080/713650008
70. Bamford C, Lamont S, Eccles M, et al. Disclosing a diagnosis of dementia: a systematic review. *Int J Geriatr Psychiatry.* 2004;19(2):151–169. doi:10.1002/gps.1050
71. Phillips J, Pond CD, Paterson NE, et al. Difficulties in disclosing the diagnosis of dementia: a qualitative study in general practice. *Br J Gen Pract.* 2012;62(601):e546–e553. doi:10.3399/bjgp12X653598
72. Walker R, Ratcliffe J, White A, et al. Dementia assessment services: what are the perceptions of older people? *Australas J Ageing.* 2018;37(1):43–47. doi:10.1111/ajag.12455
73. Badrakalimuthu V, Barclay S. Do people with dementia die at their preferred location of death? A systematic literature review and narrative synthesis. *Age Ageing.* 2014;43(1):13–19. doi:10.1093/ageing/aft151
74. McMurtray A, Saito E, Nakamoto B. Language preference and development of dementia among bilingual individuals. *Hawaii Med J.* 2009;68(9):223.
75. Cohen-Mansfield J, Golander H, Arnheim G. Self-identity in older persons suffering from dementia: preliminary results. *Soc Sci Med.* 2000;51(3):381–394. doi:10.1016/S0277-9536(99)00471-2
76. Mungas D, Cooper JK, Weiler PG, et al. Dietary preference for sweet foods in patients with dementia. *J Am Geriatr Soc.* 1990;38(9):999–1007. doi:10.1111/j.1532-5415.1990.tb04423.x
77. Ortega JV, Iwata BA, Nogales-González C, et al. Assessment of preference for edible and leisure items in individuals with dementia. *J Appl Behav Anal.* 2012;45(4):839–844. doi:10.1901/jaba.2012.45-839
78. Kremer HP, Roos RAC, Dingjan GM, et al. The hypothalamic lateral tuberal nucleus and the characteristics of neuronal loss in Huntington's disease. *Neurosci Lett.* 1991;132(1):101–104. doi:10.1016/0304-3940(91)90443-W
79. de Sant'Anna M, Vallet C, Kadouche R, et al. Computer accessibility for individuals suffering from mild to moderate Alzheimer's disease. *Eur Geriatr Med.* 2010;1(3):186–192. doi:10.1016/j.eurger.2010.04.003
80. Bohlken J, Riedel-Heller S, Steininger G, Kostev K, Michalowsky B. Trends in dementia and mild cognitive impairment prevalence and incidence in German general and specialist practices between 2015 and 2019. *J Alzheimers Dis.* 2021;79:1683–1690. doi:10.3233/JAD-201385

Patient Preference and Adherence

Dovepress

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focusing on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize clinical outcomes for existing disease

states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/patient-preference-and-adherence-journal>