






Finding genetically-supported drug targets for Parkinson's disease using Mendelian randomization of the druggable genome

Catherine S. Storm ¹, Demis A. Kia¹, Mona M. Almramhi^{1,2}, Sara Bandres-Ciga³, Chris Finan ^{4,5,6}, International Parkinson's Disease Genomics Consortium (IPDGC)*, Aron D. Hingorani ^{4,5,7} & Nicholas W. Wood ¹ 

Parkinson's disease is a neurodegenerative movement disorder that currently has no disease-modifying treatment, partly owing to inefficiencies in drug target identification and validation. We use Mendelian randomization to investigate over 3,000 genes that encode druggable proteins and predict their efficacy as drug targets for Parkinson's disease. We use expression and protein quantitative trait loci to mimic exposure to medications, and we examine the causal effect on Parkinson's disease risk (in two large cohorts), age at onset and progression. We propose 23 drug-targeting mechanisms for Parkinson's disease, including four possible drug repurposing opportunities and two drugs which may increase Parkinson's disease risk. Of these, we put forward six drug targets with the strongest Mendelian randomization evidence. There is remarkably little overlap between our drug targets to reduce Parkinson's disease risk versus progression, suggesting different molecular mechanisms. Drugs with genetic support are considerably more likely to succeed in clinical trials, and we provide compelling genetic evidence and an analysis pipeline to prioritise Parkinson's disease drug development.

¹ Department of Clinical and Movement Neurosciences, University College London Queen Square Institute of Neurology, London, UK. ² Department of Medical Laboratory Technology, Faculty of Applied Medical Sciences, King Abdulaziz University, Jeddah, Kingdom of Saudi Arabia. ³ Laboratory of Neurogenetics, National Institute on Aging, Bethesda, MD, USA. ⁴ Institute of Cardiovascular Science, Faculty of Population Health, University College London, London WC1E 6BT, UK. ⁵ University College London British Heart Foundation Research Accelerator Centre, New Delhi, India. ⁶ Department of Cardiology, Division Heart and Lungs, University Medical Center Utrecht, Heidelberglaan 100, 3584 CX Utrecht, the Netherlands. ⁷ Health Data Research UK, 222 Euston Road, London, UK. *A list of authors and their affiliations appears at the end of the paper. ✉email: n.wood@ucl.ac.uk

Parkinson's disease (PD) is a neurodegenerative movement disorder that currently has no disease-modifying treatment. Despite efforts, around 90% of drugs that enter clinical trials fail, mostly due to insufficient efficacy or safety^{1–3}. This contributes to the staggering \$1.3 billion mean price of bringing a new drug to the market¹.

Incorporating genetics in drug development could be one of the most efficient ways to improve the process, because drugs with genetic support are considerably more likely to succeed in clinical trials^{4–6}. “Druggable” genes encode proteins that have been targeted by medications or are possible to target with a small molecule or monoclonal antibody^{7,8}. While genome-wide association studies (GWAS) have effectively identified single nucleotide polymorphisms (SNPs) linked to PD risk and progression^{9–11}, the GWAS design cannot reliably pinpoint causal genes and directly inform drug development.

Mendelian randomization (MR) is a genetic technique that can predict the efficacy of a drug by mimicking a randomized controlled trial^{12–15}. SNPs associated with expression levels of a gene (expression quantitative trait loci, eQTLs) may be analogous to lifelong exposure to a medication targeting the encoded protein^{8,16}. The association between the same genetic variants and a disease (the outcome) can then be extracted from a GWAS for the outcome (Fig. 1a). The SNP-gene-expression and SNP-disease associations can be combined using MR to infer the causal effect of the exposure on the outcome. Since the exposure and outcome can be measured in two independent cohorts, openly available data from two large-scale GWASs can be used for one well-powered MR study. Because of Mendel's law of independent assortment, individuals are “randomized” at conception to have genetically higher or lower expression levels of the druggable gene (Fig. 1b). Individuals are generally unaware of their genotype, so the MR study is effectively blinded.

In this study, we use eQTLs in blood and brain tissue to predict the efficacy of over 3000 drug-targeting mechanisms in two independent PD case-control cohorts and examine several PD progression markers (Fig. 1c). Where possible, we repeat the analysis using SNPs associated with circulating levels of the encoded proteins. Using large-scale, openly available data and MR techniques, we propose a list of genetically-supported drug targets for PD, including repurposing opportunities of already-licensed or clinical-phase drugs.

Results

Mimicking medications with expression quantitative trait loci.

The druggable genome encompasses human genes that encode drug targets, including proteins targeted by approved and clinical-phase drugs, proteins similar to approved drug targets and proteins accessible to monoclonal antibodies or drug-like small molecules *in vivo*⁷. The most comprehensive version to date includes 4863 genes, and we sought to identify openly available eQTL data for these genes to mimic exposure to the corresponding medications⁷. Although the transcript level is biologically a step before the protein level, expression GWASs cover many genes across the genome and provide tissue specificity. Gene expression GWAS data thus provide a very good resource for high-level screens to develop drug-targeting hypotheses.

We used eQTL data from blood (31,684 mostly European-ancestry individuals)¹⁷ and brain tissue (1387 prefrontal cortex samples of mostly European ancestry, including 679 healthy controls, 497 schizophrenia, 172 bipolar disorder, 31 autism spectrum disorder, 8 affective disorder patients)¹⁸. We kept eQTLs with false discovery rate (FDR) < 0.05 and located within 5 kb of the associated gene to increase the specificity of the eQTL.

As such, eQTLs were available for 2786 and 2448 druggable genes in blood and brain tissue, respectively, and these were clumped at $r^2 = 0.2$. Compared to a lower clumping threshold, this increases the number of SNPs available per gene, which in turn improves the power to detect an effect and makes it possible to test for biases in the MR estimate (as discussed later). When clumping at $r^2 = 0.2$, SNPs are not strictly independent. We therefore used MR methods that incorporate a linkage disequilibrium matrix based on the 1000 genomes EUR reference panel in the MR analysis, which accounts for correlation between SNPs^{19,20}. These methods therefore take linkage disequilibrium into account.

Discovery phase identifies 31 potential drug targets to prevent PD.

The largest GWAS available for a PD trait studied disease risk in European-ancestry individuals, which we obtained from the International PD Genomics Consortium (IPDGC)⁹. Our discovery cohort consisted of 13,708 PD patients and 95,282 controls collected for a 2014 GWAS meta-analysis²¹. The MR effect estimate for each SNP (Wald ratio) was calculated, and where >1 eQTL was available per gene after clumping, Wald ratios were meta-analysed, weighted by inverse-variance (IVW). Genetically-determined expression of 31 genes (11 in blood only; 17 in brain tissue only; three in both blood and brain tissue) was significantly associated with PD risk in the discovery cohort at FDR < 0.05. All remained significant when clumping at $r^2 = 0.001$ (Supplementary Data 1).

15 potential preventative agents replicate in an independent PD case-control cohort.

We sought to replicate all genes that reached significance in the discovery phase using the Wald ratio or IVW method in an independent PD case-control cohort (Fig. 1). The replication population consisted of 8036 PD patients and 5803 controls (no overlap with the discovery cohort)⁹. The MR methods were identical to those used in the discovery phase.

Genetically-predicted expression of 15 genes (four in blood only; nine in brain tissue only; two in both tissues) replicated using the Wald ratio or IVW method (Fig. 2 and Supplementary Data 1). *BST1*, *CD38*, *CHRN1*, *CTSB*, *GPNMB*, *LGALS3*, *MAPT*, *MMRN1*, *NDUFAF2*, *PIGF*, *VKORC1* and *WNT3* reached FDR < 0.05; *ACVR2A*, *HSD3B7* and *MAP3K12* reached nominal significance. *GPNMB* and *HSD3B7* reached significance in both blood and brain tissue. Of these 15 potential drug targets to prevent PD, nine were not nominated by the PD risk GWAS meta-analysis⁹.

Three replicated genes encode targets of approved or clinical-phase drugs with an appropriate direction of effect for PD protection, presenting a possible repurposing opportunity: *CHRN1*, *NDUFAF2* and *VKORC1* (Table 1 and Supplementary Data 1). The *GPNMB* protein is a receptor targeted by glematimumab, an antibody-drug conjugate that is being evaluated for several types of cancer²². After binding to *GPNMB*, the drug is internalised by the cell and is cytotoxic. Since this mechanism of action does not reflect a change in *GPNMB* levels, we do not consider glematimumab a potential candidate for repurposing. We find that CD38-inhibitors such as daratumumab, licensed to treat multiple myeloma, and *MAP3K12*-inhibitors such as CEP-1347 may increase PD risk. Interestingly, CEP-1347 failed to modify PD progression in a phase 3 clinical trial²³, and our data may provide a genetic explanation why CEP-1347 was unsuccessful.

MR quality control suggests that *CD38*, *CTSB*, *GPNMB* and *MAP3K12* have the most robust MR evidence for PD risk.

We completed a series of quality control steps to prioritise




Fig. 1 Overview of MR and our study. **a** Genetic variants associated with the expression of a gene are called eQTLs, and they mimic life-long exposure to higher or lower levels of gene expression (the exposure). These variants affect PD (the outcome) through the exposure only, i.e. there is no horizontal pleiotropy. **b** MR is analogous to a randomized controlled trial, where individuals are randomly allocated to a genotype according to Mendel's law of independent assortment¹⁴. **c** Workflow and summarized results of our MR study. eQTL expression quantitative trait locus, MR Mendelian randomization, PD Parkinson's disease, pQTL protein quantitative trait locus.

the replicated genes. The direction of effect was consistent between the discovery and replication phases for all 15 replicated genes (Supplementary Data 2). Previous eQTL-based MR studies have reported heterogeneity in magnitude and direction of effect between tissues^{8,24}, and we found that raised *HSD3B7* expression was associated with raised PD risk in blood and reduced PD risk in brain tissue (Fig. 2). This pattern was

consistent between the discovery and replication phase. Although this may suggest opposing effects between tissues, there was only one eQTL available for *HSD3B7* in brain and two eQTLs in blood (discovery phase). Results based on one or two SNPs should be interpreted with caution, because it is not possible to perform the additional quality control discussed below.




Fig. 2 Fifteen potential preventative drug targets reach significance in two independent PD case-control cohorts. Forest plots showing the discovery-phase results for the 15 replicated genes. The centre of the error bars represents the PD odds ratio per 1-standard-deviation increase in gene expression, calculated using the Wald ratio (if 1 SNP) or IVW (if >1 SNP) and corrected for the number of genes tested. Results are colour-coded according to the tissue (red = blood, blue = brain tissue). 95% CI 95% confidence interval, FDR false discovery rate, OR odds ratio, PD Parkinson’s disease.

Table 1 Four potential drug-targeting mechanisms for PD may constitute repurposing opportunities for existing drugs.

Gene	Outcome (tissue)	Drug name	Clinical phase	Indications/Uses
<i>CHRNA1</i>	Risk (brain)	Rocuronium	Approved	Muscle relaxant in anaesthesia
<i>NDUFAF2</i>	Risk (brain)	Metformin	Approved	Type 2 diabetes mellitus, polycystic ovarian syndrome
<i>RHD</i>	Dyskinesia (brain)	Roledumab	Phase 2	Prevent alloimmunisation in Rhesus negative mothers carrying a Rhesus positive child
<i>VKORC1</i>	Risk (blood)	Warfarin	Approved	Prophylactic anticoagulation *high risk of falls in Parkinson’s disease

These drugs are either approved or in clinical trial phase, and the mechanism of action is consistent with the direction of our MR effect estimate. The second column displays the potential effect on PD and target tissue. Clinical phase and drug indication based on <https://clinicaltrials.gov/> and the British National Formulary. Direction of effect was confirmed using <https://www.drugbank.ca> or <https://www.ebi.ac.uk/chembl/> databases.

The IVW method assumes that (1) the genetic variant(s) must be associated with the exposure, (2) the genetic variant(s) must not be associated with any confounders, and (3) the genetic variant(s) must not be associated directly with the outcome. This means that the SNP should affect the outcome (PD risk) through the exposure (gene expression) only, so the y-intercept of the IVW regression is fixed at zero²⁵. This assumption is violated if there is genetic pleiotropy, where a SNP affects the outcome through an alternative pathway. This kind of pleiotropy may arise due to measured and unmeasured confounders, for example if the SNP is an eQTL for another gene that is not tested in this MR study. If pleiotropy pushes the effect in one direction, the IVW method yield a biased effect estimate. The MR-Egger method relaxes this assumption by not constraining the y-intercept. If the MR-Egger y-intercept significantly deviates from zero, this suggests that there is directional pleiotropy. This method assumes that any pleiotropic effects are independent of the gene-exposure association²⁶.

If several meta-analysis methods yield a similar result, such as the MR-Egger and maximum likelihood methods, we consider the MR result more robust^{25,27,28}. The latter allows more uncertainty in the SNP-exposure and SNP-outcome associations²⁹. These methods are only possible if >2 SNPs are available per gene, and all genes with >2 SNPs reached at least

nominal significance using the maximum likelihood method (uncorrected $p < 0.05$). The magnitude and direction of effect were largely consistent between methods, except for *BST1*. For *BST1*, the MR-Egger estimate was in the opposite direction to the IVW and maximum likelihood results (Supplementary Data 1). All genes with >2 SNPs available passed the MR-Egger intercept test except *BST1*, explaining the deviant MR-Egger estimate for this gene (Supplementary Data 2).

Nevertheless, if SNPs for a gene are pleiotropic in opposing directions, the MR-Egger y-intercept will still be zero. The Cochran’s Q and I^2 tests usefully assess overall heterogeneity between Wald ratios. *NDUFAF2*, *WNT3* and *VKORC1* did not pass the Cochran’s Q ($p < 0.05$) nor I^2 ($I^2 > 0.50$) tests (Supplementary Data 2). This means that there is significant heterogeneity in the MR result for these genes, and such heterogeneity among Wald ratios can for example happen if at least one SNP for the gene is pleiotropic³⁰.

We repeated the analysis in the discovery outcome data using only SNPs that were specifically associated with our replicated genes. In other words, we removed any SNPs associated with the expression of any other gene in the original eQTL dataset. All replicated genes remained significant in this analysis (Supplementary Data 8 and 9).

In addition, a spurious MR result may arise from a locus where the SNP-exposure and SNP-outcome associations are rooted in two distinct causal SNPs in close linkage disequilibrium³⁰. When the SNP is significantly associated with both exposure and outcome, this can be probed using colocalization analysis³¹. There is evidence that proteins with both MR and colocalization evidence are more likely to be successful drug targets³²; this may simply reinforce that GWAS-nominated drug targets are more likely to reach approval⁴. Using the discovery outcome data, we had sufficient power ($PPH3 + PPH4 \geq 0.8$) to perform a colocalization analysis for 13 genes (see ‘Methods’ and Supplementary Data 7). Of these, *ACVR2A*, *BST1*, *CHRN1*, *CTSB*, *GPNMB*, *HSD3B7*, *LGALS3*, *MAPT*, *MMRN1* and *VKORC1* had strong evidence of colocalization ($PPH4 \geq 0.75$). All genes with sufficient power colocalized in the replication data (*BST1*, *CD38*, *GPNMB*, *HSD3B7*, *MAPT*, *MMRN1*, *VKORC1* and *WNT3*). Similarly, Kia et al. recently found that eQTLs in brain tissue for *CD38* and *GPNMB* based on a different eQTL dataset colocalize with PD risk loci³³, strengthening the evidence for the encoded proteins as drug targets for PD.

Four potential targets for preventative drugs may also affect PD age at onset. Pharmacologically delaying the age of onset of a debilitating disease may have a considerable impact on both socioeconomic burden of disease and quality of life by providing disability-free years to people at risk. Evidence from polygenic risk score analyses suggests that genetic risk of PD is correlated with PD age at onset^{11,34–36}. We therefore asked whether expression of the genes reaching significance in our MR discovery phase for PD risk also predict PD age of onset. We sourced openly available summary statistics from a PD age of onset GWAS, including 17,996 patients (Fig. 1c). Based on the same analysis pipeline as the replication step for PD risk, expression of four genes predicted PD age of onset at $p < 0.05$: *BST1* in blood, *CD38* in brain tissue, *CTSB* in brain tissue and *MMRN1* in brain tissue (Fig. 3 and Supplementary Data 3). *CD38* and *MMRN1* remained significant when clumping at $r^2 = 0.001$. There were >2 SNPs available for *BST1*, *CD38* and *CTSB*, and the IVW, maximum likelihood and MR-Egger methods yielded a consistent direction of effect for these genes (Supplementary Data 1). All three genes passed the MR-Egger intercept ($p > 0.05$), Cochran’s Q ($p > 0.05$), and I^2 tests ($I^2 < 0.50$). *BST1* and *MMRN1* remained significant when removing SNPs associated with expression of any other gene in the original eQTL dataset

(Supplementary Data 8 and 9). Of the four genes, we had sufficient power ($PPH3 + PPH4 \geq 0.8$) to perform a colocalization analysis for *BST1*, and we found strong evidence of colocalization ($PPH4 \geq 0.75$; Supplementary Data 7).

If increased expression of a gene predicts reduced PD risk, this gene should be associated with a delayed age at onset. This was consistently the case for all four genes that reached significance for age at onset. Overall, these data suggest that there may be some shared molecular mechanisms driving PD risk and age at onset, yet this overlap may be incomplete.

There is little overlap between drug targets to prevent PD and reduce PD progression.

The PD risk GWAS data afford large discovery and replication cohorts, which is a great advantage in MR. Nevertheless, it is currently not possible to reliably predict PD, limiting the immediate usefulness of a drug to prevent or delay this condition. Many clinical trials for PD use progression markers such as the Unified PD Rating Scale (UPDRS) to evaluate drug efficacy, and it remains unclear how the molecular mechanisms driving PD risk relate to clinical progression. We used MR to probe whether expression of any of the 4863 druggable genes is significantly associated with PD progression, measured by the UPDRS (total and parts 1 to 4), mini-mental state examination (MMSE), Montreal cognitive assessment (MOCA), modified Schwab and England activities of daily living scale (SEADL), Hoehn and Yahr stage, dementia, depression, and dyskinesia. The MR pipeline for each progression marker was identical to the discovery phase for PD risk (Fig. 1).

We used openly available summary statistics from a GWAS for these PD progression markers in 4093 European PD patients, followed over a median of 2.97 years¹⁰. 3455 druggable genes had an eQTL available for MR using a PD progression marker (2752 in blood, 2353 in brain tissue), and eight genes reached significance across five progression outcomes (Fig. 4 and Supplementary Data 1). One of these, *RHD*, encodes the target of a clinical-phase medication with an appropriate direction of effect, possibly representing a repurposing opportunity (Table 1).

IRAK3 expression in blood was significantly associated with UPDRS parts 2 and 4, depression, and dyskinesias. *LMAN1* expression in blood reached significance for dyskinesias and UPDRS part 2. Reaching significance for several progression markers strengthens the evidence for these two genes. No genes reached significance for both PD risk and progression. Since age at motor symptom onset may be considered an early marker of




Fig. 3 Four potential preventative drugs may also affect PD age at onset. Forest plot; the centre of the error bars represents the standard-deviation change in PD age at onset per 1-standard-deviation increase in gene expression, calculated using the Wald ratio (if 1 SNP) or IVW (if >1 SNP) and colour-coded by tissue (red = blood, blue = brain tissue). A negative beta corresponds to a younger age at onset, and a positive beta corresponds to an older age at onset. 95% CI 95% confidence interval, PD Parkinson’s disease.




Fig. 4 Genetically-predicted expression of eight genes in blood or brain tissue is associated with PD progression markers. Forest plot; the centre of the error bars show the standard-deviation change in each progression marker, per 1-standard-deviation increase in gene expression, calculated using the Wald ratio (if 1 SNP) or IVW (if >1 SNP). Results are colour-coded by tissue (red = blood, blue = brain tissue) and corrected for the number of genes tested. 95% CI 95% confidence interval, DEPR depression, FDR false discovery rate, HY Hoehn and Yahr, DEPR depression, UPDRS2-4 Unified Parkinson's Disease Rating Scale parts 2 to 4.

PD progression, we also used our MR approach to assess whether genes discovered by our progression analysis causally predict age at onset. Of the genes that reached significance for a progression marker, none reached nominal significance using the age at onset data (Supplementary Data 3 and 4). *CD177*, *IRAK3*, *RHD* and *STK4* remained significant when removing SNPs associated with expression of any other gene in the original eQTL dataset (Supplementary Data 8 and 9). It was not possible to perform a reliable colocalization analysis in our progression study, since the discovered genes did not have sufficient power to do so (i.e. $PPH3 + PPH4 < 0.8$).

The direction of effect was consistent between the IVW, maximum likelihood and MR-Egger methods for all genes except *RHD*, where the MR-Egger method opposed the direction of the IVW and maximum likelihood methods. *CD177* (depression), *RHD* (dyskinesia), *PYGL* (UPDRS part 4) and *STK4* (Hoehn and Yahr) reached significance when clumping at $r^2 = 0.001$. *ADAM32* (Hoehn and Yahr), *IRAK3* (dyskinesia), *LMAN1* (UPDRS part 2), and *RHD* (dyskinesia) passed MR-Egger intercept, Cochran's Q and I^2 tests (Supplementary Data 2). Taken together, these five genes have the most robust MR evidence for modifying a PD progression marker.

Protein quantitative trait locus data provide further genetic evidence. Most clinically-used drugs target proteins, not gene expression, and genetic variants associated with protein levels (protein quantitative trait loci, pQTLs) may model drug target effects more accurately than eQTLs⁸. Even with high throughput protein assays, however, the spectrum of reliable, well-powered GWAS data on protein targets is limited. Many genetic studies on protein levels are based on plasma and lack tissue diversity^{37–39}. Of the 23 proposed targets, we found pQTLs for BST1, CD38, CTSS, GPNMB and LGALS3 for PD risk, as well as PYGL and QDPR for UPDRS part 4^{37–40}.

Our MR analysis found that BST1, CTSS and LGALS3 levels were consistently associated with PD risk ($p < 0.05$; Fig. 5 and Supplementary Data 5). The result for GPNMB (risk) and PYGL (UPDRS part 4) lost significance when using data from different pQTL studies. The direction of effect was consistent between the pQTL and eQTL results for all genes except BST1, and the MR-Egger intercept, Cochran's Q and I^2 tests suggest that the BST1 results may be biased by genetic pleiotropy (Supplementary Data 6). This illustrates the importance of MR quality control—maximizing the number of SNPs available per drug target and validation with different data types and independent replication cohorts is essential for a reliable effect estimate.

Discussion

This work explicitly seeks to identify new drug targets for PD, and we provide genetic evidence in favour of 23 potential disease-modifying drug targets. Tables 2 and 3 summarize the evidence supporting these genes. The genes were prioritised using several meta-analysis methods (IVW, MR-Egger and maximum likelihood), the MR-Egger intercept test, Cochran's Q test, the I^2 test, a pQTL study, colocalization analysis and previously published MR and colocalization evidence. This allowed us to look for pleiotropy due to both measured and unmeasured confounders²⁵. We propose six drug targets with the strongest MR evidence: *CTSS*, *GPNMB*, *CD38*, *RHD*, *IRAK3* and *LMAN1*.

We identified four genes encoding targets for existing drugs warranting further discussion (Table 1). *NDUFAF2* encodes a subunit of a target of metformin, an approved medication for type 2 diabetes mellitus. There is extensive evidence for a relationship between diabetes mellitus and PD⁴¹, and several rodent studies have investigated the potential of metformin as a neuroprotective agent^{41–43}. We found significant heterogeneity in the MR result for this gene. Although this may be because at least one SNP for this gene is pleiotropic, we speculate that this could occur if the




Fig. 5 Protein quantitative trait loci in blood provide further genetic evidence. Forest plots showing the results for all proteins and outcomes where a pQTL was available. The centre of the error bars show the (a) PD odds ratio and (b) standard-deviation change in UPDRS part 4 score, per 1-standard-deviation increase in circulating protein levels, calculated using the Wald ratio (if 1 SNP) or IVW (if >1 SNP). The “pQTL Source” column indicates which pQTL study the SNPs were derived from. 95% CI 95% confidence interval, OR odds ratio, PD Parkinson’s disease, pQTL protein quantitative trait locus, UPDRS Unified Parkinson’s Disease Rating Scale.

effect is driven by a subset of PD patients. This, however, remains a subject for future research, because the GWAS data used in this study are not stratified by any kind of PD subtype. Epidemiological studies on the relationship between long-term medication use and incidence of a disease are an invaluable contribution to evaluating preventative agents for PD. A retrospective cohort study of over 6000 patients with type 2 diabetes mellitus found that more than four years of metformin use maybe associated with a reduced PD incidence⁴⁴. Our MR study thus provides further evidence in favour of repurposing anti-diabetic drugs for PD.

Other medications may not be as suitable for repurposing. To our knowledge, there is no evidence linking PD and the drug roledumab, which is currently in a phase II clinical trial to prevent alloimmunisation in Rhesus negative mothers carrying a Rhesus positive child (NCT02287896). Our evidence suggests that *RHD* expression in brain tissue, rather than blood, is associated with PD dyskinesia. Next, *CHRNBI* encodes the beta subunit of the muscle acetylcholine receptor at the neuromuscular junction,

which is inhibited by muscle relaxants used during surgical anaesthesia. *VKORCI* encodes the catalytic subunit of the vitamin K epoxide reductase, and this enzyme is targeted by the oral anticoagulant warfarin. The key adverse effect of warfarin treatment is haemorrhage, and since PD is a movement disorder where patients experience frequent falls, any potential benefit of warfarin treatment would likely be outweighed by the added risk of haemorrhagic strokes and complications of bleeding.

The two-sample MR design allowed us to explore different tissues and PD traits, and we identified different candidates to prevent, delay onset, and slow progression of PD (Figs. 2, 3 and 4). Although we found that four of the drug targets for PD risk may also affect PD age at onset, we found very different candidates for progression. Age at motor symptom onset can be considered an early sign of PD progression, and it is striking that none of the genes that reached significance for a progression outcome reached significance in the age at onset data. These results are in line with the GWAS data, finding little overlap between loci associated with PD risk, age at onset and progression

Table 2 Evidence supporting druggable genes whose expression was significantly associated with PD risk or age at onset using the Wald ratio or IVW method.

Gene	PD outcome	Tissue	Replication	Sign. with max. lik.	Sign. with MR-Egger	MR-Egger intercept test	Cochran's Q test	I ² test	pQTL evidence	Coloc	Previously published MR or coloc support
GPNMB	Risk	Blood	✓	✓	✓	✓	✓	✓	✓	✓	MR ⁹ MR ²⁴ ; coloc ³³
	Risk	Brain	✓	✓	✓	✓	✓	✓	✓	✓	
CTSB	Risk	Brain	✓	✓	x	✓	✓	✓	✓	✓	MR ⁹
	Age at onset	Brain	✓	✓	x	✓	✓	✓	✓	✓	
VKORC1	Risk	Blood	✓	✓	x	✓	✓	✓	✓	✓	Coloc ³³
CD38	Risk	Brain	✓	✓	x	✓	✓	✓	x	x	
	Age at onset	Brain	✓	✓	x	✓	✓	✓	✓	✓	
MAP3K12	Risk	Blood	✓	✓	x	✓	✓	✓	✓	x	MR ⁹
NDUFAF2	Risk	Brain	✓	✓	x	✓	✓	✓	✓	x	
BST1	Risk	Blood	✓	✓	x	x	✓	✓	✓	✓	
	Age at onset	Blood	✓	✓	x	✓	✓	✓	✓	✓	
HSD3B7	Risk	Blood	✓	✓	✓	✓	✓	✓	✓	x	MR ²⁴
	Risk	Brain	✓	✓	✓	✓	✓	✓	✓	✓	
LGALS3	Risk	Brain	✓	✓	✓	✓	✓	✓	✓	✓	MR ⁹
ACVR2A	Risk	Brain	✓	✓	✓	✓	✓	✓	✓	✓	
CHRN1	Risk	Brain	✓	✓	✓	✓	✓	✓	✓	✓	
MAPT	Risk	Brain	✓	✓	✓	✓	✓	✓	✓	✓	MR ⁹
MMRN1	Risk	Brain	✓	✓	✓	✓	✓	✓	✓	✓	
	Age at onset	Brain	✓	✓	✓	✓	✓	✓	✓	✓	
PIGF	Risk	Brain	✓	✓	✓	✓	✓	✓	✓	✓	MR ⁹
WNT3	Risk	Blood	✓	✓	x	✓	x	x	x	x	

✓ pass, x fail, blank not possible to test, coloc colocalization, max. lik. maximum likelihood, MR Mendelian randomization, PD Parkinson's disease, pQTL protein quantitative trait locus, sign significant.

Table 3 Evidence supporting druggable genes whose expression was significantly associated with a PD progression trait using the Wald ratio or IVW method.

Gene	PD outcome	Tissue	Sign. with max. lik.	Sign. with MR-Egger	MR-Egger intercept test	Cochran's Q test	I ² test	pQTL evidence	Coloc
RHD	Dyskinesia	Brain	✓	x	✓	✓	✓	✓	
IRAK3	Dyskinesia	Blood	✓	✓	✓	✓	✓	✓	
	Depression	Blood	✓	✓	x	✓	✓	✓	
	UPDRS part 2	Blood	✓	x	x	✓	✓	✓	
	UPDRS part 4	Blood	✓	✓	x	✓	✓	✓	
ADAM32	Hoehn and Yahr	Brain	✓	x	✓	✓	✓	✓	
LMAN1	UPDRS part 2	Blood	✓	x	✓	✓	✓	✓	
	Dyskinesia	Blood	✓	✓	x	✓	✓	✓	
PYGL	UPDRS part 4	Brain						✓	
CD177	Depression	Brain							
QDPR	UPDRS part 4	Blood	✓	✓	✓	x	x	x	x
STK4	Hoehn and Yahr	Brain			x	x	x	x	

✓ pass, x fail, blank not possible to test, coloc colocalization, max. lik. maximum likelihood, MR Mendelian randomization, PD Parkinson's disease, pQTL protein quantitative trait locus, sign significant, UPDRS Unified Parkinson's Disease Rating Scale.

markers⁹⁻¹¹. This may reflect the limited sample size of current PD progression GWAS data. Nevertheless, this raises questions about what drives PD susceptibility versus progression, painting a yet unclear picture of partially overlapping molecular mechanisms.

Our candidates to slow PD progression may be of most immediate relevance, because currently PD cannot be accurately predicted. A preventative agent would need to be highly tolerable and have a very safe side effect profile, and our approach is not well suited for systematically evaluating the safety aspects of our proposed candidates in this study. To our knowledge, the data

used here are from the largest openly available progression GWAS to date. We did not find any non-overlapping PD progression GWAS with sufficient power for a replication step in our progression analysis, which would have to measure progression in a similar way to the study used here. As such, the preventative list carries more robust evidence, because each gene reached significance in two large, independent cohorts. Replication is critical to validating scientific findings and eliminating false positives, and this has been an crucial lesson for genetic research⁴⁵⁻⁴⁷. Replication is not common practice in MR yet⁴⁸, and it is a key strength of our study. Although including all samples available in

one analysis would maximise statistical power^{46,49}, using independent discovery and replication cohorts allowed us to validate our proposed drug targets. Since our overarching intention was to provide genetic evidence to improve success rates in clinical trials, we made this decision to reduce the number of false positives.

Our study has valuable advantages compared to previous MR projects studying PD using QTL data. In the latest GWAS meta-analysis for PD risk in Europeans, Nalls et al. selected SNPs associated with PD risk and used MR to identify whether any of these loci alter expression or methylation of genes within 1 Mb of the SNP⁹. This contrasts with our exposure-centred MR analysis, where we chose SNPs associated with the expression of a druggable gene, rather than the disease outcome. More recently, Baird and colleagues conducted a transcriptome-wide MR study for a series of brain diseases and found six genes whose expression in brain tissue was significantly associated with PD risk²⁴. Two of these were also discovered in our study: *GPNMB* and *HSD3B7*. The remaining four were either not part of the druggable genome, rendering the encoded proteins less actionable drug targets, or did not reach significance in our discovery or replication cohorts, illustrating the importance of replication. Furthermore, our MR study is the first to study druggable genes in the context of PD age at onset and progression.

Nevertheless, progression and age at onset studies are particularly affected by collider bias^{50–52}. For example, if expression of a gene and depression are both associated with disease risk, that gene's expression will be artificially associated with depression in a cohort containing only cases. In a progression study, genetic variants that cause disease will thus be associated with other risk factors for disease. The druggable genes we identified in our progression study did not reach significance in our risk study, so this kind of collider bias is less likely to have occurred for our candidate genes. The age at onset analysis was comparably more affected, since we tested genes that reached significance for PD risk. Overall, this emphasises the importance of MR quality control methods (including replication) for identifying reliable causal effects, representative sampling in GWAS, as well as continued development of methods to formally test for collider bias^{53,54}.

Another key limitation of this study is that MR cannot fully recapitulate a clinical trial. MR mimics lifelong, low-dose exposure to a drug and assumes a linear relationship between exposure and outcome. This differs from a clinical trial, which typically investigates comparably high doses of drug over a much shorter timeframe. The MR result may therefore not directly correspond to the effect size in practice and does not perfectly predict the effect of a drug.

In addition, the eQTL cohorts contained some non-European individuals^{17,18}, three of the pQTL studies sourced were based on Icelandic, Scottish and German cohorts^{38–40}, and the PD populations were comprised of European individuals only^{9–11}. Linkage disequilibrium patterns differ between populations, which may compromise how well our QTLs mimic drug action in the PD cohorts and introduce bias to the MR effect estimate²⁵.

It is difficult to interpret which tissue would be the most appropriate site of action. Whereas the genes that reached significance in both blood and brain tissue may have stronger MR evidence, targeting the protein of a widely expressed gene may lead to systemic side-effects. Brain tissue may be more biologically relevant for neurodegeneration, but a drug acting in the blood stream may not need to cross the blood–brain barrier to exert its effect. A limitation of using brain tissue is that gene expression is quantified post-mortem, and measured expression levels are influenced by RNA degradation occurring after cell death⁵⁵. We included both blood and brain tissue eQTLs to

capture as many genes as possible and explore two potential tissue sites of action, but we note that it is difficult to prioritise genes based on which tissue(s) they reached significance in.

Furthermore, the sample size of our blood eQTL data ($n = 31,684$) is larger than that for brain tissue eQTLs ($n = 1387$) and the blood pQTL study ($n = 750–4137$). A larger sample size allows greater power to detect QTLs, meaning there are more SNPs per gene. Nevertheless, it is unclear how well QTL data mimic medications that modulate activity levels of the protein. We are encouraged that five of the seven proteins we were able to probe using both eQTL and pQTL data were successfully validated, adding to existing evidence that regulatory variants may be used for robust causal inferences in drug target MR⁸. Nevertheless, this MR study does not provide functional evidence for the proposed drug targets, and the MR process does not replace pre-clinical evaluation of drug targets in vitro and in vivo. Genomic approaches serve as adjuncts thereto, promising to better prioritise drug targets carried forward to clinical phase trials.

A 9.6% vs. 13.8% success rate for drugs from phase 1 trials to approval may mean a \$480 million difference in the median research and development cost of bringing a new drug to the market¹. The druggable genome resource has opened up new avenues for drug target identification using existing genetic data^{7,56,57}, and if genetic evidence increases success rates even by a few percent, this could have a substantial effect on drug development costs^{4,5}. As such, MR a highly compelling, time- and cost-effective adjunct to the randomized controlled trial. We have made our code openly available for use beyond PD research (https://github.com/catherinestorm/mr_druggable_genome_pd/)⁵⁸, and we have demonstrated ways to prioritise drug targets based on genetic data. We have provided human genetic evidence of drug efficacy for PD, and we hope that these data will serve as a useful resource for prioritising drug development efforts.

Methods

All DNA positions are based on the human reference genome build hg19 (GRCh37). Data processing was completed using R software version 3.6.3⁵⁹.

Exposure data. Tissue-specific eQTL data were obtained from the eQTLGen (<https://eqtlgen.org/>) and PsychENCODE consortia (<http://resource.psychencode.org/>); full descriptions of the data are available in the original publications^{17,18}. Briefly, the eQTLGen data consisted of cis-eQTLs for 16,987 genes and 31,684 blood samples, of which most were healthy European-ancestry individuals. We downloaded the full significant cis-eQTL results (FDR < 0.05) and allele frequency information from the eQTLGen consortium on 13 May 2020.

The PsychENCODE data included 1387 prefrontal cortex, primarily-European samples (679 healthy controls, 497 schizophrenia, 172 bipolar disorder, 31 autism spectrum disorder and 8 affective disorder patients). We downloaded all significant eQTLs (FDR < 0.05) for genes with expression >0.1 fragments per kilobase per million mapped fragments (FPKM) in at least ten samples and all SNP information, accessed on 13 May 2020.

We obtained an updated version of the druggable genome containing 4863 genes from the authors of the original publication⁷, double-checking the druggability level for all genes marked as approved or in clinical trials ("druggability tier 1"). We removed non-autosomal genes, leaving 4560 druggable genes. We filtered both eQTL datasets to include SNPs 5 kb upstream of the target druggable gene start or 5 kb downstream of the target druggable gene end position.

We sought freely available pQTL data from blood or brain tissue for all druggable genes that reached significance for any PD outcome in our study. Out of 23 pQTL studies identified, four studies (1) reported significant pQTLs in individuals of European descent for any of the druggable proteins proposed by our eQTL analysis, (2) provided all the SNP information required for MR and (3) reported SNPs that were available in our PD outcome data^{37–40}. Sun and colleagues measured 3622 proteins in 3301 healthy European blood donors from the INTERVAL study and identified 1927 pQTLs for 1478 proteins. Emilsson and colleagues measured 4137 proteins in the serum of 5457 Icelanders from AGES Reykjavik study. Effect alleles and effect allele frequencies were obtained from the authors. Suhre and colleagues measured 1124 proteins in 1000 blood samples from a German population. Hillary and colleagues measured 92 proteins in the blood of 750 healthy Scottish controls.

In total, we found pQTLs that were available in the appropriate PD outcome data for seven of our druggable proteins of interest: BST1, CD38, CTSS, GPNMB, LGALS3, PYGL and QDPR. All pQTLs included in our analysis had $p < 5e-6$ in the original pQTL study. All pQTLs were found on the same chromosome as the associated gene except for: rs62143198 for PYGL, rs62143197 for QDPR, rs4253282 for GPNMB, rs2731674 for GPNMB³⁷. These latter four SNPs are therefore acting in trans.

Outcome data. All PD data were obtained from the IPDGC, and details on recruitment and quality control are available in the original publications^{9–11,21}. In the discovery phase for PD risk, we used openly available summary statistics from a 2014 case-control GWAS meta-analysis, which included 13,708 PD patients and 95,282 controls²¹.

In the replication phase for PD risk, we obtained summary statistics from 11 case-control GWAS studies included in the most recent PD risk GWAS meta-analysis from the authors⁹. The 11 studies, as named and described in the PD GWAS meta-analysis, were Spanish Parkinson's, Baylor College of Medicine/University of Maryland, McGill Parkinson's, Oslo Parkinson's Disease Study, Parkinson's Progression Markers Initiative (PPMI), Finnish Parkinson's, Harvard Biomarker Study (HBS), UK PDMED (CouragePD), Parkinson's Disease Biomarker's Program (PDBP), Tübingen Parkinson's Disease cohort (CouragePD) and Vance (dbGap phs000394). These yielded a total of 8036 PD cases and 5803 controls. We meta-analysed the data using METAL (version 2011-03-25) using default settings, weighting by sample size⁶⁰. The overall genomic inflation factor was $\lambda = 1.116$, and when scaled to 1000 cases and 1000 controls $\lambda_{1000} = 1.017$. Based on genomic inflation factors and quantile–quantile plots of the original GWASs^{9,21}, we considered our quantile–quantile plot to show adequate agreement with the expected null distribution (Supplementary Fig. 1).

For the progression marker analyses, we used summary statistics from the largest publicly available GWAS meta-analyses for PD age at onset and clinical progression^{10,11}. For age at onset, this included 17,996 PD cases, and age at onset was defined as self-reported age at motor symptom onset or PD diagnosis. The authors reported a high correlation between age at diagnosis and age at onset.

The progression GWAS meta-analysis included 4093 PD patients from 12 cohorts, followed over a median of 2.97 years (mean visits per individual over the study period: 5.44). We downloaded summary statistics for nine continuous outcomes and four binomial outcomes (<https://pdgenetics.shinyapps.io/pdprogmetagwasbrowser/>). Continuous outcomes included Hoehn and Yahr stage (PD progression rating scale), total UPDRS/Movement Disorder Society revised version total (PD progression rating scale), UPDRS parts 1 to 4 (1 = non-motor symptoms, 2 = motor symptoms, 3 = motor examination, 4 = motor complications), MOCA (cognitive impairment), MMSE (cognitive impairment) and SEADL (activities of daily living and independence). The binomial outcomes we used were dementia, depression, dyskinesia, as well as reaching Hoehn and Yahr stage 3 or more.

Mendelian randomization. MR analyses were completed using the R package “TwoSampleMR” (version 0.5.4)⁶¹, unless stated otherwise. The exposure and outcome data were loaded and harmonized using in-built functions. SNPs were then clumped at $r^2 < 0.2$ using European samples from the 1000 Genomes Project^{20,61}. Steiger filtering was used to remove genes where SNPs explained a greater proportion of variation in the outcome (PD trait) than variation in the exposure (gene expression). For the eQTL analysis, the Steiger filtering excluded 0–403 genes per outcome tested in a tissue, representing 0–15% of all genes studied per outcome tested in a tissue.

Wald ratios were calculated for all SNPs. These were meta-analysed using the IVW, MR-Egger and maximum likelihood methods, including a linkage disequilibrium matrix to account for correlation between SNPs; this function uses the R package “MendelianRandomization” version 0.4.2²⁹. Forest plots were produced using the R package “forestplot”.

Where >2 SNPs were available per exposure, we assessed whether the MR-Egger intercept significantly deviated from zero, as well as Cochran's Q and I^2 methods to test for heterogeneity between Wald ratios⁶². FDR-corrected p-values were calculated within each exposure-outcome combination to correct for multiple testing. In the discovery study for PD risk and the PD progression studies, we considered FDR < 0.05 significant. In the replication studies for PD risk and age at onset, as well as the pQTL study, we considered nominal $p < 0.05$ significant.

For genes which reached significance using the IVW method (>1 SNP available), we carried out another MR analysis, clumping at $r^2 < 0.001$. If $>1-2$ SNPs were available at this clumping threshold, Wald ratios were meta-analysed using the IVW, MR-Egger, weighted mode and weighted median methods.

Colocalization. We carried out a colocalization analysis for PD risk, age at onset and progression outcomes using the R package “coloc”³¹. We harmonized exposure and outcome datasets using the “TwoSampleMR” package. We used default priors: $p1 = 10^{-4}$, $p2 = 10^{-4}$, $p12 = 10^{-5}$. $p1$, $p2$ and $p12$ are the prior probabilities that a SNP in the tested region is significantly associated with expression of the tested gene, the tested PD outcome, or both, respectively. The colocalization yields

posterior probabilities corresponding to one of five hypotheses: PPH0, no association with either trait; PPH1, association with expression of the gene, but not the PD trait; PPH2, association with the PD trait, but not expression of the gene; PPH3, association with the PD trait and expression of the gene, with distinct causal variants; PPH4, association with the PD trait and expression of the gene, with a shared causal variant³¹. A low PPH3 and PPH4 in combination with a high PPH0, PPH1 and/or PPH2 indicates limited power in the colocalization analysis³¹. We therefore restricted our analysis to genes reaching $PPH3 + PPH4 \geq 0.8$.

Reporting summary. Further information on research design is available in the Nature Research Reporting Summary linked to this article.

Data availability

The data generated in this study and a data dictionary (Supplementary Information) are provided in the Supplementary Information. Tissue-specific eQTL data were obtained from the eQTLGen (<https://eqtlgen.org/>) and PsychENCODE consortia (<http://resource.psychencode.org/>). The druggable genome data were obtained via from the original authors; an immediately accessible version of the druggable genome is available in the supplementary materials of the original publication⁷. Subsets of the most recent PD risk GWAS were obtained from the original authors⁹. GWAS summary statistics for PD age at onset are available on the IPDGC website (<http://pdgenetics.org/resources>). Parkinson's progression GWAS data can be found here: <https://pdgenetics.shinyapps.io/pdprogmetagwasbrowser/>.

Code availability

The code used for this study is openly available on GitHub, accompanied by instructions for use and required tools (https://github.com/catherinestorm/mr_druggable_genome_pd). Data processing was completed using R software (version 3.6.3), with packages TwoSampleMR (version 0.5.4), MendelianRandomization (version 0.4.2), coloc (version 4.0.4), dplyr (version 1.0.0), readr (version 1.3.1), stringr (version 1.4.0), tidyverse (version 1.3.0), forestplot (version 1.9), plyr (version 1.8.6), devtools (version 2.3.0), remotes (version 2.1.1). The GWAS meta-analysis was completed using METAL (version 2011-03-25).

Received: 21 October 2020; Accepted: 14 September 2021;

Published online: 20 December 2021

References

- Wouters, O. J., McKee, M. & Luyten, J. Estimated research and development investment needed to bring a new medicine to market, 2009–2018. *JAMA - J. Am. Med. Assoc.* **323**, 844–853 (2020).
- Smetana, K., Siatkowski, M. & Møller, M. Trends in clinical success rates. *Nat. Rev. Drug Discov.* **15**, 379–380 (2016).
- Harrison, R. K. Phase II and phase III failures: 2013–2015. *Nat. Rev. Drug Discov.* **15**, 817–818 (2016).
- Nelson, M. R. et al. The support of human genetic evidence for approved drug indications. *Nat. Genet.* **47**, 856–860 (2015).
- King, E. A., Wade Davis, J. & Degner, J. F. Are drug targets with genetic support twice as likely to be approved? Revised estimates of the impact of genetic support for drug mechanisms on the probability of drug approval. *PLoS Genet.* **15**, 1–20 (2019).
- Hingorani, A. D. et al. Improving the odds of drug development success through human genomics: modelling study. *Sci. Rep.* **9**, 1–25 (2019).
- Finan, C. et al. The druggable genome and support for target identification and validation in drug development. *Sci. Transl. Med.* **9**, eaag1166 (2017).
- Schmidt, A. F. et al. Genetic drug target validation using Mendelian randomisation. *Nat. Commun.* <https://doi.org/10.1038/s41467-020-16969-0> (2020).
- Nalls, M. A. et al. Identification of novel risk loci, causal insights, and heritable risk for Parkinson's disease: a meta-analysis of genome-wide association studies. *Lancet Neurol.* **18**, 1091–1102 (2019).
- Iwaki, H. et al. Genetic risk of Parkinson disease and progression: an analysis of 13 longitudinal cohorts. *Neurol. Genet.* **5**, e354 (2019).
- Blauwendraat, C. et al. Parkinson's disease age at onset genome-wide association study: defining heritability, genetic loci, and α -synuclein mechanisms. *Mov. Disord.* 1–10, <https://doi.org/10.1002/mds.27659> (2019).
- Katan, M. B. Apolipoprotein E isoforms, serum cholesterol, and cancer. *Lancet* **1**, 507–508 (1986).
- Davey Smith, G. & Ebrahim, S. ‘Mendelian randomization’: can genetic epidemiology contribute to understanding environmental determinants of disease? *Int. J. Epidemiol.* **32**, 1–22 (2003).

14. Hingorani, A. & Humphries, S. Nature's randomised trials. *Lancet* **366**, 1906–1908 (2005).
15. Holmes, M. V., Ala-Korpela, M. & Davey Smith, G. Mendelian randomization in cardiometabolic disease: challenges in evaluating causality. *Nat. Rev. Cardiol.* **14**, 577–590 (2017).
16. Zhu, Z. et al. Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. *Nat. Genet.* **48**, 481–487 (2016).
17. Vösa, U. et al. Large-scale cis- and trans-eQTL analyses identify thousands of genetic loci and polygenic scores that regulate blood gene expression. *Nat. Genet.* **53**, 1300–1310 (2021).
18. Wang, D. et al. Comprehensive functional genomic resource and integrative model for the human brain. *Science* **362**, eaat8464 (2018).
19. Burgess, S., Dudbridge, F. & Thompson, S. G. Combining information on multiple instrumental variables in Mendelian randomization: Comparison of allele score and summarized data methods. *Stat. Med.* **35**, 1880–1906 (2016).
20. The 1000 Genomes Project Consortium. An integrated map of genetic variation from 1,092 human genomes. *Nature* **491**, 56–65 (2012).
21. Nalls, M. A. et al. Large-scale meta-analysis of genome-wide association data identifies six new risk loci for Parkinson's disease. *Nat. Genet.* **46**, 989–993 (2014).
22. Rose, A. A. N., Biondini, M., Curiel, R. & Siegel, P. M. Targeting GPNMB with glembatumumab vedotin: current developments and future opportunities for the treatment of cancer. *Pharmacol. Ther.* **179**, 127–141 (2017).
23. Wang, L. H. & Johnson, E. M. Mixed lineage kinase inhibitor CEP-1347 fails to delay disability in early Parkinson's disease. *Neurology* **71**, 462–463 (2008).
24. Baird, D. A. et al. Identifying drug targets for neurological and psychiatric disease via genetics and the brain transcriptome. *PLoS Genet.* **17**, 1–26 (2021).
25. Burgess, S. et al. Guidelines for performing Mendelian randomization investigations. *Wellcome Open Res.* **4**, 186 (2020).
26. Bowden, J. et al. Assessing the suitability of summary data for two-sample Mendelian randomization analyses using MR-Egger regression: the role of the I² statistic. *Int. J. Epidemiol.* **45**, 1961–1974 (2016).
27. Haycock, P. C. et al. Statistical Commentary Best (but oft-forgotten) practices: the design, analysis, and interpretation of Mendelian randomization studies 1. *Am. J. Clin. Nutr.* **103**, 965–978 (2016).
28. Slob, E. A. W. & Burgess, S. A comparison of robust Mendelian randomization methods using summary data. *Genet. Epidemiol.* 1–17, <https://doi.org/10.1002/gepi.22295> (2020).
29. Yavorska, O. O. & Burgess, S. MendelianRandomization: an R package for performing Mendelian randomization analyses using summarized data. *Int. J. Epidemiol.* **46**, 1734–1739 (2017).
30. Hemani, G., Bowden, J. & Davey Smith, G. Evaluating the potential role of pleiotropy in Mendelian randomization studies. *Hum. Mol. Genet.* **27**, R195–R208 (2018).
31. Giambartolomei, C. et al. Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet.* **10**, e1004383 (2014).
32. Zheng, J. et al. Phenome-wide Mendelian randomization mapping the influence of the plasma proteome on complex diseases. *Nat. Genet.* **52**, 1122–1131 (2020).
33. Kia, D. A. et al. Integration of eQTL and Parkinson's disease GWAS data implicates 11 disease genes. *JAMA Neurol.* <https://doi.org/10.1101/627216> (2020).
34. Escott-Price, V. et al. Polygenic risk of Parkinson disease is correlated with disease age at onset. *Ann. Neurol.* **77**, 582–591 (2015).
35. Ibanez, L. et al. Parkinson disease polygenic risk score is associated with Parkinson disease status and age at onset but not with alpha-synuclein cerebrospinal fluid levels. *BMC Neurol.* **17**, 1–9 (2017).
36. Nalls, M. A. et al. Genetic risk and age in Parkinson's disease: continuum not stratum. *Mov. Disord.* **30**, 850–854 (2015).
37. Sun, B. B. et al. Genomic atlas of the human plasma proteome. *Nature* **558**, 73–79 (2018).
38. Emilsson, V. et al. Co-regulatory networks of human serum proteins link genetics to disease. *Science* **361**, 769–773 (2018).
39. Suhre, K. et al. Connecting genetic risk to disease end points through the human blood plasma proteome. *Nat. Commun.* **8**, 14357 (2017).
40. Hillary, R. F. et al. Genome and epigenome wide studies of neurological protein biomarkers in the Lothian Birth Cohort 1936. *Nat. Commun.* **10**, 3160 (2019).
41. Polytynie, T. & Athauda, D. *Repurposing anti-diabetic drugs for the treatment of Parkinson's disease: Rationale and clinical experience. Progress in Brain Research* 252 (Elsevier B.V., 2020).
42. Rotermund, C., Machetanz, G. & Fitzgerald, J. C. The therapeutic potential of metformin in neurodegenerative diseases. *Front. Endocrinol.* **9**, 1–26 (2018).
43. Mor, D. E. et al. Metformin rescues Parkinson's disease phenotypes caused by hyperactive mitochondria. *PNAS* **117**, 26438–26447 (2020).
44. Shi, Q., Liu, S., Fonseca, V. A., Thethi, T. K. & Shi, L. Effect of metformin on neurodegenerative disease among elderly adult US veterans with type 2 diabetes mellitus. *BMJ Open* **9**, e024954 (2019).
45. Hirschhorn, J. N., Lohmueller, K., Byrne, E. & Hirschhorn, K. A comprehensive review of genetic association studies. *Genet. Med.* **4**, 45–61 (2002).
46. Chanoock, S. J. et al. Replicating genotype-phenotype associations. *Nature* <https://doi.org/10.1038/447655a> (2007).
47. Marigorta, U. M., Rodriguez, J. A., Gibson, G. & Navarro, A. Replicability and prediction: lessons and challenges from GWAS. *Trends Genet.* **34**, 504–517 (2018).
48. Burgess, S., Foley, C. N. & Zuber, V. Inferring causal relationships between risk factors and outcomes from genome-wide association study data stephen. *Annu. Rev. Genom. Hum. Genet.* 303–27, <https://doi.org/10.1146/annurev-genom-083117-021731-021731> (2018).
49. Huffman, J. E. Examining the current standards for genetic discovery and replication in the era of mega-biobanks. *Nat. Commun.* **9**, 1–4 (2018).
50. Paternoster, L., Tilling, K. & Smith, G. D. Genetic epidemiology and Mendelian randomization for informing disease therapeutics: conceptual and methodological challenges. 1–10, <https://doi.org/10.1371/journal.pgen.1006944> (2017).
51. Munafò, M. R., Tilling, K., Taylor, A. E., Evans, D. M. & Davey Smith, G. Collider scope: when selection bias can substantially influence observed associations. *Int. J. Epidemiol.* **47**, 226–235 (2018).
52. Bartlett, J. W., Harel, O. & Carpenter, J. R. Asymptotically unbiased estimation of exposure odds ratios in complete records logistic regression. *Am. J. Epidemiol.* **182**, 730–736 (2015).
53. Gkatzionis, A. & Burgess, S. Contextualizing selection bias in Mendelian randomization: How bad is it likely to be? *Int. J. Epidemiol.* **48**, 691–701 (2019).
54. Griffith, G. J. et al. Collider bias undermines our understanding of COVID-19 disease risk and severity. *Nat. Commun.* **11**, 1–12 (2020).
55. Ferreira, P. G. et al. The effects of death and post-mortem cold ischemia on human tissue transcriptomes. *Nat. Commun.* **9**, 490 (2018).
56. Hegvik, T. A. et al. Druggable genome in attention deficit/hyperactivity disorder and its co-morbid conditions. New avenues for treatment. *Mol. Psychiatry* <https://doi.org/10.1038/s41380-019-0540-z> (2019).
57. Jacobs, B. M. et al. Summary-data-based Mendelian randomization prioritizes potential druggable targets for multiple sclerosis. *Brain Commun.* **2**, 1–17 (2020).
58. Storm, C. S. Finding genetically-supported drug targets for Parkinson's disease using Mendelian randomization of the druggable genome. *GitHub Repos.* (2020).
59. R Core Team. *R: A Language and Environment for Statistical Computing* (2019).
60. Willer, C. J., Li, Y. & Abecasis, G. R. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* **26**, 2190–2191 (2010).
61. Hemani, G. et al. The MR-Base platform supports systematic causal inference across the human phenome. *Elife* **7**, e34408 (2018).
62. Del Greco M, F., Minelli, C., Sheehan, N. A. & Thompson, J. R. Detecting pleiotropy in Mendelian randomisation studies with summary data and a continuous outcome. *Stat. Med.* **34**, 2926–2940 (2015).

Acknowledgements

C.S.S. would like to thank Dr. Vishal Rawji for his invaluable support and insightful ideas about the clinical implications and communication of this study. C.S.S. is funded by Rosetrees Trust, John Black Charitable Foundation and the University College London MBPhD Programme. D.A.K. is supported by an MBPhD Award from the International Journal of Experimental Pathology. M.A. is funded by the Faculty of Applied Medical Sciences, King Abdulaziz University, Jeddah, Saudi Arabia. N.W.W. and A.D.H. are National Institute for Health Research senior investigators. N.W.W., A.D.H. and C.F. receive support from the National Institute for Health Research University College London Hospitals Biomedical Research Centre. We would like to thank all members of the International Parkinson Disease Genomics Consortium (IPDGC) and the authors of QTL projects referenced here, who make their data openly available. We thank all the patients and families whose decision to donate tissue samples make our research possible.

Author contributions

Conceptualization: C.S.S., D.A.K., M.M.A. and N.W.W.; methodology: C.S.S., D.A.K., M.M.A. and N.W.W.; investigation, formal analysis, visualization: C.S.S.; resources: S.B.C., C.F., A.D.H. and IPDGC; writing—original draft: C.S.S.; writing—review & editing: C.S.S., D.A.K., M.M.A., N.W.W., S.B.C., C.F., A.D.H. and IPDGC.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41467-021-26280-1>.

Correspondence and requests for materials should be addressed to Nicholas W. Wood.

Peer review information *Nature Communications* thanks Neil Davies, Haydeh Payami and Heiko Runz for their contribution to the peer review of this work. Peer reviewer reports are available.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021

International Parkinson's Disease Genomics Consortium (IPDGC)

Alastair J. Noyce^{8,9}, Rauan Kaiyrzhanov¹⁰, Ben Middlehurst¹¹, Demis A. Kia¹, Manuela Tan¹², Henry Houlden¹⁰, Catherine S. Storm¹³, Huw R. Morris¹², Helene Plun-Favreau¹⁰, Peter Holmans^{13,14}, John Hardy¹⁰, Daniah Trabzuni¹⁰, John Quinn¹¹, Vivien Bubb¹¹, Kin Y. Mok¹⁰, Kerri J. Kinghorn¹⁵, Nicholas W. Wood¹⁶, Patrick Lewis¹⁶, Sebastian R. Schreglmann¹⁰, Ruth Lovering¹⁷, Lea R'Bibo¹⁰, Claudia Manzoni¹⁶, Mie Rizig¹⁰, Mina Ryten¹⁰, Sebastian Guelfi¹⁰, Valentina Escott-Price¹⁸, Viorica Chelban¹⁰, Thomas Foltynie¹⁹, Nigel Williams²⁰, Karen E. Morrison²¹, Carl Clarke^{22,23}, Kirsten Harvey²⁴, Benjamin M. Jacobs⁸, Alexis Brice²⁵, Fabrice Danjou²⁵, Suzanne Lesage²⁵, Jean-Christophe Corvol²⁶, Maria Martinez^{27,28}, Claudia Schulte^{29,30}, Kathrin Brockmann^{29,30}, Javier Simón-Sánchez^{29,30}, Peter Heutink³¹, Patrizia Rizzu³², Manu Sharma³³, Thomas Gasser^{29,30}, Susanne A. Schneider³⁴, Mark R. Cookson³, Sara Bandres-Ciga³, Cornelis Blauwendraat³, David W. Craig³⁵, Kimberley Billingsley³, Mary B. Makarios³, Derek P. Narendra³⁶, Faraz Faghri^{3,37}, J. Raphael Gibbs³⁸, Dena G. Hernandez³, Kendall Van Keuren-Jensen³⁹, Joshua M. Shulman^{40,41}, Hirotaka Iwaki³, Hampton L. Leonard³, Mike A. Nalls^{3,42}, Laurie Robak⁴³, Jose Bras⁴⁴, Rita Guerreiro⁴⁴, Steven Lubbe⁴⁵, Timothy Troycoco⁴⁶, Steven Finkbeiner^{47,48,49}, Niccolo E. Mencacci⁵⁰, Codrin Lungu⁵¹, Andrew B. Singleton³, Sonja W. Scholz⁵², Xylena Reed³, Ryan J. Uitti⁵³, Owen A. Ross⁵⁴, Francis P. Grenn³, Anni Moore³, Roy N. Alcalay^{55,56}, Zbigniew K. Wszolek⁵³, Ziv Gan-Or⁵⁷, Guy A. Rouleau⁵⁷, Lynne Krohn⁵⁷, Kheireddin Mufti⁵⁷, Jacobus J. van Hilten⁵⁸, Johan Marinus⁵⁸, Astrid D. Adames-Gómez⁵⁹, Miquel Aguilar⁶⁰, Ignacio Alvarez⁶⁰, Victoria Alvarez⁶¹, Francisco Javier Barrero⁶², Jesús Alberto Bergareche Yarza⁶³, Inmaculada Bernal-Bernal⁵⁹, Marta Blazquez⁶¹, Marta Bonilla-Toribio⁵⁹, Juan A. Botía⁶⁴, María Teresa Bongiorno⁶⁰, Dolores Buiza-Rueda⁵⁹, Ana Cámara⁶⁵, Fátima Carrillo⁵⁹, Mario Carrión-Claro⁵⁹, Debora Cerdan⁶⁶, Jordi Clarimón^{67,68}, Yaroslau Compta⁶⁵, Monica Diez-Fairen⁶⁰, Oriol Dols-Icardo^{67,68}, Jacinto Duarte⁶⁶, Raquel Duran⁶⁹, Francisco Escamilla-Sevilla⁷⁰, Mario Ezquerra⁶⁵, Cici Feliz⁷¹, Manel Fernández⁶⁵, Rubén Fernández-Santiago⁶⁵, Ciara Garcia⁶¹, Pedro García-Ruiz⁷², Pilar Gómez-Garre⁵⁹, Maria Jose Gomez Heredia⁷³, Isabel Gonzalez-Aramburu⁷⁴, Ana Gorostidi Pagola⁶³, Janet Hoenicka⁷⁵, Jon Infante^{68,76}, Silvia Jesús⁵⁹, Adriano Jimenez-Escrig⁷⁷, Jaime Kulisevsky^{68,78}, Miguel A. Labrador-Espinosa⁵⁹, Jose Luis Lopez-Sendon⁷⁷, Adolfo López de Munain Arregui⁶³, Daniel Macias⁵⁹, Irene Martínez Torres⁷⁹, Juan Marín^{68,78}, Maria Jose Marti⁸⁰, Juan Carlos Martínez-Castrillo⁸¹, Carlota Méndez-del-Barrio⁵⁹, Manuel Menéndez González⁶¹, Marina Mata⁸², Adolfo Mínguez⁸², Pablo Mir⁵⁹, Elisabet Mondragon Rezola⁶³, Esteban Muñoz⁸⁰, Javier Pagonabarraga^{78,83}, Pau Pastor⁶⁰, Francisco Perez Errazquin⁷³, Teresa Periñán-Tocino⁵⁹, Javier Ruiz-Martínez⁸⁴, Clara Ruz⁶⁹, Antonio Sanchez Rodriguez⁷⁴, María Sierra⁷⁴, Esther Suarez-Sanmartin⁶¹, Cesar Tabernero⁶⁶,

Juan Pablo Tartari⁶⁰, Cristina Tejera-Parrado⁵⁹, Eduard Tolosa⁸⁰, Francesc Valldeoriola⁸⁰, Laura Vargas-González⁵⁹, Lydia Vela⁸⁵, Francisco Vives⁶⁹, Alexander Zimprich⁸⁶, Lasse Pihlstrom⁸⁷, Mathias Toft⁸⁸, Pille Taba⁸⁹, Sulev Koks^{90,91}, Sharon Hassin-Baer^{92,93}, Kari Majamaa^{94,95}, Ari Siitonen^{94,95}, Pentti Tienari^{96,97}, Njideka U. Okubadejo⁹⁸, Oluwadamilola O. Ojo⁹⁸, Chingiz Shashkin⁹⁹, Nazira Zharkinbekova¹⁰⁰, Vadim Akhmetzhanov¹⁰¹, Gulnaz Kaishybayeva¹⁰², Altynay Karimova¹⁰², Talgat Khaibullin¹⁰³ & Timothy L. Lynch^{104,105}

⁸Preventive Neurology Unit, Wolfson Institute of Preventive Medicine, QMUL, London, UK. ⁹Department of Molecular Neuroscience, UCL, London, UK. ¹⁰Department of Molecular Neuroscience, UCL Institute of Neurology, London, UK. ¹¹Institute of Translational Medicine, University of Liverpool, Liverpool, UK. ¹²Department of Clinical Neuroscience, University College London, London, UK. ¹³Biostatistics and Bioinformatics Unit, Institute of Psychological Medicine and Clinical Neuroscience, Cardiff, UK. ¹⁴MRC Centre for Neuropsychiatric Genetics & Genomics, Cardiff, UK. ¹⁵Institute of Healthy Ageing, Research Department of Genetics, Evolution and Environment, University College London, London, UK. ¹⁶University of Reading, Reading, UK. ¹⁷University College London, London, UK. ¹⁸MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff University School of Medicine, Cardiff, UK. ¹⁹UCL Institute of Neurology, London, UK. ²⁰MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff, UK. ²¹Faculty of Medicine, University of Southampton, Southampton, UK. ²²University of Birmingham, Birmingham, UK. ²³Sandwell and West Birmingham Hospitals NHS Trust, Birmingham, UK. ²⁴UCL School of Pharmacy, London, UK. ²⁵Institut du Cerveau et de la Moelle épinière, ICM, Inserm U 1127, CNRS, UMR 7225, Sorbonne Universités, UPMC University Paris 06, UMR 5 1127, AP-HP, Pitié-Salpêtrière Hospital, Paris, France. ²⁶Institut du Cerveau et de la Moelle épinière, ICM, Inserm U 1127, CNRS, UMR 7225, Sorbonne Universités, UPMC University Paris 06, UMR 5 1127, Centre d'Investigation Clinique Pitié Neurosciences CIC-1422, AP-HP, Pitié-Salpêtrière Hospital, Paris, France. ²⁷INSERM UMR 1220, Toulouse, France. ²⁸Paul Sabatier University, Toulouse, France. ²⁹Department for Neurodegenerative Diseases, Hertie Institute for Clinical Brain Research, University of Tübingen, Tübingen, Germany. ³⁰DZNE, German Center for Neurodegenerative Diseases, Tübingen, Germany. ³¹DZNE, German Center for Neurodegenerative Diseases and Department for Neurodegenerative Diseases, Hertie Institute for Clinical Brain Research, University of Tübingen, Tübingen, Germany. ³²DZNE, German Center for Neurodegenerative Diseases, Bonn, Germany. ³³Centre for Genetic Epidemiology, Institute for Clinical Epidemiology and Applied Biometry, University of Tübingen, Tübingen, Germany. ³⁴Department of Neurology, Ludwig-Maximilians-University Munich, München, Germany. ³⁵Department of Translational Genomics, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA. ³⁶Inherited Movement Disorders Unit, National Institute of Neurological Disorders and Stroke, Bethesda, MD, USA. ³⁷Department of Computer Science, University of Illinois at Urbana-Champaign, Urbana, IL, USA. ³⁸Laboratory of Neurogenetics, National Institute on Aging, National Institutes of Health, Bethesda, MD, USA. ³⁹Neurogenomics Division, TGen, Phoenix, AZ, USA. ⁴⁰Departments of Neurology, Neuroscience, and Molecular & Human Genetics, Baylor College of Medicine, Houston, TX, USA. ⁴¹Jan and Dan Duncan Neurological Research Institute, Texas Children's Hospital, Houston, TX, USA. ⁴²CEO/Consultant Data Tecnica International, Glen Echo, MD, USA. ⁴³Baylor College of Medicine, Houston, TX, USA. ⁴⁴Center for Neurodegenerative Science, Van Andel Research Institute, Grand Rapids, MI, USA. ⁴⁵Ken and Ruth Davee Department of Neurology and Simpson Querrey Center for Neurogenetics, Northwestern University Feinberg School of Medicine, Chicago, IL, USA. ⁴⁶National Institutes of Health, Bethesda, MD, USA. ⁴⁷Departments of Neurology and Physiology, University of California, San Francisco, CA, USA. ⁴⁸Gladstone Institute of Neurological Disease, San Francisco, CA, USA. ⁴⁹Taube/Koret Center for Neurodegenerative Disease Research, San Francisco, CA, USA. ⁵⁰Northwestern University Feinberg School of Medicine, Chicago, IL, USA. ⁵¹National Institutes of Health Division of Clinical Research, NINDS, National Institutes of Health, Bethesda, MD, USA. ⁵²Neurodegenerative Diseases Research Unit, National Institute of Neurological Disorders and Stroke, Bethesda, MD, USA. ⁵³Department of Neurology, Mayo Clinic, Jacksonville, FL, USA. ⁵⁴Departments of Neuroscience & Clinical Genomics, Mayo Clinic, Jacksonville, FL, USA. ⁵⁵Department of Neurology, College of Physicians and Surgeons, Columbia University Medical Center, Taub Institute for Research on Alzheimer's Disease, New York, NY, USA. ⁵⁶The Aging Brain, College of Physicians and Surgeons, Columbia University Medical Center, New York, NY, USA. ⁵⁷Montreal Neurological Institute and Hospital, Department of Neurology & Neurosurgery, Department of Human Genetics, McGill University, Montréal, QC H3A 0G4, Canada. ⁵⁸Department of Neurology, Leiden University Medical Center, Leiden, Netherlands. ⁵⁹Instituto de Biomedicina de Sevilla IBiS Hospital Universitario Virgen del Rocío/CSIC/Universidad de Sevilla, Seville, Spain. ⁶⁰Fundació Docència i Recerca Mútua de Terrassa and Movement Disorders Unit, Department of Neurology, University Hospital Mutua de Terrassa, Terrassa, Barcelona, Spain. ⁶¹Hospital Universitario Central de Asturias, Oviedo, Spain. ⁶²Hospital Universitario San Cecilio de Granada, Universidad de Granada, Granada, Spain. ⁶³Instituto de Investigación Sanitaria Biodonostia, San Sebastián, Spain. ⁶⁴Universidad de Murcia, Murcia, Spain. ⁶⁵Hospital Clinic de Barcelona, Barcelona, Spain. ⁶⁶Hospital General de Segovia, Segovia, Spain. ⁶⁷Memory Unit, Department of Neurology, IIB Sant Pau, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain. ⁶⁸Centro de Investigación Biomédica en Red en Enfermedades Neurodegenerativas (CIBERNED), Madrid, Spain. ⁶⁹Centro de Investigación Biomédica, Universidad de Granada, Granada, Spain. ⁷⁰Hospital Universitario Virgen de las Nieves, Instituto de Investigación Biosanitaria de Granada, Granada, Spain. ⁷¹Departamento de Neurología, Instituto de Investigación Sanitaria Fundación Jiménez Díaz, Madrid, Spain. ⁷²Instituto de Investigación Sanitaria Fundación Jiménez Díaz, Madrid, Spain. ⁷³Hospital Universitario Virgen de la Victoria, Malaga, Spain. ⁷⁴Hospital Universitario Marqués de Valdecilla-IDIVAL, Santander, Spain. ⁷⁵Institut de Recerca Sant Joan de Déu, Barcelona, Spain. ⁷⁶Hospital Universitario Marqués de Valdecilla-IDIVAL and University of Cantabria, Santander, Spain. ⁷⁷Hospital Universitario Ramón y Cajal, Madrid, Spain. ⁷⁸Movement Disorders Unit, Department of Neurology, IIB Sant Pau, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain. ⁷⁹Department of Neurology, Instituto de Investigación Sanitaria La Fe, Hospital Universitario y Politécnico La Fe, Valencia, Spain. ⁸⁰Hospital Clinic Barcelona, Barcelona, Spain. ⁸¹Instituto Ramón y Cajal de Investigación Sanitaria, Hospital Universitario Ramón y Cajal, Madrid, Spain. ⁸²Hospital Universitario Virgen de las Nieves, Granada, Instituto de Investigación Biosanitaria de Granada, Spain. ⁸³Department of Neurology, Hospital Universitario Infanta Sofía, Madrid, Spain. ⁸⁴Hospital Universitario Donostia, Instituto de Investigación Sanitaria Biodonostia, San Sebastián, Spain. ⁸⁵Department of Neurology, Hospital Universitario Fundación Alcorcón, Madrid, Spain. ⁸⁶Department of Neurology, Medical University of Vienna, Vienna, Austria. ⁸⁷Department of Neurology, Oslo University Hospital, Oslo, Norway. ⁸⁸Department of Neurology and Institute of Clinical Medicine, Oslo University Hospital, Oslo, Norway. ⁸⁹Department of Neurology and Neurosurgery, University of Tartu, Tartu, Estonia. ⁹⁰Centre for Molecular Medicine and Innovative Therapeutics, Murdoch University, Murdoch, 6150 Perth, Western Australia, Australia. ⁹¹The Perron Institute for Neurological and Translational Science, Nedlands, 6009 Perth, Western Australia, Australia. ⁹²The Movement Disorders Institute, Department of Neurology and Sagol Neuroscience Center, Chaim Sheba Medical Center, Tel-Hashomer, 5262101 Ramat

Gan, Israel. ⁹³Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel. ⁹⁴Institute of Clinical Medicine, Department of Neurology, University of Oulu, Oulu, Finland. ⁹⁵Department of Neurology and Medical Research Center, Oulu University Hospital, Oulu, Finland. ⁹⁶Clinical Neurosciences, Neurology, University of Helsinki, Helsinki, Finland. ⁹⁷Helsinki University Hospital, Helsinki, Finland. ⁹⁸University of Lagos, Lagos State, Nigeria. ⁹⁹Kazakh National Medical University named after Asfendiyarov, Almaty, Kazakhstan. ¹⁰⁰South Kazakhstan Medical Academy, Shymkent, Kazakhstan. ¹⁰¹Astana Medical University, Astana, Kazakhstan. ¹⁰²Scientific and Practical Center “Institute of Neurology named after Smagul Kaishibayev”, Almaty, Kazakhstan. ¹⁰³Semey Medical University, Semey, Kazakhstan. ¹⁰⁴School of Medicine and Medical Science, University College Dublin, Dublin, Ireland. ¹⁰⁵The Dublin Neurological Institute at the Mater Misericordiae University Hospital, Dublin, Ireland.