



## Original Research Article

# Dietary lycopene supplementation improves meat quality, antioxidant capacity and skeletal muscle fiber type transformation in finishing pigs

Wanxue Wen<sup>a,1</sup>, Xiaoling Chen<sup>a,1</sup>, Zhiqing Huang<sup>a,\*</sup>, Daiwen Chen<sup>a</sup>, Bing Yu<sup>a</sup>, Jun He<sup>a</sup>, Yuheng Luo<sup>a</sup>, Hui Yan<sup>a</sup>, Hong Chen<sup>b</sup>, Ping Zheng<sup>a</sup>, Jie Yu<sup>a</sup>

<sup>a</sup> Key Laboratory for Animal Disease-Resistance Nutrition of China Ministry of Education, Institute of Animal Nutrition, Sichuan Agricultural University, Chengdu, Sichuan 611130, China

<sup>b</sup> College of Food Science, Sichuan Agricultural University, Ya'an, Sichuan 625014, China

## ARTICLE INFO

## Article history:

Received 9 March 2021

Received in revised form

4 June 2021

Accepted 18 June 2021

Available online 29 September 2021

## Keywords:

Lycopene

Meat quality

Antioxidant capacity

Skeletal muscle fiber type transformation

Finishing pig

## ABSTRACT

This study aimed to investigate effects of dietary lycopene supplementation on meat quality, antioxidant ability and muscle fiber type transformation in finishing pigs. In a 70-day experiment, 18 Duroc × Landrace × Yorkshire barrows were randomly allocated to 3 dietary treatments including a basal diet supplemented with 0, 100 and 200 mg/kg lycopene, respectively. Each dietary treatment had 6 replicates with one pig each. Results showed that dietary 200 mg/kg lycopene supplementation increased muscle redness  $a^*$  value, intramuscular fat and crude protein contents, and decreased muscle lightness  $L^*$  and yellowness  $b^*$  values ( $P < 0.05$ ), suggesting that addition of 200 mg/kg lycopene to the diet of finishing pigs improved color, nutritional value and juiciness of pork after slaughter. Results also showed that dietary lycopene supplementation enhanced antioxidant capacity of finishing pigs ( $P < 0.05$ ). Moreover, dietary supplementation of 200 mg/kg lycopene significantly increased slow myosin heavy chain (MyHC) protein level and slow-twitch fiber percentage, and decreased fast MyHC protein level and fast-twitch fiber percentage ( $P < 0.05$ ), suggesting that the addition of 200 mg/kg lycopene to the diet of finishing pigs promoted muscle fiber type conversion from fast-twitch to slow-twitch. Together, we provide the first evidence that dietary 200 mg/kg lycopene supplementation improves meat quality, enhances antioxidant capacity and promotes muscle fiber type transformation from fast-twitch to slow-twitch in finishing pigs.

© 2021 Chinese Association of Animal Science and Veterinary Medicine. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co. Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Muscle fibers are the basic muscle units and are divided into type I (slow-twitch, slow oxidative), type IIa (fast-twitch, fast oxidative), type IIx (fast-twitch, intermediate) and type IIb (fast-

twitch, fast glycolytic) based on the myosin heavy chain (MyHC) polymorphism (Xu et al., 2020a). Slow-twitch fibers are richer in myoglobin, and mitochondria, and have a higher ability of oxidative metabolism compared with fast-twitch fibers (Xu et al., 2020b). In animal production, the composition and proportion of different muscle fiber types are closely related to muscle color, tenderness and flavor (Cho et al., 2019; Li et al., 2018). In the pursuit of safer, healthier, higher quality and nutritional value pork, more and more attention has been focused on improving meat quality by controlling skeletal muscle fiber. Recent studies have indicated that dietary supplementation of certain nutrients could improve meat quality by inducing skeletal muscle fiber type conversion (Zhang et al., 2015, 2019). For example, dietary resveratrol supplementation improved pork quality, and its underlying mechanism was partly due to the alteration of muscle fiber type composition (Wen et al., 2020a; Zhang et al., 2015). In addition, dietary supplementation of

\* Corresponding author.

E-mail address: [zqhuang@sicau.edu.cn](mailto:zqhuang@sicau.edu.cn) (Z. Huang).

<sup>1</sup> These authors contributed equally to this work.

Peer review under responsibility of Chinese Association of Animal Science and Veterinary Medicine.



betaine (Wen et al., 2019) and methionine (Lebret et al., 2018) improved meat quality by enhancing antioxidant capacity.

Lycopene, a member of the carotenoid family, is mainly extracted from tomatoes and guavas and has various biological functions such as having a strong antioxidant capacity, and being anti-inflammatory, anti-cancer and anti-apoptosis (Costa-Rodrigues et al., 2018). It has been reported that lycopene reduced reactive oxygen species (ROS) generation and malondialdehyde (MDA) content, and increased the enzyme activities of superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px) in L02 cells (Xu et al., 2019). Moreover, lycopene effectively decreased the MDA content in liver and in plasma, and improved the meat quality of rainbow trout (Yonar, 2012). Turkey breast muscle treated with lycopene significantly increased the ratio of unsaturated fatty acid to saturated fatty acid and improved meat quality (Skiepkó et al., 2016). In finishing pigs, dietary lycopene supplementation affected the cellular and humoral immune response (Fachinello et al., 2018). However, effects of dietary lycopene supplementation on meat quality and antioxidant capacity in finishing pigs have not been reported. It has been reported that the AMP-activated protein kinase (AMPK) signaling pathway is closely associated with muscle fiber type transformation and meat quality (Wen et al., 2020b; Xu et al., 2020b). Based on lycopene effectively activating the AMPK signaling in mice (Lin et al., 2018), we speculated that lycopene might regulate skeletal muscle fiber type transformation and improve pork quality. To verify our hypothesis, this study was performed to investigate the effects of dietary lycopene supplementation on meat quality, antioxidant capacity and skeletal muscle fiber type transformation in finishing pigs.

## 2. Material and methods

### 2.1. Ethics statement

The experimental procedures were approved by the Animal Care Advisory Committee of Sichuan Agricultural University under permit no. YYS200208.

### 2.2. Experimental design and diets

A total of 18 healthy castrated Duroc × Landrace × Yorkshire (DLY) pigs with an average body weight (BW) of  $63.89 \pm 1.15$  kg were randomly allocated to 3 dietary treatments including a basal diet supplemented with 0, 100 and 200 mg/kg lycopene, respectively. There were 6 replicates in each group and 1 pig in each replicate. The pigs were individually housed in an independent pen and in the same room. The basal diet (Table 1) was formulated based on a maize-soybean meal and met the nutrient recommendations for growing-finishing pigs by the National Research Council (NRC 2012). Lycopene (a solid powder with 10% of pure lycopene and 90% of carrier dextrin stored under low temperature conditions and kept in a dark place) used in this study was purchased from Xi'an Xiaocao Plant Technology Co., Ltd (Xi'an, China). For all groups, solid powdered diets were prepared every 2 weeks and stored in a dark room under low temperature conditions. The experiment lasted 10 weeks. All pigs had free access to water and were fed 3 times (08:30, 14:30, 20:30) per day, each time with a little excess feed in the trough after the pigs were full. Then we summed the weight of 3 meals and then recorded this as the daily feed intake. The initial and final BW of fasting pigs were measured individually at the beginning and end of the experiment, respectively, using ST200L (Changzhou Lingheng Instrument Co., Ltd.) in the morning. The average daily feed intake (ADFI), average daily weight gain (ADG) and the ratio of feed to gain (F:G) were calculated.

**Table 1**

Feed ingredients and nutrient levels of the basal diet (% air-dry basis).

Item	Content
<b>Ingredients</b>	
Corn	81.57
Soybean meal	11.48
Soybean oil	2.30
Wheat bran	2.00
L-Lysine-HCl	0.39
DL-Methionine	0.09
L-Threonine	0.13
L-Tryptophan	0.03
Choline chloride	0.10
Limestone	0.70
Dicalcium phosphate	0.68
NaCl	0.30
Vitamin premix <sup>1</sup>	0.03
Mineral premix <sup>2</sup>	0.20
<b>Total</b>	<b>100.00</b>
<b>Nutrient levels</b>	
Digestible energy, Mcal/kg	3.40
Crude protein	12.26
Calcium	0.54
Total phosphorus	0.43
Available phosphorus	0.24
Lysine	0.73
Methionine	0.26
Threonine	0.46
Tryptophan	0.13

<sup>1</sup> Vitamin premix provides the following per kilogram of complete diet: vitamin A, 9,000 IU; vitamin D<sub>3</sub>, 3,000 IU; vitamin E, 24 IU; vitamin K<sub>3</sub>, 3 mg; vitamin B<sub>12</sub>, 0.036 mg; vitamin B<sub>1</sub>, 3 mg; vitamin B<sub>6</sub>, 3.6 mg; vitamin B<sub>2</sub>, 7.5 mg; vitamin B<sub>5</sub>, 15 mg; folic acid, 1.5 mg; biotin, 0.15 mg; nicotinamide, 30 mg.

<sup>2</sup> Mineral premix provides the following per kilogram of complete diet: Se (Na<sub>2</sub>SeO<sub>3</sub>) 0.15 mg; I (KI) 0.14 mg; Zn (ZnSO<sub>4</sub>) 50 mg; Cu (CuSO<sub>4</sub> · 5H<sub>2</sub>O) 3 mg; Fe (FeSO<sub>4</sub> · H<sub>2</sub>O) 40 mg; Mn (MnSO<sub>4</sub>) 2 mg.

### 2.3. Sample collection and carcass characteristics measurement

At the end of the experiment, fasting blood samples of finishing pigs (20 mL) were collected from the jugular vein. Samples were kept at room temperature for 30 min and then centrifuged at 3,000 × g for 10 min. Serum samples were stored at −20 °C until analysis. Pigs were slaughtered by electronarcosis and exsanguination based on the standard commercial procedures. Carcass weight and tare weight were used to calculate the dressing percentage. The average backfat thickness was calculated by the average thickness of the first rib, last rib and last lumbar vertebra. The eye muscle area (EMA) was measured at the last rib using vernier calipers. The longissimus dorsi (LD) muscle from the left side of each carcass was used to measure the meat color, marbling score, pH, shear force, dripping loss, cooking loss, intramuscular fat (IMF) content and crude protein content. LD muscle from the 10th rib of the left carcass was used for RNA and protein analysis.

### 2.4. Meat quality measurement

Meat color (L\* lightness, a\* redness and b\* yellowness) was recorded at 45 min after slaughter by a portable chromameter (CR-300, Minolta, Japan) calibrated with a white tile. The marbling score at 24 h was measured using the National Pork Producer Council (NPPC, Des Moines, IA, USA). Muscle pH<sub>45min</sub> and pH<sub>24h</sub> were calculated by a pH meter (pH-STAR, SFK-Technology, Denmark) calibrated with pH 4.6 and pH 7.0 buffers. Drip loss was measured according to the method described previously (Xu et al., 2019). About 30 g of muscle (a 3 cm length cube) was weighed (W<sub>1</sub>) after 45 min postmortem, and hung in a storage bag with a fishhook at 4 °C for 24 h, after which the muscle was removed from the

fishhook, blotted dry and reweighed ( $W_2$ ). The drip loss value was expressed as follows: drip loss (%) =  $(W_1 - W_2)/W_1 \times 100$ . Cooking loss was determined as described previously (Yu et al., 2020). About 120 g of muscle was weighed ( $W_3$ ) and cooked in a steamer for 30 min, after which the muscle sample was quickly removed from the steamer, cooled for 20 min at room temperature and reweighed ( $W_4$ ). The cooking loss value was expressed as follows: cooking loss (%) =  $(W_3 - W_4)/W_3 \times 100$ . The shear force was measured using a texture analyzer (TA.XT. Plus, Stable Micro Systems, Godalming, UK) according to the manufacturer's manual.

### 2.5. Crude protein and intramuscular fat contents measurement

Crude protein and intramuscular fat contents were determined as described previously (Xu et al., 2019). About 50 g of LD muscle was sliced up, weighed ( $W_1$ ), removed into a weighing bottle, and weighed ( $W_2$ ). The weighing bottle was placed into a freeze dryer for 48 h at  $-50^\circ\text{C}$  and reweighed ( $W_3$ ). The moisture percentage was expressed as following: moisture percentage (%) =  $(W_2 - W_3)/(W_2 - W_1) \times 100$ . The dried samples were pulverized by a muller and then the contents of crude protein and intramuscular fat were measured according to methods of the Association of Analytical Chemists (AOAC; Rockville, MD, USA).

### 2.6. Immunofluorescence

The LD muscle samples from finishing pigs stored in 4% paraformaldehyde were embedded in paraffin. LD muscles were sliced into 10- $\mu\text{m}$  thick sections by a pathology microtome (Leica, China). Sections were blocked with 3% bovine serum albumin (BSA) for 30 min and then incubated with primary antibodies (1:500, slow MyHC, Abcam, Cat. No. ab11083; 1:500, fast MyHC, Abcam, Cat. No. ab91506). After being washed with PBS (pH 7.4), muscle sections were incubated with goat anti-rabbit IgG (1:300, Sevicebio, Cat. No. GB21303) or goat anti-mouse IgG (1:300, Sevicebio, Cat. No. GB25301). DAPI (Servicebio, Cat. No. G1012) was used to stain the nucleus. Images were collected by Fluorescent Microscopy (Nikon, Japan). The numbers of slow-twitch fiber and fast-twitch fiber were counted by Image-Pro Plus 6.0.

### 2.7. Antioxidant status assay

The total antioxidant capacity (T-AOC), total superoxide dismutase (T-SOD), GSH-Px and catalase (CAT) activities and the MDA content in serum, liver and LD muscle were measured by commercial kits according to the manufacturers' instructions, which were purchased from Nanjing Jiancheng Bioengineering Institute (Nanjing, China).

### 2.8. Metabolic enzyme activities assay

The activities of succinic dehydrogenase (SDH), malate dehydrogenase (MDH) and lactate dehydrogenase (LDH) were tested using kits purchased from Nanjing Jiancheng Bioengineering Institute (Nanjing, China).

### 2.9. Western blotting

Total proteins were extracted by RIPA lysis buffer (Beyotime, China) and the protein concentrations were measured by BCA protein assay kit (Pierce, Rockford, IL, USA). After denaturing the protein lysates at  $98^\circ\text{C}$  for 10 min, the proteins (20  $\mu\text{g}$ ) were separated with 8% SDS-PAGE and transferred onto PVDF

membranes (Millipore, Eschborn, Germany). Membranes were blocked with 5% BSA in 1  $\times$  TBST (20 mmol/L Tris-HCl pH 7.6, 8 g/L NaCl, and 0.1% Tween-20) and then incubated with primary antibodies at  $4^\circ\text{C}$  for 16 h. The primary antibodies used were as follows: slow MyHC (1:500, Sigma, Cat. No. M8421), fast MyHC (1:500, Sigma, Cat. No. M4276), Myoglobin (1:1,000, Cell Signaling, Cat. No. 25919), cytochrome c (CytC, 1:1,000, ProteinTech Biotechnology, Cat. No. 10993-1-AP), nuclear respiratory factor 1 (NRF1, 1:1,000, Cell Signaling, Cat. No. 46743), calcium/calmodulin-dependent protein kinase  $\beta$  (CaMKK $\beta$ , 1:1,000, Cell Signaling, Cat. No. 16810), phospho-AMPK(p-AMPK, 1:1,000, Cell Signaling, Cat. No. 2535), AMPK (1:1,000, Cell Signaling, Cat. No. 5831), sirtuin1 (Sirt1, 1:1,000, Cell Signaling, Cat. No. 8469), peroxisome proliferator activated receptor- $\gamma$  coactivator-1 $\alpha$  (PGC-1 $\alpha$ , 1:800, Cell Signaling, Cat. No. 2178) and  $\beta$ -actin (1:1,000, Cell Signaling, Cat. No. 4967) antibodies. Subsequently, specific secondary antibodies were used. BeyoECL Plus (Beyotime, China) was used to visualize the protein bands. Gel-Pro Analyzer was used to detect the protein expressions, which were normalized to  $\beta$ -actin protein.

### 2.10. Real-time quantitative PCR

Total RNA from LD muscle was extracted by RNAiso Plus reagent (TaKaRa, China). One microgram of RNA was reverse transcribed into cDNA using a HiScript III RT SuperMix for qPCR (+gDNA wiper) kit (Vazyme, Nanjing, China). Real-time quantitative PCR was used to relatively quantify mRNA by a ChamQ SYBR Color qPCR Master Mix (Vazyme). The list of primer sequences is presented in Table 2. The PCR cycling conditions were as follows: 45 cycles of  $95^\circ\text{C}$  for 15 s and  $60^\circ\text{C}$  for 15 s. The  $2^{-\Delta\Delta\text{Ct}}$  method was used to calculate the relative gene expression normalized to glyceraldehyde-3-phosphate dehydrogenase (GAPDH) mRNA.

### 2.11. Statistical analyses

All data were analyzed by one-way analysis of variance followed by Duncan's multiple-range test using SPSS 21.0 software (Chicago, IL, USA). All results were presented as means and standard error of the mean (SEM).  $P < 0.05$  was considered to be statistically significant among groups.

## 3. Results

### 3.1. Growth performance and carcass characteristics

There was no significant difference in growth performance and carcass characteristics of finishing pigs (Table 3;  $P > 0.05$ ).

### 3.2. Meat quality

As shown in Table 4, dietary lycopene supplementation improved the meat quality of finishing pigs. Compared with the control group, dietary 100 mg/kg lycopene supplementation only decreased muscle  $L^*$  value and increased crude protein content, but dietary 200 mg/kg lycopene supplementation significantly increased muscle  $a^*$  value and intramuscular fat and crude protein contents, and decreased muscle  $L^*$  and  $b^*$  values ( $P < 0.05$ ; Table 4).

### 3.3. Muscle antioxidant capacity

As shown in Table 5, dietary 100 and 200 mg/kg lycopene supplementation significantly decreased the content of MDA in liver and LD muscle. Compared with the control group, dietary

**Table 2**  
Primer sequences used in this study.

Gene	Primer	Sequence (5' to 3')	GenBank ID	Size, bp
<i>MyHC I</i>	Forward	CTTCTACAGGCCTGGGCTTAC	NM_080728	128
	Reverse	CTCCTTCTCAGACTCCGCAG		
<i>MyHC IIa</i>	Forward	TTCCAGAAGCCTAAGGTGGTC	NM_001039545	94
	Reverse	GCCAGCCAGTGATGTTGTAAT		
<i>MyHC IIx</i>	Forward	CAACCCATACGACTACGCCT	NM_030679	119
	Reverse	CATCAGAAGTGAAGCCAGAAAT		
<i>MyHC IIb</i>	Forward	CTTGCTGACTCAAGCCTGCC	NM_010855	158
	Reverse	TCGCTCCTTTTCAGACTCCG		
<i>TNNI1</i>	Forward	TGAAGCCAAATGCCTCCACAAC	NM_006529382	155
	Reverse	ACACCTTGTGCTTAGAGCCCACTA		
<i>AMPK<math>\alpha</math>1</i>	Forward	CGGCAAAGTGAAGGTTGG	NM_001167633	123
	Reverse	AGGTTCTGAATTTCTCTGCGG		
<i>AMPK<math>\alpha</math>2</i>	Forward	TGAGGTCGATATCTGGAGCTG	NM_214266	151
	Reverse	AGTGGCAACAGAACGATTGAG		
<i>Sirt1</i>	Forward	ACAGTGACAGTGGCACATGC	NM_019812	130
	Reverse	AATCCAGATCCTCCAGCACA		
<i>PGC-1<math>\alpha</math></i>	Forward	CCAGTACAACAATGAGCCTGC	NM_008904	118
	Reverse	CAATCCGCTTCATCCACG		
<i>Cyt c</i>	Forward	TGCGGAGTGTAAACTTTTCAGG	NM_001129970	191
	Reverse	TCCTTAACAGGCTAGTGAACA		
<i>NRF1</i>	Forward	CTTTGTGGTGGGAGGAATGTT	XM_005657993	77
	Reverse	AGTATGCTGGCTGACCTTGTC		
<i>TFAM</i>	Forward	GCTCTCCGTTCCAGTTTTCGG	AY923074	187
	Reverse	GGAAGTTCCCTCCACAGCTC		
<i>TFB1M</i>	Forward	GCAAGCAGTGAAGCAGCTA	NM_001128475	82
	Reverse	CAGACTGCCAGCTTTCCTTAC		
<i>CS</i>	Forward	AGCCCTCAACAGTGAAGCA	NM_026444	174
	Reverse	TCAATGGCTCCGATACTGCTG		
<i>COX1</i>	Forward	GTGAGTCAGTCACCTTGAGC	NM_001025218	180
	Reverse	TCTGGCCTACTCAGGAAGGA		
<i>SOD1</i>	Forward	AGACCTGGGCAATGTGACTG	NM_001190422	102
	Reverse	GTGCGCCAATGATGAATG		
<i>SOD2</i>	Forward	TGAACAACCTGAACGTCGTG	NM_214127.2	102
	Reverse	AGCGGTCAACTTCTCCTTGA		
<i>CAT</i>	Forward	CAGATGAAGCATTGGAAGGACC	NM_214301	83
	Reverse	TTGTCTCCTATCGGATTCCTCAG		
<i>GPX1</i>	Forward	GTGAATGGCGCAAATGCTCA	NM_214201	126
	Reverse	ATTGGCAGACACTGGAGACC		
<i>GST</i>	Forward	CCAACCCAGAAGACTGCTCA	AB000884	102
	Reverse	CATTACAGGTGGGCTCTTCGT		
<i>GR</i>	Forward	GTGAGCCGACTGAACACCAT	XM_003483635	102
	Reverse	CAGGATGTGAGGAGCTGTGT		
<i>Nrf2</i>	Forward	GCCCCTGGAAGCGTTAAAC	XM_003133500	67
	Reverse	GGAAGTATATCCCAAGGTTGT		
<i>Keap1</i>	Forward	ACGACGTGGAGACAGAAACGT	NM_001114671	56
	Reverse	GCTTCGCGGATGCTTCA		
<i>GAPDH</i>	Forward	ACTCACTTCTACCTTTGATGCT	NM_001206359	100
	Reverse	TGTTGCTGTAGCCAAATTCA		

*MyHC I* = myosin heavy chain I; *MyHC IIa* = myosin heavy chain IIa; *MyHC IIx* = myosin heavy chain IIx; *MyHC IIb* = myosin heavy chain IIb; *TNNI1* = troponin I type 1; *AMPK $\alpha$ 1* = AMP-activated protein kinase  $\alpha$ 1; *AMPK $\alpha$ 2* = AMP-activated protein kinase  $\alpha$ 2; *Sirt1* = sirtuin1; *PGC-1 $\alpha$*  = peroxisome proliferator activated receptor- $\gamma$  coactivator-1 $\alpha$ ; *Cyt c* = cytochrome c; *NRF1* = nuclear respiratory factor 1; *TFAM* = mitochondrial transcription factor A; *TFB1M* = mitochondrial transcription factor B1; *CS* = citrate synthase; *COX1* = cyclooxygenase 1; *SOD1* = superoxide dismutase 1; *SOD2* = superoxide dismutase 2; *CAT* = catalase; *GPX1* = glutathione peroxidase 1; *GST* = glutathione S-transferase; *GR* = glutathione reductase; *NRF2* = nuclear factor erythroid 2-related factor 2; *Keap1* = Kelch-like ECH-associated protein 1; *GAPDH* = glyceraldehyde-3-phosphate dehydrogenase.

100 mg/kg lycopene supplementation not only increased T-AOC and T-SOD activities in serum, but also increased the activities of T-SOD, GSH-Px and CAT in liver ( $P < 0.05$ ; Table 5). Additionally, dietary 200 mg/kg lycopene supplementation significantly increased the GSH-Px and CAT activities in serum, the T-AOC and GSH-Px activities in liver, and the T-SOD and CAT activities in LD muscle ( $P < 0.05$ ; Table 5). Further, dietary 100 mg/kg lycopene supplementation significantly increased the mRNA levels of superoxide dismutase 1 (*SOD1*), glutathione peroxidase 1 (*GPX1*), glutathione S-transferase (*GST*), glutathione reductase (*GR*) and nuclear factor erythroid 2-related factor 2 (*Nrf2*) and decreased the mRNA level of Kelch-like ECH-associated protein 1 (*Keap1*) ( $P < 0.05$ ; Fig. 1). Compared with the control group, dietary 200 mg/kg lycopene supplementation significantly increased the mRNA levels of

superoxide dismutase 1 (*SOD1*), superoxide dismutase 2 (*SOD2*), catalase (*CAT*), *GPX1*, *GST*, *GR* and *Nrf2* and decreased the mRNA level of *Keap1* ( $P < 0.05$ ; Fig. 1). These data suggested that dietary lycopene supplementation enhanced the antioxidant capacity of finishing pigs.

#### 3.4. Skeletal muscle fiber transformation

As shown in Fig. 2A, compared with the control group, dietary supplementation of 100 and 200 mg/kg lycopene significantly promoted slow MyHC protein expression and inhibited fast MyHC protein expression in LD muscle ( $P < 0.05$ ). Compared with the control group, dietary 200 mg/kg lycopene supplementation significantly increased the percentage of slow-twitch fiber and

**Table 3**  
Effect of dietary lycopene supplementation on growth performance and carcass characteristics of finishing pigs.<sup>1</sup>

Item	Lycopene, mg/kg			SEM	P-value
	0	100	200		
Initial weight, kg	64.10	63.63	63.95	1.15	0.988
Final weight, kg	136.57	138.95	135.30	2.10	0.793
ADG, kg/d	1.04	1.08	1.02	0.01	0.553
ADFI, kg/d	3.34	3.31	3.18	0.05	0.469
F:G	3.22	3.09	3.13	0.03	0.322
Carcass weight, kg	103.31	101.22	101.27	2.04	0.904
Dressing percentage, %	75.54	72.79	74.75	0.49	0.058
Carcass length, cm	111.35	111.43	109.13	0.82	0.463
Backfat thickness, cm					
First rib	3.29	3.22	3.49	0.11	0.643
Last rib	2.19	2.49	2.61	0.08	0.112
Last lumbar vertebra	1.58	1.42	1.57	0.07	0.668
Average backfat	2.35	2.38	2.55	0.07	0.522
Eye muscle area, cm <sup>2</sup>	54.74	56.94	58.64	1.47	0.587
Abdominal fat, kg	2.58	2.85	2.20	0.12	0.100
Abdominal fat index <sup>2</sup> , %	1.90	2.06	1.63	0.09	0.165

ADG = average daily gain; ADFI = average daily feed intake; F:G = feed-to-gain ratio.

<sup>1</sup> Results are presented as the mean and SEM (n = 6).

<sup>2</sup> Abdominal fat index (%) = abdominal fat weight/live weight × 100.

**Table 4**  
Effect of dietary lycopene supplementation on meat quality of finishing pigs.

Item	Lycopene, mg/kg			SEM	P-value
	0	100	200		
pH <sub>45min</sub>	6.49	6.53	6.52	0.05	0.961
pH <sub>24h</sub>	5.61	5.57	5.59	0.01	0.329
L* (lightness)	42.17 <sup>b</sup>	41.17 <sup>a</sup>	41.16 <sup>a</sup>	0.13	<0.001
a* (redness)	4.10 <sup>a</sup>	4.32 <sup>a</sup>	4.58 <sup>b</sup>	0.06	<0.001
b* (yellowness)	2.97 <sup>b</sup>	2.50 <sup>ab</sup>	2.45 <sup>a</sup>	0.09	0.029
Drip loss, %	2.58	2.82	2.75	0.05	0.199
Cooking loss, %	33.54	34.38	32.85	0.73	0.719
Shear force, kg	5.53	5.44	5.09	0.13	0.387
Marbling score	2.83	3.67	3.00	0.23	0.319
Intramuscular fat content, %	3.26 <sup>a</sup>	3.56 <sup>ab</sup>	3.98 <sup>b</sup>	0.12	0.047
Crude protein content, %	22.93 <sup>a</sup>	24.74 <sup>b</sup>	25.30 <sup>b</sup>	0.28	<0.001
Moisture, %	70.87	70.02	69.61	0.42	0.486

Results are presented as the mean and SEM (n = 6).

<sup>a, b</sup> Values within a row with different superscripts differ significantly at P < 0.05.

decreased the percentage of fast-twitch fiber (P < 0.05; Fig. 2C). Besides, dietary 200 mg/kg lycopene supplementation notably increased myoglobin protein level (P < 0.05; Fig. 2B). Furthermore, dietary 200 mg/kg lycopene supplementation significantly up-regulated mRNA expressions of *MyHC I*, *MyHC IIa*, *MyHC IIx* and troponin I type 1 (*TNNI1*), whereas it downregulated *MyHC IIb* mRNA expression (P < 0.05; Fig. 2D).

Metabolic enzymes and mitochondrial function indirectly reflect skeletal muscle fiber composition. Our results showed that dietary 200 mg/kg lycopene supplementation significantly decreased LDH activity and increased the activities of SDH and MDH in LD muscle of finishing pigs (P < 0.05; Fig. 2E–G). Our results also showed that dietary lycopene supplementation increased mRNA expressions of mitochondrial function-related genes, such as *Cytc*, *NRF1*, mitochondrial transcription factor A (*TFAM*), mitochondrial transcription factor B1 (*TFB1M*), citrate synthase (*CS*) and cyclooxygenase 1 (*COX1*) (P < 0.05; Fig. 2H). A similar result was also observed in *Cytc* protein level (P < 0.05; Fig. 2I). These results indicated that dietary lycopene

**Table 5**  
Effect of dietary lycopene supplementation on antioxidant capacity of finishing pigs.<sup>1</sup>

Item	Lycopene, mg/kg			SEM	P-value
	0	100	200		
Serum					
MDA, nmol/mL	3.52	2.72	3.10	0.21	0.350
T-AOC, U/mL	0.42 <sup>a</sup>	0.45 <sup>b</sup>	0.43 <sup>ab</sup>	0.00	0.014
T-SOD, U/mL	137.97 <sup>a</sup>	159.63 <sup>b</sup>	159.31 <sup>ab</sup>	4.52	0.011
GSH-Px, U/mL	2,888.69 <sup>a</sup>	3,142.86 <sup>ab</sup>	3,239.77 <sup>b</sup>	54.73	0.015
CAT, U/mL	3.68 <sup>a</sup>	4.15 <sup>a</sup>	5.42 <sup>b</sup>	0.25	0.001
Liver					
MDA, nmol/mg prot	3.97 <sup>b</sup>	2.72 <sup>a</sup>	2.51 <sup>a</sup>	0.25	0.018
T-AOC, U/mg prot	0.63 <sup>a</sup>	0.64 <sup>ab</sup>	0.65 <sup>b</sup>	0.00	0.006
T-SOD, U/mg prot	1,404.42 <sup>a</sup>	1,619.14 <sup>b</sup>	1,481.39 <sup>a</sup>	26.56	<0.001
GSH-Px, U/mg prot	98.92 <sup>a</sup>	118.78 <sup>b</sup>	115.10 <sup>b</sup>	2.50	<0.001
CAT, U/mg prot	7.04 <sup>a</sup>	7.94 <sup>b</sup>	7.64 <sup>ab</sup>	0.15	0.030
LD muscle					
MDA, nmol/mg prot	1.21 <sup>c</sup>	1.01 <sup>b</sup>	0.81 <sup>a</sup>	0.05	<0.001
T-AOC, U/mg prot	0.47	0.48	0.48	0.00	0.237
T-SOD, U/mg prot	49.84 <sup>a</sup>	49.82 <sup>a</sup>	54.10 <sup>b</sup>	0.60	<0.001
GSH-Px, U/mg prot	8.85	11.32	13.83	0.98	0.110
CAT, U/mg prot	0.68 <sup>a</sup>	0.86 <sup>ab</sup>	1.14 <sup>b</sup>	0.07	0.026

T-AOC = total antioxidant capacity; T-SOD = total superoxide dismutase; GSH-Px = glutathione peroxidase; CAT = catalase; MDA = malonaldehyde; prot = protein.

<sup>a, b, c</sup> Values within a row with different superscripts differ significantly at P < 0.05.

<sup>1</sup> Results are presented as the mean and SEM (n = 6).

supplementation promoted skeletal muscle fiber type conversion from fast-twitch to slow-twitch.


### 3.5. Gene and protein expression

To further investigate the underlying mechanism of lycopene on skeletal muscle fiber transformation, we measured the expression levels of AMPK signal components. The result showed that dietary 200 mg/kg lycopene supplementation significantly increased p-AMPK, NRF1, CaMKKβ, Sirt1 and PGC-1α protein levels (P < 0.05; Fig. 3A and B) and *AMPKα1*, *AMPKα2*, *Sirt1* and *PGC-1α* mRNA levels (P < 0.05; Fig. 3C).

## 4. Discussion

In this study, dietary lycopene supplementation improved meat quality, enhanced antioxidant capacity and promoted muscle fiber type transformation from fast-twitch to slow-twitch in finishing pigs. We also found that the mechanism of lycopene on skeletal muscle fiber transformation might be related to the AMPK signaling pathway.

Meat color, intramuscular fat content and nutritional value are the key indicators to assess the meat quality of livestock (Joo et al., 2013; Leseigneur-Meynier and Gandemer, 1991). As far as we know, this study is the first to investigate the effect of dietary lycopene supplementation on the meat quality of finishing pigs. Here we showed that dietary lycopene supplementation significantly increased muscle a\* value and decreased muscle b\* value and L\* value, indicating that dietary lycopene supplementation could improve meat color. Generally, crude protein and intramuscular fat contents are thought to be the main factors in assessing the nutritional value and juiciness of meat (Li et al., 2016; Zhang et al., 2021). Here, we also showed that dietary lycopene supplementation significantly increased muscle crude protein and intramuscular fat contents. Together, our data indicated that the addition of



**Fig. 1.** Effect of dietary lycopene supplementation on the mRNA levels of antioxidant enzyme related genes in longissimus dorsi (LD) muscle of finishing pigs. Real-time quantitative PCR analyzed the mRNA expressions of superoxide dismutase 1 (*SOD1*), superoxide dismutase 2 (*SOD2*), catalase (*CAT*), glutathione peroxidase 1 (*GPX1*), glutathione S-transferase (*GST*), glutathione reductase (*GR*), nuclear factor erythroid 2-related factor 2 (*Nrf2*) and Kelch-like ECH-associated protein 1 (*Keap1*). Data are presented as the mean and SEM ( $n = 6$ ). <sup>a, b, c</sup> Bar with different letter indicates significant difference ( $P < 0.05$ ).

lycopene to the diet of finishing pigs improved color, nutritional value and juiciness of pork after slaughter.


The antioxidant ability is positively correlated with a healthy body and meat quality (Grabowska et al., 2019). Our results showed that dietary lycopene supplementation significantly increased the activities of antioxidant enzymes including T-AOC, CAT, GSH-Px and T-SOD in serum, liver and LD muscle of finishing pigs, and decreased the content of MDA. The activities of antioxidant enzymes are partly dependent on the expression levels of antioxidant enzyme related genes (Wang et al., 2015). Nrf2 is reported to bind to the antioxidant-responsive element (ARE) and has the ability to control the transcriptions of antioxidant enzyme related genes (Chen et al., 2013). Keap1, a Nrf2-binding protein, is reported to prevent the translocation of Nrf2 to the nucleus (Liang et al., 2018). In this study, we showed that dietary lycopene supplementation increased *SOD1*, *SOD2*, *CAT*, *GPX1*, *GST*, *GR* and *Nrf2* mRNA levels, and decreased *Keap1* mRNA level in LD muscle of finishing pigs. These results indicated that dietary lycopene supplementation significantly enhanced the antioxidant ability of finishing pigs, which contributed to improving pork quality.

Muscle fibers account for 90% of the total skeletal muscle. The composition and proportion of different muscle fiber types directly affect muscle physiological and biochemical function and meat quality (Kim et al., 2018; Zhang et al., 2015). Skeletal muscle rich in slow-twitch fiber has a red appearance, better taste and flavor, and stronger aerobic metabolism (Zhang et al., 2019). Our results showed that dietary lycopene supplementation promoted slow MyHC and myoglobin protein expressions and increased the percentage of slow-twitch fiber, whereas it decreased the protein level of fast MyHC and the percentage of fast-twitch fiber. Additionally, dietary lycopene supplementation increased the mRNA levels of *MyHC I* and *TNNI1*, and decreased the mRNA levels of *MyHC IIa*, *MyHC IIx* and *MyHC IIb*. These data suggested that dietary lycopene supplementation promoted muscle fiber type transformation from fast-twitch to slow-twitch. Metabolism enzyme activities indirectly reflect the composition of skeletal muscle fiber type. SDH and MDH belong to oxidative enzymes and are positively associated with the content of slow oxidative fibers, whereas the glycolytic enzyme LDH is associated with the content of fast glycolytic fibers (Chen et al., 2018). In this study, we also showed that dietary lycopene supplementation


increased SDH and MDH enzyme activities, and decreased LDH activity, further supporting that dietary lycopene supplementation promoted muscle fiber type transformation from fast-twitch to slow-twitch. Mitochondrial biogenesis is a new regulator of skeletal muscle fiber type transformation. In this study, our results showed that dietary lycopene supplementation up-regulated the expressions of mitochondrial biogenesis related factors including NRF1, TFAM, TFB1M, CytC, CS and ATP synthase subunit C1 (ATP5G). Taken together, these results suggested that dietary lycopene supplementation promoted muscle fiber type transformation from fast-twitch to slow-twitch, which contributed to improving pork quality.

In this study, we showed that dietary lycopene supplementation promoted slow MyHC and myoglobin protein expressions and increased the percentage of slow-twitch fiber, and decreased the protein level of fast MyHC and the percentage of fast-twitch fiber. It is well known that the slow-twitch fiber has smaller fiber volume (area) than those of the fast-twitch fiber. So, the higher proportion of slow-twitch fiber in the skeletal muscle means a lower area of eye muscle (Zhang et al., 2019). However, it has also been reported that there is no significant correlation between eye muscle area and muscle fiber composition (Ryu et al., 2004). In this study, we also showed that dietary lycopene supplementation increased the area of the eye muscle numerically, although the difference was not statistically significant. So, the relationship between eye muscle area and muscle fiber composition needs to be further studied.

AMPK plays a key role in regulation of skeletal muscle fiber transformation (Xu et al., 2020a, 2020b). To evaluate the potential mechanisms by which lycopene induces skeletal muscle fiber transformation, we firstly measured the effect of dietary lycopene supplementation on AMPK signaling. Our data showed that dietary lycopene supplementation could activate AMPK. NRF1 is an upstream factor of CaMKK $\beta$ , which binds to CaMKK $\beta$  promoter and enhances CaMKK $\beta$  transcription, contributing to increase the expression of p-AMPK (Koh et al., 2017). Sirt1 and PGC-1 $\alpha$ , downstream regulators of AMPK are directly activated by AMPK and drive the transformation of fast glycolytic fiber to slow oxidative fiber (Kulkarni and Cantó, 2015; Li et al., 2002). Meaningfully, our results showed that dietary lycopene supplementation up-regulated the expression levels of NRF1, CaMKK $\beta$ , Sirt1 and PGC-1 $\alpha$ , suggesting that lycopene might activate AMPK through the



**Fig. 2.** Effect of dietary lycopene supplementation on muscle fiber type conversion in longissimus dorsi (LD) muscle of finishing pigs. (A) Western blotting measured the protein expression levels of slow myosin heavy chain (MyHC) and fast MyHC. (B) Western blotting measured the protein expression level of myoglobin. (C) Immunofluorescence analyzed the percentage of slow-twitch fiber and fast-twitch fiber. The white scale bars represent 100  $\mu$ m. (D) Real-time quantitative PCR analyzed the mRNA levels of *MyHC I*, *MyHC IIa*, *MyHC IIx*, *MyHC IIb* and troponin I type 1 (*TNNI1*). (E to G) The activities of succinic dehydrogenase (SDH), malate dehydrogenase (MDH) and lactate dehydrogenase (LDH) enzymes. (H) The mRNA levels of mitochondrial biogenesis related genes including cytochrome c (*Cytc*), nuclear respiratory factor 1 (*NRF1*), mitochondrial transcription factor A (*TFAM*), mitochondrial transcription factor B1 (*TFB1M*), citrate synthase (*CS*) and cyclooxygenase 1 (*COX1*). (I) The protein level of Cytc. Data are presented as the mean and SEM ( $n = 6$ ). a, b, c Bar with a different letter indicates significant difference ( $P < 0.05$ ).



**Fig. 3.** Effect of dietary lycopene supplementation on AMP-activated protein kinase (AMPK) signaling pathway in longissimus dorsi (LD) muscle of finishing pigs. (A) The protein expressions of AMP-activated protein kinase (AMPK) and phospho-AMP-activated protein kinase (p-AMPK). (B) The protein expressions of nuclear respiratory factor 1 (NRF1), calcium/calmodulin-dependent protein kinase  $\beta$  (CaMKK $\beta$ ), sirtuin1 (Sirt1) and peroxisome proliferator activated receptor- $\gamma$  coactivator-1 $\alpha$  (PGC-1 $\alpha$ ). (C) The mRNA expressions of AMP-activated protein kinase  $\alpha$ 1 (AMPK $\alpha$ 1), AMP-activated protein kinase  $\alpha$ 2 (AMPK $\alpha$ 2), sirtuin1 (Sirt1) and peroxisome proliferator activated receptor- $\gamma$  coactivator-1 $\alpha$  (PGC-1 $\alpha$ ). Data are presented as the mean and SEM ( $n = 6$ ). <sup>a, b, c</sup> Bar with different letter indicates significant difference ( $P < 0.05$ ).

NRF1/CaMKK $\beta$  axis and control muscle fiber type conversion via the AMPK signaling pathway.

In summary, this study provides evidence that dietary lycopene supplementation could enhance the antioxidant capacity of finishing pigs. We also found that dietary lycopene supplementation activated AMPK through the NRF1/CaMKK $\beta$  axis, thereby activating Sirt1 and PGC-1 $\alpha$  and inducing skeletal muscle fiber transformation from fast-twitch to slow-twitch. The regulatory functions of lycopene on antioxidant ability and muscle fiber type conversion contributed to the improvement of pork quality. Understanding the effects of dietary lycopene supplementation on meat quality, antioxidant ability and muscle fiber type transformation will lay the foundation of the application of lycopene in production.

#### Author contributions

**Wanxue Wen:** Investigation, Data curation, Formal analysis, Writing - original draft. **Xiaoling Chen:** Conceptualization, Funding acquisition, Methodology, Project administration, Supervision.

**Zhiqing Huang:** Conceptualization, Funding acquisition, Methodology, Supervision, Writing - review & editing. **Daiwen Chen:** Methodology. **Bing Yu:** Methodology. **Jun He:** Methodology. **Yuheng Luo:** Methodology. **Hui Yan:** Methodology. **Hong Chen:** Methodology. **Ping Zheng:** Methodology. **Jie Yu:** Methodology.

#### Declaration of competing interest

We declare that we have no financial and personal relationships with other people or organizations that can inappropriately influence our work, and there is no professional or other personal interest of any nature or kind in any product, service and/or company that could be construed as influencing the content of this paper.

#### Acknowledgments

This work was supported by the National Key R&D Program of China (No. 2018YFD0500403) and the Sichuan Youth Science and Technology Innovation Research Team Project (No. 2020JDTD0026).



## References

- Chen X, Guo Y, Jia G, Liu G, Zhao H, Huang Z. Arginine promotes skeletal muscle fiber type transformation from fast-twitch to slow-twitch via Sirt1/AMPK pathway. *J Nutr Biochem* 2018;61:155–62.
- Costa-Rodrigues J, Pinho O, Monteiro PRR. Can lycopene be considered an effective protection against cardiovascular disease? *Food Chem* 2018;245:1148–53.
- Cho IC, Park HB, Ahn JS, Han SH, Lee JB, Lim HT, et al. A functional regulatory variant of MYH3 influences muscle fiber-type composition and intramuscular fat content in pigs. *PLoS Genet* 2019;15:e1008279.
- Chen S, Zou L, Li L, Wu T. The protective effect of glycyrrhetic acid on carbon tetrachloride-induced chronic liver fibrosis in mice via upregulation of Nrf2. *PLoS One* 2013;8:e53662.
- Fachinello MR, Fernandes NLM, Souto ER, Santos TC, Costa AER, Pozza PC. Lycopene affects the immune responses of finishing pigs. *Ital J Anim Sci* 2018;17:666–74.
- Grabowska M, Wawrzyniak D, Rolle K, Chomczyński P, Oziewicz S, Jurga S, et al. Let food be your medicine: nutraceutical properties of lycopene. *Food Funct* 2019;10:3090–102.
- Joo ST, Kim GD, Hwang YH, Ryu YC. Control of fresh meat quality through manipulation of muscle fiber characteristics. *Meat Sci* 2013;95:828–36.
- Kim GD, Yang HS, Jeong JY. Intramuscular variations of proteome and muscle fiber type distribution in semimembranosus and semitendinosus muscles associated with pork quality. *Food Chem* 2018;244:143–52.
- Koh JH, Hancock CR, Terada S, Higashida K, Holloszy JO, Han DH. PPAR $\beta$  is essential for maintaining normal levels of PGC-1 $\alpha$  and mitochondria and for the increase in muscle mitochondria induced by exercise. *Cell Metabol* 2017;25:1176–85. e5.
- Kulkarni SS, Cantó C. The molecular targets of resveratrol. *Biochim Biophys Acta* 2015;1852:1114–23.
- Lebret B, Batonon-Alavo DI, Perruchot MH, Mercier Y, Gondret F. Improving pork quality traits by a short-term dietary hydroxy methionine supplementation at levels above growth requirements in finisher pigs. *Meat Sci* 2018;145:230–7.
- Leseigneur-Meynier A, Gandemer G. Lipid composition of pork muscle in relation to the metabolic type of the fibres. *Meat Sci* 1991;29:229–41.
- Li J, Kinoshita T, Pandey S, Ng CK, Gygi SP, Shimazaki K, et al. Modulation of an RNA-binding protein by abscisic-acid-activated protein kinase. *Nature* 2002;418:793–7.
- Li XK, Wang JZ, Wang CQ, Zhang CH, Li X, Tang CH, et al. Effect of dietary phosphorus levels on meat quality and lipid metabolism in broiler chickens. *Food Chem* 2016;205:289–96.
- Li YH, Li FN, Duan YH, Guo QP, Wen CY, Wang WL, et al. Low-protein diet improves meat quality of growing and finishing pigs through changing lipid metabolism, fiber characteristics, and free amino acid profile of the muscle. *J Anim Sci* 2018;96:3221–32.
- Liang M, Wang Z, Li H, Cai L, Pan J, He H, et al. L-Arginine induces antioxidant response to prevent oxidative stress via stimulation of glutathione synthesis and activation of Nrf2 pathway. *Food Chem Toxicol* 2018;115:315–28.
- Lin J, Xia J, Zhao HS, Hou R, Talukder M, Yu L, et al. Lycopene triggers Nrf2-AMPK cross talk to alleviate atrazine-induced nephrotoxicity in mice. *J Agric Food Chem* 2018;66:12385–94.
- Ryu YC, Rhee MS, Kim BC. Estimation of correlation coefficients between histological parameters and carcass traits of pig *Longissimus dorsi* muscle. *Asian-Australas J Anim Sci* 2004;17:428–33.
- Skiepkio N, Chwastowska-Siwiecka I, Kondratowicz J, Mikulski D. Fatty acid profile, total cholesterol, vitamin content, and TBARS value of Turkey breast muscle cured with the addition of lycopene. *Poultry Sci* 2016;95:1182–90.
- Wang B, Liu Y, Feng L, Jiang WD, Kuang SY, Jiang J, et al. Effects of dietary arginine supplementation on growth performance, flesh quality, muscle antioxidant capacity and antioxidant-related signalling molecule expression in young grass carp (*Ctenopharyngodon idella*). *Food Chem* 2015;167:91–9.
- Wen C, Chen Y, Leng Z, Ding L, Wang T, Zhou Y. Dietary betaine improves meat quality and oxidative status of broilers under heat stress. *J Sci Food Agric* 2019;99:620–3.
- Wen W, Chen X, Huang Z, Chen D, Chen H, Luo Y, et al. Resveratrol regulates muscle fiber type conversion via miR-22-3p and AMPK/SIRT1/PGC-1 $\alpha$  pathway. *J Nutr Biochem* 2020a;77:108297.
- Wen W, Chen X, Huang Z, Chen D, Zheng P, He J, et al. miR-22-3p regulates muscle fiber-type conversion through inhibiting AMPK/SIRT1/PGC-1 $\alpha$  pathway. *Anim Biotechnol* 2020b;14:1–8.
- Xu M, Chen X, Huang Z, Chen D, Chen H, Luo Y, et al. Procyanidin B2 promotes skeletal slow-twitch myofiber gene expression through the AMPK signaling pathway in C2C12 myotubes. *J Agric Food Chem* 2020a;68:1306–14.
- Xu M, Chen X, Huang Z, Chen D, Yu B, Chen H, et al. Grape seed proanthocyanidin extract promotes skeletal muscle fiber type transformation via AMPK signaling pathway. *J Nutr Biochem* 2020b;84:108462.
- Xu X, Chen X, Chen D, Yu B, Yin J, Huang Z. Effects of dietary apple polyphenol supplementation on carcass traits, meat quality, muscle amino acid and fatty acid composition in finishing pigs. *Food Funct* 2019;10:7426–34.
- Yonar ME. The effect of lycopene on oxytetracycline-induced oxidative stress and immunosuppression in rainbow trout (*Oncorhynchus mykiss*, W.). *Fish Shellfish Immunol* 2012;32:994–1001.
- Yu M, Li Z, Rong T, Wang G, Liu Z, Chen W, et al. Different dietary starch sources alter the carcass traits, meat quality, and the profile of muscle amino acid and fatty acid in finishing pigs. *J Anim Sci Biotechnol* 2020;11:78.
- Zhang C, Luo J, Yu B, Zheng P, Huang Z, Mao X, et al. Dietary resveratrol supplementation improves meat quality of finishing pigs through changing muscle fiber characteristics and antioxidative status. *Meat Sci* 2015;102:15–21.
- Zhang Y, Yu B, Yu J, Zheng P, Huang Z, Luo Y, et al. Butyrate promotes slow-twitch myofiber formation and mitochondrial biogenesis in finishing pigs via inducing specific microRNAs and PGC-1 $\alpha$  expression. *J Anim Sci* 2019;97:3180–92.
- Zhang L, Yin M, Wang X. Meat texture, muscle histochemistry and protein composition of *Eriocheir sinensis* with different size traits. *Food Chem* 2021;338:127632.