



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Medical countermeasures against henipaviruses: a review and public health perspective



Raúl Gómez Román, Nadia Tornieporth, Neil George Cherian, Amy C Shurtleff, Maina L'Azou Jackson, Debra Yeskey, Adam Hacker, Eric Mungai, Tung Thanh Le

Henipaviruses, including Nipah virus, are regarded as pathogens of notable epidemic potential because of their high pathogenicity and the paucity of specific medical countermeasures to control infections in humans. We review the evidence of medical countermeasures against henipaviruses and project their cost in a post-COVID-19 era. Given the sporadic and unpredictable nature of henipavirus outbreaks, innovative strategies will be needed to circumvent the infeasibility of traditional phase 3 clinical trial regulatory pathways. Stronger partnerships with scientific institutions and regulatory authorities in low-income and middle-income countries can inform coordination of appropriate investments and development of strategies and normative guidelines for the deployment and equitable use of multiple medical countermeasures. Accessible measures should include global, regional, and endemic in-country stockpiles of reasonably priced small molecules, monoclonal antibodies, and vaccines as part of a combined collection of products that could help to control henipavirus outbreaks and prevent future pandemics.

Introduction

Nipah and other henipaviral diseases are listed among the WHO research and development blueprint priority diseases that pose a substantial public health risk because of their epidemic or pandemic potential and the absence of specific medical countermeasures to control and mitigate them.¹ Like other viruses of the henipavirus genus, Nipah virus (NiV) and Hendra virus (HeV) are zoonotic viruses that can spillover from *Pteropus* spp bats—their natural hosts and reservoir—to other mammals including humans.^{2,3} Transmission occurs via exposure to animal or human secretions or respiratory droplets.^{2,3} Human infections with NiV were originally described as a syndrome of fever and rapid neurological decline (ie, encephalitis) following contact with pigs. Since 2001, outbreaks (of Nipah virus Bangladesh strain [NiV-B]) report prominent respiratory symptoms (eg, atypical pneumonia and severe respiratory problems, including acute respiratory distress) and human-to-human transmission.⁴ Since 1994, henipavirus spillover events have caused human outbreaks in several countries, including Australia, Bangladesh, India, Malaysia, the Philippines, and Singapore.⁵ Although these outbreaks have thus far involved fewer than 1000 confirmed cases in total, the case–fatality rate for henipavirus infections can be as high as 100% depending on the context and constraints of the health-care systems where outbreaks occur. The geographical range where the various *Pteropus* bat species thrive is extensive^{6,7} (encompassing Indo-Pacific territories across the southeast Asia and western Pacific regions) and covers half of the global population.⁸ In addition to the tragic loss of human lives, henipavirus outbreaks can generate fear, stigma, loss of livestock (pigs and horses), and have a negative economic effect on affected communities.⁵ Although the reproduction number (R_0) for HeV has not been calculated, the R_0 for NiV ranges from 0.19 to 0.59 in nosocomial settings or in the community, including corpse-to-person transmission.^{9,10}

In an outbreak situation, use of widespread sensitive rapid diagnostic tests (RDTs)¹¹ combined with appropriate prevention and non-pharmaceutical intervention control measures, can rapidly lower the R_0 , buying precious time

Key messages

- The pipeline for henipavirus countermeasures in development with clinical or preclinical data in the public domain includes at least eight small molecules, four monoclonal antibodies, and more than 15 vaccine candidates.
- Several of these potential henipavirus medical countermeasures employ molecules, concepts, or technologies that are being used, where available, to treat or prevent COVID-19.
- Access to henipavirus countermeasures will depend on various factors including cost and whether or not their use is authorised by the national regulatory authorities in henipavirus-affected countries.
- Given the sporadic and unpredictable nature of current henipavirus outbreaks, traditional phase 3 clinical trial regulatory pathways might not be feasible for vaccines; thus, national regulatory authorities might have to rely on alternative regulatory pathways such as conditional market authorisation, authorisation under exceptional circumstances, or the animal rule.
- To prepare for future henipavirus outbreaks, funding agencies, sponsors, and manufacturers of henipavirus medical countermeasures must jointly understand the regulatory requirements to apply for emergency use or relevant authorisation of henipavirus medical countermeasures in the affected countries where these measures are ultimately needed.
- Lessons learned from the current COVID-19 pandemic provide a platform to connect with several stakeholders and to better prepare for future virus threats, including those posed by henipaviruses.

Lancet Infect Dis 2022; 22: e13–27

Published Online
November 1, 2021
[https://doi.org/10.1016/S1473-3099\(21\)00400-X](https://doi.org/10.1016/S1473-3099(21)00400-X)

Coalition for Epidemic Preparedness Innovations (CEPI), Oslo, Norway
(R Gómez Román PhD, N Tornieporth MD, N G Cherian MBBBS, A C Shurtleff PhD, M L'Azou Jackson MSc, D Yeskey PharmD, A Hacker PhD, E Mungai MSc, T T Le MSc); University of Applied Sciences & Arts, Hanover, Germany (N Tornieporth)

Correspondence to:
Mr Tung Thanh Le, Coalition for Epidemic Preparedness Innovations (CEPI), Marcus Thranes Gate 2, 0473 Oslo, Norway
tung.le@cepi.net

to deploy medical countermeasures. Vaccines, antivirals, and other medical countermeasures must be developed and made available for use worldwide, especially to populations at the highest risk of henipavirus infection.

Internationally coordinated development of medical countermeasures against henipaviruses has mainly focused on NiV, but important progress has also been made in the development of such measures against HeV. Medical countermeasures against other henipaviruses have not been described. This Review will focus on publicly available evidence of potential medical countermeasures at any stage of development against both of these viruses. The aim is to provide a synthesis of available knowledge to aid epidemic preparedness. Potential costs and equitable access by all henipavirus-affected countries, including low-income and middle-income countries, are considered.

Scope and search strategy

We reviewed information available in the public domain on therapeutics, monoclonal antibodies, and vaccine candidates evaluated for prophylaxis or protection, primarily against NiV and HeV infection, with a small amount of information found for other henipaviruses such as Cedar henipavirus, Ghanaian bat henipavirus, and Mojiang henipavirus. A targeted literature search was done in the English language following PRISMA guidelines and using “Nipah” OR “Hendra” OR “henipaviruses” as keywords with additional MeSH terms and selection criteria described in detail in the appendix p 1. Our search strategy did not include diagnostics. Although diagnostics are classified as medical countermeasures, they are mentioned here only in the context of other medical countermeasures. A detailed landscape analysis of the henipavirus diagnostic pipeline has been published elsewhere.¹²

Evidence for medical countermeasures

Small molecules and antivirals

No approved therapy exists for henipavirus encephalitis, and a target product profile for potential NiV therapeutics is currently being developed by WHO.¹³ Where available, intensive supportive therapy for severe respiratory and neurological complications is the current standard of care for individuals infected with henipavirus.^{14–17} Further evidence is required to generate substantive guidance protocols for pre- and post-exposure prophylaxis against henipavirus encephalitis. Available evidence regarding potential therapeutics against henipavirus encephalitis is displayed in table 1. Most of these compounds are broadly active antiviral therapeutics targeting a range of RNA and DNA viruses, rather than specifically developed as henipavirus medical countermeasures.

Although some of the compounds listed in table 1 are approved for other indications, none are yet WHO-prequalified for global procurement. Although a few of these have been shown to inhibit viral replication of

henipaviruses *in vitro* and *in vivo*, there is still a paucity of evidence in animal models, partly due to the restricted access to biosafety level 4 (BSL4) laboratories required to do experiments. Thus, several research groups are doing *in-silico* analyses⁴¹ or biosafety level 2 pseudotyped virus assays⁴² to design and screen compounds for the potential treatment of henipavirus encephalitis before animal and clinical testing. Although animal challenge studies have mainly been against NiV, the evidence available for remdesivir, favipiravir, ribavirin, and griffithsin suggests that these agents could also offer broad protection against other henipaviruses (table 1). Only remdesivir and VIKI-dPEG-Toco are supported by non-human primate challenge data in African green monkeys, whereas all other experiments have been done in smaller animals such as mice, ferrets, or Syrian golden hamsters. Remdesivir has been shown to be protective against NiV challenge in African green monkeys when daily, intravenous treatment was initiated within 24 h of exposure to NiV and continued for 12 days.¹⁹ Favipiravir has been shown to be protective in Syrian golden hamsters when given immediately after NiV infection.²¹ Although these compounds have shown protection against challenge or post-exposure in animal models, demonstration of a therapeutic effect following symptom onset is still missing and merits further research.

For further development of these drugs as potential henipavirus medical countermeasures, additional pre-clinical and clinical studies are needed to determine the protective efficacy of the treatment regimen when initiated more than 24 h post-virus challenge, or even after symptom onset. Such studies will help define scenarios for prophylactic medical countermeasure use, as well as use of medical countermeasures as a post-infection intervention. The potentially narrow treatment window (within 24 h of exposure), timing of deployment, administration, and appropriate volumes of these products need to be considered for stockpiling. Notably, NiV can present with relapsing encephalitis,⁴³ and such cases might also benefit from antiviral prophylaxis.

Ribavirin (approved for hepatitis C virus) remains the only therapeutic with supportive evidence from a human outbreak;¹⁸ mortality was reduced by 36% when ribavirin was used empirically in Malaysia during the outbreak of Nipah there in September, 1998–June, 1999.¹⁸ However, this evidence is limited due to the open-label nature of the study, the use of data from historical control patients who declined treatment or received standard treatment before ribavirin was available, and the simultaneous provision of intensive supportive care to patients. Ribavirin has since been used on a compassionate basis during the 2018 Indian outbreak of Nipah encephalitis^{16,44} in which its use led to a possible reduction in viral load.⁴⁵ Soluble Ephrin-B2 receptors (EFBN2) have been shown to inhibit binding of NiV and HeV to target cells *in vitro*,⁴⁶ but to our knowledge no EFBN2 analogues have been developed and specifically tested as potential therapeutics against henipaviruses.

See Online for appendix

	Classification	Available evidence so far (route of therapeutic administration)	Target	Mode of action	Finding	Development or licensure status and indication
Ribavirin ¹⁸	Antiviral	Open-label clinical trial in humans (oral and intravenous)	NiV	Viral replication inhibition	Reduced mortality of treated patients compared with non-treated patients (by 36%)* in Malaysian outbreak (1998)	Approved for hepatitis C virus and respiratory syncytial virus in several countries
Remdesivir ¹⁹	Antiviral	African green monkeys (intravenous)	NiV	Nucleotide analogue prodrug, inhibits viral replication	Protection against viral challenge	Approved for COVID-19 by the US FDA. Emergency use authorisation for COVID-19 in Australia, Bangladesh, India, Singapore, Japan, Taiwan, and the European Union ²⁰
Favipiravir ²¹	Antiviral	Syrian golden hamsters (oral and subcutaneous)	NiV, HeV	Viral RNA-dependent RNA polymerase inhibitor	Protection against viral challenge	Approved for influenza A in Japan, and for COVID-19 in several countries. Emergency use authorisation for COVID-19 in India ²²
Chloroquine ^{23,24}	Antimalarial	Syrian golden hamsters (intraperitoneal) and ferrets (route not stated)	NiV, HeV	Inhibition of F protein maturation	Inhibition of viral replication in vitro. No conclusive evidence in vivo challenge—in combination with ribavirin	Approved for malaria in several countries
Heparin ²⁵	Anticoagulant	Syrian golden hamsters (subcutaneous)	NiV, HeV	Inhibits cell-mediated viral trans-infection by binding to heparan sulfate	Inhibition of viral trans-infection in vitro. Heparin treatment restricts NiV infection in Syrian golden hamsters	Approved for coagulopathies in several countries. Experimental: preclinical (Syrian golden hamster study)
Rintatolimid ²⁶	Interferon inducer	Syrian golden hamsters (subcutaneous and intraperitoneal)	NiV	Induces IFN- α and IFN- β production, inhibition of viral replication	Inhibition of viral replication and protection against viral challenge	Approved for chronic fatigue syndrome in Argentina. Experimental (phase 1 and 2 trials) for HIV and chronic fatigue syndrome ²⁷
Griffithsin ²⁸	Antiviral lectin	Syrian golden hamsters (intranasal)	NiV	Inhibits viral entry, replication and syncytia formation	Reduced viral replication in vitro and provides partial protection against viral challenge	Experimental: phase 1 trials for HIV ²⁹
VIKI-dPEG4-Toco, VIKI-PEG4-choj ^{30,31}	Viral fusion inhibitory peptide	African green monkeys and Syrian golden hamsters (intratracheal and intranasal)	NiV	Inhibition of F protein fusion and cell entry	Partial protection against viral challenge	Experimental: preclinical
Gliotoxin ³²	Mycotoxin	In vitro	NiV, HeV	Viral RNA-dependent RNA polymerase inhibitor	Inhibition of infection and replication; Cytotoxic but possible topical applications	Experimental: exploratory
Bortezomib ³³	Anticancer	In vitro	NiV	Proteasome inhibitor	Inhibition of viral budding	Approved for multiple myeloma and mantle cell lymphoma by the US FDA
Balapiravir, R1479 ³⁴	Antiviral	In vitro	NiV, HeV	Viral RNA-dependent RNA polymerase inhibitor	Inhibition of viral replication	Experimental, discontinued in phase 1 trials for dengue virus and hepatitis C virus ³⁵
Lumicitabine, ALS-8112 ³⁶	Antiviral	In vitro	NiV	Nucleotide analogue prodrug, inhibits viral replication	Inhibition of recombinant and wild-type NiV replication, and reduced NiV infectious virus titre	Experimental: phase 1 and phase 2 trials for respiratory syncytial virus ³⁷
CH25H ³⁸	Antiviral	In vitro	NiV	Intravenous-stimulated genes: catalyses oxidation of cholesterol to 25-hydroxycholesterol	Inhibition of viral membrane fusion and NiV replication	Experimental: exploratory
KIN1408 ³⁹	Antiviral	In vitro	NiV	Immunomodulation of interferon regulatory factor 3	Inhibition of viral replication and decreased viral load in vitro	Experimental: exploratory
AB00991123, AB00992391, and AB00993210 ⁴⁰	Antivirals	In vitro	NiV	Sulfonamide compounds, unknown	Inhibition of NiV-induced cytopathic effect and virus replication	Experimental: exploratory

HeV=Hendra virus. NiV=Nipah virus. *Patients also received intensive supportive treatment, and comparators were historical control patients.

Table 1: Available evidence on potential small molecules against henipavirus encephalitis

A major bottleneck in moving promising compounds to clinical development is access to BSL4 facilities to conduct proof-of-concept challenge studies. Other key aspects for repurposed therapeutics include the need for further evidence on oral, intravenous, or intranasal

routes of administration, the associated costs of intravenous administration, and the need for hospital infrastructure and supportive treatment. Regulatory requirements for repurposing approved medications, such as additional clinical trials for novel indications and

reformulation of the product, must also be taken into account.

Monoclonal antibodies

Monoclonal antibodies (mAbs) could be a viable medical countermeasure option for deployment under compassionate use for prophylaxis, and this might remain the case until a vaccine is available for emergency use or licensed. Similar to small molecules and antivirals, mAbs can potentially be used for both pre-exposure and post-exposure prophylaxis and fill in a critical gap before vaccine availability. mAbs may even be deployed in an emergency situation even after a vaccine becomes available, given that many of the vaccines follow a two-dose regimen and would require several days to elicit protective immunity. Four main mAb projects constitute the current landscape of mAbs under development against henipaviruses (table 2). These mAbs have also been used as reagents to characterise antigens and inform vaccine design strategies.

mAb m102.4 is the furthest in development and is the only NiV mAb with published phase 1 data. It has been shown to protect African green monkeys against challenge with both Malaysia and Bangladesh NiV strains.^{47–49} m102.4 has been administered under compassionate use as post-exposure prophylaxis for HeV to 14 individuals in Australia and the USA.⁵⁰ The phase 1 trial in Australia included 40 healthy adults, who received escalating single doses from 1 mg/kg to 20 mg/kg, or two 20 mg/kg infusions, 72 h apart, or placebo.⁵⁰ No serious adverse events or adverse events leading to discontinuation were observed in this study. The most commonly reported adverse events included mild to moderate infections and infestations and headaches, occurring at a similar frequency in active treatment and placebo recipients. Another mAb, h5B3.1 (ie, the humanised version of mAb 5B3 originally derived from mouse hybridoma 5B3), neutralises HeV and NiV Bangladesh and Malaysia strains and inhibits viral fusion *in vitro*.⁵¹ In NiV and HeV post-exposure prophylaxis models in ferrets, mAb h5B3.1 protected animals against disseminated disease when administered

intraperitoneally on days 3 and 5 post-infection.⁵² In 2020, mAb HENV-26 and mAb HENV-32, two naturally occurring human mAbs, were isolated from a donor immunised with a veterinary HeV protein subunit vaccine (Equivac; Zoetis, Rhodes, NSW, Australia) administered under compassionate use. These mAbs were shown to neutralise both HeV and NiV Bangladesh and Malaysia strains.⁵³ In a NiV post-exposure prophylaxis model in ferrets, these mAbs also protected animals against disseminated disease when administered intraperitoneally on days 3 and 5 post-infection. Although their different epitope specificity would suggest that these mAbs could have synergistic properties if administered as a mAb cocktail, the mAbs were only tested separately in virus neutralisation assays and protective efficacy experiments. Furthermore, despite binding to NiV and HeV G proteins, the mAbs did not bind to G proteins from other henipaviruses such as Ghanaian bat henipavirus or Cedar henipavirus.⁵³

The current engagement from academic, public, and biotechnology product developers is important for progression of the existing mAb pipeline. However, to further accelerate the development process and ensure that final products reach the target population, a clear strategy is required on the following four areas: scenario planning for product development and regulatory pathways in relation to disease epidemiology; further participation from industry, especially product developers with experience in the development of medical countermeasures against other pathogens, to lead and support henipavirus medical countermeasure development efforts; technical transfer and engagement of local manufacturers in endemic countries; and further progression of mAb development to clinical trials and potential licensure.

Vaccines

A draft Nipah vaccine target product profile published by WHO⁵⁵ stipulates that a NiV vaccine be used for reactive immunisation (ie, active immunisation of at-risk individuals in the area of an ongoing outbreak) and “in conjunction with other control measures to curtail or end an outbreak”.⁵⁵ In an outbreak scenario, key preferred

	Evidence available	Current development stage	Target	Developer
mAb 102.4	Human (phase 1 trial), ferrets, and African green monkeys ^{47–50}	Phase 1	HeV or NiV G glycoprotein	Henry M Jackson Foundation for the Advancement of Military Medicine (Bethesda, MD, USA)
mAb 5B3, mAb h5B3.1	Mice and ferrets ^{51,52}	Preclinical	HeV or NiV pre-fusion F glycoprotein	University of Washington (Seattle, WA, USA) and Uniformed Services University (Bethesda, MD, USA)
mAb HENV-26, mAb HENV-32	Ferrets ⁵³	Preclinical	HeV or NiV G glycoprotein (receptor-binding protein)	Vanderbilt Vaccine Center (Nashville, TN, USA)
Anti-G mAb, anti-F mAb	Hamsters ⁵⁴	Preclinical	NiV G and F glycoprotein	INSERM (Paris, France), Université Claude Bernard Lyon 1 (Lyon, France), and Institut Pasteur (Paris, France)

HeV=Hendra virus. mAb=monoclonal antibody. NiV=Nipah virus.

Table 2: Preclinical and clinical evidence for candidate monoclonal antibodies against henipaviruses

characteristics of a NiV vaccine include an ability to elicit protective immunity rapidly (preferably within 2 weeks) after a single dose, acceptable safety profile, high efficacy (>90%), thermostability (2–8°C), and an ability to confer immunity and protection against Malaysia and Bangladesh strains of NiV.

Table 3 shows the current landscape of the henipavirus vaccine pipeline. There are more than 40 candidates in development with data available in the public domain; however, eight of these are intended primarily for veterinary use.^{68,73,77} Nearly half (n=19) of henipavirus vaccine candidates are based on viral vector platforms, 17 are protein subunits or virus-like particles formulated in various adjuvants, and two are based on mRNA technology. The main targets of henipavirus vaccines in development are the surface G glycoprotein or the fusion F protein. All listed vaccines are highly immunogenic and elicit henipavirus binding antibodies, neutralising antibodies, or both, with at least 15 candidates conferring various levels of protection against challenge with

homologous or heterologous henipavirus strains in African green monkeys, Syrian golden hamsters, or ferrets (table 3).

The recombinant, soluble HeV G glycoprotein candidate is the only vaccine candidate that has progressed to phase 1 clinical trials.^{91,101} This candidate builds on existing knowledge gained from using the same soluble protein in several pre-clinical and veterinary studies,^{79,80,82,83,85} including pivotal data used to develop the Equivac HeV vaccine approved for veterinary use in horses.⁸⁵ In addition to good manufacturing practices and good clinical practices required to develop the soluble HeV G product for human use, a key difference between the vaccine intended for humans and its previous versions is the adjuvant formulation in Alhydrogel (Croda, USA) at a 1:10 ratio.⁹¹

It is too early to ascertain whether any of these candidate vaccines will achieve licensure and meet the preferred product characteristics described in the draft WHO target product profile for a NiV vaccine.⁵⁵ The low temperature requirement could create challenges for

	Vaccine regimen and administration route	Animal models used	Henipavirus challenge strain	Reference
Viral vectors				
Recombinant vaccinia viruses (modified vaccinia virus Ankara) expressing NiV-M or HeV F, G, or N	Single dose (intraperitoneal)	Mice	None	56
Vaccinia virus vector (NYVAC) expressing NiV-M G or F	2 doses, 1 month apart (subcutaneous)	Syrian golden hamsters	NiV-M	57
Canarypox vector (ALVAC) expressing NiV-M G or F	2 doses, 14 days apart (intramuscular)	Pigs*	NiV-M	58,59
Venezuelan equine encephalitis virus expressing HeV or NiV-M G or F	3 doses on weeks 0, 5, and 18 (footpad inoculation)	Mice	None	60
Replication-defective VSV-DG vector expressing NiV-M G or F	Single dose (intranasal or intramuscular)	Mice	None	61
Newcastle disease virus vector expressing NiV-M F or G	2 doses, 4 weeks apart (intramuscular)	Mice, pigs*	None	62
Single-cycle replication VSV-DG vector expressing NiV-B G and/or F	Single dose (intramuscular)	Ferrets	NiV-M	63†
Adeno-associated virus vector expressing NiV-M G	Single dose (intramuscular)	Syrian golden hamsters	NiV-M and HeV	64
Measles virus vaccine vectors (HL and Ed strains) expressing NiV-M G	2 doses, 21 or 28 days apart (intraperitoneal [Syrian golden hamsters] or subcutaneous [African green monkeys])	Syrian golden hamsters, African green monkeys	NiV-M	65
Live-attenuated rVSV-ZEBOV-GP vector expressing NiV-M G, F, or N	Single dose (intraperitoneal)	Syrian golden hamsters	NiV-M	66‡
Single-cycle replication VSV-DG vector expressing NiV-M G or F	Single dose (intramuscular)	Syrian golden hamsters	NiV-M	67
Live-attenuated and beta-propiolactone-inactivated VSV or rabies virus vaccine vectors expressing codon-optimised HeV G	Single dose (live) or three doses (beta-propiolactone), on weeks 0, 2, and 3 (intramuscular)	Mice*	None	68
Live-attenuated rVSV-ZEBOV-GP vector expressing NiV-M G	Single dose (intramuscular)	African green monkeys	NiV-M	69‡
Live-attenuated rVSV-ZEBOV-GP vector expressing NiV-M G	Single dose (intraperitoneal)	Syrian golden hamsters	NiV-M	70‡
Canarypox vector (ALVAC) expressing HeV G or F	2 doses, 21 days apart (intramuscular)	Syrian golden hamsters and ponies (horses)	None	71
Non-replicating VSV-DG vectors expressing NiV-M G and/or F	Single dose (intranasal or intracranial)	Mice	None	72
Live-attenuated and beta-propiolactone-inactivated rabies virus vaccine vector expressing NiV-B G	Single dose (live) or 2 doses (beta-propiolactone), 28 days apart (intramuscular)	Mice	None	73
Single-cycle replication VSV-DG vector expressing NiV-B G and/or F	Single dose (intramuscular)	African green monkeys	NiV-B	74†
Chimpanzee adenovirus vector expressing NiV-B G	Single or two doses, 28 days apart (intramuscular)	Syrian golden hamsters	NiV-M, NiV-B, and HeV	75
Modified vaccinia virus Ankara expressing NiV-M sG or G	Single or 2 doses, 21 days apart (intraperitoneal or intramuscular)	IFNAR -/- mice	None	76
Recombinant rabies viruses Evelyn-Rokitnicki-Abelseth strain, rERAG _{333E} expressing NiV-M G or F	2 doses, 8 weeks apart (oral)	Mice and pigs*	None	77
Bovine herpes virus 4 or canarypox vectors (ALVAC) expressing NiV-M G or F	2 doses, 3 weeks apart (intramuscular)	Pigs*	None	5,78

(Table 3 continues on next page)

	Vaccine regimen and administration route	Animal models used	Henipavirus challenge strain	Reference
(Continued from previous page)				
Protein subunits§				
sG _{NiV-M} or sG _{HeV} in CSIRO triple adjuvant (Montanide/QuilA/DEAE-dextran)	3 doses, 2 weeks apart (subcutaneous)	Cats	NiV-M	79
Recombinant soluble HeV G glycoprotein in CpG plus Alhydrogel adjuvant	2 doses, 21 days apart (intramuscular)	Cats	NiV-M	80
Soluble trimeric forms of HeV and NiV-M F proteins (sF _{GM1}) in Sigma Adjuvant System adjuvant	4 doses, each 30 days apart (intraperitoneal or subcutaneous)	Mice	None	81
Recombinant soluble HeV G glycoprotein in CpG and Alhydrogel adjuvant	2 doses, 21 days apart (intramuscular)	African green monkeys	NiV M	82
Recombinant soluble HeV G glycoprotein in Alhydrogel and CpG adjuvant	2 doses, 20 days apart (subcutaneous)	Ferrets	NiV B	83
Recombinant soluble HeV G glycoprotein in Alhydrogel with or without CpG adjuvant	2 doses, 21 days apart (intramuscular)	African green monkeys	HeV	84
Recombinant soluble HeV G glycoprotein (produced in 293 or Chinese hamster ovary cells) in a proprietary adjuvant (Zoetis, Inc)	2–5 doses, weeks 0 and 3, then at 6 months and then yearly (intramuscular)	Horses*	HeV	85,86
Recombinant soluble HeV G glycoprotein in a proprietary adjuvant (Zoetis, Inc)	2 doses, 21 days apart (intramuscular)	Pigs*	NiV-M and HeV	87
Recombinant soluble HeV G glycoprotein in alhydrogel + CpG adjuvant	2 doses, 20 days apart (subcutaneous)	Ferrets	HeV	88
Molecular clamp-stabilised F protein (mcsF)	2 doses, 3 weeks apart (intramuscular)	Pigs*	NiV-M	5
Multiple pre-fusion-stabilised F and oligomeric G proteins derived from NiV-M and formulated in aluminum hydroxide	2 doses, 3 weeks apart (intramuscular)	Mice	None	89
Monovalent, bivalent, and tetravalent Fc-linked G proteins from NiV-M, HeV, GhV, and MojV formulated in CpG and Alhydrogel	2 doses, 3 weeks apart (intramuscular)	Mice	None	90
Recombinant soluble HeV G glycoprotein, produced in HEK-293 cells, formulated in Alhydrogel	Single dose or 2 doses, 4 weeks apart (intramuscular)	African green monkeys	HeV (Brisbane) and NiV B	91
Virus-like particles				
Virus-like particles containing NiV-M M, G, and F¶	3 doses on days 0, 15, and 29 (subcutaneous)	Mice	None	92
Virus-like particles containing NiV-M M, F, and G, formulated in various adjuvants (alum, monophosphoryl lipid A, and CpG)¶	Single dose or 3 doses on days 0, 21, and 42 (intramuscular)	Syrian golden hamsters	NiV-M	93
Virus-like particles containing NiV-M M and F or G in Sigma Adjuvant System	3 doses on weeks 0, 3, and 5 (intramuscular)	Rabbits	None	94
Virus-like particles containing NiV-M F and G and HeV M	3 doses on weeks 0, 3, and 6 (intraperitoneal)	Mice	None	95
Cellular debris				
Pellets and supernatants from sF9 cells expressing recombinant NiV-M F and G proteins in a baculovirus system	2 doses, 3 weeks apart (intramuscular and intraperitoneal)	Mice	None	96
DNA				
Plasmids encoding codon-optimised NiV-M F and/or G	2 doses, 4 weeks apart (intramuscular)	Mice	None	97
Plasmids encoding NiV-M F and/or G	Single dose (intramuscular) followed by electroporation	Mice	NiV-M pseudovirus	98
mRNA				
HeV G codon-optimised mRNA in liquid nanoparticles	Single dose (intramuscular)	Syrian golden hamsters	NiV-M	99
mRNA-1215, mRNA encoding NiV-M F and G in liquid nanoparticles	To be determined	Undisclosed preclinical development	To be determined	100

GhV=Ghanaian bat henipavirus. GP=glycoprotein. HeV=Hendra virus. MojV=Mojiang henipavirus. NiV=Nipah virus. NiV-B=Nipah virus Bangladesh. NiV-M=Nipah virus Malaysia. rVSV=recombinant vesicular stomatitis virus. VSV=vesicular stomatitis virus. ZEBOV=Zaire Ebola virus. *These vaccines are intended primarily for veterinary use. †The single-cycle replication VSV-DG vector expressing NiV-B G and/or F is identical in these studies. ‡The live-attenuated rVSV-ZEBOV-GP vector expressing NiV-M G is identical in these studies. §The soluble HeV G protein is identical in all studies using different adjuvant formulations. ¶The virus-like particles are identical in these studies using different formulations.

Table 3: Chronological overview of henipavirus vaccines in development, by platform

some vaccine candidates; however, this problem is currently being addressed with COVID-19 vaccines and is not necessarily a barrier, as even a small amount of ultracold chain capacity can be established rapidly during vaccine deployment in response to an outbreak.¹⁰² The single-dose vaccine regimen and cross-protection characteristics as described in the WHO target product profile might be possible to achieve. For example, a single dose of the ChAdOx-NiV Bangladesh vaccine candidate cross-protects Syrian golden hamsters against challenge with HeV and the NiV-Malaysia strain.⁷⁵

The global effort to develop COVID-19 vaccines might pose a substantial risk to henipavirus vaccine development capacity, including delays in animal studies and reduced manufacturing capacity and funding. However, COVID-19 vaccine development has also accelerated many of the vaccine platform technologies that could be applied to henipavirus vaccine development and rapid response. Among these, the henipavirus mRNA vaccine candidates are likely to benefit from the experience in COVID vaccine development, scale-up manufacturing, regulatory approval, and deployment.

Projected costs of henipavirus medical countermeasures in a post-COVID-19 era

The current henipavirus medical countermeasure pre-clinical and clinical landscape includes nine small molecules, four mAbs, and 15 vaccine candidates. Included candidates are those with clinical data in humans or proof-of-concept data in ferrets, Syrian golden hamsters, or African green monkeys. This list of candidates excludes those with only exploratory in-vitro data, or pre-clinical data without a known henipavirus challenge study. The development pipeline is heavily shifted towards vaccines because these are potentially the most effective public health intervention. The main funders of currently active henipavirus vaccine development projects are listed in appendix p 3. In addition to these, the National Institute of Allergy and Infectious Diseases has supported and continues to fund the development of henipavirus mAbs and other medical countermeasures.^{103–105} Furthermore, as part of its new strategic priorities, the Coalition for Epidemic Preparedness Innovations (CEPI) aims to support the development of a fully licensed vaccine or prophylactic mAb against Nipah by 2027.¹⁰⁶

Despite this progress, the development of vaccines and treatments against infectious diseases is a lengthy, costly, and risky process. The development of medical countermeasures against COVID-19 has proved that at least the speed of product development can be partially overcome with broad mobilisation of public and philanthropic funding, and engagement of scientists, policymakers, epidemiologists, public and private vaccine developers, manufacturers, and regulators involved in emergency use processes. However, this advanced speed comes with additional costs due to a greater number of projects entering the pipeline, and at-risk costs needed for large-scale manufacturing of various leading products in parallel before conclusion of the clinical testing.¹⁰⁷ COVID-19 medical countermeasure investments will probably increase the speed of product development and the number of medical countermeasure platform technologies against other pathogens, including those against henipavirus. Table 4 estimates the costs of potential henipavirus medical countermeasures in a post-COVID-19 era. Vaccines and small molecule regimens will continue to be more affordable per person than mAbs. However, vaccine impact at large, including vaccine effectiveness and cost-effectiveness of potential medical countermeasures, will need to be determined in a post-authorisation and post-licensure era.

Small molecules

For small molecules, we project costs ranging from US\$14 to \$600 per person per regimen (table 4). Advantages are associated with repurposing commercially available products for new indications, including the cost benefits of limited additional investments needed to ramp up supply, or in the cases of remdesivir and favipiravir, the leverage scale from demand generated by the current

COVID-19 indication. Furthermore, commercialised drugs are likely to have a reliable supply chain of generic versions at substantially reduced costs, with the potential for a good investment case for pre-exposure or post-exposure prophylaxis for high-risk populations in affected areas. An example of a prophylaxis campaign during an outbreak would be the use of oseltamivir (Tamiflu; F Hoffmann-La Roche AG, Basel, Switzerland) during the 2009 influenza A (H1N1) pandemic.¹¹² For henipavirus outbreaks, however, further supportive evidence in animal models and clinical trials will be needed in support of a mass prophylaxis concept. The route of administration will be especially relevant in an outbreak scenario—for example, daily intravenous administration of remdesivir would be limited by cost, infrastructure, and the need for hospital stays.

mAbs

For mAbs, we project costs of more than \$1000 per person per regimen (table 4). Further funding will be required to advance henipavirus mAb programmes into early-stage and late-stage clinical development. Delivery of mAbs, especially via intravenous infusion, requires capacity and infrastructure, which could prove challenging in some outbreak settings. Subcutaneous administration of mAbs against COVID-19 is being tested and submitted for licensure.¹¹³ If successful, this concept will fundamentally improve access to mAb treatments. Casirivimab and imdevimab (Regeneron [Tarrytown, NY, USA]) and bamlanivimab (Eli Lilly [Indianapolis, IN, USA]) COVID-19 mAbs have rapidly completed phase 1 trials and received emergency use listings as treatment against mild and moderate COVID-19,¹¹⁴ setting the precedent for future emulsion in a henipavirus outbreak. However, further data are needed regarding the added benefit to children and adolescents, and the prophylactic indication for mAbs.^{114,115}

Vaccines

The cost of henipavirus vaccines in an epidemic or pandemic setting remains unclear; however, experience from COVID-19 vaccines using similar platforms suggests the price of vaccines could range between \$4 per dose for viral vector vaccines to \$37 per dose for mRNA vaccines.¹¹¹ These figures do not include delivery costs, which could add substantially to the immunisation costs depending on vaccine logistics, infrastructure, and regimens.¹¹⁶ The cost per dose of a henipavirus vaccine in the scenario of a small number of outbreaks and a modest stockpile could also be higher as a result of the lower market potential. Although this price estimate is a wide range, vaccines might still be reasonably priced by comparison with other medical countermeasures such as mAbs. Public funding for research and development and advance purchase agreements could keep the cost affordable.

	Reference price* (US\$)	Reference cost unit	Potential dose regimen	Estimated cost of regimen per person (US\$)
Small molecules or therapeutics†				
Remdesivir	53 per vial (Cipla Limited, Mumbai, India)	100 µg vial	Ebola regimen; intravenous: 200 mg (loading dose) + 10 mg for 10 days	\$640
Favipiravir	<0.50 per tablet (Mylan Laboratories Limited, Hyderabad, India)	200 mg tablet	Influenza regimen: oral: 1800 mg twice daily (loading dose) + 800 mg twice daily for 14 days	\$31
Ribavirin	<1 per tablet	200 mg tablet (oral) or 10 mg vial (intravenous)	Chong et al ¹⁸ Nipah regimen; oral: 2 g (loading dose) + 1 g for 10 days Intravenous: 30 mg/kg (loading dose) + 16 mg/kg every 6 h for 4 days + 8 mg/kg every 8 h for 3 days	\$20–60 (oral); \$14–26 (intravenous)
Monoclonal antibodies‡				
Monoclonal antibodies produced in mammalian cell lines	1250–2100 per vial	700 mg–2.4 g per dose	COVID-19-like regimens; single intravenous infusion	\$1250–2100 (intravenous) ^{108–110}
Vaccines§				
Viral vectors, protein subunits in adjuvant, and mRNA in liquid nanoparticles	4–37 per dose	Variable concentration per dose	Single or two-dose regimen	\$8–74 ¹¹¹

*Prices based on publicly available information of generic product costs, or products developed using similar platforms. †Costs exclude expenses related to hospital care infrastructure and administration, supportive therapy, and disposables, etc. ‡Reference data for monoclonal antibodies are based on data available in the public domain for bamlanivimab (Eli Lilly, Indianapolis, IN, USA) and casirivimab plus imdevimab (REGN-COV2 antibody cocktail [Regeneron, Tarrytown, NY, USA]).^{96,97–99} §Costs exclude expenses related to vaccine delivery, transport, refrigeration, and administration.

Table 4: Projected costs of henipavirus medical countermeasures

Further investments in diagnostics and virological surveillance

Laboratory-based diagnostic testing, contact tracing, and proactive quarantine and treatment of suspected cases, often collectively referred to as “test, track/trace, and treat” strategies, are indicated at the onset of an outbreak and can quickly isolate and treat patients to halt further spread of henipaviruses.^{117,118} Currently, however, no near-patient or point-of-care tests are currently used for Nipah or Hendra viruses, which is likely to be a major limitation for the development of other medical countermeasures. Incentivisation is therefore needed to commercialise diagnostic assays and to adapt them for early detection of henipavirus encephalitis in suspected epidemic or pandemic settings.

Although numerous in-house serological methods (ELISA) and nucleic acid amplification techniques (PCR) for NiV detection exist globally,¹² harmonisation of validation methods, standards, and reagents to facilitate development of commercial PCR kits is needed. A reverse transcription-loop-mediated isothermal amplification (RT-LAMP) assay for the NiV *N* gene has recently been developed and is significantly more sensitive than RT-PCR, demonstrating potential for a quicker and simple rapid diagnostic test for use in outbreak settings.¹¹⁹ Key barriers to the development of henipavirus diagnostic tests include: the scarcity of widely available sera from henipavirus survivors, which might be necessary to accelerate the evaluation, validation, and licensure of serologic diagnostics for human use; and the BSL4 laboratory requirements for diagnostics dependent on virus isolation and wild-type virus neutralisation assays.

Notably, all experimental knowledge on Nipah virus acquired over the past two decades is derived from only two virus strains (199902916 Malaysia and 200401066 Bangladesh).¹²⁰ Investments in surveillance, virus isolation, and sequencing can further our understanding of the impact of Nipah virus strain variation on medical countermeasure efficacy.

Regulatory considerations

Modelling to evaluate the feasibility of conducting a phase 3 NiV vaccine trial in Bangladesh suggests that under the present low incidence scenario, traditional randomised, controlled efficacy trials are not feasible.¹²¹ Alternative licensure pathways in lieu of a traditional phase 3 vaccine efficacy trial, as described in the following paragraphs, should be considered.

Regulatory pathways

Various pathways exist for the licensure of new medical countermeasures or medical countermeasures for life-threatening diseases (table 5). These options include accelerated approval (by the Food and Drugs Administration [FDA]¹²² or the animal efficacy rule¹²⁶ in the USA), and conditional approval or exceptional circumstances in the European Union (EU).¹²³ These regulatory pathways might be suitable for the licensure of medical countermeasures against viruses that have unpredictable outbreak patterns and are not conducive to phase 3 human clinical efficacy trials.¹³⁵ However, NiV outbreaks are very unlikely to occur where the US FDA or the European Medicines Agency (EMA) have jurisdiction. Therefore, further development of the regulatory procedures in potentially affected countries is paramount.

	Data package could include	Examples of approved vaccines for other indications
Current scenario: $R_0 < 1$		
Accelerated approval (US FDA), ¹²² conditional marketing authorisation or exceptional circumstances approval (EMA), ¹²³ or similar mechanisms	Phase 1 data, phase 2 data*, phase 3 trial feasibility assessment (results from mathematical modelling), assay validation data, surrogate endpoint or correlates of protection data, passive transfer or adoptive transfer studies, bridging data for licensure, and post-approval confirmatory studies to demonstrate clinical benefit	Approvals based on surrogate endpoints; ¹²⁴ conditional marketing authorisation for Ervebo (recombinant vesicular stomatitis virus-Zaire Ebola virus) ¹²⁵
Animal rule (US FDA) ¹²⁶ and exceptional circumstances (EMA) ¹²³	Natural history study data and challenge data, additional requirements requested by national regulatory authorities for licensure, and post-approval confirmatory studies to demonstrate clinical benefit (if possible to conduct)	Approvals based on the animal rule; ¹²⁷ exceptional circumstances: Zabdeno (Ad26.ZEBOV [adenovirus type 26 vector-based vaccine, expressing a Zaire Ebola virus glycoprotein]) ¹²⁸ and Mvabea (MVA-BN-Filo [modified vaccinia Ankara vector-based vaccine, encoding glycoproteins from the Zaire Ebola virus]) ¹²⁹
Other, depending on specific national regulatory authority legislation. See Directorate General of Drug Administration, (Bangladesh) ¹³⁰ and the Central Drugs Standard Control Organization (India) ¹³¹ as examples	To be defined by each national regulatory authority	Vary by different national regulatory authority
Alert phase: $R_0 \sim 1$		
Rapid response: potential shift to PHEIC and pandemic scenarios
Higher incidence scenarios: $R_0 > 1$		
Accelerated approval (US FDA), ¹²² conditional marketing authorisation (EMA), ¹²³ national regulatory authority emergency use authorisation, WHO EUL, ^{132,133} and other	Phase 1 data, phase 2 data, and assay validation data†	Ebola vaccines: Ervebo ^{125,134}
EMA=European Medicines Agency. EUL=emergency use listing. PHEIC=public health emergency of international concern. US FDA=US Food and Drug Administration. *Phase 2 clinical trial material can become so-called "outbreak-ready" for an investigational stockpile. †Investigational stockpile deployed.		
Table 5: Potential regulatory pathways to pursue henipavirus vaccine authorisation		

In the absence of specific regulatory mechanisms in affected countries, this article will reference the US FDA and the EMA regulatory procedures. The US accelerated approval pathway allows for surrogate or clinical intermediate endpoints and is used for therapies for serious conditions "as soon as it can be concluded that [their] benefits justify their risks".¹³⁶ For henipaviruses, an example of a surrogate endpoint could be a protective titre of virus-neutralising antibodies. Accelerated approval would only be granted with a post-marketing commitment to demonstrate efficacy in a well-controlled clinical trial at the time of an outbreak. Given the epidemiology of NiV, timely fulfilment of this commitment would be problematic.

The animal efficacy rule (US FDA) entails using animal efficacy data instead of human clinical data.¹²⁶ To support licensure through this route, the efficacy of a medical countermeasure can be demonstrated using a disease endpoint in a relevant animal model that enables the selection of an effective dose and regimen in humans. The animal efficacy rule approval pathway is only available in a situation where licensure through the traditional or accelerated approval pathway is not feasible. Generating the data required for approval via this pathway is challenging. So far, there has been only one precedent for a vaccine approval using the animal rule

(BioThrax [Emergent BioSolutions, Rockville, MD, USA] for post-exposure prophylaxis following suspected or confirmed anthrax exposure). The process and data requirements were substantial, taking at least 5 years from conception of the regulatory pathway to approval.^{137,138} To gain approval via the animal rule, data must be generated to demonstrate a reasonably good understanding of the pathophysiology of the disease in a chosen animal model, and preferably in more than one animal species.¹³⁹ Well-designed natural history studies must be done, with solid evidence that the models can recapitulate key aspects of human disease. There have been relatively few human cases of henipavirus encephalitis to enable description of the full human clinical and pathological basis of the disease; therefore, animal disease comparisons might not reveal all features of human disease. Finally, whether or not the available strains of henipavirus encephalitis (as well as the challenge dose and method of administration) used in the animal challenge studies are epidemiologically relevant remains unclear. Hence, using the the animal rule to obtain approval for a vaccine against NiV may be challenging.

In the European Union (EU), the conditional marketing authorisation¹²³ has some similarities to the US accelerated approval procedure. The following

criteria must be met: the benefit–risk balance must be positive; the product must fulfil an unmet medical need; the benefit of making the product immediately available must be greater than the risk of additional data still being required; and comprehensive data post-authorisation must be provided in a timely manner. As is the case for the accelerated approval pathway, fulfilment of the latter is problematic given the epidemiology of NiV.

EU legislation also allows marketing authorisation to be granted in the absence of comprehensive data under exceptional circumstances.¹²³ Unlike conditional marketing authorisation, where marketing approval is granted in the likelihood that the sponsor will provide such data post-approval within an agreed timeframe, the EMA can grant authorisation under exceptional circumstances when comprehensive data cannot be obtained even after authorisation. For example, approval under exceptional circumstances was granted in July, 2020, for Zabdeno¹²⁸ and Mvabea¹²⁹ (both Janssen-Cilag International NV, Beerse, Belgium) against Ebola virus in individuals 1 year of age or older. The fact that Zabdeno and Mvabea received approval under exceptional circumstances in the EU but is yet to be approved in the USA implies that the EU exceptional circumstances regulatory route may be less challenging than the animal rule in the USA.

Early discussions with the US FDA and EMA are critical to establish specific data requirements and map the most appropriate route to approval for the developer. Importantly, the aforementioned US and EU regulatory routes are defined within specific legislation and require robust technical capabilities, especially with regard to the review and acceptance of animal efficacy data. Further work will be needed to evaluate the extent to which existing legislation in henipavirus-affected countries provides an appropriate level of flexibility to allow the use of animal efficacy data as the basis for regulatory approval. If such mechanisms are not available, the development of similar legislation should be considered by those countries where future henipavirus outbreaks could occur.

Pre-licensure mechanisms: emergency use of vaccines

National regulatory authorities might consider the authorisation of vaccines in their jurisdiction by allowing the use of an investigational product in emergency situations ($R_0 > 1$), such as in the event of a henipavirus pandemic, or if a public health emergency of international concern (PHEIC) is declared by WHO. Preparedness efforts can speed up the time from the declaration of a PHEIC to the licensure of vaccines for emergency use. The WHO emergency use listing procedure^{132,133,140} has enabled rapid deployment of COVID-19 vaccines to a broad range of countries.

All countries with potential for a henipavirus outbreak must have emergency legislation to enable the rapid

deployment of vaccines in the event an emergency use listing is granted by WHO. Since there might be a delay between the emergence of a henipavirus outbreak and declaration of a PHEIC, national regulatory authorities should also have legislation to enable the deployment of vaccines under their own emergency measures.

Data packages for vaccine candidates with some clinical data could be formally submitted to regulators in those countries where future outbreaks might occur and an authorisation could be requested based on existing clinical and non-clinical data, supplemented by data from the vaccine platform technology. These data could undergo a regulatory assessment based on anticipating the local benefit–risk in the country of deployment in the event of an outbreak. This would allow research to continue and additional data to be generated to support licensure. Such a licensed vaccine could be rapidly deployed in an attempt to control the emerging outbreak. Effectiveness and pharmacovigilance data could be collected in the real-world setting to confirm vaccine efficacy and safety. Although there is no current regulatory mechanism to support this, we advocate for an open debate on the feasibility of this approach. Case studies from the deployment of COVID-19 vaccines licensed in China, India, and Russia before the availability of phase 3 efficacy data, as well as learnings from undertaking clinical vaccine trials during the 2014–16 west African Ebola outbreak, could contribute to the framing of innovative regulatory frameworks in response to future pandemics.

A minimal dataset will be reviewed by regulatory authorities to enable rapid access and deployment of a vaccine during an emergency, irrespective of the regulatory mechanism used. As such, the ultimate objective of early deployment should be to continue to generate and collect the appropriate level of data to confirm effectiveness and achieve full licensure of the vaccine.

In the event that vaccines against NiV, HeV, or both do achieve licensure, it will also be important to advocate for implementation of accelerated regulatory approaches to enable the use of so-called core dossiers based on the pre-pandemic strains^{141–144} and manufacturing technology platform experience to rapidly enable a strain change in the event that the henipavirus outbreak contains mutations that make the licensed vaccines less efficacious due to insufficient cross-reactivity.

Ensuring equitable access

Crucially, affected populations must be able to access medical countermeasures irrespective of cost. Four countries (Bangladesh, India, Malaysia, and the Philippines) historically affected by henipaviruses are low-income or middle-income countries¹⁴⁵ whereas two (Australia and Singapore) are high-income countries. Australia is the first country with a HeV vaccine and a documented, in-country stockpile of a henipavirus mAb, m102·4, which has been administered for compassionate

use.⁵⁰ With scientific expertise in HeV, BSL4 capacity, and a renowned medical countermeasure consortia,¹⁴⁶ Australia could be a leading stakeholder in ensuring access through large-scale manufacturing and technology transfer of mAbs in the western Pacific region. In countries like Bangladesh, where Nipah outbreaks have predominantly affected poor populations in rural areas and health-care expenditures are generally out-of-pocket expenses,¹⁴⁷ central and local governments can prepare for future outbreaks by adapting strategies to reduce out-of-pocket expenses¹⁴⁸ or by devising mechanisms to share the costs of medical countermeasure procurement, deployment, and delivery, avoiding any undue delays to populations in need, regardless of their ability to pay for treatment.

Strategies to reduce the cost of medical countermeasure production while increasing the number of market players would help to increase access to medical countermeasures for low-income and middle-income countries; however, this approach is challenging in the face of a somewhat small henipavirus medical countermeasure portfolio. There might be little financial incentive perceived by multinational pharmaceutical companies, which have the most competitive resources and manufacturing capacity to invest in medical countermeasures.^{149,150} Several initiatives have been set up to address this gap—for example, CEPI has invested more than \$100 million in the pre-clinical and clinical development of four NiV vaccine candidates, including a protein subunit vaccine,⁹¹ a measles vector-based vaccine,⁶⁴ a vesicular stomatitis virus vector-based vaccine,^{66,69,70} and a chimpanzee adenovirus vector vaccine.⁷⁵ NiV investigational vaccine stockpiles and other medical countermeasures, together with operations for intervention in the field, should be in place before a major epidemic or pandemic.¹⁵¹ However, more financial investment and continued participation from governments and the pharmaceutical private sector will be required to achieve this goal. The public, philanthropic, and private funding of active henipavirus vaccine development projects (appendix p 3) could help to decouple the price of the product from the cost of research and development, making it potentially more affordable. Efforts to increase supply and local manufacturing are also needed to potentially reduce prices and spur innovation. Funders must therefore include low-income and middle-income countries and strengthen their capacities in the crucial stages of vaccine development such as manufacturing, for example by establishing mechanisms to transfer technology and know-how to emerging developers in low-income and middle-income countries.¹⁵²

Compared with vaccines, small molecules and antivirals are easier to generically produce. The COVID-19 pandemic has demonstrated that in some cases, prices can be set fairly low by overcoming intellectual property issues.¹⁵³ Therefore, both upstream and downstream strategies that address such issues across all the medical countermeasures will be vital for accessibility.

Funding agencies, sponsors, and manufacturers of henipavirus medical countermeasures must share the responsibility of understanding regulatory requirements to apply for emergency use or relevant authorisation of henipavirus medical countermeasures in affected countries. Poor knowledge of country-specific regulatory pathways can potentially create authorisation delays and an access roadblock for affected countries. Communication between product developers and the appropriate national regulatory authorities is essential to obtain clarity and guidance on local clinical trial requirements, regulatory legislation, and import requirements. Regulatory pathways and national regulatory authority capabilities vary between henipavirus-affected countries; thus, early engagement with regulators, preferably as soon as a target product profile is conceived with defined target populations, is essential to identify country-specific considerations.¹³

Coordination and cooperation among all stakeholders is crucial to ensure equitable access to henipavirus medical countermeasures. Although the price of the various measures could be reduced, supply might still be low in the case of a large outbreak or pandemic, as is the case with COVID-19.¹⁵⁴ Improved mechanisms to facilitate mobilisation, coordination, and cooperation across all the stakeholders are required.

Conclusion

In this Review we have presented an overview of several medical countermeasures under development that have the potential to control henipavirus outbreaks. The pipeline of such measures is diverse (and mostly in the pre-clinical stage), with vaccines leading both in the number of candidates and the lowest anticipated cost per person per regimen. Investment in a combined portfolio of several medical countermeasures, including surveillance systems, should be part of a coordinated, multilateral strategy for epidemic and pandemic preparedness given the unpredictability of outbreaks and the high case–fatality rate. Given such risk, ethical considerations must feature prominently when planning clinical trials and establishing the trial design before the outbreak. Active exchange of data between developers of human and animal medical countermeasures should be encouraged. Regulatory agencies across different nations will require convincing data packages to approve the start of human clinical testing. Animal efficacy data collected in well-designed, high-quality studies could enable the start of these clinical studies and the collection of further data to support licensure via non-traditional regulatory pathways. For these efforts to be effective, multilateralism will be necessary. Multilateral strategies should be based on scenario planning and include an outbreak response plan that clearly delineates responsibilities for the coordinated, sequential deployment of medical countermeasures in the event of a henipavirus outbreak. The current COVID-19 pandemic

opens a timely and unique opportunity to implement lessons learned from SARS-CoV-2 and apply them to preparedness efforts against henipaviruses and other pathogens of pandemic potential.

Contributors

RGR, NT, NGC, ACS, MLJ, DY, EM, and TTL contributed to conceptualisation of the study. RGR did the literature search for the vaccines section, NGC did the search for the small molecules section, and TTL did the search for the monoclonal antibodies section. RGR verified the data in the small molecules section, NGC verified the data in the monoclonal antibodies section, and TTL verified the data in the vaccines section. RGR, NGC, ACS, MLJ, DY, AH, EM, and TTL contributed to drafting of the manuscript. NT, MLJ, DY, and AH critically reviewed the scientific content. All authors reviewed and approved the final version.

Declaration of interests

NT is an independent consultant to the Coalition for Epidemic Preparedness Innovations (CEPI). All other authors are employees of CEPI. CEPI is supporting the research and development of a diverse portfolio of vaccine candidates (including vaccines against Nipah virus) based on a range of vaccine approaches.

Acknowledgments

We thank Amol Chaudhari, Gabrielle Breugelmanns, and Georges Thiry for reviewing the manuscript, and Julia Granerod for providing scientific writing support.

References

- WHO. Prioritizing diseases for research and development in emergency contexts. 2021. <https://www.who.int/activities/prioritizing-diseases-for-research-and-development-in-emergency-contexts> (accessed April 28, 2021).
- Lee B, Rota PA. Henipavirus: ecology, molecular virology, and pathogenesis. New York, NY: Springer, 2012.
- Halpin K, Rota P. A review of Hendra virus and Nipah virus infections in man and other animals. In: Sing A, ed. Zoonoses – infections affecting humans and animals: focus on public health aspects. Dordrecht: Springer, 2015: 997–1012.
- Hauser N, Gushiken AC, Narayanan S, Kotttilil S, Chua JV. Evolution of Nipah virus infection: past, present, and future considerations. *Trop Med Infect Dis* 2021; **6**: 24.
- Coalition for Epidemic Preparedness Innovations. Nipah@20. Proceedings from Nipah Virus International Conference, Singapore, 9 to 11 December 2019. 2019. https://cepi.net/wp-content/uploads/2020/06/2019-CEPI-Duke-WHO-NIAID-Nipah-Conference_FINAL.pdf (accessed April 28, 2021).
- WHO. Nipah virus: risk of importation to the countries of EMR. *Wkly Epidemiol Monitor* 2018; **11**: 23.
- Enchery F, Horvat B. Understanding the interaction between henipaviruses and their natural host, fruit bats: paving the way toward control of highly lethal infection in humans. *Int Rev Immunol* 2017; **36**: 108–21.
- WHO. Population data by WHO region. Global health observatory repository data. 2016. <https://apps.who.int/gho/data/view.main.POP2020?lang=en> (accessed April 28, 2021).
- Nikolay B, Salje H, Hossain MJ, et al. Transmission of Nipah virus—14 years of investigations in Bangladesh. *N Engl J Med* 2019; **380**: 1804–14.
- Sazzad HMS, Hossain MJ, Gurley ES, et al. Nipah virus infection outbreak with nosocomial and corpse-to-human transmission, Bangladesh. *Emerg Infect Dis* 2013; **19**: 210–17.
- Koczula KM, Gallotta A. Lateral flow assays. *Essays Biochem* 2016; **60**: 111–20.
- Mazzola LT, Kelly-Cirino C. Diagnostics for Nipah virus: a zoonotic pathogen endemic to Southeast Asia. *BMJ Glob Health* 2019; **4** (suppl 2): e001118.
- WHO. Nipah research & development (R&D) roadmap. 2018. [https://www.who.int/publications/m/item/nipah-research-and-development-\(r-d\)-roadmap](https://www.who.int/publications/m/item/nipah-research-and-development-(r-d)-roadmap) (accessed April 28, 2021).
- BMJ. BMJ best practice. Henipaviruses. 2021. <https://bestpractice.bmj.com/topics/en-gb/1607/management-approach> (accessed April 28, 2021).
- Institute of Epidemiology Disease Control And Research. National Guideline for management, prevention and control of Nipah virus infection. 2020. <https://iedcr.gov.bd/publication/guidelines/10adb7d5-b36b-4e87-a37c-7fc981f4f888> (accessed June 2, 2021).
- Central team Ministry of Health and Family Welfare, Government of India. Clinical management protocol for Nipah virus disease. 2018. <http://cghealth.nic.in/cghealth17/Information/content/NipahVirus/Treatment.pdf> (accessed June 2, 2021).
- Indian Medical Association–Tamil Nadu State Branch. Surveillance, prevention and control of Nipah virus infection: a practical handbook. 2014. <https://imatin.com/wp-content/uploads/2018/05/surveillance-prevention-and-control-of-nipah-virus-infection-a-practical-handbook.pdf> (accessed June 2, 2021).
- Chong HT, Kamarulzman A, Tan CT, et al. Treatment of acute Nipah encephalitis with ribavirin. *Ann Neurol* 2001; **49**: 810–13.
- Lo MK, Feldmann F, Gary JM, et al. Remdesivir (GS-5734) protects African green monkeys from Nipah virus challenge. *Sci Transl Med* 2019; **11**: eaau9242.
- Reuters. FACTBOX—Countries where remdesivir is approved or supported for treating COVID-19. 2021. <https://www.reuters.com/article/healthcoronavirus-gilead-remdesivir-idUSL1N2HD1UUX> (accessed April 28, 2021).
- Dawes BE, Kalveram B, Ikegami T, et al. Favipiravir (T-705) protects against Nipah virus infection in the hamster model. *Sci Rep* 2018; **8**: 7604.
- Agrawal U, Raju R, Udawadia ZF. Favipiravir: a new and emerging antiviral option in COVID-19. *Med J Armed Forces India* 2020; **76**: 370–36.
- Freiberg AN, Worthy MN, Lee B, Holbrook MR. Combined chloroquine and ribavirin treatment does not prevent death in a hamster model of Nipah and Hendra virus infection. *J Gen Virol* 2010; **91** (Pt 3): 765–72.
- Pallister J, Middleton D, Cramer G, et al. Chloroquine administration does not prevent Nipah virus infection and disease in ferrets. *J Virol* 2009; **83**: 11979–82.
- Mathieu C, Dhondt KP, Chalons M, et al. Heparan sulfate-dependent enhancement of henipavirus infection. *mBio* 2015; **6**: e02427.
- Georges-Courbot MC, Contamin H, Faure C, et al. Poly(I)-poly(C12U) but not ribavirin prevents death in a hamster model of Nipah virus infection. *Antimicrob Agents Chemother* 2006; **50**: 1768–72.
- Hemispherx Biopharma. Executive informational overview. Nov 27, 2016. <https://cdn2.hubspot.net/hubfs/150154/docs/Hemispherx-Executive-Informational-Overview-11-27-16.pdf> (accessed April 28, 2021).
- Lo MK, Spengler JR, Krumpke LRH, et al. Griffithsin inhibits Nipah virus entry and fusion and can protect Syrian golden hamsters from lethal Nipah virus challenge. *J Infect Dis* 2020; **221** (suppl 4): S480–92.
- ClinicalTrials.gov. Studies found for griffithsin. 2021. <https://www.clinicaltrials.gov/ct2/results?recrs=&cond=&term=griffithsin&cntry=&state=&city=&dist> (accessed April 28, 2021).
- Mathieu C, Porotto M, Figueira TN, Horvat B, Moscona A. Fusion inhibitory lipopeptides engineered for prophylaxis of Nipah virus in primates. *J Infect Dis* 2018; **218**: 218–27.
- Porotto M, Rock B, Yokoyama CC, et al. Inhibition of Nipah virus infection in vivo: targeting an early stage of paramyxovirus fusion activation during viral entry. *PLoS Pathog* 2010; **6**: e1001168.
- Aljofan M, Sganga ML, Lo MK, et al. Antiviral activity of gliotoxin, gentian violet and brilliant green against Nipah and Hendra virus in vitro. *Virol J* 2009; **6**: 187.
- Wang YE, Park A, Lake M, et al. Ubiquitin-regulated nuclear-cytoplasmic trafficking of the Nipah virus matrix protein is important for viral budding. *PLoS Pathog* 2010; **6**: e1001186.
- Hotard AL, He B, Nichol ST, Spiropoulou CF, Lo MK. 4'-azidocytidine (R1479) inhibits henipaviruses and other paramyxoviruses with high potency. *Antiviral Res* 2017; **144**: 147–52.
- ClinicalTrials.gov. A study of balapiravir in patients with dengue virus infection. 2021. <https://clinicaltrials.gov/ct2/show/NCT01096576> (accessed April 28, 2021).

- 36 Lo MK, Amblard F, Flint M, et al. Potent in vitro activity of beta-D-4'-chloromethyl-2'-deoxy-2'-fluorocytidine against Nipah virus. *Antiviral Res* 2020; **175**: 104712.
- 37 ClinicalTrials.gov. 7 studies found for lumicitabine. 2021. <https://www.clinicaltrials.gov/ct2/results?recrs=&cond=&term=lumicitabine&cntry=&state=&city=&dist> (accessed April 28, 2021).
- 38 Liu SY, Aliyari R, Chikere K, et al. Interferon-inducible cholesterol-25-hydroxylase broadly inhibits viral entry by production of 25-hydroxycholesterol. *Immunity* 2013; **38**: 92–105.
- 39 Pattabhi S, Wilkins CR, Dong R, et al. Targeting innate immunity for antiviral therapy through small molecule agonists of the RLR pathway. *J Virol* 2015; **90**: 2372–87.
- 40 Tigabu B, Rasmussen L, White EL, et al. A BSL-4 high-throughput screen identifies sulfonamide inhibitors of Nipah virus. *Assay Drug Dev Technol* 2014; **12**: 155–61.
- 41 Sen N, Kanitkar TR, Roy AA, et al. Predicting and designing therapeutics against the Nipah virus. *PLoS Negl Trop Dis* 2019; **13**: e0007419.
- 42 Porotto M, Orefice G, Yokoyama CC, et al. Simulating henipavirus multicycle replication in a screening assay leads to identification of a promising candidate for therapy. *J Virol* 2009; **83**: 5148–55.
- 43 Tan CT, Goh KJ, Wong KT, et al. Relapsed and late-onset Nipah encephalitis. *Ann Neurol* 2002; **51**: 703–08.
- 44 Banerjee S, Niyas VKM, Soneja M, et al. First experience of ribavirin postexposure prophylaxis for Nipah virus, tried during the 2018 outbreak in Kerala, India. *J Infect* 2019; **78**: 491–503.
- 45 Times of India. Ribavirin effective in treating Nipah, say Pune scientists. 2018. <https://timesofindia.indiatimes.com/india/ribavirin-effective-in-treating-nipah-say-pune-scientists/articleshow/64456461.cms> (accessed July 1, 2021).
- 46 Bonaparte MI, Dimitrov AS, Bossart KN, et al. Ephrin-B2 ligand is a functional receptor for Hendra virus and Nipah virus. *Proc Natl Acad Sci USA* 2005; **102**: 10652–57.
- 47 Bossart KN, Geisbert TW, Feldmann H, et al. A neutralizing human monoclonal antibody protects african green monkeys from hendra virus challenge. *Sci Transl Med* 2011; **3**: 105ra3.
- 48 Bossart KN, Zhu Z, Middleton D, et al. A neutralizing human monoclonal antibody protects against lethal disease in a new ferret model of acute nipah virus infection. *PLoS Pathog* 2009; **5**: e1000642.
- 49 Geisbert TW, Mire CE, Geisbert JB, et al. Therapeutic treatment of Nipah virus infection in nonhuman primates with a neutralizing human monoclonal antibody. *Sci Transl Med* 2014; **6**: 242ra82.
- 50 Playford EG, Munro T, Mahler SM, et al. Safety, tolerability, pharmacokinetics, and immunogenicity of a human monoclonal antibody targeting the G glycoprotein of henipaviruses in healthy adults: a first-in-human, randomised, controlled, phase 1 study. *Lancet Infect Dis* 2020; **20**: 445–54.
- 51 Dang HV, Chan YP, Park YJ, et al. An antibody against the F glycoprotein inhibits Nipah and Hendra virus infections. *Nat Struct Mol Biol* 2019; **26**: 980–87.
- 52 Mire CE, Chan YP, Borisevich V, et al. A cross-reactive humanized monoclonal antibody targeting fusion glycoprotein function protects ferrets against lethal Nipah virus and Hendra virus infection. *J Infect Dis* 2020; **221** (suppl 4): S471–79.
- 53 Dong J, Cross RW, Doyle MP, et al. Potent henipavirus neutralization by antibodies recognizing diverse sites on Hendra and Nipah virus receptor binding protein. *Cell* 2020; **183**: 1536–50.e17.
- 54 Guillaume V, Contamin H, Loth P, et al. Antibody prophylaxis and therapy against Nipah virus infection in hamsters. *J Virol* 2006; **80**: 1972–78.
- 55 WHO. WHO target product profile for Nipah virus vaccine. 2017. https://www.who.int/blueprint/priority-diseases/key-action/Nipah_virus_vaccineTPP.pdf?ua=1 (accessed April 28, 2021).
- 56 Tamin A, Harcourt BH, Ksiazek TG, Rollin PE, Bellini WJ, Rota PA. Functional properties of the fusion and attachment glycoproteins of Nipah virus. *Virology* 2002; **296**: 190–200.
- 57 Guillaume V, Contamin H, Loth P, et al. Nipah virus: vaccination and passive protection studies in a hamster model. *J Virol* 2004; **78**: 834–40.
- 58 Weingartl HM, Berhane Y, Caswell JL, et al. Recombinant Nipah virus vaccines protect pigs against challenge. *J Virol* 2006; **80**: 7929–38.
- 59 Tamin A, Harcourt BH, Lo MK, et al. Development of a neutralization assay for Nipah virus using pseudotype particles. *J Virol Methods* 2009; **160**: 1–6.
- 60 Defang GN, Khetawat D, Broder CC, Quinnan GV Jr. Induction of neutralizing antibodies to Hendra and Nipah glycoproteins using a Venezuelan equine encephalitis virus in vivo expression system. *Vaccine* 2010; **29**: 212–20.
- 61 Chattopadhyay A, Rose JK. Complementing defective viruses that express separate paramyxovirus glycoproteins provide a new vaccine vector approach. *J Virol* 2011; **85**: 2004–11.
- 62 Kong D, Wen Z, Su H, et al. Newcastle disease virus-vectored Nipah encephalitis vaccines induce B and T cell responses in mice and long-lasting neutralizing antibodies in pigs. *Virology* 2012; **432**: 327–35.
- 63 Mire CE, Versteeg KM, Cross RW, et al. Single injection recombinant vesicular stomatitis virus vaccines protect ferrets against lethal Nipah virus disease. *Viol J* 2013; **10**: e353.
- 64 Ploquin A, Szecsi J, Mathieu C, et al. Protection against henipavirus infection by use of recombinant adeno-associated virus–vector vaccines. *J Infect Dis* 2013; **207**: 469–78.
- 65 Yoneda M, Georges-Courbot MC, Ikeda F, et al. Recombinant measles virus vaccine expressing the Nipah virus glycoprotein protects against lethal Nipah virus challenge. *PLoS One* 2013; **8**: e58414.
- 66 DeBuysscher BL, Scott D, Marzi A, Prescott J, Feldmann H. Single-dose live-attenuated Nipah virus vaccines confer complete protection by eliciting antibodies directed against surface glycoproteins. *Vaccine* 2014; **32**: 2637–44.
- 67 Lo MK, Bird BH, Chattopadhyay A, et al. Single-dose replication-defective VSV-based Nipah virus vaccines provide protection from lethal challenge in Syrian hamsters. *Antiviral Res* 2014; **101**: 26–29.
- 68 Kurup D, Wirblich C, Feldmann H, Marzi A, Schnell MJ. Rhabdovirus-based vaccine platforms against henipaviruses. *J Virol* 2015; **89**: 144–54.
- 69 Prescott J, DeBuysscher BL, Feldmann F, et al. Single-dose live-attenuated vesicular stomatitis virus-based vaccine protects African green monkeys from Nipah virus disease. *Vaccine* 2015; **33**: 2823–29.
- 70 DeBuysscher BL, Scott D, Thomas T, Feldmann H, Prescott J. Peri-exposure protection against Nipah virus disease using a single-dose recombinant vesicular stomatitis virus-based vaccine. *NPJ Vaccines* 2016; **1**: 16002.
- 71 Guillaume-Vasselin V, Lemaitre L, Dhondt KP, et al. Protection from Hendra virus infection with canarypox recombinant vaccine. *NPJ Vaccines* 2016; **1**: 16003.
- 72 van den Pol AN, Mao G, Chattopadhyay A, Rose JK, Davis JN. Chikungunya, influenza, Nipah, and Semliki Forest chimeric viruses with vesicular stomatitis virus: actions in the brain. *J Virol* 2017; **91**: e02154–60.
- 73 Keshwara R, Shiels T, Postnikova E, et al. Rabies-based vaccine induces potent immune responses against Nipah virus. *NPJ Vaccines* 2019; **4**: 15.
- 74 Mire CE, Geisbert JB, Agans KN, et al. Use of single-injection recombinant vesicular stomatitis virus vaccine to protect nonhuman primates against lethal Nipah virus disease. *Emerg Infect Dis* 2019; **25**: 1144–52.
- 75 van Doremalen N, Lambe T, Sebastian S, et al. A single-dose ChAdOx1-vectored vaccine provides complete protection against Nipah Bangladesh and Malaysia in Syrian golden hamsters. *PLoS Negl Trop Dis* 2019; **13**: e0007462.
- 76 Kalodimou G, Veit S, Jany S, et al. A soluble version of Nipah virus glycoprotein G delivered by vaccinia virus MVA activates specific CD8 and CD4 T Cells in mice. *Viruses* 2019; **12**: 26.
- 77 Shuai L, Ge J, Wen Z, Wang J, Wang X, Bu Z. Immune responses in mice and pigs after oral vaccination with rabies virus vectored Nipah disease vaccines. *Vet Microbiol* 2020; **241**: 108549.
- 78 Pedrera M, Macchi F, McLean RK, et al. Bovine herpesvirus-4 vectored delivery of Nipah virus glycoproteins enhances T cell immunogenicity in pigs. *Vaccines* 2020; **8**: 115.
- 79 Mungall BA, Middleton D, Crameri G, et al. Feline model of acute Nipah virus infection and protection with a soluble glycoprotein-based subunit vaccine. *J Virol* 2006; **80**: 12293–302.
- 80 McEachern JA, Bingham J, Crameri G, et al. A recombinant subunit vaccine formulation protects against lethal Nipah virus challenge in cats. *Vaccine* 2008; **26**: 3842–52.

- 81 Chan YP, Lu M, Dutta S, et al. Biochemical, conformational, and immunogenic analysis of soluble trimeric forms of henipavirus fusion glycoproteins. *J Virol* 2012; **86**: 11457–71.
- 82 Bossart KN, Rockx B, Feldmann F, et al. A Hendra virus G glycoprotein subunit vaccine protects African green monkeys from Nipah virus challenge. *Sci Transl Med* 2012; **4**: 146ra07.
- 83 Pallister JA, Klein R, Arkininstall R, et al. Vaccination of ferrets with a recombinant G glycoprotein subunit vaccine provides protection against Nipah virus disease for over 12 months. *Virol J* 2013; **10**: e237.
- 84 Mire CE, Geisbert JB, Agans KN, et al. A recombinant Hendra virus G glycoprotein subunit vaccine protects nonhuman primates against Hendra virus challenge. *J Virol* 2014; **88**: 4624–31.
- 85 Middleton D, Pallister J, Klein R, et al. Hendra virus vaccine, a one health approach to protecting horse, human, and environmental health. *Emerg Infect Dis* 2014; **20**: 372–79.
- 86 Tan R, Hodge A, Klein R, et al. Virus-neutralising antibody responses in horses following vaccination with Equivac(R) HeV: a field study. *Aust Vet J* 2018; **96**: 161–66.
- 87 Pickering BS, Hardham JM, Smith G, et al. Protection against henipaviruses in swine requires both, cell-mediated and humoral immune response. *Vaccine* 2016; **34**: 4777–86.
- 88 Pallister J, Middleton D, Wang LF, et al. A recombinant Hendra virus G glycoprotein-based subunit vaccine protects ferrets from lethal Hendra virus challenge. *Vaccine* 2011; **29**: 5623–30.
- 89 Loomis RJ, Stewart-Jones GBE, Tsybovsky Y, et al. Structure-based design of Nipah virus vaccines: a generalizable approach to paramyxovirus immunogen development. *Front Immunol* 2020; **11**: 842.
- 90 Li Y, Li R, Wang M, et al. Fc-based recombinant henipavirus vaccines elicit broad neutralizing antibody responses in mice. *Viruses* 2020; **12**: 480.
- 91 Geisbert TW, Bobb K, Borisevich V, et al. A single dose investigational subunit vaccine for human use against Nipah virus and Hendra virus. *NPJ Vaccines* 2021; **6**: 23.
- 92 Walpita P, Barr J, Sherman M, Basler CF, Wang L. Vaccine potential of Nipah virus-like particles. *PLoS One* 2011; **6**: e18437.
- 93 Walpita P, Cong Y, Jahrling PB, et al. A VLP-based vaccine provides complete protection against Nipah virus challenge following multiple-dose or single-dose vaccination schedules in a hamster model. *NPJ Vaccines* 2017; **2**: 21.
- 94 Schmidt R, Beltzig LC, Sawatsky B, et al. Generation of therapeutic antisera for emerging viral infections. *NPJ Vaccines* 2018; **3**: 42.
- 95 Stroh E, Fischer K, Schwaiger T, et al. Henipavirus-like particles induce a CD8 T cell response in C57BL/6 mice. *Vet Microbiol* 2019; **237**: 108405.
- 96 Wang XJ, Hu S, Ge JY, Wang QH, Qin LT, Bu ZG. [Study of fusion protein and attachment glycoprotein of Nipah virus expressed in recombinant baculovirus]. *Sheng Wu Gong Cheng Xue Bao* 2006; **22**: 418–24.
- 97 Wang X, Ge J, Hu S, et al. Efficacy of DNA immunization with F and G protein genes of Nipah virus. *Ann N Y Acad Sci* 2006; **1081**: 243–45.
- 98 Nie J, Liu L, Wang Q, et al. Nipah pseudovirus system enables evaluation of vaccines in vitro and in vivo using non-BSL-4 facilities. *Emerg Microbes Infect* 2019; **8**: 272–81.
- 99 Lo MK, Spengler JR, Welch SR, et al. Evaluation of a single-dose nucleoside-modified messenger RNA vaccine encoding Hendra virus-soluble glycoprotein against lethal Nipah virus challenge in Syrian hamsters. *J Infect Dis* 2020; **221** (suppl 4): S493–98.
- 100 Moderna. Press releases. Moderna provides business update and announces three new development programs in infectious disease vaccines. 2021. <https://investors.modernatx.com/news-releases/news-release-details/moderna-provides-business-update-and-announces-three-new> (accessed April 28, 2021).
- 101 Coalition for Epidemic Preparedness Innovations. CEPI-funded Nipah virus vaccine candidate first to reach phase 1 clinical trial. 2020. https://cepi.net/news_cepi/cepi-funded-nipah-virus-vaccine-candidate-first-to-reach-phase-1-clinical-trial/ (accessed April 28, 2021).
- 102 Jusu MO, Glauser G, Seward JF, et al. Rapid establishment of a cold chain capacity of –60°C or colder for the STRIVE Ebola vaccine trial during the Ebola outbreak in Sierra Leone. *J Infect Dis* 2018; **217** (suppl 1): S48–55.
- 103 NIH. NIH RePORTER. Project details. Project 2 – Vanderbilt University. 2021. <https://reporter.nih.gov/project-details/9897483> (accessed April 28, 2021).
- 104 NIH. NIH RePORTER. Project details. Advancement of vaccines and therapies for henipaviruses. 2021. <https://reporter.nih.gov/project-details/9897467> (accessed April 28, 2021).
- 105 NIH. NIH RePORTER. Project details. Rapid development of vaccines for emerging viruses. 2021. <https://reporter.nih.gov/project-details/10273019> (accessed April 28, 2021).
- 106 Coalition for Epidemic Preparedness Innovations (CEPI). The urgency of now. 2021. https://cepi.net/wp-content/uploads/2021/03/CEPI_3.5_billion_investment_case_10032021.pdf (accessed April 28, 2021).
- 107 Lurie N, Saville M, Hatchett R, Halton J. Developing Covid-19 vaccines at pandemic speed. *N Engl J Med* 2020; **382**: 1969–73.
- 108 Clinical Trials Arena. Eli Lilly's monoclonal antibody bamlanivimab value allows payer grace for Covid-19. 2020. <https://www.clinicaltrialsarena.com/comment/eli-lillys-monoclonal-antibody-bamlanivimab/#:~:text=Lilly%20will%20be%20supplying%20300%2C000.press%20release%20and%20CEO%20note> (accessed April 28, 2021).
- 109 Eli Lilly. News release. Lilly announces 650,000 additional doses of neutralizing antibody bamlanivimab (LY-CoV555) purchased by U.S. Government to treat COVID-19. 2020. <https://investor.lilly.com/news-releases/news-release-details/lilly-announces-650000-additional-doses-neutralizing-antibody> (accessed April 28, 2021).
- 110 Johnson CY, McGinley L. Experimental drug given to Trump to treat covid-19 wins FDA clearance. The Seattle Times. 2020. <https://www.seattletimes.com/nation-world/experimental-drug-given-to-trump-to-treat-covid-19-wins-fda-clearance/> (accessed April 28, 2021).
- 111 Statista. The cost per jab of Covid-19 vaccine candidates. 2020. <https://www.statista.com/chart/23658/reported-cost-per-dose-of-covid-19-vaccines/> (accessed April 28, 2021).
- 112 Lee VJ, Yap J, Cook AR, et al. Oseltamivir ring prophylaxis for containment of 2009 H1N1 influenza outbreaks. *N Engl J Med* 2010; **362**: 2166–74.
- 113 Regeneron. Press Release. Phase 3 prevention trial showed 81% reduced risk of symptomatic Sars-cov-2 infections with subcutaneous administration of Regen-cov™ (casirivimab with imdevimab). 2021. <https://investor.regeneron.com/news-releases/news-release-details/phase-3-prevention-trial-showed-81-reduced-risk-symptomatic-sars> (accessed April 28, 2021).
- 114 Cohen MS. Monoclonal antibodies to disrupt progression of early Covid-19 infection. *N Engl J Med* 2021; **384**: 289–91.
- 115 Wolf J, Abzug MJ, Wattier RL, et al. Initial guidance on use of monoclonal antibody therapy for treatment of COVID-19 in children and adolescents. *J Pediatric Infect Dis Soc* 2021; **10**: 629–34.
- 116 Vaughan K, Ozaltin A, Mallow M, et al. The costs of delivering vaccines in low- and middle-income countries: findings from a systematic review. *Vaccine X* 2019; **2**: 100034.
- 117 Arunkumar G, Chandni R, Mourya DT, et al. Outbreak investigation of Nipah virus disease in Kerala, India, 2018. *J Infect Dis* 2019; **219**: 1867–78.
- 118 Sahay RR, Yadav PD, Gupta N, et al. Experiential learnings from the Nipah virus outbreaks in Kerala towards containment of infectious public health emergencies in India. *Epidemiol Infect* 2020; **148**: e90.
- 119 Ma L, Chen Z, Guan W, Chen Q, Liu D. Rapid and specific detection of all known Nipah virus strains' sequences with reverse transcription-loop-mediated isothermal amplification. *Front Microbiol* 2019; **10**: 418.
- 120 Gurley ES, Spiropoulou CF, de Wit E. Twenty years of Nipah virus research: where do we go from here? *J Infect Dis* 2020; **221** (suppl 4): S359–62.
- 121 Nikolay B, Lipsitch M, Rahman M, et al. Assessing the feasibility of Nipah vaccine efficacy trials based on previous outbreaks in Bangladesh. *medRxiv* 2020; published online Dec 8. <https://doi.org/10.1101/2020.12.06.20244871> (preprint).
- 122 US Food and Drug Administration. Accelerated approval. 2018. <https://www.fda.gov/patients/fast-track-breakthrough-therapy-accelerated-approval-priority-review/accelerated-approval> (accessed April 28, 2021).
- 123 European Medicines Agency. Conditional marketing authorisation. 2021. <https://www.ema.europa.eu/en/human-regulatory/marketing-authorisation/conditional-marketing-authorisation> (accessed April 28, 2021).

- 124 US Food and Drug Administration. CDER drug and biologic accelerated approvals based on a surrogate endpoint. As of December 31, 2020. 2020. <https://www.fda.gov/media/88907/download> (accessed April 28, 2021).
- 125 European Medicines Agency. Ervebo. 2021. <https://www.ema.europa.eu/en/medicines/human/EPAR/ervebo> (accessed April 28, 2021).
- 126 US Food and Drug Administration. Animal rule approvals. 2021. <https://www.fda.gov/drugs/nda-and-bla-approvals/animal-rule-approvals> (accessed April 28, 2021).
- 127 US Food and Drug Administration. CDER drug and biologic animal rule approvals. 2021. <https://www.fda.gov/media/145827/download> (accessed April 28, 2021).
- 128 European Medicines Agency. Zabdeno. 2021. <https://www.ema.europa.eu/en/medicines/human/EPAR/zabdeno> (accessed April 28, 2021).
- 129 EMA. Mvabea. 2021. <https://www.ema.europa.eu/en/medicines/human/EPAR/mvabea> (accessed April 28, 2021).
- 130 Directorate General of Drug Administration (DGDA). 2013. <https://www.dgda.gov.bd/> (accessed April 28, 2021).
- 131 Central Drugs Standard Control Organization. 2021. <https://cdsco.gov.in/opencms/opencms/en/Home/> (accessed April 28, 2021).
- 132 WHO. Regulation and prequalification. Emergency use listing. 2021. <https://www.who.int/teams/regulation-prequalification/eul> (accessed April 28, 2021).
- 133 WHO. WHO publishes emergency use listing procedure and roadmap to make new medical products more readily available during health emergencies. 2021. <https://www.who.int/news/item/09-01-2020-emergency-use-listing-procedure-and-roadmap-health-emergencies> (accessed April 28, 2021).
- 134 Wolf J, Bruno S, Eichberg M, et al. Applying lessons from the Ebola vaccine experience for SARS-CoV-2 and other epidemic pathogens. *NPJ Vaccines* 2020; 5: 51.
- 135 Schrauf S, Tschisnarov R, Tauber E, Ramsauer K. Current efforts in the development of vaccines for the prevention of Zika and Chikungunya virus infections. *Front Immunol* 2020; 11: 592.
- 136 FDA Guidance for Industry. Expedited programs for serious conditions – drugs and biologics. <https://www.fda.gov/files/drugs/published/Expedited-Programs-for-Serious-Conditions-Drugs-and-Biologics.pdf> (accessed Oct 19, 2021).
- 137 Beasley DWC, Brasel TL, Comer JE. First vaccine approval under the FDA Animal Rule. *NPJ vaccines* 2016; 1: 16013.
- 138 US Food and Drug Administration. Summary basis for regulatory action template for BioThrax. 2015. <https://wayback.archive-it.org/7993/20170723023407/https://www.fda.gov/downloads/BiologicsBloodVaccines/Vaccines/ApprovedProducts/UCM474886.pdf> (accessed April 28, 2021).
- 139 Snoy PJ. Establishing efficacy of human products using animals: the US food and drug administration's "animal rule". *Vet Pathol* 2010; 47: 774–78.
- 140 WHO. Emergency use listing procedure. Version 13. December 2020. https://extranet.who.int/pqweb/sites/default/files/documents/EUL-FINAL-13_12_2020.pdf (accessed April 28, 2021).
- 141 Medicines & Healthcare products Regulatory Agency. Guidance on strain changes in authorised COVID-19 vaccines. 2021. <https://www.gov.uk/government/publications/access-consortium-guidance-on-strain-changes-in-authorised-covid-19-vaccines/guidance-on-strain-changes-in-authorised-covid-19-vaccines> (accessed April 28, 2021).
- 142 US Food and Drug Administration. Emergency use authorization for vaccines to prevent COVID-19. Guidance for industry. Document issued on Feb 22, 2021. <https://www.fda.gov/media/142749/download> (accessed April 28, 2021).
- 143 European Medicines Agency. Reflection paper on the regulatory requirements for vaccines intended to provide protection against variant strain(s) of SARS-CoV-2. 2021. <https://www.ema.europa.eu/en/documents/scientific-guideline/reflection-paper-regulatory-requirements-vaccines-intended-provide-protection-against-variant-en.pdf> (accessed April 28, 2021).
- 144 WHO. Addendum to considerations for evaluation of COVID-19 vaccines for prequalification or emergency use listing considerations for evaluation of modified COVID-19 vaccines. Points to consider for manufacturers of COVID-19 vaccines. Version 12. March, 2021. 2021. https://extranet.who.int/pqweb/sites/default/files/documents/Addendum_Evaluation_Modified_Covid-19%20Vaccine.pdf (accessed April 28, 2021).
- 145 Wellcome Trust. Low- and middle-income countries. 2021. <https://wellcome.org/grant-funding/guidance/low-and-middle-income-countries> (accessed April 28, 2021).
- 146 DMTC. DMTC Annual Report 2020. MCM segment. 2020. https://dmtc.com.au/wp-content/uploads/2020/12/MCM-Segment-DMTC-Annual-Report_2020.pdf (accessed April 28, 2021).
- 147 Mahumud RA, Sarker AR, Sultana M, Islam Z, Khan J, Morton A. Distribution and determinants of out-of-pocket healthcare expenditures in Bangladesh. *J Prev Med Public Health* 2017; 50: 91–99.
- 148 WHO. Out-of-pocket payments, user fees and catastrophic expenditure. https://www.who.int/health_financing/topics/financial-protection/out-of-pocket-payments/en/ (accessed June 2, 2021).
- 149 Crager SE. Improving global access to new vaccines: intellectual property, technology transfer, and regulatory pathways. *Am J Public Health* 2018; 108: S414–20.
- 150 Medecins Sans Frontieres Access Campaign. Giving developing countries the best shot: an overview of vaccine access and R&D. 2020. <https://msfaccess.org/giving-developing-countries-best-shot-overview-vaccine-access-and-rd> (accessed April 28, 2021).
- 151 Coalition for Epidemic Preparedness Innovations. Why we exist. 2021. <https://cepi.net/about/whyweexist/> (accessed April 28, 2021).
- 152 PATH. The Meningitis Vaccine Project: a groundbreaking partnership. 2015. <https://www.path.org/articles/about-meningitis-vaccine-project/> (accessed April 28, 2021).
- 153 Gilead. Voluntary licensing agreements for remdesivir. 2021. <https://www.gilead.com/purpose/advancing-global-health/covid-19/voluntary-licensing-agreements-for-remdesivir> (accessed April 28, 2021).
- 154 WHO. In the COVID-19 vaccine race, we either win together or lose together. Feb 10, 2021. <https://www.who.int/news/item/10-02-2021-in-the-covid-19-vaccine-race-we-either-win-together-or-lose-together> (accessed Oct 12, 2021).

© 2021 Elsevier Ltd. All rights reserved.