© The Author(s) 2021. Published by Oxford University Press on behalf of The Japanese Society for Immunology.

Advance Access publication 13 December 2021 This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (https://creativecommons.org/licenses/by-nc/4.0/), which permits noncommercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Corrigendum

Corrigendum to: Deficiency of TGR5 exacerbates immune-mediated cholestatic hepatic injury by stabilizing the β-catenin destruction complex

Jianhua Rao^{1,2,3}, Chao Yang^{1,2,3}, Shikun Yang^{1,2,3}, Hao Lu^{1,2,3}, Yuanchang Hu^{1,2,3}, Ling Lu^{1,2,3}, Feng Cheng^{1,2,3} and Xuehao Wang^{1,2,3}

¹Hepatobiliary Center, The First Affiliated Hospital of Nanjing Medical University, Nanjing 210029, China. ²Key Laboratory of Liver Transplantation, Chinese Academy of Medical Sciences, Nanjing 210029, China. ³NHC Key Laboratory of Living Donor Liver Transplantation, Nanjing 210029, China.

International Immunology, Volume 32, Issue 5, 1 May 2020, Pages 321–334, https://doi.org/10.1093/intimm/dxaa002

The authors recently realized that the two images in Fig. 1C that show staining of control liver tissue (labeled "Ctrl-1" and "Ctrl-2") were in fact from one subject; CHI-1 and CHI-2 were from two different CHI patients. The image "Ctrl-2" has been corrected in the revised Fig. 1, which now shows control tissue from two different subjects. This correction does not alter the overall conclusions of the study.

