



Erratum

Erratum to “The Dermatological Aspects of hEDS in Women”.
[International Journal of Women's Dermatology 7/3 (2021) 285-289]



Cynthia O. Edimo BS^{a,*}, Julia R. Wajsberg BA^{a,1}, Sammi Wong BS^{a,1},
Zachary P. Nahmias MD^b, Bernadette A. Riley DO^c

^a New York Institute of Technology College of Osteopathic Medicine, Old Westbury, NY, United States

^b FAAD, Samaritan Medical Center, Watertown, NY, United States

^c FACOFP, New York Institute of Technology, College of Osteopathic Medicine, Ehlers–Danlos Syndrome/Hypermobility Treatment Center, Old Westbury, NY, United States

The publisher regrets the [Figures 3 and 4](#) were interchanged in the published version. The correct figures and the corresponding caption is listed below:



Figure 3. Wound dehiscence: Note the evidence of former sutures at the lower border, now 3 weeks after the injury and treatment with antibiotics. Scars tend to stretch further in the 6 months after closure (Kang, et al., n.d. Orringer: Fitzpatrick's Dermatology, Ninth Edition. Copyright McGraw-Hill Education. All rights reserved).

DOI of original article: [10.1016/j.ijwd.2021.01.020](https://doi.org/10.1016/j.ijwd.2021.01.020)

* Corresponding author.

E-mail address: cedimo@nyit.edu (C.O. Edimo).

¹ Co-second authors.

<https://doi.org/10.1016/j.ijwd.2021.07.010>

2352-6475/© 2021 Women's Dermatologic Society. Published by Elsevier Inc. All rights reserved.



Figure 4. Clinical picture of piezogenic pedal papules. (Photo courtesy of Zachary P. Nahmias, MD.)

The publisher would like to apologise for any inconvenience caused.