

LETTER TO THE EDITOR

Open Access



Identification of tumor tissue-derived DNA methylation biomarkers for the detection and therapy response evaluation of metastatic castration resistant prostate cancer in liquid biopsies

Thomas Dillinger^{1,2}, Raheleh Sheibani-Tezerji^{1,2} , Walter Pulverer³, Ines Stelzer^{1,4}, Melanie R. Hassler^{2,5}, Janine Scheibelreiter², Carlos Uziel Pérez Malla¹, Madeleine Kuroll¹, Sandra Domazet¹, Elisa Redl², Sarah Ely^{1,6}, Stefanie Brezina⁷, Andreas Tiefenbacher^{1,2}, Katharina Rebhan⁵, Nicolai Hübner⁵, Bernhard Grubmüller⁵ , Markus Mitterhauser^{1,8} , Marcus Hacker⁸ , Andreas Weinhaeuser³ , Judit Simon^{1,4} , Markus Zeitlinger^{1,6} , Andrea Gsur⁷ , Gero Kramer⁵, Shahrokh F. Shariat^{5,9,10,11,12,13,14,15} , Lukas Kenner^{2,9,16,17} and Gerda Egger^{1,2,9*}

Main text

Prostate cancer (PCa) is among the most common cancers in men worldwide [1] and comprises a highly heterogeneous disease, which ranges from indolent localized cancer to aggressive high-risk stages, including metastatic hormone sensitive or hormone refractory PCa. Due to the limited specificity and sensitivity of current biomarkers such as PSA [2], there is an urgent need for better biomarkers that can reliably differentiate benign from malignant prostate conditions, localized from metastatic, as well as aggressive from indolent disease. Furthermore, the development of predictive biomarkers that allow for better patient stratification and of biomarkers for early monitoring of treatment response is of utmost importance [3].

Treatment options for PCa are mostly based on non-targeted therapies and include radical prostatectomy,

hormonal treatment using androgen deprivation therapy (ADT), androgen receptor (AR) signaling-targeting agents, chemotherapy or radiation, depending on disease state and risk classification [4, 5]. Despite good initial response to ADT, tumors eventually progress to metastatic castration resistant PCa (mCRPC). In this setting, recent treatments including poly(ADP-ribose) polymerase inhibitors (PARPi) for tumors harboring mutations in DNA repair genes, or radionuclide therapy using ¹⁷⁷Lu-PSMA for PSMA positive cancers have shown promising results [5].

Liquid biopsy assays, analyzing circulating free tumor DNA (ctDNA) or circulating tumor cells (CTCs) in plasma or other body fluids have proven as a useful source for biomarkers and have already entered the clinics for companion diagnostic use [6]. Aside from genetic alterations, epigenetic tumor-specific changes including DNA methylation are measurable in ctDNA and CTCs and their potential as diagnostic, prognostic and predictive epigenetic biomarkers has been demonstrated in a large number of studies although only few have made it into clinical practice, yet [7].

*Correspondence: gerda.egger@meduniwien.ac.at

⁹ Comprehensive Cancer Center, Medical University of Vienna, Vienna, Austria

Full list of author information is available at the end of the article



In this study, we investigated the suitability of DNA methylation-based biomarkers for non-invasive PCa diagnostics. Based on experiments and *in silico* analyses we identified two DNA methylation signatures, which could be used as minimal-invasive markers in liquid biopsies for the detection of methylated ctDNA. These signatures allowed for the classification of mCRPC with high specificity and sensitivity and were able to distinguish responders from non-responders following different treatment modalities. Importantly, several individual marker genes had prognostic potential for radiographic progression free survival independent of other clinical variables.

Results and discussion

Identification of tissue-specific DNA methylation markers

Differences in DNA methylation profiles between normal and PCa tissue were shown in several studies to reveal potential DNA methylation-based biomarkers for PCa detection [8]. In order to define suitable epigenetic markers for non-invasive diagnostics, we combined experimental and *in silico* data to derive 92 methylation markers that were significantly differentially methylated between tumor and normal adjacent tissues, which we subsequently tested on ctDNA of PCa patients (Fig. 1A).

First, we used the Infinium HumanMethylation450 BeadChip array and probed DNA isolated from six localized PCa tissues compared to six adjacent normal tissues (Table S1). From these analyses, 7 genes including *SERPIN1*, *ACSS3*, *SCGB3A1*, *NKX2-6*, *HOXA7*, *CRABP2* and *DHRS4L2* were found significantly hypermethylated in the tumor samples compared to normal adjacent tissues (Fig. S1A). Methylation levels of those genes continually rose from benign to PCa to metastatic tumors, as analyzed in two published datasets (Fig. S1B) [8, 9]. Along these lines, we could confirm the tumor-specific hypermethylation of all seven markers with high significance

in the publicly available cancer genome atlas PCa dataset TCGA-PRAD (Fig. S1C). In addition, we inferred 85 methylation markers from the TCGA-PRAD methylation data including PCa ($n=498$) and adjacent normal tissue ($n=54$). We selected suitable regions for marker evaluation containing at least three neighboring CpG sites, which were significantly differentially methylated ($p < 0.05$), showed low methylation in the controls (below 20%) and a methylation difference of at least 15% between controls and tumor. Out of the total 92 candidate regions, 80 were located in promoter regions (-1500bp upstream to +1000bp downstream of the transcription start site) of the respective genes, with 65 out of the 80 located in CpG islands. The remaining 12 candidate regions were located in intergenic regions or gene bodies (Table S2).


Identification of a methylation classifier for metastatic castration resistant prostate cancer in liquid biopsies of patients

To test the 92 identified methylation markers for their suitability to detect PCa-specific DNA methylation of ctDNA in plasma samples of patients, we used a high-throughput methylation sensitive restriction enzyme (MSRE) assay [10]. This method is based on the selective digestion of unmethylated DNA, and can be applied in a multiplex setting detecting DNA amounts as low as 10 copies and 0.1-1% of methylated DNA in an unmethylated background.

First, we analyzed a total of 174 plasma samples, including patients with benign conditions ($n=48$), localized PCa ($n=65$) and mCRPC ($n=61$) (Table S1). One sample from the benign cohort was removed from further analysis due to poor performance in the MSRE-qPCR. Relative methylation values were calculated based on percentage of methylation ratios (PMR), where samples were normalized for input DNA and

(See figure on next page.)

Fig. 1 Identification and liquid biopsy testing of DNA methylation-based markers. **A** Experimental Workflow. Flow chart indicates biomarker identification, validation and analysis in different patient cohorts and publicly available datasets. **B** Unsupervised clustering of PMR-values resulting from MSRE-qPCR analysis of ccfDNA isolated from plasma of indicated patient groups for 92 marker candidates ($n=47$ benign, $n=65$ localized PCa, $n=61$ mCRPC, $n=1$ PBMC). **C** DNA methylation levels of ccfDNA isolated from plasma in patients with localized PCa or mCRPC versus controls for three signature genes. Differences between the three groups were assessed using one-way ANOVA (*CHST11*, *PCDHGC4* $n=47$ benign, $n=65$ localized PCa, $n=61$ mCRPC, *CUGBP2* $n=46$ benign, $n=62$ localized PCa, $n=54$ mCRPC; **** $p < 0.0001$). **D** ROC-curve analysis based on the three gene signature as in (C) for classification of mCRPC samples compared to benign and localized PCa patients combined, calculated with different prediction algorithms, ((Bayesian) Compound Covariate Predictor (BCCP, CCP) and Diagonal Linear Discriminant Analysis (DLDA)) using recursive feature elimination ($n=112$ benign + localized PCa, $n=61$ mCRPC). **E** Unsupervised clustering of PMR-values resulting from MSRE-qPCR of ccfDNA isolated from plasma of mCRPC patients before and after treatment, or healthy PBMC controls for 92 marker candidates ($n=17$ responders, $n=12$ non-responders, $n=5$ PBMC). **F** ROC-curve analysis based on a 3-gene signature (*AKR1B1*, *KLF8*, *LDAH*) for classification of responders versus non-responders. Calculations were performed using BCCP, CCP and DLDA with recursive feature elimination using BRB array tools software. **G** Methylation levels of the three signature genes for individual responders and non-responders pre- and post-treatment ($n=17$ responder, $n=12$ non-responder ** $p < 0.01$, * $p < 0.05$, ns $p > 0.05$; two-way ANOVA). **H** Kaplan-Meier-Analysis for radiographic progression-free survival (rPFS) of three signature genes using post-treatment samples (*LDAH*, *KLF8* $n=17$ responder, $n=12$ non-responder; *AKR1B1* $n=15$ responder, $n=10$ non-responder; p values shown on each plot calculated with Mantel-Cox-test, censored subjects indicated on plots by strokes)



two control assays indicating 100% methylation. Unsupervised clustering revealed high levels of methylation in most mCRPC plasma samples, whereas normal and localized PCa plasma samples showed very low methylation values and were clustering together with the PBMC control (Fig. 1B).

Next, the 92 analyzed marker candidates were used for prediction model calculations by inputting PMR values for class prediction models based on different algorithms including Diagonal Linear Discriminant Analysis (DLDA), Nearest Centroid Predictor, k-Nearest-Neighbor Predictor, Support Vector Machines and (Bayesian) Compound Covariate Predictor (BCCP/CCP). We used a cutoff p value of $p \leq 0.01$ and 10-fold cross-validation. When comparing the benign with the mCRPC sample group, 83 out of the 92 markers were calculated to accurately classify between 93 and 96% of the samples to the correct group, depending on the used prediction model (Table S3). Areas under the curve (AUC) also depended on the used model and showed values of 0.968 (CCP), 0.972 (DLDA) and 0.966 (BCCP) (Fig. S2A). Similarly, comparing the localized PCa to mCRPC cohort, 83 out of the tested 92 marker candidates classified between 87 and 96% of the samples to the correct group with AUCs of 0.956 (CCP), 0.958 (DLDA) and 0.949 (BCCP) (Fig. S2B and Table S4). While no accurate prediction model could be inferred for benign versus localized PCa in general, comparisons between benign samples and PCa with a Gleason Score of 9 or above, resulted in AUCs ranging from 0.828 (CCP), 0.794 (DLDA) and 0.8 (BCCP) with 79 to 89% correct classification for 12 of the 92 analyzed marker genes (Fig. S2C, D and Table S5). Although we also observed a significant increase of cfDNA in mCRPC samples compared to benign and localized PCa samples, no correlation of DNA methylation and overall cfDNA concentration was detectable (Fig. S2E, F). Furthermore, DNA methylation outperformed overall cfDNA concentration measures in ROC analyses (Fig. S2G).

To identify a minimal set of markers, which allow for an accurate detection of mCRPC, we performed signature classifier calculations using recursive feature elimination. This resulted in a set of three marker candidates including *CHST11*, *CUGBP2* and *PCDHGC4* that accurately differentiated mCRPC from the combined group of localized PCa and benign patients (Fig. 1C). Depending on the prediction model used, this gene signature classified between 92% and 95% of the samples to the correct group with AUCs of 0.963 (CCP), 0.978 (DLDA) and 0.953 (BCCP) (Fig. 1D, Table S6). Individual genes showed AUCs of 0.982 (*CHST11*), 0.632 (*CUGBP2*) and 0.906 (*PCDHGC4*) (Fig. S2H).

Together, these data suggest that mCRPC can be identified based on methylation signatures with high accuracy, whereas organ-confined PCa with Gleason scores lower than 9 cannot be differentiated from benign samples, most likely due to limited amounts of ctDNA, which was also described in other studies analyzing DNA methylation in localized PCa patients using digital droplet PCR [11, 12]. Generally, ctDNA shows smaller fragment sizes as compared to cfDNA shed from normal cells [13, 14]. When performing fragment analysis of a subset of benign, localized PCa and mCRPC plasma samples ($n=20$ per group), we observed a significant shift of the mean cfDNA fragment size from 175 bp in benign and localized PCa (range 168 - 183bp) to 168 bp in mCRPC (range 145 - 179 bp) samples (Figure S2I).

Thus, our markers might be suitable to identify high risk patients, who have already developed micrometastases, which cannot be detected by regular computed tomography (CT). Interestingly, hypermethylation of *CHST11* has been found in breast cancer cell lines originating from luminal cells, whereas basal-like breast cancer cell lines showed rather hypomethylation [15]. This might also apply to PCa, which most frequently originates from luminal cells.

DNA methylation markers differentiate treatment responsive from non-responsive patients

Therapy options for mCRPC are diverse and evaluation of response to different treatments is thus essential for treatment decision making [4, 5]. To evaluate the potential of our identified marker genes in monitoring treatment response in mCRPC patients, we performed MSRE-qPCR analyses in liquid biopsies from mCRPC patients responsive ($n=17$) or non-responsive ($n=12$) to different therapies. Therapy response was based on increasing/decreasing PSA blood levels after therapy. Liquid biopsies were taken before treatment start and following therapy (Table S7).

We detected a trend towards higher methylation in non-responder patients before as well as after treatment, whereas responder patients showed reduced methylation after treatment compared to pre-treatment samples (Fig. 1E)

Using recursive feature elimination, signature classifier prediction models calculated a gene signature of three genes (*AKR1B1*, *LDAH*, *KLF8*) to distinguish responders from non-responders following treatment. Dependent on the prediction algorithm used, this signature correctly classified 83 - 90% of patients with AUCs of 0.902 (CCP), 0.892 (DLDA) and 0.863 (BCCP) (Fig. 1F, G and Table S8). Individual genes resulted in AUCs of 0.931 (*AKR1B1*), 0.765 (*KLF8*) and 0.980 (*LDAH*) (Fig. S2J). No significant

Table 1 Univariate and multivariate cox regression analysis based on methylation of individual genes in plasma post treatment

Univariate				Multivariate			
Model Tested	Variable	HR [95% CI]	P-value	Model Tested	Variable	HR	P-value
Overall Survival	CRABP2	4.013 [1.069, 15.06]	0.0395	Progression free survival (Radiographic recurrence)	ALOX5	3.7244	0.021
	TNFAIP8	0.036 [0.002, 0.582]	0.0191		Abiraterone	0.2594	0.02
Progression -free survival (Radiographic recurrence)	ALOX5	4.225[1.47, 12.15]	0.007	C2ORF88	4.85	0.006	
	APC	4.269 [1.553, 11.73]	0.0049	Abiraterone	0.214	0.008	
	C2ORF88	4.528 [1.579, 12.99]	0.005	CHST11	7.97	0.0004	
	CHST11	8.79 [2.975, 25.97]	8.43e-05	Abiraterone	0.2944	0.05	
	CPVL	3.867 [1.118, 13.37]	0.0326	CPVL	4.25	0.026	
	CRABP2	12.56 [2.441, 64.63]	0.002	Abiraterone	0.22	0.006	
	DNAH10	4.5326 [1.55, 13.26]	0.006	CRABP2	6	0.04	
	HOCX4,5,6	2.829 [1.034, 7.741]	0.0429	Abiraterone	0.3	0.07	
	KLF8	2.977 [1.043, 8.496]	0.0415	DNAH10	3.3	0.045	
	KLHL34	5.9 [1.751, 19.87]	0.004	Abiraterone	0.4	0.11	
	KLK10	5.402 [1.862, 15.67]	0.002	KLHL34	4.4	0.024	
	LDAH	3.422[1.14, 10.27]	0.028	Abiraterone	0.3	0.042	
	NKX2-6	3.673 [1.327, 10.17]	0.0123	KLK10	5.056	0.005	
	NOL3	6.89 [2.353, 20.17]	0.0004	Abiraterone	0.2529	0.02	
	PCDHGC4	4.225 [1.47, 12.15]	0.0075	NOL3	6.66	0.001	
	PROM1	3.447[1.191, 9.974]	0.023	Abiraterone	0.25	0.02	
	RGS20	4.69 [1.644, 13.36]	0.003	PCDHGC4	3.72	0.021	
	SCGB3A1	7.03 [2.361, 20.93]	0.0005	Abiraterone	0.26	0.02	
	SERPINB1	2.97 [10.978, 9.043]	0.0109	PROM1	3.2	0.04	
	SPARC	3.33 [1.148, 9.669]	0.0269	Abiraterone	0.24	0.012	
TMLHE	3.081 [1.1, 8.626]	0.0322	SCGB3A1	6.66	0.001		
TPM4	5.402 [1.862, 15.67]	0.002	Abiraterone	0.26	0.03		
ZNF154	4.347 [1.519, 12.44]	0.005	SERPINB1	3.33	0.04		
Abiraterone	0.2337 [0.082, 0.668]	0.0067	Abiraterone	0.2	0.006		
PSA post	1.002 [1.001, 1.004]	0.0003	TPM4	5.0559	0.005		
			Abiraterone	0.253	0.02		
			ZNF154	4.85	0.006		
			Abiraterone	0.198	0.007		
			APC	3.1	0.046		
			PSA post	1	0.003		
			CHST11	6.6	0.002		
			PSA post	1	0.01		
			CRABP2	6.8	0.036		
			PSA post	1	0.002		
			DNAH10	3.6	0.026		
			PSA post	1	0.002		
			KLK10	3.7	0.03		
			PSA post	1	0.005		
			NOL3	5.1	0.007		
			PSA post	1	0.006		
			RGS20	3.6	0.02		
			PSA post	1	0.002		
			SCGB3A1	5.3	0.003		
			PSA post	1	0.006		
			SPARC	3.4	0.04		
			PSA post	1	0.0005		
			TPM4	5.3	0.003		
			PSA post	1	0.001		
			ZNF154	3.5	0.027		
			PSA post	1	0.002		

difference of overall cfDNA concentrations was evident for the different patient groups before and after treatment (Fig. S2K).

When comparing post-treatment plasma samples of responders and non-responders to abiraterone acetate treatment only, we were able to correctly classify 84% of the patients based on 13 marker genes (Table S9).

In summary, we defined a methylation classifier for measuring treatment response to different therapeutic regimens based on ctDNA methylation of 3 distinct marker genes. Therapy-response monitoring is an important task for the clinical management of mCRPC. With a continuously growing number of available therapeutic options for mCRPC patients [16] it is of high importance to choose proper treatments and adjust in a timely manner upon failed response. Thus, we suggest that our identified methylation markers could allow for early detection of non-responders and indicate patient prognosis, which could allow for patient stratification to adjust treatment and prepare appropriate countermeasures.

Prognostic potential of DNA methylation markers for progression-free survival

To test whether our methylation markers could predict clinical outcomes such as radiographic progression-free survival (rPFS), we performed survival statistics using methylation levels of our candidate genes in plasma post treatment. Kaplan-Meier survival analysis revealed significant associations of methylation and decreased rPFS for several marker candidates including the three signature genes (*AKR1B1*, *KLF8*, *LDAH*) (Fig. 1H and S3).

Next, we performed univariate and covariate cox regression analyses to determine the prognostic value of our markers for overall survival (OS) and rPFS. For OS, methylation of *CRABP2* and *TNEAIP8* were significant prognostic factors (HR 4.013 and 0.036; *p* values 0.0395 and 0.0191) in plasma of patients following treatment in univariate analysis, but did not remain significant in co-variate models including treatment or PSA levels as variables (Table 1).

For rPFS, methylation of 23 individual marker genes was a negative prognostic factor for disease recurrence in univariate cox regression analysis, with hazard ratios (HR) ranging from 2.829 (*HOCX4,5,6*, *p* = 0.04) to 12.56 (*CRABP2*, *p* = 0.002).

Since most of the responder patients underwent abiraterone acetate treatment, covariate cox regression was performed. Despite abiraterone acetate treatment showing significant predictive value for radiographic recurrence (HR 0.2337, *p* = 0.0067), methylation of 15

marker candidates remained an independent significant predictor with HRs ranging from 3.2 (*PROM1*, *p* = 0.04) to 7.968 (*CHST11*, *p* = 0.0004) (Table 1). Notably, 11 of the marker genes also remained significant in co-variate analysis adjusting for PSA as a variable (HR 3.1–6.8) suggesting that our markers are suitable for prognosis independent of PSA levels. This finding might have important implications for detecting tumors that progress regardless of PSA increase.

Taken together our findings reveal the potential of DNA methylation-based marker genes to monitor treatment response in mCRPC patients at an early stage following therapy administration. Importantly, using radiographic recurrence as an endpoint, we reveal several methylation markers that could predict tumor progression following treatment. Due to the small sample size and heterogenous treatment history of our study population, it will be important to validate our findings in an independent homogenous cohort in future studies.

Conclusions

To this date, mCRPC remains an incurable disease. A variety of therapeutic approaches for the clinical management of mCRPC have been developed recently, and it is essential to closely monitor therapy response in those patients to maximize their survival and quality of life. Our study presents several DNA methylation-based biomarkers with the potential to detect metastatic disease and to monitor treatment response and predict disease progression in liquid biopsies of patients with advanced cancer. Aside from their usability as cancer specific biomarkers, some of our candidates might also have important biological functions for prostate cancer [17–24]. As a next step, it will be important to test the performance of our methylation markers in prospective clinical trials including mCRPC patients undergoing different treatment regimes. We envision the development of a multiplex MSRE-qPCR kit including our 3-gene signatures for routine testing of therapy response of patients with advanced PCa in clinical labs in the future.

Abbreviations

PCa: Prostate cancer; PSA: Prostate specific antigen; BPH: Benign prostatic hyperplasia; ADT: Androgen deprivation therapy; CRPC: Castration resistant prostate cancer; mCRPC: Metastatic castration resistant prostate cancer; ctDNA: Cell-free circulating tumor DNA; CTCs: Circulating tumor cells; FFPE: Formalin fixed paraffin embedded; PSMA: Prostate specific membrane antigen; PMR: Percentage methylated reference; ccfDNA: Circulating cell-free DNA; Tm: Melting temperature; MSRE-qPCR: Methylation sensitive restriction enzyme – quantitative PCR; ms-qPCR: Methylation-specific quantitative PCR; PBMC: Peripheral blood mononuclear cells; sQ: Sample quantity; DLDA: Diagonal linear discriminant analysis; CCP: Compound covariate predictor; BCCP: Bayesian compound covariate predictor; rPFS: Radiographic progression-free survival; AUC: Area under the curve; ROC: Receiver operating characteristics; CT: Computed tomography; PET: Positron emission tomography.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12943-021-01445-0>.

Additional file 1: Table S1. Patient cohorts and characteristics. **Table S2.** List of biomarker assays for MSRE analysis of liquid biopsies. **Table S3.** Classification results of benign versus mCRPC. **Table S4.** Classification results of primary PCa versus mCRPC. **Table S5.** Classification results of benign versus primary PCa GSc 9+. **Table S6.** Classification results of benign+primary PCa versus mCRPC. **Table S7.** Clinical data of mCRPC responder and non-responder patients. **Table S8.** Classification results of mCRPC non-responder post-treatment vs. mCRPC responder post-treatment. **Table S9.** Classification results of mCRPC non-responder post-treatment vs. mCRPC responder post-treatment PMR-values from patients that underwent Abiraterone-acetate treatment. **Table S10.** Primers used for ms-qPCR assays

Additional file 2: Figure S1. Validation of maker candidates. **Figure S2.** ROC analysis and cfDNA concentration. **Figure S3.** Survival analysis of responder and non-responder patients.

Additional file 3: Materials and Methods.

Acknowledgements

We thank Helga Schachner, Margit Schmeidl and Jasmin Huber for technical support.

Availability data and materials

The results within this publication are in part based upon data generated by the TCGA Research Network: <https://www.cancer.gov/tcga>. In addition, the datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

G.E. and T.D. conceived and designed the study and drafted the manuscript. T.D., W.P., MR.H., J.Sch., M.K., S.D. and E.R. performed experiments. T.D., W.P., R.S.-T., I.S. and CU.P.M. performed data and statistical analyses. S.E., N.H., B.G. collected patient samples and data. S.B., A.G. provided samples and clinical data. A.T., K.R. provided patient data. L.K. performed pathological assessment. A.W., J.S., M.Z., S.F.S. supervised parts of the study. M.H. and M.M. provided resources. The author(s) read and approved the final manuscript.

Funding

This study was supported by the Austrian National Bank (OeNB) Jubileefund (Nr. 13061) and by a Marie Curie Actions- International Re-integration Grant (IRG) (FP7-People-IRG-2008; Nr. 230984).

Declarations

Ethics approval and consent to participate

All patients included in this study were treated in the General Hospital of Vienna and were included upon informed consent. All study protocols for the analysis of patient tissues and liquid biopsies were approved by the ethics committee of the Medical University of Vienna (Study numbers: 1492/2012, 1224/2014, 1932/2018).

Consent for publication

All authors agree to publication of the article.

Competing interests

The authors declare that they have no conflicts of interest.

Author details

¹Ludwig Boltzmann Institute Applied Diagnostics, Vienna, Austria. ²Department of Pathology, Medical University of Vienna, Vienna, Austria. ³Health & Environment Department, Molecular Diagnostics, AIT-Austrian Institute of Technology GmbH, Vienna, Austria. ⁴Department of Health Economics, Center for Public Health, Medical University of Vienna, Vienna, Austria. ⁵Department of Urology, Medical University Vienna, Vienna, Austria. ⁶Department

of Clinical Pharmacology, Medical University of Vienna, Vienna, Austria.

⁷Institute of Cancer Research, Department of Medicine I, Medical University of Vienna, Vienna, Austria. ⁸Division of Nuclear Medicine, Department of Biomedical Imaging and Image-Guided Therapy, Vienna, Austria. ⁹Comprehensive Cancer Center, Medical University of Vienna, Vienna, Austria. ¹⁰Department of Urology, University of Texas Southwestern Medical Center, Dallas, TX, USA. ¹¹Department of Urology, Weill Cornell Medical College, New York, NY, USA. ¹²Department of Urology, Second Faculty of Medicine, Charles University, Prague, Czech Republic. ¹³Karl Landsteiner Institute of Urology and Andrology, Vienna, Austria. ¹⁴Division of Urology, Department of Special Surgery, Jordan University Hospital, The University of Jordan, Amman, Jordan. ¹⁵European Association of Urology Research Foundation, Arnhem, The Netherlands. ¹⁶Unit of Laboratory Animal Pathology, University of Veterinary Medicine, Vienna, Austria. ¹⁷Christian Doppler Laboratory for Applied Metabolomics, Medical University of Vienna, Vienna, Austria.

Received: 2 August 2021 Accepted: 22 October 2021

Published online: 03 January 2022

References

- Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global Cancer Statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2021;71:209–49.
- Potosky AL, Feuer EJ, Levin DL. Impact of screening on incidence and mortality of prostate cancer in the United States. *Epidemiol Rev*. 2001;23:181–6.
- Sumanasuriya S, Omlin A, Armstrong A, Attard G, Chi KN, Bevan CL, et al. Consensus statement on circulating biomarkers for advanced prostate cancer. *Eur Urol Oncol*. 2018;1:151–9.
- Parker C, Castro E, Fizazi K, Heidenreich A, Ost P, Procopio G, et al. *clinicalguidelines@esmo.org* EGCEa: prostate cancer: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2020;31:1119–34.
- Rebello RJ, Oing C, Knudsen KE, Loeb S, Johnson DC, Reiter RE, et al. Prostate cancer. *Nat Rev Dis Primers*. 2021;7:9.
- Ignatiadis M, Sledge GW, Jeffrey SS. Liquid biopsy enters the clinic - implementation issues and future challenges. *Nat Rev Clin Oncol*. 2021;18:297–312.
- Lam D, Clark S, Stirzaker C, Pidsley R. Advances in prognostic methylation biomarkers for prostate cancer. *Cancers*. 2020;12(10):2993.
- Kobayashi Y, Absher DM, Gulzar ZG, Young SR, McKenney JK, Peehl DM, et al. DNA methylation profiling reveals novel biomarkers and important roles for DNA methyltransferases in prostate cancer. *Genome Res*. 2011;21:1017–27.
- Aryee MJ, Liu W, Engelmann JC, Nuhn P, Gurel M, Haffner MC, et al. DNA methylation alterations exhibit intraindividual stability and interindividual heterogeneity in prostate cancer metastases. *Sci Transl Med*. 2013;5:169ra110.
- Beikircher G, Pulverer W, Hofner M, Noehammer C, Weinhaeusel A. Multiplexed and sensitive DNA methylation testing using methylation-sensitive restriction enzymes "MSRE-qPCR". *Methods Mol Biol*. 2018;1708:407–24.
- Bjerre MT, Norgaard M, Larsen OH, Jensen SO, Strand SH, Ostergren P, et al. Epigenetic analysis of circulating tumor DNA in localized and metastatic prostate cancer: evaluation of clinical biomarker potential. *Cells*. 2020;9.
- Hennigan ST, Trostel SY, Terrigino NT, Voznesensky OS, Schaefer RJ, Whitlock NC, et al. Low abundance of circulating tumor DNA in localized prostate cancer. *JCO Precis Oncol*. 2019;3:PO.19.00176.
- Chen E, Cario CL, Leong L, Lopez K, Marquez CP, Chu C, et al. Cell-free DNA concentration and fragment size as a biomarker for prostate cancer. *Sci Rep*. 2021;11:5040.
- Mouliere F, Chandrananda D, Piskorz AM, Moore EK, Morris J, Ahlborn LB, et al. Enhanced detection of circulating tumor DNA by fragment size analysis. *Sci Transl Med*. 2018;10(466):eaat4921.
- Herman D, Leakey TI, Behrens A, Yao-Borengasser A, Cooney CA, Jousheghany F, et al. CHST11 gene expression and DNA methylation in breast cancer. *Int J Oncol*. 2015;46:1243–51.

16. Powers E, Karachaliou GS, Kao C, Harrison MR, Hoimes CJ, George DJ, et al. Novel therapies are changing treatment paradigms in metastatic prostate cancer. *J Hematol Oncol.* 2020;13:144.
17. Currall BB, Chen M, Sallari RC, Cotter M, Wong KE, Robertson NG, et al. Loss of LDAH associated with prostate cancer and hearing loss. *Hum Mol Genet.* 2018;27:4194–203.
18. Ilijazi D, Pulverer W, Ertl IE, Lemberger U, Kimura S, Abufaraj M, et al. Discovery of molecular DNA methylation-based biomarkers through genome-wide analysis of response patterns to BCG for bladder cancer. *Cells.* 2020;9(8):1839.
19. Khayami R, Hashemi SR, Kerachian MA. Role of aldo-keto reductase family 1 member B1 (AKR1B1) in the cancer process and its therapeutic potential. *J Cell Mol Med.* 2020;24:8890–902.
20. Lee HM, Hwang KA, Choi KC. Diverse pathways of epithelial mesenchymal transition related with cancer progression and metastasis and potential effects of endocrine disrupting chemicals on epithelial mesenchymal transition process. *Mol Cell Endocrinol.* 2017;457:103–13.
21. Moller M, Strand SH, Mundbjerg K, Liang G, Gill I, Haldrup C, et al. Heterogeneous patterns of DNA methylation-based field effects in histologically normal prostate tissue from cancer patients. *Sci Rep.* 2017;7:40636.
22. Pique L, de Paz Martinez A, Pineyro D, Martinez-Cardus A, Castro de Moura M, Llinas-Arias P, et al. Epigenetic inactivation of the splicing RNA-binding protein CELF2 in human breast cancer. *Oncogene.* 2019;38:7106–12.
23. Vega-Benedetti AF, Loi E, Moi L, Blois S, Fadda A, Antonelli M, et al. Clustered protocadherins methylation alterations in cancer. *Clin Epigenetics.* 2019;11:100.
24. Xu N, Wu YP, Ke ZB, Liang YC, Cai H, Su WT, et al. Identification of key DNA methylation-driven genes in prostate adenocarcinoma: an integrative analysis of TCGA methylation data. *J Transl Med.* 2019;17:311.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

