




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Letter to the Director

About “Absence of Relevant Clinical Effects of SARS-CoV-2 on the Affinity of Hemoglobin for O₂ in Patients with COVID-19”


To the Director,

The SARS-CoV-2 virus damages the lungs and also the erythrocytes which often results in arterial hypoxia and severe anemia. Both factors typically cause a right shift of the hemoglobin oxygen dissociation curve (ODC) in man due to an increase in the concentration of 2,3-bisphosphoglycerate in the red cells. Consequently oxygen discharge to the consuming tissues is improved but the uptake in the lungs is hindered. Surprisingly, however, investigations in COVID-19 patients detected no changes or even a left shift of the ODC (reviewed in Ref. 1). Notably, COVID-19 effects on the ODC may depend on the severity of the illness: In mechanically ventilated patients with profound anemia² a clear left shift was observed. In contrast Gille et al.³ detected no significant change in less severe cases without marked anemia, although the number of measurements might be too low for a decisive conclusion in our opinion.⁴ In the recent publication of Pascual-Guardia et al. in this journal⁵ the patients do on average not seem severely ill. Mechanical ventilation is not mentioned and anemia is mostly moderate with mean Hb concentrations of 13.3 ± 1.8 SD g/dl in arterial samples from 139 patients, while unfortunately values are lacking for the 215 patients with venous samples. Notably, the authors did not detect a significant overall change in oxygen affinity but a very large scattering. Specifically, the difference between measured and calculated (from different standard curves) SO₂ in ca. 35 venous samples is larger than two times the standard deviation (approx. $\pm 6\%$), in 22 of these samples the measured values exceed the calculated ones. The latter corresponds to a marked reduction of the standard P₅₀ by at least 5 mmHg compared to the normal value of approximately 27 mmHg. In view of these large deviations in some patients and the potential effect of disease severity on ODC discussed above, it would be of considerable interest to check whether P₅₀ depends on changes of the Hb-concentration or other indicators of severe illness (e. g. mechanical ventilation) in this patient cohort. One could expect that the results for critical illness with heavy anemia will be similar to those reported by Vogel et al.,² and thus explain the large scatter in the study by Pascual-Guardia et al.⁵ In this context it is interesting that erythrocytes with injured membranes can often but not always be detected in lung biopsies from COVID-19 patients (W. Bloch, unpublished results). Without

an intact membrane the intraerythrocytic pH and therefore the oxygen affinity must be increased because of a disappearing Donnan equilibrium. Another cause for a left shift of the ODC in this scenario might be a loss of 2,3-bisphosphoglycerate (2,3-BPG) to the plasma.

A notable drawback of all papers cited here is that 2,3-BPG has not been measured in COVID-19 patients. The likely reason is that formerly used test kits are no longer available. In view of the current pandemic, there is now an urgent need for the reintroduction of 2,3-BPG measurements.

Funding

No funders pertaining to this article.

Conflict of interests

The authors declare no conflict of interests.

References

- Böning D, Kuebler WM, Bloch W. The oxygen dissociation curve of blood in COVID-19. *Am J Physiol Lung Cell Mol Physiol.* 2022;321:L349–357. <http://dx.doi.org/10.1152/ajplung.00079.2021>.
- Vogel DJ, Formenti F, Retter AJ, Vasques F, Camporota L. A left shift in the oxyhaemoglobin dissociation curve in patients with severe coronavirus disease 2019 (COVID-19). *Br J Haematol.* 2020;191:390–3.
- Gille T, Sese L, Aubourg E, Fabre EE, Cymbalista F, Ratnam KC, et al. The affinity of hemoglobin for oxygen is not altered during COVID-19. *Front Physiol.* 2021;12:578708.
- Böning D, Bloch W, Kuebler W. The pathophysiology of COVID-19 and SARS-CoV-2 infection. Reply to Gille et al. *Am J Physiol Lung Cell Mol Physiol.* 2022;322:L176–177. <http://dx.doi.org/10.1152/ajplung.00461.2021>.
- Pascual-Guardia S, Ferrer A, Diaz O, Caguana AO, Tejedor E, Bellido-Calduch S, et al. Absence of relevant clinical effects of SARS-CoV-2 on the affinity of hemoglobin for O₂ in patients with COVID-19. *Arch Bronconeumol (Engl Ed).* 2021;57:757–63.

Dieter Böning^{a,*}, Wilhelm Bloch^b, Wolfgang M. Kuebler^a

^a *Institut für Physiologie, Campus Mitte, Charité – Universitätsmedizin Berlin, Virchowweg 6, 10117 Berlin, Germany*

^b *Institut für Kreislaufforschung und Sportmedizin, Deutsche Sporthochschule Köln, Am Sportpark Müngersdorf 6, 50933 Köln, Germany*

Corresponding author.

E-mail address: dieter.boening@charite.de (D. Böning).