



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Letter to the Editor Regarding: Acute Coronary Syndrome in the COVID-19 Pandemic: Reduced Cases and Increased Ischaemic Time by Sutherland et al. *Heart Lung Circ.* 2022;31(1):69-76



To the Editor,

Sutherland and colleagues recently reported a reduction in patients admitted with acute coronary syndrome (ACS) and an increase in total ischaemic time during the first and second wave of the COVID-19 outbreak in Melbourne [1]. They also observed a 20% rebound increase in ACS presentations following the relaxation of public health restrictions in November–December 2020 compared to the same period of 2019 and suggested that this could represent long-term sequelae of untreated ACS including reinfarction and heart failure. However, several additional mechanisms could play a role.

Daily Life Triggers

A trend of reduced hospitalisations for all ACS types during the pandemic has been observed worldwide [1–3]. Besides social distancing, stay-at-home orders, and fear of acquiring the infection, attenuated exposure to well-recognised ACS triggers, particularly reduced air pollution, and decreased physical and work activities, has been linked to this phenomenon [3,4]. The alleviation of restrictions after the pandemic waves restored the level of exposure to ACS triggers which could have contributed to a gradual increase in ACS hospitalisations [4]. Moreover, fear of lack of medical care, lockdown stress, anger, loneliness, job loss, financial stress, and binge smoking have been suggested as ACS triggers due to the ramification of the pandemic [5]. Along this line, a post-restriction increase in hospitalisations for ST-segment elevation myocardial infarction has been documented in Israel [2].

Population Vulnerability

Anti-pandemic measures favour sedentary behaviour, physical inactivity, unhealthy nutritional habits, weight gain,

and increased alcohol consumption, as confirmed by recent meta-analyses [6,7]. Consequent negative effects on metabolism likely worsened the population's cardiovascular health and, given their presence for over a year now, could have increased the number of vulnerable coronary patients [4]. Accordingly, increased population cardiovascular vulnerability coupled with enhanced exposure to ACS triggers could be important mechanisms of a rebound increase in ACS.

Viktor Čulić, MD, PhD^{a,b*}
Ahmed Alturki, MD^c
Riccardo Proietti, MD, PhD^d

^aDepartment of Cardiology and Angiology, University Hospital Centre Split, Split, Croatia

^bUniversity of Split School of Medicine, Split, Croatia
^cDivision of Cardiology, McGill University Health Centre Montreal, Quebec, Canada

^dLiverpool Centre for Cardiovascular Science, University of Liverpool, Liverpool, UK

*Corresponding author at: Department of Cardiology and Angiology, University Hospital Centre Split, Šoltanska 1, 21000 Split, Croatia
Email: viktor.culic@st.t-com.hr

Received 11 October 2021; Accepted 7 November 2021; online published-ahead-of-print xxx

References

- [1] Sutherland N, Dayawansa NH, Filipopoulos B, Vasanthakumar S, Narayan O, Ponnuthurai FA, et al. Acute coronary syndrome in the COVID-19 pandemic: reduced cases and increased ischaemic time. *Heart Lung Circ.* 2022;31:69–76. <https://doi.org/10.1016/j.hlc.2021.07.023>.
- [2] Fardman A, Oren D, Berkovitch A, Segev A, Levy Y, Beigel R, et al. Post Covid-19 acute myocardial infarction rebound. *Can J Cardiol.* 2020;36:1832.e15–6.

- [3] Rattka M, Dreyhaupt J, Winsauer C, Stuhler L, Baumhardt M, Thiessen K, et al. Effect of the COVID-19 pandemic on mortality of patients with STEMI: a systematic review and meta-analysis. *Heart*. 2021. <https://doi.org/10.1136/heartjnl-2020-318360>.
- [4] Čulić V, AlTurki A, Proietti R. Covid-19 pandemic and possible rebound phenomenon in incidence of acute myocardial infarction. *Can J Cardiol*. 2021;37(8):1294.
- [5] Hammoudeh A, Abu-Hantash H, Tabbalat R, Al-Mousa E, Madanat E, Al-Muhaisen R, et al. The Covid-19 pandemic and triggered acute myocardial infarction among non-infected individuals. *Int J Clin Cardiol*. 2020;7:185.
- [6] Bakaloudi DR, Barazzoni R, Bischoff SC, Breda J, Wickramasinghe K, Chourdakis M. Impact of the first COVID-19 lockdown on body weight: a combined systematic review and a meta-analysis. *Clin Nutr*. 2021. <https://doi.org/10.1016/j.clnu.2021.04.015>.
- [7] Bakaloudi DR, Jeyakumar DT, Jayawardena R, Chourdakis M. The impact of COVID-19 lockdown on snacking habits, fast-food and alcohol consumption: a systematic review of the evidence. *Clin Nutr*. 2021. <https://doi.org/10.1016/j.clnu.2021.04.020>.