

RESEARCH ARTICLE

Suicidal behaviors among Bangladeshi university students: Prevalence and risk factors

M. Rasheduzzaman^{1,2}, Firoj al-Mamun^{1,3}, Ismail Hosen^{1,3}, Tahmina Akter^{1,4}, Moazzem Hossain⁵, Mark D. Griffiths⁶*, Mohammed A. Mamun^{1,3}✉*

1 CHINTA Research Bangladesh, Savar, Dhaka, Bangladesh, **2** Institute of Social Welfare and Research, University of Dhaka, Dhaka, Bangladesh, **3** Department of Public Health and Informatics, Jahangirnagar University, Savar, Dhaka, Bangladesh, **4** Department of Epidemiology, Bangladesh University of Health Sciences, Dhaka, Bangladesh, **5** Institute of Allergy and Clinical Immunology of Bangladesh, Savar, Dhaka, Bangladesh, **6** Psychology Department, Nottingham Trent University, Nottingham, United Kingdom

✉ These authors contributed equally to this work.

* mamunphi46@gmail.com (MAM); mark.griffiths@ntu.ac.uk (MDG)



OPEN ACCESS

Citation: Rasheduzzaman M, al-Mamun F, Hosen I, Akter T, Hossain M, Griffiths MD, et al. (2022) Suicidal behaviors among Bangladeshi university students: Prevalence and risk factors. PLoS ONE 17(1): e0262006. <https://doi.org/10.1371/journal.pone.0262006>

Editor: Muhammed Elhadi, University of Tripoli, LIBYA

Received: March 6, 2021

Accepted: December 15, 2021

Published: January 13, 2022

Copyright: © 2022 Rasheduzzaman et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) files.

Funding: The present study did not get any financial supports. Besides, the authors involved in this research communication do not have any relationships with other people or organizations that could inappropriately influence (bias) the findings.

Competing interests: The authors of the research work do not have any conflict of interest.

Abstract

Background

Bangladeshi university students are considered to be highly suicide-prone compared to other populations and cohorts. However, no prior epidemiological studies have assessed the suicidality (i.e., past-year suicidal ideation [SI], lifetime suicide plan [SP], and lifetime suicide attempt [SA]) among Bangladeshi students, including the variables such as past-year stressful life events and family mental health history. This is arguably a major knowledge gap in the country. Therefore, the present study investigated the prevalence and associated risk factors for suicidal behaviors among Bangladeshi university students.

Methods

A cross-sectional study was conducted utilizing a convenience sampling method among a total of 1844 university students between October and November 2019. Data were collected based on the information related to socio-demographics, perceived health-related questions, past-year stressful life events, family mental health history, and suicidal behaviors (i.e., SI, SP, and SA). Chi-square tests and binary logistic regressions were used to analyze the data utilizing SPSS statistical software.

Results

The prevalence of past-year suicidal ideation, lifetime suicide plans, and suicide attempts were 13.4%, 6.0%, and 4.4%, respectively. Females reported significantly higher suicidal behavior than males (i.e., 20.6% vs. 10.2% SI; 9% vs. 4.6% SP; and 6.4% vs. 3.6% SA). Risk factors for SI were being female, year of academic study, residing in an urban area, using psychoactive substances, experiencing both past year physical and mental illness, experiencing any type of stressful past-year life events, experiencing campus ragging (i.e., senior students abusing, humiliating and/or harassing freshers or more junior students),

experiencing family mental illness history, and having family suicide attempt history. SP was associated with several factors including being female, year of academic study, using psychoactive substance, experiencing both past-year physical and mental illness, and experiencing any type of stressful past-year life events. Risk factors for SA were being female, year of academic study, using psychoactive substances, experiencing past-year mental illness, experiencing any type of stressful past-year life events, and having family suicide attempt history.

Conclusions

University students appear to be a vulnerable group for experiencing suicidal behaviors. The present findings warrant rigorous action and early intervention programs such as counseling and other mental health professional services by university authorities. Longitudinal studies are highly recommended involving countrywide representative samples.

1 Introduction

Suicidal behaviors can be defined as individuals experiencing repeated thoughts of killing themselves life (suicidal ideation), planning to kill themselves (suicide plan), and actual efforts to kill themselves (suicide attempt), while suicide refers to actually killing themselves. Suicidal behaviors are frequently accompanied by overwhelming hopelessness, depression, or self-destructive behavior (parasuicidal behaviors) [1]. According to a recent meta-analysis, the prevalence of suicidal ideation worldwide is reported to be 10.62% for past-year, 6.14% for lifetime suicide plan, and 3.22% for lifetime suicide attempt [2]. However, suicide has become a global public health problem and accounts for nearly 800,000 deaths among all age groups every year [3]. Of these suicides, 79% of all cases occur in low-income and middle-income countries like Bangladesh [3]. Suicide mostly affects the 15-29-year age group (many of whom are likely to be students) and is the second-highest cause of death after unintentional injury-related deaths from accidents [3].

The present study was carried out in Bangladesh, and a recent Bangladeshi retrospective study reported that individuals aged below 30 years account for almost 61% of the total suicide deaths [4]. Similarly, a few recent retrospective studies using media reports have also explored Bangladeshi students' suicide vulnerability. For instance, one study reported five student suicides within a ten-day period at the University of Dhaka [5], and another study reported 13 Bangladeshi medical sciences student suicides in a 23-month period [6]. Moreover, another study reported 56 Bangladeshi students' suicide cases from January 2018 to June 2019 [7].

In Bangladesh, the number of university students has steadily increased over the past few years, but university facilities and subsequent career infrastructure do not meet many students' needs [8]. Furthermore, there are many problems related to campus and academic life in Bangladesh (i.e., lack of proper accommodation, campus ragging (i.e., senior students abusing, humiliating and/or harassing freshers or more junior students), political violence, poor environment and academic facilities, economic hardship due to living costs) [9, 10]. Along with the aforementioned issues, there are psychological stressors related to the lack of job security and career progression after graduation in Bangladesh [8]. These issues are highly associated with mental health suffering, and recent studies have reported that more than half of Bangladeshi students have mental health issues [9, 11, 12], where similar mental health suffering was

noted among the job-seeking graduates [8]. Based on these findings, it is evident that the current Bangladeshi students appear to be at high risk of mental health disorders due to the aforementioned academic and job-related problems. These mental health disorders also contribute to suicide and suicidal behaviors by mediating both distal and proximal suicide risk factors [13–15]. Other risk factors for suicide and suicide-related behaviors include suffering from physical diseases [16–18], stressful life events [14, 19–22], having a family history of mental disorders and suicide [23–26].

A recent meta-analysis claimed that expression of suicidal behaviors (i.e., suicidal ideation) is one of the prominent predictors of suicide completion [27], and successful suicides are often preceded by up to 20 previous attempts suggested by the World Health Organization [3]. But evidence-based data on suicidal behaviors (i.e., suicidal ideation, suicide plans, and suicide attempts) are needed for suicide prevention programs to inform policy-based legislation and public health strategies, public and physician education, and general awareness [3]. Although suicide is one of the preventable public health problems, it has not been effectively addressed in Bangladesh because there is less awareness regarding suicide prevention [7, 22, 28]. Consequently, epidemiological data is much needed for suicide prevention activities in Bangladesh. Therefore, the present study explored suicidal behaviors among Bangladeshi university students and examined associated risk factors (socio-demographics, personal health-related behaviors and traumatic events, and family mental illness and suicide history).

2 Methods

2.1 Study procedure and participants

A cross-sectional study was conducted among undergraduate students at the University of Dhaka, Bangladesh (mean age = 20.92 years; $SD \pm 1.72$ years) during October and November 2019. The data were collected through a 'paper-and-pencil' survey administered during lectures across all departments of the university by the research team. A convenience sampling technique was used to collect data from participants. Approximately 2,000 students were approached to participate in the survey, with 1897 agreeing to take part (94.6% response rate). Inclusion criteria for the study were (i) being a student of the university and (ii) being present in the class during data collection. Participants were excluded if they were not currently students at the university or were graduate students of the university. After removing the incomplete questionnaires, data from 1844 participants remained for final analysis. Prior to survey completion, study-related issues were introduced, and the research team briefed participants about the whole survey, including the terminology used. The survey took approximately 35 minutes to complete.

2.2 Measures

2.2.1 Sociodemographic factors. This survey included questions relating to sociodemographic variables such as age, gender, and whether the participants came from a rural or urban area.

2.2.2 Perceived health-related questions. Self-rated health status, that is, suffering from any type of past-year physical illnesses (e.g., diabetes, asthma, chronic pain, dengue, etc.) and past-year mental health illness (e.g., mood disorders, anxiety disorders, psychotic disorders, trauma-related disorders, etc.) was assessed based on a previous study conducted in Bangladeshi context [29]. Additionally, students were asked if they currently smoked cigarettes and engaged in any other psychoactive substance use (e.g., alcohol, cannabis, illicit drugs, non-medical use of prescription drugs) using a binary response option (i.e., yes/no).

2.2.3 Past-year stressful life events. Past-year stressful life events (i.e., if they had a failure in the examination, if they had relationship complexities, if they were bullied on campus [ragging], if they had family problems, and if they had other problems) were assessed utilizing a binary response ‘yes/no’ response.

2.2.4 Family mental health history. The history of family mental illness (if any of the family members had any mental illness), suicide completion (if any family members had actually committed suicide), and suicide attempt (if any family members attempted suicide) were assessed using a binary response (‘yes/no’) for each of these three variables.

2.2.5 Suicidal behaviors. To assess suicidal behaviors (i.e., suicidal ideation, suicide plans, and suicide attempts), questions used in previous studies were utilized (i.e., binary ‘yes/no’ responses). Participants were asked if they had ever thought about committing suicide during the past year (past-year suicidal ideation; SI), whether such thoughts were persistent across their lifetime, whether they had ever made suicide plans to kill themselves (lifetime suicide plan; SP), and whether they had ever attempted suicide during their lifetime (lifetime suicidal attempt; SA) [30–32].

2.3 Ethical considerations

The study followed the medical ethical guidelines of Helsinki Declaration, 1975. The study was reviewed and approved by the ethics board of the Institutional Review Board of the Institute of Allergy and Clinical Immunology of Bangladesh (IACIB), Dhaka, Bangladesh [Reference Number: IRBIACIB/CEC/07201903]. All participants signed an informed consent form prior to participating in the study, and were assured that their data would be anonymous and confidential. They were also informed about the nature, purpose, and procedure of the study, as well as being informed about the right to withdraw their data at any time from the study.

2.4 Statistical analysis

This study utilized Statistical Package for Social Science (SPSS) version 22.0 for the data analysis. The analysis included descriptive and inferential statistics such as frequencies, percentages, and means. First-order analysis, including chi-squares and binary logistic regression, also utilized SPSS. All of the variables were added in the unadjusted model (univariate analysis) and then the adjusted model (multivariate analysis) only included the significant variables in the unadjusted model. The unadjusted model was applied for a single predictor and adjusted model was responsible for more than one predictor and where past-year suicidal ideation, lifetime suicide plan, and lifetime suicide attempts were considered as the dependent variables. Odds ratios were used as a measure of risk association, confidence intervals were used as a measure of estimation/precision, and significance levels ($p < 0.05$) were used as a measure of statistical significance.

3 Results

3.1 Characteristics of the participants

The participants’ characteristics are shown in [Table 1](#). The sample comprised 70% males, 84.9% came from a village area, 16.7% were cigarette smokers, 3.3% were psychoactive substance users, 10.4% had suffered from physical illnesses in the past year, and 8.4% had experienced mental health psychological suffering. The number of females was less in the present study simply because there was a much smaller proportion of females enrolled at the university. Results also indicated that in the past year, 31.4% had experienced stressful life events, 11.8% had failed examinations, 13.1% had relationship difficulties, 29.2% experienced ragging

Table 1. Distribution of the variables with suicidal behaviors.

Variables	Total; n (%)	Past-year suicidal ideation (N = 247; 13.4%)			Life-time suicide plans (N = 110; 6.0%)			Life-time suicide attempts (N = 82; 4.4%)		
		Yes; n (%)	χ^2 test value	p-value	Yes; n (%)	χ^2 test value	p-value	Yes; n (%)	χ^2 test value	p-value
Socio-demographic factors										
Gender										
Female	567; 30.7%	117; 20.6%	36.997	<0.001	51; 9.0%	13.395	<0.001	36; 6.4%	7.018	0.008
Male	1277; 69.3%	130; 10.2%			59; 4.6%			46; 3.6%		
Year of study										
4 th year	303; 16.5%	65; 21.5%	22.986	0.069	36; 11.9%	23.446	<0.001	29; 9.6%	23.167	<0.001
3 rd year	507; 27.6%	50; 9.9%			25; 4.9%			19; 3.8%		
2 nd year	519; 28.2%	68; 13.1%			21; 4.0%			20; 3.9%		
1 st year	511; 27.8%	63; 12.3%			28; 5.5%			14; 2.7%		
Permanent residence										
Rural	1544; 84.9%	192; 12.4%	9.320	<0.001	94; 6.0%	0.169	0.681	72; 4.6%	0.578	0.447
Urban	277; 15.1%	53; 19.1%			15; 5.4%			10; 3.6%		
Personal health-related variables										
Cigarette smoker										
Yes	308; 16.7%	51; 16.6%	3.296	0.069	32; 10.4%	12.879	<0.001	29; 9.4%	21.425	<0.001
No	1535; 83.3%	195; 12.7%			78; 5.1%			53; 3.5%		
Psychoactive substance user										
Yes	60; 3.3%	23; 38.3%	33.249	<0.001	17; 28.3%	55.316	<0.001	17; 28.3%	83.218	<0.001
No	1784; 96.7%	224; 12.6%			93; 5.2%			65; 3.6%		
Past-year physical health illness										
Yes	191; 10.4%	66; 34.6%	82.239	<0.001	43; 22.5%	104.012	<0.001	31; 16.2%	69.566	<0.001
No	1653; 89.6%	181; 10.9%			67; 4.1%			51; 3.1%		
Past-year mental health illness										
Yes	154; 8.4%	70; 45.5%	148.751	<0.001	59; 38.3%	313.214	<0.001	47; 30.5%	268.472	<0.001
No	1689; 91.6%	177; 10.5%			51; 3.0%			35; 2.1%		
Past-year stressful life events										
Any types of stressful life events during past-year										
Yes	573; 31.4%	155; 27.1%	134.623	<0.001	86; 15.0%	118.712	<0.001	62; 10.8%	80.228	<0.001
No	1250; 68.6%	89; 7.1%			24; 1.9%			19; 1.5%		
Examination failure										
Yes	217; 11.8%	65; 30.0%	58.134	<0.001	33; 15.2%	37.450	<0.001	26; 12.0%	33.136	<0.001
No	1627; 88.2%	182; 11.2%			77; 4.7%			56; 3.4%		
Relationship difficulties										
Yes	242; 13.1%	76; 31.4%	77.111	<0.001	53; 21.9%	126.103	<0.001	41; 17.0%	102.933	<0.001
No	1602; 86.9%	171; 10.7%			57; 3.6%			41; 2.6%		
Campus ragging										
Yes	120; 6.5%	35; 29.2%	27.523	<0.001	15; 12.5%	9.771	<0.002	6; 5.0%	0.092	0.762
No	1724; 93.5%	212; 12.3%			95; 5.5%			76; 4.4%		
Family problems										
Yes	213; 11.6%	66; 31.0%	64.238	<0.001	40; 18.8%	70.492	<0.001	36; 16.9%	87.838	<0.001
No	1631; 88.4%	181; 11.1%			70; 4.3%			46; 2.8%		
Others										
Yes	25; 1.4%	2; 8.0%	0.637	0.4257	1; 4.0%	0.175	0.676	1; 4.0%	0.012	0.912
No	1818; 98.6%	245; 13.5%			109; 6.0%			81; 4.5%		
Family history of psychiatric suffering										

(Continued)

Table 1. (Continued)

Variables	Total; n (%)	Past-year suicidal ideation (N = 247; 13.4%)			Life-time suicide plans (N = 110; 6.0%)			Life-time suicide attempts (N = 82; 4.4%)		
		Yes; n (%)	χ^2 test value	p-value	Yes; n (%)	χ^2 test value	p-value	Yes; n (%)	χ^2 test value	p-value
Family mental illness history										
Yes	229; 12.4%	75; 32.8%	84.925	<0.001	42; 18.3%	72.426	<0.001	31; 13.5%	50.651	<0.001
No	1612; 87.6%	171; 10.6%			67; 4.2%			51; 3.2%		
Family suicide history										
Yes	48; 2.6%	17; 35.4%	20.603	<0.001	11; 22.9%	25.246	<0.001	6; 12.5%	7.514	0.006
No	1796; 97.4%	230; 12.8%			99; 5.5%			76; 4.2%		
Family suicide attempt history										
Yes	99; 5.4%	38; 38.45%	56.314	<0.001	21; 21.2%	43.355	<0.001	18; 18.2%	46.408	<0.001
No	1745; 94.6%	209; 12.0%			89; 5.1%			64; 3.7%		

<https://doi.org/10.1371/journal.pone.0262006.t001>

by other students on campus, 31.0% had experienced family problems, and 8.0% reported experiencing other events (e.g., having personal items stolen [money, smartphone], being in or witnessing a road traffic accident, being humiliated by another person, being beaten up by another person, witnessing others' injuries and deaths, etc.). Finally, participants reported a history of family mental illness (12.4%), family suicide completion (2.6%) and family suicide attempts (5.4%) (Table 1).

3.2 Prevalence of suicidal behaviors

The present study found that 13.4% of the total participants had past-year suicidal ideation (SI), whereas 6.0% reported having made lifetime suicide plans (SP), and 4.4% had at least one-lifetime suicide attempt (SA) (Table 1).

3.3 Association between socio-demographics and suicidal behaviors

Results demonstrated that in relation to gender, females had higher rate of experiencing suicidal behaviors compared to males for SI (20.6% vs. 10.2%; $\chi^2 = 36.997$, $p < 0.001$), SP (9% vs. 4.6%; $\chi^2 = 13.395$, $p < 0.001$) and SA (6.4% vs. 3.6%; $\chi^2 = 7.018$, $p = 0.008$). Fourth-year students had significantly higher SP (11.9% vs. 5.5%, 4% and 4.9%; $p < 0.001$) and SA (9.6% vs. 2.7%, 3.9%, and 3.8%; $p < 0.001$) compared to first-year, second-year, and third-year year students respectively. Students from urban areas had higher SI compared to rural areas students (19.1% vs. 12.4%; $p < 0.001$; Table 1).

3.4 Association between health-related variables and suicidal behaviors

Results indicated that cigarette smoking was not associated with SI, but was significantly associated with SP (10.4% vs. 5.1%; $\chi^2 = 12.879$, $p < 0.001$) and SA (9.4% vs. 3.5%; $\chi^2 = 21.425$, $p < 0.001$). Psychoactive substance users reported a higher significant rate of all suicidal behaviors compared to non-users. Similarly, participants with a past-year health suffering (both physical and psychological) reported significantly higher levels of all types of suicidal behaviors than those who had no health suffering (Table 1).

3.5 Association between past-year stressful life events and suicidal behaviors

Participants with a history of any type of past-year stressful life event (compared to those that did not) had significantly higher levels of SI (27.1% vs. 7.1%; $\chi^2 = 134.623$, $p < 0.001$), SP

(15.0% vs. 1.9%; $\chi^2 = 118.712, p < 0.001$) and SA (10.8% vs. 1.5%; $\chi^2 = 80.228, p < 0.001$). Similarly, past-year stressful life events were significantly associated with suicidal behaviors. This including examination failure (SI: $\chi^2 = 58.134, p < 0.001$; SP: $\chi^2 = 37.450, p < 0.001$; and SA: $\chi^2 = 33.136, p < 0.001$), relationship difficulties (SI: $\chi^2 = 77.111, p < 0.001$; SP: $\chi^2 = 126.103, p < 0.001$; and SA: $\chi^2 = 102.933, p < 0.001$), family problems (SI: $\chi^2 = 64.238, p < 0.001$; SP: $\chi^2 = 70.492$; and $p < 0.001$; and SA: $\chi^2 = 87.838, p < 0.001$) and being ragged by other students on campus (SI: $\chi^2 = 27.523, p < 0.001$; and SP: $\chi^2 = 9.771, p < 0.001$ –although it was not associated with SA: $\chi^2 = 0.092, p = 0.762$) (**Table 1**).

3.6 Association between family mental health history and suicidal behaviors

Participants with a family history of psychiatric illness (compared to those who did not) had significantly higher levels of SI (32.8% vs. 10.6%; $\chi^2 = 84.925, p < 0.001$), SP (18.3% vs. 4.2%; $\chi^2 = 72.426, p < 0.001$) and SA (13.5% vs. 3.2%; $\chi^2 = 50.651, p < 0.001$). Similarly, participants with a suicide-related family history also had higher levels of all types of suicide behaviors compared to those that did not [i.e., suicide completion ($\chi^2 = 20.603, p < 0.001$; $\chi^2 = 25.246, p < 0.001$; and $\chi^2 = 7.514, p = 0.006$ for SI, SP and SA respectively) and suicide attempt ($\chi^2 = 56.314, p < 0.001$; $\chi^2 = 43.355, p < 0.001$; and $\chi^2 = 46.408, p < 0.001$ for SI, SP and SA respectively)] (**Table 1**).

3.7 Risk factors for suicidal ideation

Table 2 shows the risk factors for suicidal ideation utilizing multivariate analysis (Nagelkerke's $R^2 = 0.259$). The significant predictors were gender (using male as reference; AOR = 2.257, 95% CI = 1.60–3.17), year of academic study (using first-year as reference; AOR = 0.53, 95% CI = 0.34–0.83), residence (using living in an urban area as reference, AOR = 0.61, 95% CI = 0.42–0.90), past-year physical illness (using no physical illness as reference, AOR = 1.80, 95% CI = 1.19–2.73), past-year mental illness (using no mental illness as reference, AOR = 2.69, 95% CI = 1.73–4.22), any type of past-year stressful life events (using no past-year stressful life events as reference, AOR = 2.20, 95% CI = 1.45–3.34), family mental illness history (using no family mental illness history as reference, AOR = 1.56, 95% CI = 1.05–2.33), family suicide attempt history (using no family suicide attempt as reference, AOR = 2.07, 95% CI = 1.22–3.49) (**Table 2**).

3.8 Risk factors for suicide planning

Table 3 shows the risk factors for suicide planning utilizing multivariate analysis (Nagelkerke's $R^2 = 0.384$). The significant predictors were gender (using male as reference, AOR = 2.03, 95% CI = 1.21–3.42), year of academic study (using first-year as reference, AOR = 0.52, 95% CI = 0.27–0.98), psychoactive substance user (using no psychoactive substance use as reference, AOR = 2.74, 95% CI = 1.07–7.02), past-year physical illness (using no past-year physical illness as reference, AOR = 2.09, 95% CI = 1.22–3.58), past-year mental illness (using no past-year mental illness as reference, AOR = 7.74, 95% CI = 4.50–13.32), any type of stressful past-year life events (using no type of stressful past-year events as reference, AOR = 3.03, 95% CI = 1.62–5.68) (**Table 2**).

3.9 Risk factors for suicide attempts

Table 4 shows the risk factors for suicide attempts utilizing multivariate analysis (Nagelkerke's $R^2 = 0.379$). The significant risk factors were gender (using male as reference, AOR = 2.02, 95% CI = 1.11–3.67), year of academic study (using first-year as reference, AOR = 0.34, 95%

Table 2. Logistic regression analysis of the variables associated with suicidal ideation.

Variables	Unadjusted model			Adjusted model (-2 Log likelihood = 1137.861; Nagelkerke's R ² = 0.259)		
	Odds ratio (OR)	95% Confidence Interval (CI)	p-value	Adjusted odds ratio (AOR)	95% Confidence Interval (CI)	p-value
Gender						
Female	2.29	1.74–3.01	<0.001	2.25	1.60–3.17	<0.001
Male	Reference			Reference		
Year of study						
1 st year	0.51	0.35–0.75	<0.001	0.53	0.34–0.83	0.014
2 nd year	0.55	0.38–0.80		0.63	0.41–0.97	
3 rd year	0.40	0.26–0.59		0.51	0.32–0.81	
4 th year	Reference			Reference		
Permanent residence						
Rural	0.59	0.42–0.83	0.002	0.61	0.42–0.90	0.014
Urban	Reference			Reference		
Cigarette smoker						
Yes	1.36	0.97–1.90	0.070	1.35	0.85–2.15	0.191
No	Reference			Reference		
Psychoactive substance user						
Yes	4.32	2.52–7.42	<0.001	2.13	1.00–4.55	0.049
No	Reference			Reference		
Past-year physical illness						
Yes	4.29	3.06–6.00	<0.001	1.80	1.19–2.73	0.005
No	Reference			Reference		
Past-year mental illness						
Yes	7.11	5.00–10.13	<0.001	2.69	1.72–4.22	<0.001
No	Reference			Reference		
Any type of stressful life event during past year						
Yes	4.83	3.64–6.42	<0.001	2.20	1.45–3.34	<0.001
No	Reference			Reference		
Examination failure						
Yes	3.39	2.44–4.71	<0.001	1.34	0.89–2.01	0.153
No	Reference			Reference		
Relationship difficulties						
Yes	3.83	2.79–5.24	<0.001	1.18	0.77–1.81	0.444
No	Reference			Reference		
Campus ragging						
Yes	2.93	1.93–4.46	<0.001	1.71	1.01–2.89	0.044
No	Reference			Reference		
Family problems						
Yes	3.59	2.58–4.99	<0.001	1.21	0.78–1.86	0.381
No	Reference			Reference		
Other problems						
Yes	0.55	0.13–2.38	0.432	0.79	0.17–3.55	0.761
No	Reference			Reference		
Family mental illness history						
Yes	4.10	2.98–5.64	<0.001	1.56	1.05–2.33	0.027
No	Reference			Reference		
Family suicide history						
Yes	3.73	2.03–6.85	<0.001	1.32	0.63–2.75	0.456
No	Reference			Reference		

(Continued)

Table 2. (Continued)

Variables	Unadjusted model			Adjusted model (-2 Log likelihood = 1137.861; Nagelkerke's R ² = 0.259)		
	Odds ratio (OR)	95% Confidence Interval (CI)	p-value	Adjusted odds ratio (AOR)	95% Confidence Interval (CI)	p-value
Family suicide attempt history						
Yes	4.57	2.97–7.03	<0.001	2.07	1.22–3.49	0.006
No	Reference			Reference		

<https://doi.org/10.1371/journal.pone.0262006.t002>

CI = 0.15–0.74), psychoactive substance user (using no psychoactive substance use as reference, AOR = 3.62, 95% CI = 1.33–9.86), past-year mental illness (using no past-year substance use as reference, AOR = 8.71, 95% CI = 4.72–16.07), any type of past-year stressful life events (using no past-year stressful life events as reference, AOR = 2.15, 95% CI = 1.01–4.43), and family suicide attempt history (using no family suicide attempt history as reference, AOR = 2.38, 95% CI = 1.13–5.03) (Table 4).

4 Discussion

In the present study, findings indicated that the prevalence rate among Bangladeshi students for (i) past-year suicidal ideation (SI) was 13.4%, (ii) lifetime suicide plans (SP) was 6.0%, and (iii) lifetime suicide attempt (SA) was 4.4% respectively. In other Bangladeshi studies, the rate of past-year suicidality among university students was reported to be 28.5% in a multi-institutional study [11], 12.4% among dental students [10], and 17.7% among university entrance test exam students [33]. Compared to prior Bangladeshi studies, it is evident that the reported suicidal ideation in the present study appears to be lower.

A study of 19 colleges comprising 13,984 first-year students across eight countries (i.e., Australia, Belgium, Germany, Mexico, Northern Ireland, South Africa, Spain, and the United States) reported prevalence rates of 17.2% for past-year SI prevalence, 17.5% for lifetime SP, and 4.3% for lifetime SA [34]. Another study examining adolescents from 32 low-income and middle-income countries, reported a pooled past-year SI prevalence rate of 12.2% for males (11.7%–12.7%) and 16.2% for females (15.6%–16.7%) [30], compared to a past-year prevalence rate of 18.2% among Ghanaian high school students (N = 1,984 [31]). However, a recent meta-analysis among 36 studies comprising college students (N = 634,662 students: 15 undergraduate samples, four graduate samples, 11 mixed undergraduate/graduate samples, and six not reported) estimated prevalence rates of 10.62% for past-year SI (9.10% to 12.25%), 6.14% for lifetime SP (4.78% to 7.75%) and 3.22% for lifetime SA (2.16% to 4.46%) [2]. Based on the aforementioned suicidal behaviors prevalence rates, it can be concluded that the present sample had a higher prevalence of suicidal behaviors for SI (13.4% vs. 10.62%) and SA (4.4% vs. 3.22%), and an equivalent prevalence rate for SP (6.0% vs. 6.14%). These higher rates may be particularly due to the university itself because previous research in Bangladesh examining actual suicides (rather than suicidal behaviors more generally) at the same university as the present study (i.e., University of Dhaka) reported five suicidality cases within a 10-day period [5].

Globally, gender differences on suicidal death and suicidal behaviors have been consistent (i.e., the female suicide rate is lower than males, but they experience a higher prevalence of suicide-related behaviors—such as SA—compared to males) [35]. Compared to findings globally (i.e., more SA among females) and Bangladeshi suicide trends (i.e., more suicides among females), the present study's findings are consistent (i.e., females had higher prevalence rates among all types of suicidal behavior). In addition, depending on the reasons for suicide, the difference between males and females has been found to be higher due to economic problems,

Table 3. Logistic regression analysis of the variables associated with suicide plans.

Variables	Unadjusted model			Adjusted model (-2 Log likelihood = 545.245; Nagelkerke's R ² = 0.384)		
	Odds ratio (OR)	95% Confidence Interval (CI)	p-value	Adjusted odds ratio (AOR)	95% Confidence Interval (CI)	p-value
Gender						
Female	2.04	1.38–3.00	<0.001	2.03	1.21–3.42	0.007
Male	Reference			Reference		
Year of study						
1 st year	0.43	0.25–0.72	<0.001	0.52	0.27–0.98	0.012
2 nd year	0.31	0.17–0.54		0.32	0.16–0.64	
3 rd year	0.38	0.22–0.65		0.51	0.27–0.99	
4 th year	Reference			Reference		
Permanent Residence						
Rural	1.12	0.64–1.97	0.681	1.31	0.66–2.62	0.430
Urban	Reference			Reference		
Cigarette smoker						
Yes	2.16	1.40–3.33	<0.001	1.59	0.82–3.09	0.167
No	Reference			Reference		
Psychoactive substance user						
Yes	7.18	3.94–13.08	<0.001	2.74	1.07–7.02	0.035
No	Reference			Reference		
Past-year physical illness						
Yes	6.87	4.52–10.45	<0.001	2.09	1.22–3.58	0.007
No	Reference			Reference		
Past-year mental illness						
Yes	19.94	13.00–30.60	<0.001	7.74	4.50–13.32	<0.001
No	Reference			Reference		
Any type of stressful life events during past-year						
Yes	9.02	5.66–14.35	<0.001	3.03	1.62–5.68	<0.001
No	Reference			Reference		
Examination failure						
Yes	3.61	2.33–5.58	<0.001	1.11	0.63–1.94	0.711
No	Reference			Reference		
Relationship difficulties						
Yes	7.60	5.07–11.37	<0.001	1.51	0.86–2.65	0.151
No	Reference			Reference		
Campus ragging						
Yes	2.45	1.37–4.37	0.002	0.96	0.45–2.06	0.935
No	Reference			Reference		
Family problems						
Yes	5.15	3.39–7.84	<0.001	1.19	0.68–2.09	0.540
No	Reference			Reference		
Other problems						
Yes	0.65	0.08–4.87	0.678	0.99	0.11–8.44	0.997
No	Reference			Reference		
Family mental illness history						
Yes	5.17	3.42–7.83	<0.001	1.39	0.79–2.45	0.252
No	Reference			Reference		
Family suicide history						
Yes	5.09	2.52–10.29	<0.001	1.41	0.57–3.47	0.455
No	Reference			Reference		

(Continued)

Table 3. (Continued)

Variables	Unadjusted model			Adjusted model (-2 Log likelihood = 545.245; Nagelkerke's R ² = 0.384)		
	Odds ratio (OR)	95% Confidence Interval (CI)	p-value	Adjusted odds ratio (AOR)	95% Confidence Interval (CI)	p-value
Family suicide attempt history						
Yes	5.01	2.95–8.48	<0.001	1.75	0.88–3.48	0.110
No	Reference			Reference		

<https://doi.org/10.1371/journal.pone.0262006.t003>

relationship problems, and educational failure [36]. Studies have also reported that relationship complexities are the primary cause of suicide among females, whereas economic concerns and illness are the major causes of suicide among males [36, 37]. Moreover, other biological and/or psychological factors, including coping style, impulsivity, and personality, may influence gender differences in suicidal behaviors.

It should also be noted that the adjusted model in the present study provides a more accurate depiction of the risk factors associated with suicidal behaviors than the unadjusted model. Moreover, the present study found higher prevalence rates of all suicidal behaviors among psychoactive substance users (i.e., alcohol, cannabis, illicit drugs, non-medical use of prescription drugs), and cigarette smoking was significantly associated with both suicide planning and suicide attempts (but not suicidal ideation). Previous research indicates that substance abuse can have a wide range of direct and indirect effects on both physical and mental health. As reported in a recent systematic review [38], there are significant associations between all types of substance use and suicidal behaviors. These effects often depend upon the drug specification, amount of use, frequency of use, personal health capabilities, and other factors. However, the present study did not consider these factors [38]. Therefore, further studies are needed to examine these specific relationships and factors between substance use and suicidality.

Strong relationships between physical illnesses and extreme mental health conditions (i.e., suicide and suicidality) are well-established [16]. Physical illnesses (e.g., high blood pressure, heart attacks, strokes, arthritis, chronic headaches, other chronic pain, respiratory conditions and bronchial asthma, diabetes, arthritis, hypothyroidism, etc.) can predispose individuals to mental illnesses by mediating abnormal and imbalanced secretions of neurotransmitters (e.g., serotonin, dopamine, norepinephrine, etc.) that make individuals more suicide-prone (even in the absence of any mental disorders; [16]), have also been reported in the Bangladeshi literature [18]. Individuals with mental disorders (with or without physical illnesses) are also at high risk of suicide-related behaviors and have been reported globally [16, 27]. In Bangladesh, recent retrospective studies reported that up to 60% of individuals with SI experience depression and other disorders such as schizophrenia, bipolar disorders, obsessive-compulsive disorder, generalized anxiety disorder, personality disorders, anxiety disorder, panic disorder, and conversion disorder [18, 39, 40]. Consistent with the prior studies, this study found a higher risk of suicidal behaviors of these participants with either mental health problems or physical illnesses.

Negative and traumatic life experiences such as criminal victimization, interpersonal violence (e.g., being raped, sexually molested, physically assaulted, physically abused as a child, seriously neglected as a child, threatened with a weapon, held captive or kidnapped), non-interpersonal violence (e.g., suffering great shock, life-threatening accidents, fire/flood/natural disasters, and witnessing bad injuries/deaths), domestic violence, childhood abuse and neglect, torture, sexual traumatization, natural disasters, and holocausts, are highly associated with suicidal behaviors and suicide contribution [21, 23], also the findings of the present study support this.

Table 4. Logistic regression analysis of the variables associated with suicide attempt.

Variables	Unadjusted model			Adjusted model (-2 Log likelihood = 438.198; Nagelkerke's R ² = 0.379)		
	Odds ratio (OR)	95% Confidence Interval (CI)	p-value	Adjusted odds ratio (AOR)	95% Confidence Interval (CI)	p-value
Gender						
Female	1.81	1.16–2.84	0.009	2.02	1.11–3.67	0.020
Male	Reference			Reference		
Year of study						
1 st year	0.26	0.13–0.51	<0.001	0.34	0.15–0.74	0.035
2 nd year	0.37	0.21–0.68		0.48	0.23–0.98	
3 rd year	0.36	0.20–0.67		0.50	0.24–1.03	
4 th year	Reference			Reference		
Permanent Residence						
Rural	1.29	0.66–2.54	0.448	1.85	0.80–4.28	0.147
Urban	Reference			Reference		
Cigarette smoker						
Yes	2.90	1.81–4.64	<0.001	1.86	0.89–3.92	0.099
No	Reference			Reference		
Psychoactive substance user						
Yes	10.44	5.65–19.30	<0.001	3.62	1.33–9.86	0.012
No	Reference			Reference		
Past-year physical illness						
Yes	6.08	3.78–9.78	<0.001	1.46	0.78–2.72	0.232
No	Reference			Reference		
Past-year mental illness						
Yes	20.74	12.84–33.50	<0.001	8.71	4.72–16.07	<0.001
No	Reference			Reference		
Any type of stressful life events during past-year						
Yes	7.87	4.66–13.30	<0.001	2.15	1.05–4.43	0.036
No	Reference			Reference		
Examination failure						
Yes	3.83	2.35–6.25	<0.001	1.30	0.70–2.44	0.400
No	Reference			Reference		
Relationship difficulties						
Yes	7.80	4.94–12.32	<0.001	1.67	0.87–3.21	0.121
No	Reference			Reference		
Campus ragging						
Yes	1.14	0.48–2.67	0.762	0.43	0.15–1.18	0.102
No	Reference		Reference			
Family problems						
Yes	7.00	4.40–11.12	<0.001	1.62	0.88–2.99	0.121
No	Reference			Reference		
Other problems						
Yes	0.89	0.11–6.68	0.913	1.47	0.17–12.73	0.725
No	Reference			Reference		
Family mental illness history						
Yes	4.78	2.99–7.66	<0.001	1.24	0.64–2.37	0.517
No	Reference			Reference		
Family suicide history						
Yes	3.23	1.33–7.83	0.009	0.69	0.23–2.08	0.521
No	Reference			Reference		

(Continued)

Table 4. (Continued)

Variables	Unadjusted model			Adjusted model (-2 Log likelihood = 438.198; Nagelkerke's R ² = 0.379)		
	Odds ratio (OR)	95% Confidence Interval (CI)	p-value	Adjusted odds ratio (AOR)	95% Confidence Interval (CI)	p-value
Family suicide attempt history						
Yes	5.83	3.30–10.29	<0.001	2.38	1.13–5.03	0.022
No	Reference			Reference		

<https://doi.org/10.1371/journal.pone.0262006.t004>

Several possible pathways between exposure to traumatic events have been suggested, including the mediating role of post-traumatic stress disorder (PTSD) symptoms, depression, psychiatric comorbidity, and dissociation, as well as the impact upon personality and cognitive development [19–21]. As consistent with the aforementioned literature, Bangladeshi studies also suggest that students' negative events (i.e., lack of proper accommodation, campus ragging, and political violence, etc.) mediate common psychological problems such as depression, anxiety, and stress [9], and these disorders contribute proximal suicide risk factors [22]. In the present study, stressful life-events (e.g., examination failure, relationship difficulties, campus ragging, family problems, etc.) were highly associated with all suicidal behaviors, although campus ragging did was not a risk factor for SA. In addition, experiencing a self-reported physical and mental illness were significantly associated with SI, SP and SA (except physical comorbidities) in the present study. This finding can be explained by the relationship between physical illness, mental illness, and suicidal behavior where physical co-morbidities can trigger psychiatric disorder alongside feelings of hopelessness or helplessness, a dramatic change in personality or appearance and/or irrational or bizarre behaviour. It has also been reported that psychiatric disorders are estimated to be responsible for a large proportion of suicides [41, 42].

This study also found the importance of the family history of mental illness, and suicide and suicidal behaviors in the association of all suicidal behaviors. As reported previously, both fatal and non-fatal suicidal behaviors of offspring are consistently associated with a history of affective and mood disorders, substance abuse, internal family conflicts, inappropriate parent-child relationships, history of suicide completion, and suicide attempts within the family [23–26]. Previous Bangladeshi findings are the same (e.g., 16.5% of individuals with SI had a family SA history [43]).

The present study has a number of limitations including (i) it being a cross-sectional study, (ii) assessing mental health illness and physical health illness using self-report, and (iii) a limited number of variables being examined and the omission of potentially important variables (e.g., family income, relationship status, childhood maltreatment, etc.). Moreover, assessing only a single university in Bangladesh limits the generalizability of the findings for other universities inside or outside of the country. Therefore, future (preferably longitudinal) research using countrywide representative student samples is needed to establish causal pathways between the variables examined in the present study. Despite these limitations, the study presented novel data concerning students' suicidal behaviors using a relatively large sample which will hopefully facilitate suicide prevention initiatives to be implemented by university authorities as well as further studies in the country.

5 Concluding remarks

Based on the present research (and elsewhere [9, 10]), campus-related issues such as ragging (among freshers) and examination failure (among final-year students) are prominent problems that should also be taken into account when developing suicide prevention programs on campus. However, other issues such as relationship complexities, family problems, and

psychoactive substance abuse also require consideration in such programs. Additionally, providing a student-friendly campus environment with appropriate psychological support (i.e., gatekeeper training, mental health support programs, etc.) is recommended based on the present findings.

Supporting information

S1 Data.

(SAV)

Acknowledgments

The authors also like to acknowledge that the project was run by the Undergraduate Research Organization, presently which will be introduced as the CHINTA Research Bangladesh. In addition, the authors would like to mention that, from this project, a paper was published with distinct aims and objectives that does not overlap with the present study [44].

Author Contributions

Conceptualization: Mohammed A. Mamun.

Data curation: M. Rasheduzzaman.

Formal analysis: M. Rasheduzzaman, Firoj al-Mamun, Ismail Hosen, Tahmina Akter, Mohammed A. Mamun.

Investigation: M. Rasheduzzaman, Moazzem Hossain.

Methodology: M. Rasheduzzaman, Firoj al-Mamun, Ismail Hosen, Tahmina Akter, Moazzem Hossain, Mark D. Griffiths, Mohammed A. Mamun.

Project administration: M. Rasheduzzaman, Moazzem Hossain, Mohammed A. Mamun.

Resources: M. Rasheduzzaman, Moazzem Hossain, Mark D. Griffiths, Mohammed A. Mamun.

Supervision: Moazzem Hossain, Mark D. Griffiths, Mohammed A. Mamun.

Validation: M. Rasheduzzaman, Firoj al-Mamun, Ismail Hosen, Tahmina Akter, Moazzem Hossain, Mark D. Griffiths, Mohammed A. Mamun.

Visualization: M. Rasheduzzaman, Firoj al-Mamun, Ismail Hosen, Tahmina Akter, Moazzem Hossain, Mark D. Griffiths, Mohammed A. Mamun.

Writing – original draft: M. Rasheduzzaman, Mohammed A. Mamun.

Writing – review & editing: Firoj al-Mamun, Ismail Hosen, Tahmina Akter, Mark D. Griffiths, Mohammed A. Mamun.

References

1. Castle K, Kreipe R. Suicidal Behavior. In: Garfunkel LC, Kaczorowski JM, Christy C, editors. *Pediatric Clinical Advisor (Second Edition)*. Second Edition. Philadelphia: Mosby; 2007. pp. 544–545. <https://doi.org/10.1016/B978-032303506-4.10315-3>
2. Mortier P, Cuijpers P, Kiekens G, Auerbach RP, Demyttenaere K, Green JG, et al. The prevalence of suicidal thoughts and behaviours among college students: a meta-analysis. *Psychol Med*. Cambridge University Press; 2018; 48: 554–565. <https://doi.org/10.1017/S0033291717002215> PMID: 28805169
3. World Health Organization. Suicide—key facts [Internet]. 2019 [cited 10 Nov 2019]. Available: <https://www.who.int/news-room/fact-sheets/detail/suicide>.

4. Shah MMA, Ahmed S, Arafat SMY. Demography and risk factors of suicide in Bangladesh: a six-month paper content analysis. *Psychiatry J. Hindawi*; 2017; 2017: e3047025. <https://doi.org/10.1155/2017/3047025> PMID: 29130035
5. Arafat SMY, Mamun MAA. Repeated suicides in the University of Dhaka (November 2018): Strategies to identify risky individuals. *Asian J Psychiatr. Elsevier*; 2019; 39: 84–85. <https://doi.org/10.1016/j.ajp.2018.12.014> PMID: 30594879
6. Mamun MA, Misti JM, Griffiths MD. Suicide of Bangladeshi medical students: Risk factor trends based on Bangladeshi press reports. *Asian J Psychiatr. Elsevier*; 2020; 48: 101905. <https://doi.org/10.1016/j.ajp.2019.101905> PMID: 31865199
7. Mamun MA, Siddique AB, Sikder MT, Griffiths MD. Student suicide risk and gender: A retrospective study from Bangladeshi press reports. *Int J Ment Health Addict. Springer*; 2020; Epub ahead of print. <https://doi.org/10.1007/s11469-020-00267-3>
8. Rafi MA, Mamun MA, Hsan K, Hossain M, Gozal D. Psychological implications of unemployment among Bangladesh Civil Service job seekers: a pilot study. *Front Psychiatry. Frontiers Media SA*; 2019; 10: 578. <https://doi.org/10.3389/fpsy.2019.00578> PMID: 31456705
9. Mamun MA, Hossain MS, Griffiths MD. Mental health problems and associated predictors among Bangladeshi students. *Int J Ment Health Addict. Springer*; 2019; Epub ahead of print. <https://doi.org/10.1007/s11469-019-00144-8>
10. Bhuiyan MAH, Griffiths MD, Mamun MA. Depression literacy among Bangladeshi pre-university students: Differences based on gender, educational attainment, depression, and anxiety. *Asian J Psychiatr. 2020*; 50: 101944. <https://doi.org/10.1016/j.ajp.2020.101944> PMID: 32106072
11. Sakib N, Islam M, Al Habib MS, et al. Depression and suicidality among Bangladeshi students: Subject selection reasons and learning environment as potential risk factors. *Perspect Psychiatr Care. Wiley Online Library*; 2021; 57(3):1150–1162. <https://doi.org/10.1111/ppc.12670> PMID: 33135191
12. Faruk MO, Mamun MA, Siddique AB, Griffiths MD. Risk factors for depression and anxiety disorders among Bangladeshi dental students: a cross-sectional survey study. *Int J Ment Health Addict. Springer*; 2021; Epub ahead of print. <https://doi.org/10.1007/s11469-021-00603-1>
13. Milner A, Witt K, LaMontagne AD, Niedhammer I. Psychosocial job stressors and suicidality: a meta-analysis and systematic review. *Occup Environ Med. BMJ Publishing Group Ltd*; 2018; 75: 245–253. <https://doi.org/10.1136/oemed-2017-104531> PMID: 28851757
14. Yozwiak JA, Lentzsch-Parcells CM, Zapolski TCB. Suicide and suicidal ideation among college students. *Int J Disabil Hum Dev. Walter de Gruyter*; 2012; 11: 185–189. <https://doi.org/10.1515/ijdh-2012-0042>
15. Mamun MA, Griffiths MD. A rare case of Bangladeshi student suicide by gunshot due to unusual multiple causalities. *Asian J Psychiatr. Elsevier*; 2020; 49: 101951. <https://doi.org/10.1016/j.ajp.2020.101951> PMID: 32078949
16. Scott KM, Hwang I, Chiu WT, Kessler RC, Sampson NA, Angermeyer M, et al. Chronic physical conditions and their association with first onset of suicidal behavior in the world mental health surveys. *Psychosom Med. American Psychosomatic Society*; 2010; 72: 712–719. <https://doi.org/10.1097/PSY.0b013e3181e3333d> PMID: 20498290
17. Juurlink DN, Herrmann N, Szalai JP, Kopp A, Redelmeier DA. Medical illness and the risk of suicide in the elderly. *Arch Intern Med. American Medical Association*; 2004; 164: 1179–1184. <https://doi.org/10.1001/archinte.164.11.1179> PMID: 15197042
18. Ahmad M, Hossain MZ. Hanging as a method of suicide: Retrospective analysis of postmortem cases. *J Armed Forces Med Coll. Armed Forces Medical College Bangladesh*; 2010; 6: 37–39. <https://doi.org/10.3329/jafmc.v6i2.7273>
19. Panagioti M, Gooding P, Tarrier N. Post-traumatic stress disorder and suicidal behavior: A narrative review. *Clin Psychol Rev. Elsevier*; 2009; 29: 471–482. <https://doi.org/10.1016/j.cpr.2009.05.001> PMID: 19539412
20. Panagioti M, Gooding PA, Tarrier N. A meta-analysis of the association between posttraumatic stress disorder and suicidality: the role of comorbid depression. *Compr Psychiatry. Elsevier*; 2012; 53: 915–930. <https://doi.org/10.1016/j.comppsy.2012.02.009> PMID: 22483367
21. Belik SL, Cox BJ, Stein MB, Asmundson GJG, Sareen J. Traumatic events and suicidal behavior: Results from a national mental health survey. *J Nerv Ment Dis. Wolters Kluwer*; 2007; 195: 342–349. <https://doi.org/10.1097/01.nmd.0b013e318060a869> PMID: 17435485
22. Mamun MA, Griffiths MD. PTSD-related suicide six years after the Rana Plaza collapse in Bangladesh. *Psychiatry Res. Elsevier*; 2020; 287: 112645. <https://doi.org/10.1016/j.psychres.2019.112645> PMID: 31685284
23. Carballo JJ, Harkavy-friedman J, Burke AK, Sher L, Baca-garcia E, Sullivan GM, et al. Family history of suicidal behavior and early traumatic experiences: Additive effect on suicidality and course of bipolar

- illness? *J Affect Disord.* Elsevier; 2008; 109: 57–63. <https://doi.org/10.1016/j.jad.2007.12.225> PMID: 18221790
24. Wagner BM, Silverman MAC, Martin CE. Family factors in youth suicidal behaviors. *Am Behav Sci.* Sage Publications; 2003; 46: 1171–1191. <https://doi.org/10.1177/0002764202250661>
 25. Goodwin RD, Beautrais AL, Fergusson DM. Familial transmission of suicidal ideation and suicide attempts: evidence from a general population sample. *Psychiatry Res.* Elsevier; 2004; 126: 159–165. <https://doi.org/10.1016/j.psychres.2004.02.010> PMID: 15123395
 26. Wang Y, Chen S, Xu Z, Shen Z, Wang Y, He X, et al. Family history of suicide and high motor impulsivity distinguish suicide attempters from suicide ideators among college students. *J Psychiatr Res.* Elsevier; 2017; 90: 21–25. <https://doi.org/10.1016/j.jpsychires.2017.02.006> PMID: 28214643
 27. Hubers AAM, Moaddine S, Peersmann SHM, Stijnen T, Van Duijn E, Van der Mast RC, et al. Suicidal ideation and subsequent completed suicide in both psychiatric and non-psychiatric populations: a meta-analysis. *Epidemiol Psychiatr Sci.* Cambridge University Press; 2018; 27: 186–198. <https://doi.org/10.1017/S2045796016001049> PMID: 27989254
 28. Mashreky SR, Rahman F, Rahman A. Suicide kills more than 10,000 people every year in Bangladesh. *Arch Suicide Res.* Taylor & Francis; 2013; 17: 387–396. <https://doi.org/10.1080/13811118.2013.801809> PMID: 24224672
 29. Hossain S, Anjum A, Hasan MT, Uddin ME, Hossain MS, Sikder MT. Self-perception of physical health conditions and its association with depression and anxiety among Bangladeshi university students. *J Affect Disord.* Elsevier; 2020; 263: 282–288. <https://doi.org/10.1016/j.jad.2019.11.153> PMID: 31818790
 30. McKinnon B, Gariépy G, Sentenac M, Elgar FJ. Adolescent suicidal behaviours in 32 low-and middle-income countries. *Bull World Health Organ.* World Health Organization; 2016; 94: 340–350F. <https://doi.org/10.2471/BLT.15.163295> PMID: 27147764
 31. Asante KO, Kugbey N, Osafo J, Quarshie EN-B, Sarfo JO. The prevalence and correlates of suicidal behaviours (ideation, plan and attempt) among adolescents in senior high schools in Ghana. *SSM-population Heal.* Elsevier; 2017; 3: 427–434. <https://doi.org/10.1016/j.ssmph.2017.05.005> PMID: 29349236
 32. Jahan S, Araf K, Gozal D, Griffiths MD, Mamun MA. Depression and suicidal behaviors among Bangladeshi mothers of children with Autistic Spectrum Disorder: a comparative study. *Asian J Psychiatr.* Elsevier; 2020; 51: 101994. <https://doi.org/10.1016/j.ajp.2020.101994> PMID: 32146143
 33. Mamun MA, Misti JM, Hosen I, Al Mamun F. Suicidal behaviors and university entrance test-related factors: A Bangladeshi exploratory study. *Perspect Psychiatr Care.* Wiley Online Library; 2021; Epub ahead of print. <https://doi.org/10.1111/ppc.12783> PMID: 33834493
 34. Mortier P, Auerbach RP, Alonso J, Bantjes J, Benjet C, Cuijpers P, et al. Suicidal Thoughts and Behaviors Among First-Year College Students: Results From the WMH-ICS Project. *J Am Acad Child Adolesc Psychiatry.* American Academy of Child & Adolescent Psychiatry; 2018; 57: 263-273.e1. <https://doi.org/10.1016/j.jaac.2018.01.018> PMID: 29588052
 35. Mendez-Bustos P, Lopez-Castroman J, Baca-García E, Ceverino A. Life cycle and suicidal behavior among women. *Sci World J.* Hindawi; 2013; 2013: e485851. <https://doi.org/10.1155/2013/485851> PMID: 23533350
 36. Oner S, Yenilmez C, Ozdamar K. Sex-related differences in methods of and reasons for suicide in Turkey between 1990 and 2010. *J Int Med Res.* SAGE Publications Sage UK: London, England; 2015; 43: 483–493. <https://doi.org/10.1177/0300060514562056> PMID: 25995209
 37. Oner S, Yenilmez C, Ayranci U, Gunay Y, Ozdamar K. Sexual differences in the completed suicides in Turkey. *Eur Psychiatry.* Elsevier; 2007; 22: 223–228. <https://doi.org/10.1016/j.eurpsy.2007.01.002> PMID: 17344031
 38. Breet E, Goldstone D, Bantjes J. Substance use and suicidal ideation and behaviour in low-and middle-income countries: a systematic review. *BMC Public Health.* BioMed Central; 2018; 18: 549. <https://doi.org/10.1186/s12889-018-5425-6> PMID: 29699529
 39. Shah MMA, Sajib MWH, Arafat SMY. Demography and risk factor of suicidal behavior in Bangladesh: A cross-sectional observation from patients attending a suicide prevention clinic of Bangladesh. *Asian J Psychiatr.* Elsevier; 2018; 35: 4–5. <https://doi.org/10.1016/j.ajp.2018.04.035> PMID: 29715650
 40. Feroz AHM, Islam SMN, Reza S, Rahman AKMM, Sen J, Mowla M, et al. A community survey on the prevalence of suicidal attempts and deaths in a selected rural area of Bangladesh. *J Med.* 2012; 13: 3–9. <https://doi.org/10.3329/jom.v13i1.10042>
 41. Fegg M, Kraus S, Graw M, Bausewein C. Physical compared to mental diseases as reasons for committing suicide: a retrospective study. *BMC Palliat Care.* BioMed Central; 2016; 15: 14. <https://doi.org/10.1186/s12904-016-0088-5> PMID: 26860949

42. MacLean J, Kinley DJ, Jacobi F, Bolton JM, Sareen J. The relationship between physical conditions and suicidal behavior among those with mood disorders. *J Affect Disord.* Elsevier; 2011; 130: 245–250. <https://doi.org/10.1016/j.jad.2010.10.028> PMID: 21078525
43. Arafat SMY, Akter H, Mali B. Psychiatric morbidities and risk factors of suicidal ideation among patients attending for psychiatric services at a tertiary teaching hospital in Bangladesh. *Asian J Psychiatr.* Elsevier; 2018; 34: 44–46. <https://doi.org/10.1016/j.ajp.2018.04.020> PMID: 29635222
44. Rasheduzzaman M, al Mamun F, Faruk MO, Hosen I, Mamun MA. Depression in Bangladeshi university students: The role of sociodemographic, personal, and familial psychopathological factors. *Perspect Psychiatr Care.* Wiley Online Library; 2021; 57(4): 1585–1594. <https://doi.org/10.1111/ppc.12722> PMID: 33442872