



Published in final edited form as:

Stress Health. 2019 October ; 35(4): 542–548. doi:10.1002/smi.2887.

Understanding mind–body disciplines: A pilot study of paced breathing and dynamic muscle contraction on autonomic nervous system reactivity

Michael S. Chin^{1,2}, Stefanos N. Kales^{3,4}

¹Division of General Internal Medicine and Public Health, Vanderbilt University School of Medicine, Nashville, Tennessee

²Vanderbilt Occupational Health, Vanderbilt University Medical Center, Nashville, Tennessee

³Environmental and Occupational Medicine and Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts

⁴Occupational Medicine, Cambridge Health Alliance, Harvard Medical School, Cambridge, Massachusetts

Abstract

Mind–body disciplines such as yoga, Tai Chi, and Qigong have been demonstrated to activate the parasympathetic nervous system, but it remains unclear how these practices achieve these results, whether by breathing, movement, or some combination. This pilot study establishes a model to examine the individual and combined effects of paced breathing and rhythmic skeletal muscle contraction on the activation of the parasympathetic system during a cognitive stressor. Male participants were randomly assigned to one of four preconditioning groups: (a) paced breathing alone, (b) alternating upper extremity muscle contractions, (c) paced breathing synchronized with alternating contractions, or (d) a neutral control task. Autonomic response was assessed by heart rate variability during a standardized cognitive stressor. The alternating contraction group had 71.7% higher activation of parasympathetic signal over respiration alone ($p < .001$). Alternating contractions synchronized with breathing demonstrated 150% higher parasympathetic activation than control ($p < .0001$). Comparing the contraction alone and synchronized groups, the synchronized group demonstrated 45.9% higher parasympathetic response during a cognitive stressor ($p < .001$). In conclusion, paced breathing synchronized with rhythmic muscle contraction leads to more resilient activation of the parasympathetic response than either alternating contractions or breathing alone, which may help explain the stress reducing benefits of mind–body disciplines.

Correspondence: Michael S. Chin, MD, MPH, Vanderbilt Occupational Health, 1211 21st Avenue South, Suite 640, Nashville, TN 37212. mchin@mail.harvard.edu.

CONFLICT OF INTEREST

The authors have declared that they have no conflict of interest.

Keywords

biological mechanisms of stress; cardiovascular reactivity; psychophysiology; respite/recovery resilience; stress; mindfulness; stress management

1 | INTRODUCTION

Recent focus on stress reduction and physiological resiliency has led to further interest in the autonomic nervous system's adaptation to stress. Many popular mind–body practices such as yoga, Tai Chi, and Qigong have been shown to activate the parasympathetic nervous system (Goyal et al., 2014; Sullivan et al., 2018; Walther, Lacker, & Ehlert, 2018). Paced respiration and dynamic tension through rhythmic skeletal muscle contraction are two core components common to yoga, Tai Chi, and Qigong; however, it remains unclear by which mechanisms these training disciplines activate the parasympathetic nervous system. Therefore, without understanding the interaction between breathing and movement, it is challenging to optimize the benefits of these practices. This pilot study establishes a model to examine the individual and combined effects of paced breathing and rhythmic skeletal muscle contraction.

Paced breathing is a technique with a proposed mechanism of action that is mediated through the activation of the body's parasympathetic nervous system through entrainment effects between respiratory rate and heart rate (Jerath, Edry, Barnes, & Jerath, 2006; Nijjar et al., 2014). Although the exact respiratory frequency may vary slightly from person to person, studies have demonstrated that the resonant breathing frequency is around 0.1 Hz, or one breath cycle every 10 s (Lehrer, Vaschillo, & Vaschillo, 2000; Lin, Tai, & Fan, 2014).

Less recognized and understood, concurrent evidence also suggests that voluntary rhythmic muscle contraction alone could be an alternative way of stimulating the parasympathetic nervous system. In a study on rhythmic contraction alone, Lehrer et al. demonstrated that when large muscle groups alternated rhythmically between contraction and relaxation, subjects experienced a similar parasympathetic entrainment effect observed in the earlier paced breathing studies (Lehrer, Vaschillo, Trost, & France, 2009; Vaschillo, Vaschillo, Pandina, & Bates, 2011). However, the autonomic effects of performing rhythmic synchronized breathing and muscle contraction tasks simultaneously have not been studied systematically. Understanding the synergistic effect between these components may help to explain the observed benefits many practitioners experience after various types of mind–body training.

Additionally, practice of mind–body disciplines has been demonstrated to improve cognitive function. Three meta-analyses have shown that training in yoga, Tai-chi, and Qigong led to improved cognitive function in older populations (Chan, Deng, Wu, & Yan, 2019; Wayne et al., 2014; Wu et al., 2019). However, the mechanism of how these practices achieve these effects has not been established. Considering that these disciplines are partly meditative arts, short-term studies have suggested that meditation can have near immediate positive effects on task performance (Chan, Immink, & Lushington, 2017; Chan, Lushington, & Immink, 2018; Colzato, Sellaro, Samara, & Hommel, 2015; Colzato, van der Wel, Sellaro, & Hommel, 2016).

Using established heart rate variability (HRV) measures to determine the autonomic state, the first aim of this study will address whether entrainment with rhythmic muscle contraction synchronized with respiration at a resonant frequency can produce greater activation of the parasympathetic system more effectively than either technique alone when challenged with a cognitively induced sympathetic state. Second, this study will evaluate whether the type of preconditioning affects task performance.

2 | METHODS

2.1 | Participants

Because HRV changes have been shown to vary differently between males and females due to hormonal differences, only males were recruited for this pilot study to minimize variation (Koenig & Thayer, 2016; Sato & Miyake, 2004). Forty-eight healthy male participants, ages 18–55, were recruited and consented from Harvard T.H. Chan School of Public Health- and Harvard Medical School-affiliated student programs, fellowships, and training residencies. An initial telephone screening interview was performed for all participants. Individuals with any history of restrictive or obstructive lung disease, hypertension, or taking any medications that could alter blood pressure were excluded. Although participants using prescription stimulants were excluded from the study, caffeine consumption was not specifically restricted as it has been shown that withdrawal effects on HRV may exist for habitual caffeine users (Zimmermann-Viehoff et al., 2016). The study protocol was reviewed and approved by the IRB of the Harvard T.H. Chan School of Public Health.

The participants were randomized prior to the testing session to one of four preconditioning groups using the online Research Randomizer tool (www.randomizer.com). Preconditioning groups were (a) diaphragmatic breathing at 0.1 Hz (inhale nose 5 s, exhale mouth 5 s) for 5 min, (b) rhythmic contraction and relaxation of arm muscles by grasping a tennis ball at 0.1 Hz for 5 min (alternating contractions in left and right arms every 5 s), (c) performing synchronized diaphragmatic breathing and contractions at 0.1 Hz for 5 min, and (d) a control group assigned to read four adapted articles from Scientific American that were rated as emotionally neutral for 5 min (van den Broek, Lorch, Linderholm, & Gustafson, 2001).

2.2 | Testing procedure

All testing occurred over a 30-min session with only the experimenter and participant present in a quiet room between the daytime hours of 09:00 and 16:00. After written informed consent was obtained, a Polar H7 heart rate monitor (Polar Electro Oy, Kempele, Finland) was then placed at the participant's mid-chest level. The Polar chest strap systems have already been validated as a reliable research device measuring heart rate with accuracy comparable with electrocardiograms used in clinical medicine by measuring R-R intervals at a sampling rate of 1000 Hz (Barbosa, da Silva, de Azevedo, Pastre, & Vanderlei, 2016; Giles, Draper, & Neil, 2016). The R-R interval is the beat-to-beat interval as measured between peaks of consecutive QRS complexes on electrocardiogram. A recording application on iPad (Apple Inc., Cupertino, CA, United States) recorded the R-R intervals from the chest strap, which were analysed offline.

Participants were instructed to sit upright quietly for 5 min while reading the study instructions, and then, baseline heart rate, cuff blood pressure, and respiration rate were obtained. A 5-min period of the assigned preconditioning task was then performed with a graphical timer application on iPad was used to visually cue the initiation and completion of each cycle of breathing or contraction, depending on the group assignment.

Immediately after the preconditioning task, a computerized version of the Stroop test (<http://cognitivedfun.net>) was then run for 5 min to provoke a sympathetic response. The Stroop test has been demonstrated to produce a mild sympathetic response measured by HRV through dissonant executive task function (Salahuddin, Cho, Jeong, & Kim, 2007; Visnovcova et al., 2014). The program automatically performed the test by presenting words written in colour text, and participants were asked to indicate the colour of the word (and not its meaning) by key stroke as fast as possible while minimizing errors. A reaction time for each word pair was recorded by the computer program. The premise of the Stroop test is that incongruent pairs have longer reaction times when compared with congruent pairs (Dyer, 1973). As a marker for performance, a reaction time gap was calculated for each participant from the difference between congruent and incongruent pair reaction times.

For validation of Stroop engagement across groups, at the end of this task period, a 5-min questionnaire (Short Flow State Scale) was then administered to assess degree of task immersion for all groups (Jackson, Martin, & Eklund, 2008). Flow state is the degree of perceived immersion in the task. The Short Flow State Scale has been validated for evaluation of performance engagement. Respiration rates were monitored during all phases of testing to ensure they were within the 9–24 cycles per minute range required for high frequency (HF) to correspond accurately to vagal tone (Laborde, Mosley, & Thayer, 2017).

2.3 | HRV analysis

High frequency (HF) and low frequency (LF) HRV components are reflective of parasympathetic and sympathetic activation, respectively. As a corollary, the ratio LF/HF is generally regarded as the overall sympathovagal balance and degree of autonomic excitement (Shaffer & Ginsberg, 2017). Recorded R-R intervals from the Polar H7 chest strap were downloaded and analysed using Kubios HRV Premium software (Kubios Oy, Kuopio, Finland). The software converted the R-R intervals into frequency domain indices: LF power (ms^2), HF power (ms^2), and LF/HF ratio. Each of the indices was calculated over 2-min intervals based on recommendations from published standards. It is generally accepted that 1 min is needed to assess the HF component of HRV whereas approximately 2 min are needed to address the LF component (Laborde et al., 2017).

For each participant, the baseline and Stroop HRV recordings were processed by Kubios HRV Premium. Automated artifact correction was performed for all recordings prior to analysis. One hundred twenty-second sampling periods were utilized to derive LF, HF, and LF/HF using fast Fourier transformation spectrum method (Figure 1). LF and HF bands were standardly defined as 0.04–0.15 and 0.15–0.4 Hz, respectively, and absolute power for each band was analysed in normalized units, LF or HF divided by total power (Malliani, Lombardi, & Pagani, 1994). The Stroop HRV measurements were normalized to each

participant's baseline. Normalized Stroop HRV indices for each preconditioning group were compared by one-way ANOVA with Tukey's post-hoc test.

3 | RESULTS

All enrolled participants ($n = 48$) completed the study sessions without any adverse events. Participants' average age, resting blood pressure, heart and spontaneous respiration rates are summarized in Table 1. As a result of randomization, 12 subjects were assigned to the control group, and 12 subjects were assigned to each preconditioning group.

3.1 | Parasympathetic response (HF)

ANOVA for HF demonstrated significant changes between groups ($F(3,44) = 32.85$, $p < .0001$; Figure 2a). During administration of the Stroop test, there was no significant difference between breathing alone group and reading control. The alternating contraction group had 71.7% higher activation of HF over respiration alone ($p < .001$). Alternating contractions synchronized with breathing demonstrated 150% higher HF activation than control ($p < .0001$). Between contraction alone and combined contraction groups, the combined group demonstrated 45.9% higher HF response ($p < .001$).

3.2 | Sympathetic response (LF)

ANOVA for LF demonstrated significant changes between groups ($F(3,44) = 8.258$, $p < .001$; Figure 2b). Sympathetic activation during administration of the Stroop test was not different between the breathing alone group and reading control. The alternating the contraction group and synchronized groups had 33.5% and 45.2% lower LF HRV activation than control, respectively ($p < .01$ and $p < .0001$). There was no significant difference in LF response between the contraction alone and synchronized groups.

3.3 | Sympathovagal response (LF/HF)

ANOVA for LF/HF demonstrated significant changes between groups ($F(3,44) = 10.62$, $p < .0001$; Figure 2c). Sympathovagal balance results mirrored the sympathetic activation. There was no significant LF/HF ratio difference between the breathing alone group and the reading controls. Similar to LF, both contraction groups had significantly lower LF/HF over control, respectively ($p < .01$ and $p < .001$). Again, similar to LF, there was no significant difference in LF/HF response between the contraction alone and synchronized groups.

3.4 | Cognitive performance and engagement

There were no differences for Stroop test reaction time among groups ($F(3,44) = 1.359$, $p > .05$). Similarly, there were no significant differences between Short State Flow State scores among the four preconditioning groups ($F(3,44) = 1.172$, $p > .05$; data not shown).

4 | DISCUSSION

Despite the widespread adoption of various mind-body practices such as yoga, Tai Chi, and Qigong, there still lacks consensus regarding how these disciplines produce their powerful benefits (Riley & Park, 2015). The Respiratory Vagal Stimulation Model hypothesizes

that these disciplines confer benefits through attentively regulated breathing (Gerritsen & Band, 2018). However, this theory ignores another key component of these mind–body disciplines, muscle contraction. Our pilot study provides a simplified model to study the combined effects between two fundamental, common elements of these practices: respiration and muscle contraction. Our results suggest that cyclic respiration synchronized with alternating muscle contraction may be one underlying phenomenon that confers increased parasympathetic balance.

Contrary to the common preconception that breathing practices lead to stress reduction, it is interesting to note that priming with paced respiration alone did not confer lasting increased HRV reactivity during the stress event. Although current research has commonly associated mindful breathing practices with reduction in stress, these results are usually after weeks of training and may be due to post-event stress recovery rather than pre-event priming (Goyal et al., 2014; Wolever et al., 2012). Our results suggest that there may be a distinction between pre-event priming for a stressful event and stress mitigation, wherein entrainment of the autonomic system with synchronized respiration and contraction lead to a more resilient response than either component alone.

The powerful priming effect observed when combining cyclic respiration with muscle contraction supports the “bottom-up” polyvagal theory proposed by Stephen Porges (Sullivan et al., 2018). This theory hypothesizes that autonomic regulation is possible through interoception (i.e., an awareness of the internal state of the body’s systems) and self-regulatory skills (Gard, Noggle, Park, Vago, & Wilson, 2014). The finding that entrainment with breathing synchronized rhythmic muscle contraction activates the parasympathetic relaxation response more effectively than either one alone could be explained by this process of enhanced interoception. Indeed, movement disciplines, such as the Russian martial art Systema, may have empirically developed an optimized priming process through highly coordinated rhythmic movement and respiration exercises (Vasiliev, Meredith, & Ryabko, 2006).

When assessing level of task immersion through the flow questionnaire, we observed that there were no differences in level of task engagement among all four groups. This is not unexpected as engagement is dependent on many individual factors such as level of proficiency, personal interest, as well as personality characteristics (Keller & Bless, 2008).

4.1 | Study limitations

Our pilot study represents a relatively small sample. Studies have demonstrated that estrogen levels may affect HRV during the stress response (Koenig & Thayer, 2016; Sato & Miyake, 2004; Thayer, Ahs, Fredrikson, Sollers, & Wager, 2012). Due to these concerns, we restricted our participant enrollment to male participants to minimize hormonal variability, and therefore, we may have limited generalizability to females. Future studies should validate our findings in a female population.

4.2 | Future directions

Although no differences in flow engagement were found between the preconditioning groups, we propose that future studies should examine how these synchronized respiration

and contraction patterns may enhance cognitive performance under stress by varying the autonomic response. These findings may be highly relevant for individuals who need to make quick and accurate decisions in high stress environments. Real-time management of stress could lead improved performance under pressure as suggested by a large-scale study of emergency medical responders, where increased perceived anxiety was correlated with patient safety events and compromised decision making (Guise et al., 2017).

5 | PRACTICAL IMPLICATIONS

- Alternating isometric muscle contraction can help to prepare for a stressful event.
- Synchronizing isometric muscle contraction with paced breathing leads to an even more resilient relaxation response in preparation for a stressful event.
- Using rhythmic breathing alone before a stressful event does not lead to a lasting relaxation response.

6 | CONCLUSIONS

Entrainment using paced breathing synchronized with rhythmic skeletal muscle contraction at 0.1 Hz produces a more resilient parasympathetic response than either breathing or muscle contraction alone during a stressful task. Preconditioning type does not appear to lead to significant differences in reaction time performance or task flow immersion.

ACKNOWLEDGMENTS

MC was partly supported by a training grant from the Harvard Education and Research Center for Occupational Safety and Health (T42 OH008416). This study was funded by an award from the Harvard Chan-NIEHS Center for Environmental Health (NIEHS Grant P30 ES000002). The authors would like to thank Ms. Ann Backus for her assistance with study participant recruitment. MC would like to recognize the mentorship of Mr. Vladimir Vasiliev, whose teachings on Systema breathwork and Russian martial arts served as the original inspiration for this study.

REFERENCES

- Barbosa MP, da Silva NT, de Azevedo FM, Pastre CM, & Vanderlei LC (2016). Comparison of Polar(R) RS800G3 heart rate monitor with Polar(R) S810i and electrocardiogram to obtain the series of RR intervals and analysis of heart rate variability at rest. *Clinical Physiology and Functional Imaging*, 36(2), 112–117. 10.1111/cpf.12203 [PubMed: 25348547]
- Chan JSY, Deng K, Wu J, & Yan JH (2019). Effects of meditation and mind-body exercises on older adults' cognitive performance: A meta-analysis. *The Gerontologist*. 10.1093/geront/gnz022
- Chan RW, Immink MA, & Lushington K (2017). The influence of focused-attention meditation states on the cognitive control of sequence learning. *Consciousness and Cognition*, 55, 11–25. 10.1016/j.concog.2017.07.004 [PubMed: 28750361]
- Chan RW, Lushington K, & Immink MA (2018). States of focused attention and sequential action: A comparison of single session meditation and computerised attention task influences on top-down control during sequence learning. *Acta Psychologica*, 191, 87–100. 10.1016/j.actpsy.2018.09.003 [PubMed: 30240891]
- Colzato LS, Sellaro R, Samara I, & Hommel B (2015). Meditation-induced cognitive-control states regulate response-conflict adaptation: Evidence from trial-to-trial adjustments in the Simon task. *Consciousness and Cognition*, 35, 110–114. 10.1016/j.concog.2015.04.012 [PubMed: 25982057]
- Colzato LS, van der Wel P, Sellaro R, & Hommel B (2016). A single bout of meditation biases cognitive control but not attentional focusing: Evidence from the global-local task. *Consciousness and Cognition*, 39, 1–7. 10.1016/j.concog.2015.11.003 [PubMed: 26637968]

- Dyer FN (1973). The Stroop phenomenon and its use in the study of perceptual, cognitive, and response processes. *Memory & Cognition*, 1(2), 106–120. 10.3758/bf03198078 [PubMed: 24214501]
- Gard T, Noggle JJ, Park CL, Vago DR, & Wilson A (2014). Potential self-regulatory mechanisms of yoga for psychological health. *Frontiers in Human Neuroscience*, 8, 770. 10.3389/fnhum.2014.00770 [PubMed: 25368562]
- Gerritsen RJS, & Band GPH (2018). Breath of life: The respiratory vagal stimulation model of contemplative activity. *Frontiers in Human Neuroscience*, 12, 397. 10.3389/fnhum.2018.00397 [PubMed: 30356789]
- Giles D, Draper N, & Neil W (2016). Validity of the Polar V800 heart rate monitor to measure RR intervals at rest. *European Journal of Applied Physiology*, 116(3), 563–571. 10.1007/s00421-015-3303-9 [PubMed: 26708360]
- Goyal M, Singh S, Sibinga EM, Gould NF, Rowland-Seymour A, Sharma R, ... Haythornthwaite JA (2014). Meditation programs for psychological stress and well-being: A systematic review and meta-analysis. *JAMA Internal Medicine*, 174(3), 357–368. 10.1001/jamainternmed.2013.13018 [PubMed: 24395196]
- Guise JM, Hansen M, O'Brien K, Dickinson C, Meckler G, Engle P, ... Jui J (2017). Emergency medical services responders' perceptions of the effect of stress and anxiety on patient safety in the out-of-hospital emergency care of children: A qualitative study. *BMJ Open*, 7 (2), e014057. 10.1136/bmjopen-2016-014057
- Jackson SA, Martin AJ, & Eklund RC (2008). Long and short measures of flow: The construct validity of the FSS-2, DFS-2, and new brief counterparts. *Journal of Sport & Exercise Psychology*, 30(5), 561–587. 10.1123/jsep.30.5.561 [PubMed: 18971512]
- Jerath R, Edry JW, Barnes VA, & Jerath V (2006). Physiology of long pranayamic breathing: Neural respiratory elements may provide a mechanism that explains how slow deep breathing shifts the autonomic nervous system. *Medical Hypotheses*, 67(3), 566–571. 10.1016/j.mehy.2006.02.042 [PubMed: 16624497]
- Keller J, & Bless H (2008). Flow and regulatory compatibility: An experimental approach to the flow model of intrinsic motivation. *Personality and Social Psychology Bulletin*, 34(2), 196–209. 10.1177/0146167207310026 [PubMed: 18212330]
- Koenig J, & Thayer JF (2016). Sex differences in healthy human heart rate variability: A meta-analysis. *Neuroscience and Biobehavioral Reviews*, 64, 288–310. 10.1016/j.neubiorev.2016.03.007 [PubMed: 26964804]
- Laborde S, Mosley E, & Thayer JF (2017). Heart Rate Variability and Cardiac Vagal Tone in Psychophysiological Research – Recommendations for Experiment Planning, Data Analysis, and Data Reporting. *Frontiers in Psychology*, 8, 213. 10.3389/fpsyg.2017.00213 [PubMed: 28265249]
- Lehrer P, Vaschillo E, Trost Z, & France CR (2009). Effects of rhythmical muscle tension at 0.1Hz on cardiovascular resonance and the baroreflex. *Biological Psychology*, 81(1), 24–30. 10.1016/j.biopsycho.2009.01.003 [PubMed: 19428965]
- Lehrer PM, Vaschillo E, & Vaschillo B (2000). Resonant frequency biofeedback training to increase cardiac variability: Rationale and manual for training. *Applied Psychophysiology and Biofeedback*, 25(3), 177–191. 10.1023/A:1009554825745 [PubMed: 10999236]
- Lin IM, Tai LY, & Fan SY (2014). Breathing at a rate of 5.5 breaths per minute with equal inhalation-to-exhalation ratio increases heart rate variability. *International Journal of Psychophysiology*, 91(3), 206–211. 10.1016/j.ijpsycho.2013.12.006 [PubMed: 24380741]
- Malliani A, Lombardi F, & Pagani M (1994). Power spectrum analysis of heart rate variability: A tool to explore neural regulatory mechanisms. *British Heart Journal*, 71(1), 1–2. 10.1136/hrt.71.1.1
- Nijjar PS, Puppala VK, Dickinson O, Duval S, Duprez D, Kreitzer MJ, & Benditt DG (2014). Modulation of the autonomic nervous system assessed through heart rate variability by a mindfulness based stress reduction program. *International Journal of Cardiology*, 177(2), 557–559. 10.1016/j.ijcard.2014.08.116 [PubMed: 25179555]
- Riley KE, & Park CL (2015). How does yoga reduce stress? A systematic review of mechanisms of change and guide to future inquiry. *Health Psychology Review*, 9(3), 379–396. 10.1080/17437199.2014.981778 [PubMed: 25559560]

- Salahuddin L, Cho J, Jeong MG, & Kim D (2007). Ultra short term analysis of heart rate variability for monitoring mental stress in mobile settings. *Conference Proceedings: Annual International Conference of the IEEE Engineering in Medicine and Biology Society, 2007*, 4656–4659. 10.1109/IEMBS.2007.4353378
- Sato N, & Miyake S (2004). Cardiovascular reactivity to mental stress: Relationship with menstrual cycle and gender. *Journal of Physiological Anthropology and Applied Human Science*, 23(6), 215–223. 10.2114/jpa.23.215 [PubMed: 15599065]
- Shaffer F, & Ginsberg JP (2017). An overview of heart rate variability metrics and norms. *Frontiers in Public Health*, 5, 258. 10.3389/fpubh.2017.00258 [PubMed: 29034226]
- Sullivan MB, Erb M, Schmalzl L, Moonaz S, Noggle Taylor J, & Porges SW (2018). Yoga therapy and polyvagal theory: The convergence of traditional wisdom and contemporary neuroscience for self-regulation and resilience. *Frontiers in Human Neuroscience*, 12, 67. 10.3389/fnhum.2018.00067 [PubMed: 29535617]
- Thayer JF, Ahs F, Fredrikson M, Sollers JJ 3rd, & Wager TD(2012). A meta-analysis of heart rate variability and neuroimaging studies: Implications for heart rate variability as a marker of stress and health. *Neuroscience and Biobehavioral Reviews*, 36(2), 747–756. 10.1016/j.neubiorev.2011.11.009 [PubMed: 22178086]
- van den Broek P, Lorch RF Jr., Linderholm T, & Gustafson M (2001).The effects of readers' goals on inference generation and memory for texts. *Memory & Cognition*, 29(8), 1081–1087. 10.3758/BF03206376 [PubMed: 11913743]
- Vaschillo EG, Vaschillo B, Pandina RJ, & Bates ME (2011). Resonances in the cardiovascular system caused by rhythmical muscle tension. *Psychophysiology*, 48(7), 927–936. 10.1111/j.1469-8986.2010.01156.x [PubMed: 21143610]
- Vasiliev V, Meredith S, & Ryabko M (2006). *Let every breath: Secrets of the Russian Breath Masters: Vasiliev*. Toronto, Canada: Russian Martial Art.
- Visnovcova Z, Mestanik M, Javorka M, Mokra D, Gala M, Jurko A, ... Tonhajzerova I (2014). Complexity and time asymmetry of heart rate variability are altered in acute mental stress. *Physiological Measurement*, 35(7), 1319–1334. 10.1088/0967-3334/35/7/1319 [PubMed: 24854052]
- Walther A, Lacker TJ, & Ehlert U (2018). Everybody was Kung-Fu fighting—The beneficial effects of Tai Chi Qigong and self-defense Kung-Fu training on psychological and endocrine health in middle aged and older men. *Complementary Therapies in Medicine*, 36, 68–72. 10.1016/j.ctim.2017.11.021 [PubMed: 29458935]
- Wayne PM, Walsh JN, Taylor-Piliae RE, Wells RE, Papp KV, Donovan NJ, & Yeh GY (2014). Effect of tai chi on cognitive performance in older adults: Systematic review and meta-analysis. *Journal of the American Geriatrics Society*, 62(1), 25–39. 10.1111/jgs.12611 [PubMed: 24383523]
- Wolever RQ, Bobinet KJ, McCabe K, Mackenzie ER, Fekete E, Kusnick CA, & Baime M (2012). Effective and viable mind-body stress reduction in the workplace: A randomized controlled trial. *Journal of Occupational Health Psychology*, 17(2), 246–258. 10.1037/a0027278 [PubMed: 22352291]
- Wu C, Yi Q, Zheng X, Cui S, Chen B, Lu L, & Tang C (2019). Effects of mind-body exercises on cognitive function in older adults: A meta-analysis. *Journal of the American Geriatrics Society*, 67(4), 749–758. 10.1111/jgs.15714 [PubMed: 30565212]
- Zimmermann-Viehoff F, Thayer J, Koenig J, Herrmann C, Weber CS, & Deter HC (2016). Short-term effects of espresso coffee on heart rate variability and blood pressure in habitual and non-habitual coffee consumers—A randomized crossover study. *Nutritional Neuroscience*, 19(4), 169–175. 10.1179/1476830515Y.0000000018 [PubMed: 25850440]




FIGURE 1.

Examples of heart rate variability (HRV) data after frequency analysis. After R-R interval data were processed in Kubios HRV Premium, fast Fourier transformation spectral analysis was performed separating the HRV power into low frequency (LF) and high frequency (HF) components. For comparison, the autonomic state during the Stroop test was normalized to baseline reading for each subject




FIGURE 2.

(a) Parasympathetic response (high frequency [HF]). There was no significant difference in HF between breathing alone group and reading control during the Stroop test. Alternating contraction group had 71.7% higher activation of HF heart rate variability (HRV) over respiration alone ($p < .001$). Alternating contractions synchronized with breathing demonstrated 150% higher parasympathetic activation than control ($p < .0001$). Between the contraction alone and synchronized groups, the synchronized group demonstrated 45.9% higher parasympathetic response ($p < .001$). (b) Sympathetic response (low frequency [LF]). LF activation during administration of the Stroop test did not demonstrate a difference between breathing alone group and reading control. The alternating contraction group and synchronized groups had 33.5% and 45.2% lower LF activation over control respectively ($p < .01$ and $p < .0001$). There was no significant difference in LF response between contraction alone and synchronized groups. (c) Sympathovagal balance (LF/HF). There was no significant LF/HF ratio difference between breathing alone group and reading control. Both contraction groups had significantly less sympathetic activation over control respectively ($p < .01$ and $p < .001$). There was no significant difference in LF response between contraction alone and synchronized groups

TABLE 1

Baseline characteristics

Group	Mean age	Mean systolic BP	Mean diastolic BP	Mean HR	Mean R-R
Breath Only n=12	29.7 (22–40)	121.8 (97–136)	76.2 (70–88)	69.0 (52–95)	16.9 (10–25)
Contraction Only n=12	26.7 (22–33)	121.5 (104–136)	73.2 (61–86)	69.9 (57–82)	18.8 (15–24)
Breath and Contraction n=12	32.3 (22–38)	120.1 (107–134)	74.8 (68–88)	67.3 (55–83)	16.4 (9–26)
Control n=12	30.3 (22–46)	122.7 (102–145)	79.7 (63–96)	63.7 (53–81)	16.2 (8–25)

Abbreviations: BP = Blood Pressure, HR = Heart Rate, RR = Respiration Rate