



A PRACTICAL GUIDE TO OPTIMIZING THE BENEFITS OF POST-STROKE SPASTICITY INTERVENTIONS WITH BOTULINUM TOXIN A: AN INTERNATIONAL GROUP CONSENSUS

Gerard E. FRANCISCO, MD¹, Alexander BALBERT, PhD², Ganesh BAVIKATTE, MD, FRCP, FEBPRM³, Djamel BENSMAIL, MD⁴, Stefano CARDA, MD, PhD⁵, Thierry DELTOMBE, MD⁶, Nathalie DRAULANS, MD⁷, Steven ESCALDI, DO⁸, Raphaël GROSS, MD, PhD⁹, Jorge JACINTO, MD¹⁰, Nicholas KETCHUM, MD¹¹, Franco MOLteni, MD¹², Susana MORALEDA, MD¹³, Michael W. O'DELL, MD¹⁴, Rajiv REEBYE, MD, FRCP¹⁵, Patrik SÄTERÖ, MD¹⁶, Monica VERDUZCO-GUTIERREZ, MD¹⁷, Heather WALKER, MD¹⁸ and Jörg WISSEL, MD, FRCP¹⁹

From the ¹University of Texas Health Science Center McGovern Medical School, Department of Physical Medicine and Rehabilitation, and The Institute for Rehabilitation and Research (TIRR) Memorial Hermann Hospital, Houston, Texas, USA, ²Department of Adaptive Physical Training, Ural University of Physical Education, Sverdlovsk regional hospital for war veterans, Yekaterinburg, Russia, ³Department of Rehabilitation Medicine, The Walton Centre National Health Service Trust, Lower Lane, Liverpool, UK, ⁴Raymond Poincaré Hospital, Assistance publique - Hôpitaux de Paris, University of Versailles Saint-Quentin, Garches, France, ⁵Service of Neuropsychology and Neurorehabilitation, Department of Clinical Neurosciences, Lausanne University Hospital (CHUV), Lausanne, Switzerland, ⁶Medecine Physique & Readaptation, Centre Hospitalier Universitaire UCL Namur site Godinne, Yvoir, Belgium, ⁷Libra Rehabilitation & Audiology, Eindhoven, The Netherlands, ⁸JFK-Johnson Rehabilitation Institute, New Jersey, USA, Rutgers – Robert Wood Johnson Medical School, Hackensack Meridian Health School of Medicine at Seton Hall University, New Jersey, USA, ⁹Nantes Université, Centre Hospitalier Universitaire de Nantes, Movement - Interactions - Performance (MIP), Nantes, France, ¹⁰Alcoitão Medical Rehabilitation Centre, Rua Conde Barão, Alcabideche, Portugal, ¹¹Medical College of Wisconsin, Department of Physical Medicine and Rehabilitation, Milwaukee, Wisconsin, USA, ¹²Valduce Hospital "Villa Beretta" Rehabilitation Center, Costa Masnaga, Italy, ¹³Physical Medicine & Rehabilitation Department, La Paz University Hospital, Madrid, Spain, ¹⁴Weill Cornell Medicine, New York, USA, ¹⁵Department of Medicine, Division of Physical Medicine and Rehabilitation, University of British Columbia, Vancouver BC Canada, ¹⁶Rehdo Rehabilitation Centre, Gothenburg, Sweden, ¹⁷Joe R. & Teresa Lozano Long School of Medicine at UT Health San Antonio, San Antonio, Texas, USA, ¹⁸Medical University of South Carolina Health Rehabilitation Hospital of Charleston an Affiliate of Encompass Health, Medical University of South Carolina, Charleston, South Carolina, USA and ¹⁹Neurological Rehabilitation and Physical Therapy, Vivantes Hospital Spandau, Berlin, Germany

This consensus paper is derived from a meeting of an international group of 19 neurological rehabilitation specialists with a combined experience of more than 250 years (range 4–25 years; mean 14.1 years) in treating post-stroke spasticity with botulinum toxin A. The group undertook critical assessments of some recurring practical challenges, not yet addressed in guidelines, through an extensive literature search. They then discussed the results in the light of their individual clinical experience and developed consensus statements to present to the wider community who treat such patients. The analysis provides a comprehensive overview of treatment with botulinum toxin A, including the use of adjunctive therapies, within a multidisciplinary context, and is aimed at practicing clinicians who treat patients with post-stroke spasticity and require further practical guidance on the use of botulinum toxin A. This paper does not replicate information published elsewhere, but instead aims to provide practical advice to help optimize the use of botulinum toxin A and maximize clinical outcomes. The recommendations for each topic are summarized in a series of statements. Where published high-quality evidence exists, the recommendations reflect this. However, where evidence is not yet conclusive, the group members issued statements and, in some cases, made recommendations based on their clinical experience.

Key words: spasticity; botulinum toxin; consensus guideline.

Accepted Sep 29, 2020; pub ahead of print Oct 15, 2020

J Rehabil Med 2021; 53: jrm00134

LAY ABSTRACT

A group of doctors from around the world, who are experts in treating muscle stiffness and spasm (also called spasticity), reviewed the current scientific evidence supporting the effectiveness of using botulinum toxin injections in treatment of spasticity that results from a stroke. When evidence is not available, they discussed and agreed on the best way to treat spasticity using botulinum toxin. The recommendations made by these expert doctors can be used by less-experienced doctors as a guide to how best to use botulinum toxin injection in treating spasticity after a stroke.

Correspondence address: Gerard E. Francisco, McGovern Medical School, Department of Physical Medicine & Rehabilitation, TIRR Memorial Hermann Hospital, Physical Medicine and Rehabilitation, Houston, USA. E-mail: gerard.e.francisco@uth.tmc.edu

Botulinum toxin A (BoNT-A) has been in clinical use for treating post-stroke spasticity for approximately 30 years and is the accepted standard of care for focal post-stroke spasticity (1). It is currently known that BoNT-A treatment is safe and effective for use in both upper and lower limb spasticity, where it can result in both active and passive functional gains (2). Furthermore, BoNT-A is a first-line pharmacological treatment in the management of post-stroke focal and multi-focal spasticity, which, along with a multidisciplinary team (MDT) approach, should be part of a rehabilitation programme to promote optimal clinical effect (3–5). In addition, the Royal College of Physicians' (RCP) guidelines for management of adult spasticity using BoNT-A

(6) recommend that patient selection and management should be based on individualized criteria, resulting in a patient-centred approach to management.

Despite the ever-expanding literature base on this topic, it is clear that further clinical research is necessary to increase understanding and fill gaps in post-stroke spasticity treatment protocols. The group felt that, in the intervening time, there remained a need to provide practical advice on how best to tailor treatment regimens using BoNT-A for individual patients in order to optimize care.

Although BoNT-A is an established treatment for focal spasticity, there is little consensus on how to improve efficacy, and there is a need to increase prescribers' confidence in its use, share current best practice, and identify reasons for sub-optimal responses (e.g. injection technique, dosing, muscle selection).

The group agreed 3 key areas in which additional practical guidance and/or personal training and supervision is required: (i) individualized approach to spastic upper limb in stroke; (ii) optimal injection technique and preparation of the toxin; and (iii) adjunctive treatments. They examined the evidence for each topic, obtained from literature searches using the College of Physicians and Surgeons of British Columbia review, Medline, CINAHL and PubMed databases. They subsequently met on 2 separate occasions for full-day discussions to agree consensus statements on the topics. The gaps in the literature were filled with the knowledge acquired from the combined clinical experience of the group (Table I). Some topics, such as the assessment of spasticity and measurement of the effects of the treatment, have not been addressed by this review, as the authors felt that they have been adequately evaluated elsewhere. However, the authors agree that the Ashworth/Modified Ashworth scale, in spite of its limitations, is currently the preferred scale for assessment of spasticity, due to its simplicity of use. Whilst there is increasing interest in the Tardieu scale, due to its consideration of the velocity-dependent nature of the condition, it is more complex to implement and, therefore, more difficult to interpret. The measurement of treatment effects has not been included in this review, as it has been addressed in many guidelines (6).

INDIVIDUALIZED APPROACH TO THE TREATMENT OF SPASTICITY IN STROKE

The need for an individualized approach is well established (2, 7, 8). While randomized controlled trials (RCTs) have become the accepted gold standard for directing treatment, their strict protocols may not allow an individualized approach or treatment interventions that are applicable to real-life clinical scenarios.

Goal-setting

Patients and caregivers have wide-ranging goals and expectations from treatment, and optimal management plans should accommodate these variables whenever possible.

Goal-setting, as measured by the Goal Attainment Scaling (GAS), in which individual goals for intervention are tailored to individual patients (9, 10), has merit. It sets targets for intervention and provides a standardized measure of outcome (11). GAS has been shown to be sensitive to changes in areas of symptoms/impairment and function/participation, following focal interventions, which are not detected by more global measures (12). Approximately half of the group involved in this analysis stated that they use GAS always or often.

However, consideration must also be paid to how goals are defined and prioritized. For example, a goal may be "reduction in pain", but we must define whether this is a reduction in *intensity or duration* of pain. In addition, while a patient may consider being able to sign his or her name or shake hands as desirable functional goals, the physician might regard the manual dexterity improvement as insufficient if they cannot unscrew a bottle cap.

It is undecided if a new treatment schedule should start with a modest list of goals that are gradually extended, or if the patient and physician should strive from the outset to accomplish an ambitious list of desired outcomes.

Throughout the continuum of care, goals need to be re-evaluated and re-prioritized as it becomes clear what can be achieved, considering the outcome and the limitations of treatment.

Optimal dose of botulinum toxin A

The optimal dose of BoNT-A per injection session is not clear. Although Summaries of Product Characteristics (SPC) give recommendations regarding dosage, it is not known if these are optimal. Clinical experience and recent guideline updates (3) have identified opportunities to tailor treatment to increase the benefits for patients. Several patient/practitioner surveys conducted during the last few years have highlighted a requirement for more tailored treatment options and more flexibility in dose (and/or injection intervals) than those currently approved (13–15).

Increasing the toxin dose can reduce the Modified Ashworth Scale (MAS) score (3, 16–18). Most physicians who administer BoNT to patients with post-stroke spasticity believe that greater flexibility regarding dosing and treatment intervals for injections might benefit some patients (8,19). Approximately 60% of physicians would use higher doses if there were no label restrictions (20).

Table I. Consensus statements from the international group of experts

Statements	Key literature, selected clinical studies and reviews
An individualized approach to spastic upper limb in stroke	
1. A patient-centred collaborative approach should be taken towards management of post-stroke spasticity, and physicians should agree goals with patients and care-givers	(1-3)
<ul style="list-style-type: none"> • A new treatment schedule should start with a modest list of goals, which are reassessed and extended over time as the results of treatment become apparent • Consider using GAS 	(9-12)
2. For patients with multifocal spasticity the approved doses of BoNT-A may not be sufficient to fulfil their needs, in which case the goals should be reviewed and re-prioritized considering patient needs and expectations	(3, 16, 17, 21)
3. Injectors can treat more disabling clinical patterns/aim for more relevant patients' goals, safely and effectively, when employing higher dosages that have demonstrated efficacy and safety in published studies	(17, 21, 27-29)
<ul style="list-style-type: none"> • Treat according to patient's needs/expectations, respecting the patient's desires and the clinician's evaluation, to achieve an optimal response • Management decisions should be based on the individualized experience of the injector and the patient and on available resources 	(3, 13-15, 30-32)
4. Consider a flexible approach towards deciding when to re-inject with botulinum toxin	
<ul style="list-style-type: none"> • Physicians should strive to maintain the efficacy of botulinum toxin and to prevent complications by reinjecting before the emergence of problems with function and/or comfort, even if this is earlier than 12 weeks (but after peak effects have been observed) 	
5. Although no clear relationship has been established between varying dilution and safety or efficacy, the manipulation of dilutions can be considered for different muscles/conditions to enhance the local effect	
<ul style="list-style-type: none"> • Overly large volumes can have a counter-productive effect in small muscles, since damage to the fascia facilitates toxin leakage from the intended site. In these cases, smaller injection volumes at higher concentration should be considered 	
6. In the case of a sub-optimal response, physicians should consider the following at the next visit:	
<ul style="list-style-type: none"> • Review SMARTness of treatment goals • Review patient/carer awareness/involvement in goal-setting • Review injection technique/targeting used • Review correlation between patterns treated and goals set • Increase the total body dose, in order to reach maximal dose/muscle or inject more clinical patterns when some patterns were left out or doses per muscle were sub-maximal, due to total dose limitations • Reduce the treatment interval when duration was insufficient and doses were maximal or adverse events are feared • Review BoNT-A dilution used • Review the adjunctive rehabilitation programme (modalities/intensity) 	
Injection technique	
1. Storage	
<ul style="list-style-type: none"> • The literature suggest that reconstituted BoNT maintains efficacy for longer than the manufacturers suggest • The group recommends that each TOXIN is stored and prepared according to the instructions on the SPC 	(33-36)
2. Reconstitution and aspiration	
<ul style="list-style-type: none"> • Please refer to each product's summary of product characteristics (SmPC) 	
3. Dilution	
<ul style="list-style-type: none"> • The consensus of the literature is that, although there is some evidence to the contrary, most studies do not support hypothesis that higher volumes lead to increased clinical effect in terms of reduction of spasticity 	(3, 7, 24)
4. Analgesia	
<ul style="list-style-type: none"> • Consider analgesia according to the needs of the patient and level of discomfort during injections: skin cooling seems to be the most effective intervention • An empathetic injector can also help reduce discomfort • The group recommends that analgesia, if available, and necessary, is used during the procedure. 	(52-54) (62, 68-72)
5. Injection guidance	
<ul style="list-style-type: none"> • Many studies show that instrumental guidance (EMG, ES or US) can improve accuracy of injections • There is no clear evidence that any of these is superior over the others • Physicians should use one or a combination of techniques according to their availability and experience 	(1, 3, 46, 61, 63, 74-78) (97-103)
6. Endplate targeting	
<ul style="list-style-type: none"> • Endplate targeting is theoretically desirable. However, there is limited evidence, to date, to support the clinical value of endplate targeting 	(86-87, 89-96))
7. Conversion ratio	
<ul style="list-style-type: none"> • Although doses of the different toxin formulations are not interchangeable, sometimes there is a need to change formulations for non-clinical reasons • A wide range of dose conversion ratios between onabotulinumtoxinA or incobotulinumtoxinA and abobotulinumtoxinA have been investigated in clinical trials, but no definitive conclusions have emerged • Where possible, dose conversion between these formulations should be avoided and each toxin titrated to the individual patient • The consensus group members were willing to use a conversion ratio of 1:1 between incobotulinumtoxinA and onabotulinumtoxinA, if faced with a need for change 	

Table I. Cont.**Adjunctive treatment (listed alphabetically)**

1. Casting/splinting/taping	See Table II
<ul style="list-style-type: none"> • Critical need for high-evidence research on effect of casting/splinting/taping post-BoNT-A and long-term benefits in this population • Current evidence lacks controlled research designs, robust sample sizes and sensitive outcome measures • Selective groups of stroke survivors have benefited from casting as an adjunctive therapy post BoNT-A 	(106)
<ul style="list-style-type: none"> • Future studies are required to assess the impact of casting on upper and lower limb function; also inhibitive casting, short duration and serial casting as an adjunctive therapy • The group recommends the use of casting and/or taping and/or splinting if available, especially if there is a high risk of soft-tissue shortening. 	(123–126, 128)
2. Constraint-induced movement therapy (CIMT)	(109, 129)
<ul style="list-style-type: none"> • There is good evidence for the use of CIMT in post-stroke rehabilitation • It is not proven if it has a specific role to play after BoNT-A injection • The group recommends that physical and occupational therapy, particularly CIMT, is to be used after toxin injection. 	
3. Extracorporeal shock wave treatment (ESWT)	
<ul style="list-style-type: none"> • ESWT should be considered as an adjunctive therapy or in combination with BoNT-A injections • ESWT can be performed at elbow flexors: forearm and triceps surae • Middle belly (1500 I, 0.030 mJ/mm²) seems appropriate • Best indications <ul style="list-style-type: none"> • In multi-pattern patients who required (very) high doses (in isolation) 	
4. Functional electrical stimulation (FES)	(133, 134)
<ul style="list-style-type: none"> • Although there is no definitive published evidence, it is likely that FES provides some degree of additional effect physiological and, possibly, clinical benefit in conjunction with botulinum toxin • The magnitude of the clinical benefit is unclear and this should be counterbalanced against financial constraints and organizational input on clinical decision-making • The ideal FES settings and protocol are also unclear (Hz, best time after injection to initiate stimulation, length and frequency of sessions), distinguish adjunct vs therapeutic • The published protocols may be of value • The mandate for further research in this area is fairly high, given the cost of BoNT and the intervention and the potential to augment the effect with an alternative, inexpensive and readily available modality <ul style="list-style-type: none"> • However, the research must be adequately powered to make a definitive conclusion on benefit, and ideally would include at least 3 arms (control and 2 fairly different FES protocols) • The group recommends that electrical stimulation of the injected muscles can be used, if available. 	
5. Self-rehabilitation	
<ul style="list-style-type: none"> • There is some evidence in the lower limb where task-oriented exercise focusing on balance control, transfers, gait, strengthening and stretching has been shown to be useful in improving gait in stroke • It must be remembered that, in lower limb, we are addressing active function, but in upper limb we need to consider both active and passive function; therefore, extrapolation of results from lower to upper limb may not always be appropriate • Since there are positive benefits seen with repetition and task-oriented exercise, home-based and tele-rehabilitation, it is recommended that patients are trained to follow a self-rehabilitation programme for spasticity (lower limb) and post-stroke recovery (in general) to supplement their clinician-administered physiotherapy 	

GAS: Goal Attainment Scaling; BoNT-A: botulinum toxin A; SMART specific, measurable, achievable, realistic/relevant and timed; SPC: summary of product characteristics; EMG: electromyogram; ES: electrical stimulation; US: ultrasound.

The use of higher than approved doses of BoNT-A has been investigated in the TOWER study (21). This prospective, open-label, single-arm, multicentre, dose-titration study investigated the safety and efficacy of increased incobotulinumtoxinA total-body doses (up to 800 U); this is a higher limit than studied previously. Doses of up to 800 U allowed the treatment of an increasing number of spasticity patterns at a single treatment session without compromising safety or tolerability, enabling patients to achieve more goals. A dose of 800 U of incobotulinumtoxinA enabled more than 97% of patients with cerebral-origin spasticity to receive simultaneous treatment of the upper and lower limb, and more patients achieved 3 or 4 goals (87%) with 800 U than with the lower dose (25% in the 400 U group). The higher dose also increased the number of patients with no quality of life (QoL) impairment on the EuroQoL-5D (EQ-5D) scale. Importantly, in the TOWER study there

was no development of clinical non-responsiveness and no development of neutralizing antibodies. The majority (82%) of the clinicians in the consensus group stated that they were comfortable using 800 U of incobotulinumtoxinA in routine clinical practice.

A recent study (18) assessed the effects of repeated injections of abobotulinumtoxinA in doses of 500, 1,000, or 1,500 over a period of one year. Although the incidence of treatment-emergent adverse events (TEAEs) decreased across cycles, the investigators' evaluation identified 2 cases (constipation, diplopia) in which possible remote spread of toxin effect could not be ruled out and 8 patients seroconverted for neutralizing antibodies up to the study end.

Whilst it could be hypothesized that higher-than-approved doses of onabotulinumtoxinA would also be safe and effective, the group felt that it was important to first demonstrate this in clinical trials.

Initiation and frequency of botulinum toxin A administration

There are also considerations about the optimal time to start treatment and treatment intervals. In lower limb studies in patients with stroke and traumatic brain injury, there is some evidence that earlier treatment of patients with BoNT-A in the post-stroke period achieves better outcomes than is seen in those treated later (17, 22–24). Wein et al. have compared treatment at <24 months with >24 months post-stroke and show greater improvement in MAS in the <24-month post-stroke group, but comment that this is perhaps due to the use of higher doses and more muscles being injected during the open-label phase (Wein et al., personal communication). In an exploratory study, early abobotulinumtoxinA treatment significantly delayed time to reach re-injection criteria compared with placebo in patients with post-stroke upper limb spasticity (ULS) (25).

Duration of effect of BoNT-A can influence the choice of treatment intervals. The duration of effect of BoNT-A can vary between patients, depending on numerous parameters (e.g. clinical condition, age of patient) (13). Although the duration of action is correlated to the amount injected at lower doses, at higher doses of BoNT-A, the duration of action is thought to saturate at approximately 3 months (26).

Flexibility in treatment protocols is determined by individual product licenses, which often lag behind real-world experience (17, 21, 27–29). It may be better to select treatment intervals based on individual patient needs rather than pre-defined regimens.

In a survey of post-stroke spasticity patients (13), while 55% of patients were re-injected at 13–14 weeks or later, 79% actually preferred shorter injection intervals (35.5% at 11–12 weeks). Patient satisfaction decreased as the effects of the previous injection wore off. Thus, patients live with a roller-coaster effect, having treatment that declines in effectiveness with time and then increases again after another injection, suggesting that attempts should be made towards maintaining a near steady-state level of patient satisfaction. To date, unlike in cervical dystonia (CD) (30–32), there are no prospective safety and tolerability data available to support flexible injection intervals in spasticity. However, more than half (53%) of the consensus group members stated that they were confident to re-inject patients when they reported that the maximum effects have worn off, irrespective of common practice (12-week intervals).

TOXIN PREPARATION AND INJECTION TECHNIQUE

A number of different methods and techniques are commonly employed to perform BoNT-A injections

and there is much variability in practice for preparing and handling the toxin and in techniques used for localization, analgesia, dilution and muscle targeting. The practical recommendations made by the group are summarized below. The authors recommend that personal hands-on training and supervision is mandatory for inexperienced clinicians.

Storage

Before reconstitution, the manufacturers' recommendations regarding storage should be followed (as detailed in the package inserts) (33–36).

The group recommends that each toxin is stored and prepared according to the SPC instructions.

Reconstitution and aspiration

Manufacturers' package inserts recommend that BoNT-A should be reconstituted with preservative-free normal saline. Usual practice is to shake gently and avoid aggressive agitation, since one study suggested that mechanical effects of reconstitution were potentially damaging to toxin structure and efficacy (37). However, further studies have shown that constant agitation of the reconstituted solution did not decrease efficacy up to 42 days (38) and vigorous vs gentle agitation made no impact on efficacy (39).

A more recent study concluded that shaking and use of small-bore needles can damage the toxin, as measured by the response time of mouse hemidiaphragm to reduction in force of muscle contraction (40). Since aspiration leaves a small volume of toxin in the vial, use of small-bore needles can be helpful in increasing aspiration efficiency (balanced with the theoretical risk of toxin damage). Some suggest removing the vial stopper to access the maximum volume of toxin, but this risks vial breakage and contamination of the solution, especially if multi-use is planned (41). Residual volume in a capped vial is 0.127 ml, equivalent to almost 13 U per 100 U vial (in a 100 U/1 ml dilution). Alternatively, by using a 2" long 21G needle, it is possible to limit loss to 2.3–4.6 U/vial (assuming 100 U/1 ml dilution) (42). The group was supportive of this approach, awaiting further evidence.

Dilution

A wide range of dilution ratios may be used according to circumstances for common postures of the upper limb in post-stroke spasticity (3, 24).

Animal studies have shown that injections of higher volumes can increase denervation (43, 44). However, the situation is not so clear in humans due to unavailability of adequately powered studies. Higher volumes

have been reported more effective in children with cerebral palsy (45). One study has considered both volume and end-plate targeting of the biceps brachii in 21 adults with spastic hemiplegia following stroke (46). Results showed that high-volume/non-targeted and low-volume/targeted were superior to low-volume/non-targeted. Higher volume and lower concentration has also shown a trend for better efficacy in patients with upper limb spasticity of various aetiologies (47). In a study of the upper limb, despite a trend for more improvement using higher volumes, the improvement was not statistically significant (48); a lack of significance was also seen in 3 studies of the lower limb, 2 of which investigated patients with spastic equinovarus foot (49–51).

The group advised that it may also be necessary to take individual muscle structure into account when injecting, and approximately half stated that they changed dilution to achieve an optimal effect. However, further recommendations are needed to identify the number and location of injection sites (7) required to achieve maximum toxin effect.

Analgesia

EMLA (eutectic mixture of local anesthetics, Aspen Pharma, Ireland) cream and ice packs are equally effective in reducing injection pain at the gastrocnemius, but ice is an easier and cheaper option (52). Vapocoolant has been found to be ineffective (55); however, the time lapse between use of vapocoolant and the injection itself could be a factor. Other authors found (on forehead injections or gastrocnemius for electromyography (EMG)) that vapocoolant is more effective than EMLA (53, 54).

An older study evaluated whether muscle temperature affects BoNT-A uptake (at least *in vitro*) (55) but firm conclusions cannot be drawn.

In considering more generalized anaesthesia, nitrous oxide was shown to be more effective than rectal midazolam in a double-blind, randomized placebo-controlled trial in 50 children with cerebral palsy (CP). Nitrous oxide had the advantage of less post-procedural sedation (56). However, a more recent study in children with CP, analgesia combining nitrous oxide and EMLA cream was less effective than expected for BoNT-A injections (57).

General anaesthesia (GA) can be used in patients with CP and adults, especially in those selected cases who show lack of cooperation, have severe conditions requiring extensive injections, or who have severe cognitive deficit (58).

To reduce injection discomfort, anaesthesia through skin cooling seems the most effective intervention. An empathetic injector may also help to alleviate discomfort (59).

The group recommends that analgesia, if available and required, is used during the procedure.

Injection guidance

Four main injection guidance techniques have been evaluated in clinical trials: needle placement under anatomical guidance (AG), electrical stimulation (ES), electromyography (EMG), and direct visualization by ultrasonography (US). Other techniques have been studied (fluoroscopy, computerized tomography (CT) scan, etc.), but data are sparse.

A number of descriptive studies have considered how to define anatomical landmarks that locate and target motor endplates (60–63).

Palpation, or manual identification of a muscle using anatomical landmarks is not accurate for all muscles; this has been demonstrated in both cadavers and patients (stroke and CP) (64–67).

The advantages of US over other guidance techniques include: direct and continuous visualization of the needle, the target, structures to be avoided and the injectate spread within the target structure. US has identified forearm muscles at distinctly different sites from those identified by anatomical guidance (62, 68, 69). Considering the results of injections, both US and EMG have been demonstrated to lead to better results compared with AG in wrist and finger flexor spasticity (70). US guidance seems to lead to better results on wrist and finger spasticity compared with AG (71). However, a recent study found no differences between US, ES and AG (72).

Supportive literature from lower limb studies show that the accuracy of injections in the gastroc-soleus muscle is higher with ES guidance than with AG. However, US guidance seems to lead to better results on triceps surae spasticity compared with AG and EMG (65).

Most studies show that instrumental guidance (EMG, ES or US) can improve accuracy of injections (73). Although there is no clear evidence that any of these is superior, the group strongly recommends the use of EMG and/or ES and/or US injection guidance to optimize muscle localization. Injection using only anatomical landmarks should be used only when other techniques are unavailable.

End-plate targeting

End-plate targeting considers the end-plate topography, but is also related to injection technique. Nerve and endplate distribution in some human muscles is quite well described (63, 74–77). It has been observed that the area of highest endplate density is an inverted V-shaped band 1 cm in width between the lower third and

upper two-thirds of the muscle belly in human biceps brachii muscles (3, 61). A high-volume dilution (20 U/ml) and an endplate-targeted injection are superior to a low-volume, endplate non-targeted injection, when injecting the biceps brachii (46).

Although some muscles have well-defined motor endplate topography, others, such as the soleus and gastrocnemius, which have diffuse, or ill-defined endplates, may require a more even spread of injection, using multiple injections and larger injection volumes (1). Im et al. (78) demonstrated that a mid-belly injection in the gastrocnemius is no different in efficacy from injection in more anatomically endplate dense regions.

The consensus group acknowledged that end-plate targeting may be helpful for superficial muscles, such as the biceps or gastrocnemius, but none of the group is actively using this technique due to increased time to perform this procedure and the resultant increased pain for the patient.

Conversion ratio

Manufacturers' product labels state that different formulations of toxin are not interchangeable or bioequivalent and conversion ratios have not been systematically studied between formulations. However, since real-world practice sometimes requires switching formulations, attempts have been made to investigate conversion ratios in both clinical studies and in practice. Conducted in variable patient populations, study results are wide-ranging and careful interpretation is recommended.

Conversion ratios of 1:1 to 1:11, between onabotulinum toxin-A and abobotulinum toxin-A, have been used in clinical studies (79–88). Although many authors considered 1:3 or 1:4 to be appropriate (89–95), some evidence suggests that conversion ratios differ between muscles (96).

Studies have shown a conversion ratio of 1:1 for incobotulinum toxin-A to onabotulinum toxin-A in CD and blepharospasm (97–103). This is reflected in the European SPC for incobotulinumtoxinA (35). In focal dystonia, a study showed that switching from

abobotulinum toxin-A to incobotulinum toxin-A with a dose ratio of 4:1 was effective and well-tolerated (104).

Although no conversion ratio can be recommended at this time, if faced with a need to change formulations for practical reasons, the consensus group was comfortable using a conversion ratio of 1:1 between incobotulinumtoxinA and onabotulinumtoxinA; there was no agreement on a conversion ratio between onabotulinumtoxinA and abobotulinumtoxinA.

CONCOMITANT AND ADJUNCTIVE THERAPIES

BoNT-A injections are a part of spasticity management. The group agreed that BoNT-A injections should always be followed with a physiotherapy programme. Table II represents the techniques that are currently used by group members. However, we need to establish the optimal adjunctive therapy to be used alongside BoNT-A.

Demetrios et al., in a Cochrane review, found only 3 RCTs (105–107) that assessed rehabilitation after BoNT-A (108).

Two recent reviews looked at adjunctive therapies post-BoNT-A in post-stroke spasticity (109, 110). Mills et al. performed a systematic literature review and concluded that there is high-level evidence suggesting that adjunct therapies may improve outcomes following BoNT-A injection. However, no results have been confirmed by independent replication; all interventions would benefit from further study.

Kinnear et al. also emphasized that the effectiveness of these therapies is uncertain, with the 95% CI sometimes spanning zero (110).

Casting, taping and splinting

Available literature on casting, taping and splinting for spasticity management post-stroke were reviewed, but due to the limited number of papers this review concentrates on casting (as there were more studies). A recent review (111) of the current evidence for

Table II. Adjunctive therapies: current use by group ($n = 19$)

Technique/modality used with BoNT-A	Never 0% of the time, n (%)	Rare 1–25% of the time, n (%)	Sometimes 25–49% of the time, n (%)	Most 50–79% of the time, n (%)	Almost always 80–100% of the time, n (%)
Skilled physiotherapist	0 (0.0)	0 (0.0)	2 (10.5)	9 (47.4)	8 (42.1)
Casting/splinting/taping	0 (0.0)	9 (47.4)	8 (42.1)	2 (10.5)	0 (0.0)
Orthosis	0 (0.0)	2 (10.5)	9 (47.4)	8 (42.1)	0 (0.0)
Functional electrical stimulation	0 (0.0)	12 (63.2)	7 (36.8)	0 (0.0)	0 (0.0)
Extracorporeal shock wave treatment*	18 (94.7)	1 (5.3)	0 (0.0)	0 (0.0)	0 (0.0)
Neuromuscular stimulation	12 (63.2)	6 (31.6)	1 (5.3)	0 (0.0)	0 (0.0)
Robotics	6 (31.6)	8 (42.1)	5 (26.3)	0 (0.0)	0 (0.0)
Self-rehabilitation	0 (0.0)	0 (0.0)	2 (10.5)	3 (15.8)	14 (73.7)

*For most members of the group, extracorporeal shock wave therapy (ESWT) was not readily available or they were unfamiliar with the technique. BoNT-A: botulinum toxin A.

casting as an adjunct therapy following botulinum toxin injection for adult limb spasticity concluded that adjunct casting of the lower limbs may improve outcomes following BoNT injections. The group also considered studies on casting/splinting and taping together, focusing on RCTs that had the highest impact and the highest PEDro scores (scale developed by the Physiotherapy Evidence Database to determine the quality of clinical trials: high quality=6–10; fair quality=4–5; poor quality= ≤ 3) (112). The relevant studies are summarized in Table III. Four RCTs studied BoNT-A plus casting/taping/splinting in lower limb with positive results (113–116). Two studies (117, 118) investigated BoNT-A plus/minus taping in lower limb which a favourable trend towards the use of taping, but no long-term benefits. A retrospective study (119)

concluded that serial casting may be an appropriate intervention following BoNT-A injection to prevent equinovarus deformity and improve quality of walking in chronic stroke patients. Another RCT (120) reviewed the use of BoNT-A plus casting/taping/splinting in the upper limb and found that adhesive taping was more effective than daily manual stretching combined with passive articular mobilization and palmar splint.

A common criticism of many papers is that they provide insufficient detail about injection technique (manual palpation, EMG, nerve stimulator or US guidance), the type of casting/splinting/taping or orthotics used. More robust studies are needed comparing BoNT-A and adjunctive therapy with casting/taping/splinting post BoNT-A with BoNT-A treatment alone. Future studies should also indicate the specific timeline and technique of casting/taping/

Table III. Summary of casting/taping/splinting studies plus botulinum toxin A (BoNT-A) in spasticity

Review article	Summary
Mills PB et al., 2016 (109)	
<i>Randomized controlled trials (RCTs) for BoNT-A plus casting/taping/splinting in lower limb</i>	
Carda S et al., 2011 (113)	Investigated the effect of different adjunctive treatments after BoNT-A 69 chronic hemiplegic adult patients with spastic equinus foot Following BoNT-A injection into plantar flexors, patients were randomly assigned to either taping, casting or stretching for one week and with stretching and gait training for the following week Combining BoNT-A to the ankle plantar flexors with casting or taping gives better and longer lasting results than stretching alone PEDro 7
Farina S et al., 2008 (114)	13 stroke patients, night cast for 4 months Prolonged stretching of spastic muscles after BoNT-A injection affords long-lasting therapeutic benefit enhancing the effects of toxin alone PEDro 5
Baricich A et al., 2008 (115)	The group treated with electrical stimulation performed better at t1 on the MAS. The taping and electrical stimulation groups performed better in all outcome measures at t3. The taping group performed better, mainly for maximum ankle dorsiflexion angle in stance. The stretching group showed a less durable result, with some worsening at the t3 evaluation compared with the assessment performed before treatment. PEDro 6
Verplancke D et al., 2005 (116)	Serial casting combined with BoNT-A reduces the development of calf contracture after severe head injury 35 patients Three patient groups: cast plus saline, cast plus BoNT-A, and control Casting alone in these patients was sufficient PEDro 8
<i>BoNT-A plus/minus taping in lower limb</i>	
Karadag-Saygi E E et al., 2010 (117)	Double-blind RCT Taped for 2 weeks Positive result in decrease in plantar flexor spasticity post-2 weeks, but no long-term effect PEDro 7
Reiter F et al., 1998 (118)	Single-blinded RCT Low-dose BoNT-A to lower limb plantar flexors Improvement in MAS, but not gait parameters No long-term follow-up PEDro 5
<i>Retrospective study</i>	
Yasar E et al., 2010 (119)	Serial casting may be an appropriate intervention following BoNT-A injection to prevent equinovarus deformity and improve quality of walking in chronic stroke patients. Role of casting and splinting are important topics that require further research. PEDro cannot be calculated (study not an RCT)
<i>RCTs for BoNT-A plus casting/taping/splinting in upper limb</i>	
Santamato A et al., 2015 (120)	Adhesive taping more effective than daily manual stretching combined with passive articular mobilization and palmar splint. PEDro 8
Lai JM et al., 2016 (105)	Individuals with spastic elbow flexors post-stroke who can tolerate 6–8 h dynamic splinting per night for 4 months may have some improvements in active elbow extension and MAS compared with BoNT-A alone. PEDro score 4

RCT: randomized clinical trial; PEDro; Physiotherapy Evidence Database score.

splinting following injection of BoNT-A. Nevertheless, based on clinical experience, the group recommends the use of casting and/or taping and/or splinting, if available, especially if there is a high risk of soft-tissue shortening.

Constraint-induced movement therapy

One study examined constraint-induced movement therapy (CIMT) in post-stroke patients in combination with BoNT (106). The PEDro score of this study was 6. Thirty-two patients were assigned to 2 groups and treated with BoNT plus CIMT or BoNT plus conventional rehabilitation (2 h per day, 3 days per week). In both groups, spasticity was better at 4 weeks and 3 months. There was a significant difference in the CIMT group at 6 months in terms of spasticity measured by MAS; utilization of the affected arm with a Motor Activity Log; and arm mobility assessed by the Action Research Arm Test. A weakness of this study was that there was no CIMT alone cohort.

The authors suggest that improvements could be attributable to several factors: improvement in strength and co-ordination in the affected upper extremity as a result of spasticity reduction and repetitive training, a change in learned non-use behaviours, or use-dependent cortical changes after the combination of BoNT-A and CIMT. Currently it is not known whether user-dependent cortical reorganization can occur in chronic patients with significant spasticity. Further research exploring CNS changes accompanying the observed motor gains is warranted (106).

The group recommends that physical and occupational therapy, particularly CIMT, is provided after toxin injection.

Extracorporeal shock wave treatment

The conclusion from the literature is that extracorporeal shock wave therapy (ESWT) reduces spasticity (alone and in combination with BoNT-A injections). A small meta-analysis of 5 studies showed that spasticity (as measured by MAS) improved immediately and at 4 weeks after ESWT compared with baseline (121). The usual target is middle belly or musculotendinous junction (1,500–3,200 I, 0.030 to 0.1 mJ/mm² reported).

A recent randomized trial (122) investigating whether ESWT is non-inferior to BoNT-A for the treatment of post-stroke upper limb spasticity found that ESWT and BoNT-A caused similar reduction in spasticity of the wrist and elbow flexors (MAS); however, ESWT yielded greater improvement in wrist and elbow passive range of motion (PROM), and upper extremity Fugl-Meyer Assessment (UE-FMA) score. One RCT vs placebo showed that BoNT-A plus ESWT is more effective than BoNT-A plus electrical stimulation (ES) in the forearm (123). The mean outcome

measure (MAS, SFS and VAS) in patients decreased in the BoNT-A plus ES group.

Used alone, ESWT can reduce spasticity at forearm and triceps surae following stroke (124–126). Three sessions provided a longer effect (16 vs 8–12 weeks) and greater hand function-wrist control improvement than one session (125). The mechanism behind ESWT is unknown, but it appears to be unrelated to a decrease in spinal excitability (127). The effects last up to 12–16 weeks (although no study with longer follow-up has been conducted). It appears to be well tolerated, with few or no adverse effects.

One publication in CP (128) compared BoNT-A injection plus 3 ESWT sessions with BoNT-A alone. At one-month, significant differences were found between groups in the injected muscles percentage of hardness ($p=0.021$) and the MAS ($p=0.001$), supporting the hypothesis that the combined effects of BoNT-A and ESWT derive from their respective action on neurological and non-neural rheological components in spastic muscles.

The group concluded that further standardization of treatment protocols including optimal sites, treatment intervals and intensities need to be established. In addition, long-term follow-up studies are needed to understand the mechanism of action and resultant muscle changes with repeated sessions.

Functional electrical stimulation

There are several advantages to combining functional electrical stimulation (FES) or electrical stimulation (ES) and BoNT treatment (129). Postulated mechanisms include: FES increases synaptic activity and increases BoNT-A uptake (short-term), FES increases mechanical spread of toxin (short-term); there is a direct effect of FES on spasticity (short- and long-term); or FES increases strength in antagonists, while BoNT-A decrease tone in agonists (long-term).

Two reviews (129, 130) have concluded that, overall, the studies have small sample sizes and varying quality, with PEDro scores ranging from 6 to 10. Thus, they lack the power needed to draw firm conclusions on the effect of FES as an adjunctive treatment with BoNT-A.

A recently published systematic review (109) provides a starting point for FES review of 8 clinical trials that have examined electrical stimulation in stroke in upper and/or lower limb spasticity. The authors concluded that “When compared with BoNT-A injection alone, we found evidence suggesting that the adjunct use of ... electrical stimulation... (Level I)... resulted in improvement in Modified Ashworth Scores by at least one grade”.

It is not always possible to translate findings in one muscle to another at a different anatomical location, but, since we are considering effects on basic nerve

functioning, it is likely that findings observed in one body region will be applicable to other areas. So, given the paucity of upper limb studies, reference should also be made to lower limb investigations.

It is not clear whether ES increases BoNT-A uptake or whether it modulates neural activity in the spinal cord or the brain. BoNT-A is taken up by neurones by binding to synaptic vesicle protein SV2. The intention of most of these studies would appear to be increasing toxin uptake by the SV2 receptors.

There is a need to define better the timing and duration of ES after BoNT-A injections. In their systematic review of electrical stimulation, Mills et al. (109) point out that the variables for the intervention (frequency, current pulse duration, intensity) as well as duration of treatment differed between the studies; as such, a meta-analysis could not be performed. Dosing ranged between 30 and 60 min per session, delivered between once daily to 6 times daily for 3 days up to 12 weeks.

Clinical outcomes have been mostly modified Ashworth scores, while physiological studies have considered drop in amplitude of compound muscle action potential.

The group recommends that ES of the injected muscles can be used, if available.

Other concomitant therapies

Two narrative reviews of repetitive transcranial magnetic stimulation (rTMS) and transcranial direct current stimulation (tDCS) as spasticity treatment have recently been published (130,131). However, the group concluded that it is too early to include these techniques as adjunctive therapy in stroke.

Most relevant studies of robotics have been conducted in the lower limb. A recent study (132) was conducted in 17 post-stroke patients who participated in daily rehabilitation sessions using the NEUROExos Elbow Module exoskeleton (BioRobotics Institute of Scuola Superiore, Sant'Anna, Italy). Results showed that the robotic exoskeleton can be safely used for post-stroke spasticity rehabilitation and that intensive early rehabilitation treatment may prevent the spasticity occurrence at a later stage.

The group concluded that it is too early to make recommendations on the use of robotics in rehabilitation, due to the paucity of evidence. However, as repetition is important in rehabilitation, robotics are likely to gain more importance in the future.

Self-rehabilitation

A Cochrane review (133) of tele-rehabilitation in stroke stated that there is insufficient evidence to reach conclusions about its effectiveness after stroke. A systematic and meta-analysis of randomized trials (134)

showed similar results for Barthel Index, Berg Balance, and Fugl-Meyer scoring in home-based rehabilitation vs conventional rehabilitation, but the studies lacked spasticity outcome measures.

Another study considering home-based tele-supervising rehabilitation (30 min per day for 5 days per week) on physical function in stroke survivors (135), but again not using spasticity outcome measures, concluded that home-based tele-supervising rehabilitation is most likely as effective as the conventional outpatient rehabilitation for improving functional recovery in stroke survivors.

An RCT (136) in 35 outpatients used a 10-m timed walk, the "Timed Up and Go" test, distance covered in 6 min over an ecological circuit, and the stair test. The results strongly suggested that a standardized self-rehabilitation programme constitutes a useful adjunct to BoNT-A injections in order to improve gait-related activities.

However, a survey (137) found that physical therapy professionals are less accepting (than physiotherapy students, professionals and physicians) of the need to engage patients with post-stroke hemiparesis into Guided Self-Rehabilitation Contracts (137) (designed to increase their exercise intensity and responsibility level). There is a need to investigate this topic further using spasticity outcome measures in the context of BoNT-A use.

Since positive benefits are seen with repetition and task-oriented exercise, home-based and tele-rehabilitation, it is recommended that patients are trained to follow a self-rehabilitation programme for spasticity (lower limb) and post-stroke recovery (in general) to supplement their clinician-administered physiotherapy.

CONCLUSION

The use of BoNT-A and innovative techniques has facilitated a more individualized approach to treatment of post-stroke spasticity, which provides physicians with the opportunity to optimize outcomes and address multiple goals. Table I outlines the consensus view of best practice on the optimal use of BoNT-A within a multidisciplinary context, including use of adjunctive therapies that are commonly employed with BoNT A. BoNT-A usage and choice of adjunctive procedures should be made according to individual needs and treatment goals.

ACKNOWLEDGEMENTS

The ToxNet Group is supported by an unrestricted educational grant from Merz. Medical writing support was provided by Brenda McCleary and Jan Hawthorn, who were paid from this grant.

All authors developed the concept for the paper, critically appraised the literature and contributed to the discussion to reach a consensus. The content of the manuscript was directed and reviewed by the authors, who approved the final version before submission.

REFERENCES

- Wissel J, Ward AB, Ertzgaard P, Bensmail D, Hecht M, Lejeune T, et al. European consensus table on the use of botulinum toxin type A in adult spasticity. *J Rehabil Med* 2009; 41: 13–25.
- Esquenazi A, Alfaro A, Ayyoub Z, Charles D, Dashtipour K, Graham GD, et al. OnabotulinumtoxinA for lower limb spasticity: guidance from a Delphi panel approach. *PM R* 2017; 9: 960–968.
- Simpson DM, Hallett M, Ashman E. Practice guideline update summary: botulinum neurotoxin for the treatment of blepharospasm , cervical dystonia , adult spasticity and headache: report of the Guideline Development Subcommittee of the American Academy of Neurology. *Am Acad Neurol* 2016; 86: 1818–1826.
- Weinstein CJ, Stein J, Arena R, American Heart Association Stroke Council on Cardiovascular and Stroke, Nursing Council on Clinical Cardiology, and Council on Quality Care and Outcomes Research. Guidelines for adult stroke rehabilitation and recovery: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2016; 47: e98–e169.
- Teasell R, Foley N, Pereira S, Sequeira K, Miller T. Evidence to practice: botulinum toxin in the treatment of spasticity post stroke. *Top Stroke Rehabil* 2012; 19: 115–121.
- Royal College of Physicians, British Society of Rehabilitation Medicine, The Chartered Society of Physiotherapy, Association of Chartered Physiotherapists in Neurology and the RC of OT. Spasticity in adults: management using botulinum toxin. National guidelines. RCP 2018. [Accessed 2018 May 11] Available from: <https://www.rcplondon.ac.uk/guidelines-policy/spasticity-adults-management-using-botulinum-toxin>.
- Simpson DM, Patel AT, Alfaro A, Ayyoub Z, Charles D, Dashtipour K, et al. OnabotulinumtoxinA injection for poststroke upper-limb spasticity: guidance for early injectors from a Delphi panel process. *PMR* 2017; 9: 136–148.
- Wissel J. Towards flexible and tailored botulinum neurotoxin dosing regimens for focal dystonia and spasticity – insights from recent studies. *Toxicon* 2018; 147: 100–106.
- Scottish Intercollegiate Network. Management of patients with stroke: rehabilitation, prevention and management of complications, and discharge planning. A National Clinical Guideline. NHS Evid 2010; SIGN (June). Available from: www.sign.ac.uk/guidelines/fulltext/.
- Holliday RC, Cano S, Freeman JA, Playford ED. Should patients participate in clinical decision making? An optimised balance block design controlled study of goal setting in a rehabilitation unit. *J Neurol Neurosurg Psychiatry* 2007; 78: 576–580.
- Ashford S, Turner-Stokes L. Goal attainment scaling for spasticity management using botulinum toxin. *Physiother Res Int* 2006; 11: 24–34.
- McCrorry P, Turner-Stokes L, Baguley IJ, De Graaff S, Katrak P, Sandanam J, et al. Botulinum toxin a for treatment of upper limb spasticity following stroke: a multi-centre randomized placebo-controlled study of the effects on quality of life and other person-centred outcomes. *J Rehabil Med* 2009; 41: 536–544.
- Bensmail D, Hanschmann A, Wissel J. Satisfaction with botulinum toxin treatment in post-stroke spasticity: results from two cross-sectional surveys (patients and physicians). *J Med Econ* 2014; 17: 618–625.
- Poliziani M, Koch M, Liu X. Striving for more good days: patient perspectives on botulinum toxin for the treatment of cervical dystonia. *Patient Prefer Adherence* 2016; 10: 1601–1608.
- Sethi KD, Rodriguez R, Olayinka B. Satisfaction with botulinum toxin treatment: a cross-sectional survey of patients with cervical dystonia. *J Med Econ* 2012; 15: 419–423.
- Yablon SA, Brin MF, Vandenburgh AM, Zhou J, Garabedian-Ruffalo SM, Abu-Shakra S, et al. Dose response with onabotulinumtoxinA for post-stroke spasticity: a pooled data analysis. *Mov Disord* 2011; 26: 209–215.
- Pittock SJ, Moore AP, Hardiman O, Ehler E, Kovac M, Bojakowski J, et al. A double-blind randomised placebo-controlled evaluation of three doses of botulinum toxin type A (Dysport®) in the treatment of spastic equinovarus deformity after stroke. *Cerebrovasc Dis* 2003; 15: 289–300.
- Gracies JM, O'Dell M, Vecchio M, Hedera P, Kocer S, Rudzinska-Bar M, et al. Effects of repeated abobotulinumtoxinA injections in upper limb spasticity. *Muscle Nerve* 2018; 57: 245–254.
- Trompetto C, Marinelli L, Mori L, Puce L, Pelosin E, Serrati C, et al. Do flexible inter-injection intervals improve the effects of botulinum toxin A treatment in reducing impairment and disability in patients with spasticity? *Med Hypotheses* 2017; 102: 28–32.
- Harriss J, Simon O, Li J, Ellers-Lenz B, Roche N, Cantú-Brito C, et al. Physician characteristics in an international, non-interventional study of botulinum toxin formulations in treatment-naïve patients with spasticity (SPACE). *Ann Phys Rehabil Med* 2014; 57: e47.
- Wissel J, Bensmail D, Ferreira JJ, Molteni F, Satkunam L, Moraleda S, et al. Safety and efficacy of incobotulinumtoxinA doses up to 800 U in limb spasticity the TOWER study. *Neurology* 2017; 88: 1321–1328.
- Burbaud P, Wiart L, Dubos JL, Gaujard E, Debelleix X, Joseph PA, et al. A randomised, double blind, placebo controlled trial of botulinum toxin in the treatment of spastic foot in hemiparetic patients. *J Neurol Neurosurg Psychiatry* 1996; 61: 265–269.
- Hesse S, Krajnik J, Luecke D, Jahnke MT, Gregoric M, Mauritz KH. Ankle muscle activity before and after botulinum toxin therapy for lower limb extensor spasticity in chronic hemiparetic patients. *Stroke* 1996; 27: 455–460.
- Kaji R, Osako Y, Suyama K, Maeda T, Uechi Y, Iwasaki M. Botulinum toxin type A in post-stroke upper limb spasticity. *Curr Med Res Opin* 2010; 26: 1983–1992.
- Rosales RL, Balcaitiene J, Berard H, Maisonnobe P, Goh KJ, Kumthornthip W, et al. Early abobotulinumtoxinA (Dysport®) in post-stroke adult upper limb spasticity: ONTIME pilot study. *Toxins (Basel)* 2018; 10: 1–12.
- Dressler D, Saberi FA, Barbosa ER. Botulinum toxin: mechanisms of action. *Arq Neuropsiquiatr* 2005; 63: 180–185.
- Poewe W, Deuschl G, Nebe A, Feifel E, Wissel J, Benecke R, et al. What is the optimal dose of botulinum toxin A in the treatment of cervical dystonia? Results of a double blind, placebo controlled, dose ranging study using Dysport. German Dystonia Study Group. *J Neurol Neurosurg Psychiatry* 1998; 64: 13–17.
- Schramm A, Ndayisaba JP, Auf Dem Brinke M, Hecht M, Herrmann C, Huber M, et al. Spasticity treatment with onabotulinumtoxin A: data from a prospective German real-life patient registry. *J Neural Transm* 2014; 121: 521–530.
- Hyman N, Glickman S, Sayer A, Richardson A, Dott C, Barnes M, et al. Botulinum toxin (Dysport®) treatment of hip adductor spasticity in multiple sclerosis: a prospective, randomised, double blind, placebo controlled, dose ranging study. *J Neurol Neurosurg Psychiatry* 2000; 68: 707–712.
- Comella CL, Jankovic J, Truong DD, Hanschmann A, Grafe S. Efficacy and safety of incobotulinumtoxinA (NT 201, XEOMIN®, botulinum neurotoxin type A, without accessory proteins) in patients with cervical dystonia. *J Neurol Sci* 2011; 308: 103–109.
- Evidente VGH, Fernandez HH, Ledoux MS, Brashear A, Grafe S, Hanschmann A, et al. A randomized, double-blind study of repeated incobotulinumtoxinA (Xeomin®) in cervical dystonia. *J Neural Transm* 2013; 120: 1699–1707.
- Truong DD, Gollomp SM, Jankovic J, Lewitt PA, Marx M, Hanschmann A, et al. Sustained efficacy and safety of repeated incobotulinumtoxinA (Xeomin®) injections in blepharospasm. *J Neural Transm* 2013; 120: 1345–1353.

33. Allergan Inc. IC. Package insert Botox 2010. [Accessed 2020 Oct] Available from: <https://media.allergan.com/actavis/actavis/media/allergan-pdf-documents/product-prescribing/20190620-BOTOX-100-and-200-Units-v3-00USP1145-v2-0MG1145.pdf>.
34. Beauford-Ipsen Ltd U. Package insert Dysport 2009. [Accessed 2020 Oct] Available from https://www.ipsen.com/websites/Ipsen_Online/wp-content/uploads/2020/07/10002305/DYS-US-004998_Dysport-PI-July-2020.pdf.
35. Medicines.org.uk. Summary of Product Characteristics. [Accessed 2018 May 11]. Available from: <https://www.medicines.org.uk/emc/product/4609/sumpc>.
36. Lanzhou C. Package Insert Prosigne 2010. [Accessed 2018 May 11] Available from: <https://www.btxa.com/upload/page/3/document/55123b7d8a853.pdf>.
37. Toth SI, Smith LA, Ahmed SA. Extreme sensitivity of botulinum neurotoxin domains towards mild agitation. *J Pharm Sci* 2009; 98: 3302–3311.
38. Shome D, Nair AG, Kapoor R, Jain V. Botulinum toxin A: is it really that fragile a molecule? *Dermatologic Surg* 2010; 36 Suppl 4: 2106–2110.
39. Kazim NA, Black EH. Botox: shaken, not stirred. *Ophthal Plast Reconstr Surg* 2008; 24: 10–12.
40. Dressler D, Altenmueller E, Bhidayasiri R, Bohlega S, Chana P, Chung TM, et al. Strategies for treatment of dystonia. *J Neural Transm* 2016; 123: 251–258.
41. Dykstra DD, Wieting JM, McGuire J, Kowalkowski T. Maximizing extraction of botulinum toxin type A from vials. *Arch Phys Med Rehabil* 2002; 83: 1638–1640.
42. Niamtu J. Neurotoxin waste from drawing product through the vial stopper. *J Clin Aesthet Dermatol* 2014; 7: 33–37.
43. Kim SH, Hwang JH, Jeong ST, Lee YT, Lee PKW, Suh YL, et al. Effect of muscle activity and botulinum toxin dilution volume on muscle paralysis. *Dev Med Child Neurol* 2003; 45: 200–206.
44. Shaari CM, Sanders I. Quantifying how location and dose of botulinum toxin injections affect muscle paralysis. *Muscle Nerve* 1993; 16: 964–969.
45. Hu G-C, Chuang Y-C, Liu J-P, Chien K-L, Chen Y-M, Chen Y-F. Botulinum toxin (Dysport) treatment of the spastic gastrocnemius muscle in children with cerebral palsy: a randomized trial comparing two injection volumes. *Clin Rehabil* 2009; 23: 64–71.
46. Gracies JM, Lugassy M, Weisz DJ, Vecchio M, Flanagan S, Simpson DM. Botulinum toxin dilution and endplate targeting in spasticity: a double-blind controlled study. *Arch Phys Med Rehabil* 2009; 90: 9–16.e2.
47. Barnes M, Schnitzler A, Medeiros L, Aguilar M, Lehnert-Batar A, Minnasch P. Efficacy and safety of NT 201 for upper limb spasticity of various etiologies - a randomized parallel-group study. *Acta Neurol Scand* 2010; 122: 295–302.
48. Francisco G, Boake C, Vaughn A. Botulinum toxin in upper limb spasticity after acquired brain injury: a randomized trial comparing dilution techniques. *Am J Phys Med Rehabil* 2002; 81: 355–363.
49. Lee J, Sung I, Yoo J, Park E, Park S. Effects of different dilutions of botulinum toxin type A treatment for children with cerebral palsy with spastic ankle plantarflexor: a randomized controlled trial. *J Rehabil Med* 2009; 41: 740–745.
50. Lee L-R, Chuang Y-C, Yang B-J, Hsu M-J, Liu Y-H. Botulinum toxin for lower limb spasticity in children with cerebral palsy: a single-blinded trial comparing dilution techniques. *Am J Phys Med Rehabil* 2004; 83: 766–773.
51. Pascual-Pascual SI, Pascual-Castroviejo I, Ruiz PJG. Treating spastic equinus foot from cerebral palsy with botulinum toxin type a: what factors influence the results?: An analysis of 189 consecutive cases. *Am J Phys Med Rehabil* 2011; 90: 554–563.
52. Fung S, Phadke CP, Kam A, Ismail F, Boulias C. Effect of topical anesthetics on needle insertion pain during botulinum toxin type a injections for limb spasticity. *Arch Phys Med Rehabil* 2012; 93: 1643–1647.
53. Irkoren S, Ozkan HS, Karaca H. A clinical comparison of EMLA cream and ethyl chloride spray application for pain relief of forehead botulinum toxin injection. *Ann Plast Surg* 2015; 75: 272–274.
54. Moon YE, Kim SH, Choi WH. Comparison of the effects of vapocoolant spray and topical anesthetic cream on pain during needle electromyography in the medial gastrocnemius. *Arch Phys Med Rehabil* 2013; 94: 919–924.
55. Poulain B, de Paiva A, Dolly JO, Weller U, Tauc L. Differences in the temperature dependencies of uptake of botulinum and tetanus toxins in Aplysia neurons. *Neurosci Lett* 1992; 139: 289–292.
56. Zier JL, Rivard PF, Krach LE, Wendorf HR. Effectiveness of sedation using nitrous oxide compared with enteral midazolam for botulinum toxin A injections in children. *Dev Med Child Neurol* 2008; 50: 854–858.
57. Brochard S, Blajan V, Lempereur M, Garlandezec R, Houx L, Le Moine P, et al. Determining the technical and clinical factors associated with pain for children undergoing botulinum toxin injections under nitrous oxide and anesthetic cream. *Eur J Paediatr Neurol* 2011; 15: 310–315.
58. Forrester M, Srinivasan J, Mhrshahi S, Waugh M, O'Flaherty S, Rice J, et al. Concious sedation or general anaesthetic for intramuscular botulinum toxin injections in children – a two-centre cross-sectional prospective audit. *Eur J Paediatr Neurol* 2012; 16: 215–217.
59. Picelli A, Vallies G, Chemello E, Gavras A, Castellazzi P, Meschieri A, et al. Influence of physician empathy on the outcome of botulinum toxin treatment for upper limb spasticity in patients with chronic stroke: a cohort study. *J Rehabil Med* 2017; 49: 410–415.
60. Harrison TP, Sadnicka A, Eastwood DM. Motor points for the neuromuscular blockade of the subscapularis muscle. *Arch Phys Med Rehabil* 2007; 88: 295–297.
61. Amiralii A, Mu L, Gracies JM, Simpson DM. Anatomical localization of motor endplate bands in the human biceps brachii. *J Clin Neuromuscul Dis* 2007; 9: 306–312.
62. Bickerton LE, Agur AMR, Ashby P. flexor digitorum superficialis: locations of individual muscle bellies for botulinum toxin injections. *Muscle Nerve* 1997; 20: 306–312.
63. An XC, Lee JH, Im S, Lee MS, Hwang K, Kim HW, et al. Anatomic localization of motor entry points and intramuscular nerve endings in the hamstring muscles. *Surg Radiol Anat* 2010; 32: 529–537.
64. Schnitzler A, Roche N, Denormandie P, Lautridou C, Paratte B, Genet F. Manual needle placement: accuracy of botulinum toxin a injections. *Muscle Nerve* 2012; 46: 531–534.
65. Picelli A, Tamburin S, Bonetti P, Fontana C, Barausse M, Dambruoso F, et al. Botulinum toxin type a injection into the Gastrocnemius muscle for spastic Equinus in adults with stroke. *Am J Phys Med Rehabil* 2012; 91: 957–964.
66. Chin TYP, Duncan JA, Johnstone BR, Graham HK. Management of the upper limb in cerebral palsy. *J Pediatr Orthop B* 2005; 14: 389–404.
67. Yang EJ, Rha D wook, Yoo JK, Park ES. Accuracy of manual needle placement for gastrocnemius muscle in children with cerebral palsy checked against ultrasonography. *Arch Phys Med Rehabil* 2009; 90: 741–744.
68. Delagi EF, Perotto A. Clinical electromyography of the hand. *Arch Phys Med Rehabil* 1976; 57: 66–69.
69. Kristi Henzel M, Munin MC, Niyonkuru C, Skidmore ER, Weber DJ, Zafonte RD. Comparison of surface and ultrasound localization to identify forearm flexor muscles for botulinum toxin injections. *PMR* 2010; 2: 642–646.
70. Picelli A, Lobba D, Midiri A, Prandi P, Melotti C, Baldessarelli S, et al. Botulinum toxin injection into the forearm muscles for wrist and fingers spastic overactivity in adults with chronic stroke: a randomized controlled trial comparing three injection techniques. *Clin Rehabil* 2014; 28: 232–242.
71. Santamato A, Micello MF, Panza F, Fortunato F, Baricich A, Cisari C, et al. Can botulinum toxin type A injection technique influence the clinical outcome of patients with

- post-stroke upper limb spasticity? A randomized controlled trial comparing manual needle placement and ultrasound-guided injection techniques. *J Neurol Sci* 2014; 347: 39–43.
72. Zeuner KE, Knutzen A, Kühl C, Möller B, Hellriegel H, Margraf NG, et al. Functional impact of different muscle localization techniques for Botulinum neurotoxin A injections in clinical routine management of post-stroke spasticity. *Brain Inj* 2017; 31: 75–82.
 73. Grigoriu AI, Dinomais M, Rémy-Néris O, Brochard S. Impact of injection-guiding techniques on the effectiveness of botulinum toxin for the treatment of focal spasticity and dystonia: a systematic review. *Arch Phys Med Rehabil* 2015; 96: 2067–2078.
 74. Van Campenhout A, Bar-On L, Desloovere K, Huenaearts C, Molenaers G. Motor endplate-targeted botulinum toxin injections of the gracilis muscle in children with cerebral palsy. *Dev Med Child Neurol* 2015; 57: 476–483.
 75. Warden JM, Roberts SL, Chang Y, Baker R, Boulias C, Ismail F, et al. Neuromuscular partitioning of subscapularis based on intramuscular nerve distribution patterns: Implications for botulinum toxin injections. *Arch Phys Med Rehabil* 2014; 95: 1408–1415.
 76. Ye JF, Lee JH, An XC, Lin CH, Yue B, Han SH. Anatomic localization of motor entry points and accurate regions for botulinum toxin injection in the flexor digitorum superficialis. *Surg Radiol Anat* 2011; 33: 601–607.
 77. Parratte B, Tatu L, Vuillier F, Diop M, Monnier G. Intramuscular distribution of nerves in the human triceps surae muscle: anatomical bases for treatment of spastic drop foot with botulinum toxin. *Surg Radiol Anat* 2002; 24: 91–96.
 78. Im S, Park JH, Son SK. Does botulinum toxin injection site determine outcome in post-stroke plantarflexion spasticity? Comparison study of two injection sites in the gastrocnemius muscle: a randomized double-blind controlled trial. *Clin Rehabil* 2014; 28: 604–613.
 79. Sampaio C, Costa J, Ferreira JJ. Clinical comparability of marketed formulations of botulinum toxin. *Mov Disord* 2004; 9 Suppl 8: S129–S1136.
 80. Rosales RL, Bigalke H, Dressler D. Pharmacology of botulinum toxin: differences between type A preparations. In: *Eur J Neurol* 2006; 13 Suppl 1: 2–10.
 81. Wohlfarth K, Sycha T, Ranoux D, Naver H, Caird D. Dose equivalence of two commercial preparations of botulinum neurotoxin type A: time for a reassessment? *Curr Med Res Opin* 2009; 25: 1573–1584.
 82. Shin J, Jeon C, Ki W, YD, Kim. Clinical Comparability of Dysport and Botox in essential blepharospasm. *J Korean Ophthalmol Soc* 2009; 50: 334–335.
 83. Mohammadi B, Abdoulrahmani Balouch S, Dengler R, Kollwe K. Long-term treatment of spasticity with botulinum toxin type A: an analysis of 1221 treatments in 137 patients. *Neurol Res* 2010; 32: 309–313.
 84. Keren-Capelovitch T, Jarus T, Fattal-Valevski A. Upper extremity function and occupational performance in children with spastic cerebral palsy following lower extremity botulinum toxin injections. *J Child Neurol* 2010; 25: 694–700.
 85. Yun J, Kim J, Kim H, Chung, Kim J, Cho J, et al. Dysport and botox at a ratio of 2.5: 1 units in cervical dystonia: a double-blind, randomized study. *Mov Disord* 2015; 30: 206–213.
 86. Wohlfarth K, Göschel H, Frevert J, Dengler R, Bigalke H. Botulinum A toxins: units versus units. *Naunyn-Schmiedeberg Arch Pharmacol* 1997; 355: 335–340.
 87. Marchetti A, Magar R, Findley L, Larsen JP, Pirtosek Z, Ráuzižka E, et al. Retrospective evaluation of the dose of dysport and botox in the management of cervical dystonia and blepharospasm: the REAL DOSE study. *Mov Disord* 2005; 20: 937–944.
 88. Kollwe K, Mohammadi B, Dengler R, Dressler D. Hemifacial spasm and reinnervation synkinesias: long-term treatment with either Botox® or Dysport®. *J Neural Transm* 2010; 117: 759–763.
 89. Marion MH, Sheehy M, Sangla S, Soulayrol S. Dose standardisation of botulinum toxin. *J Neurol Neurosurg Psychiatry* 1995; 59: 102–103.
 90. Whurr R, Nye C, Lorch M. Meta-analysis of botulinum toxin treatment of spasmodic dysphonia: a review of 22 studies. *Int J Lang Commun Disord* 1998; 33 (Suppl): 327–329.
 91. Sampaio C, Ferreira JJ, Simões F, Rosas MJ, Magalhães M, Correia AP, et al. DYSBOT: a single-blind, randomized parallel study to determine whether any differences can be detected in the efficacy and tolerability of two formulations of botulinum toxin type A—dysport and botox—assuming a ratio of 4: 1. *Mov Disord* 1997; 12: 1013–1018.
 92. Odergren T, Hjaltason H, Kaakkola S, Solders G, Hanko J, Fehling C, et al. A double blind, randomised, parallel group study to investigate the dose equivalence of Dysport® and Botox® in the treatment of cervical dystonia. *J Neurol Neurosurg Psychiatry* 1998; 64: 6–12.
 93. Ranoux D, Gury C, Fondarai J, Mas JL, Zuber M. Respective potencies of botox and dysport: a double blind, randomised, crossover study in cervical dystonia. *J Neurol Neurosurg Psychiatry* 2002; 72: 459–462.
 94. Brockmann K, Schweitzer K, Beck G, Wächter T. Comparison of different preparations of botulinumtoxin a in the treatment of cervical dystonia. *Neurol Asia* 2012; 17: 15–119.
 95. Poewe W. Respective potencies of botox and dysport: a double blind, randomised, crossover study in cervical dystonia. *J Neurol Neurosurg Psychiatry* 2002; 72: 430.
 96. Tapias G, Garcia-Romero M, Crespo C, Cuesta M, Forne C, Pascual-Pascual SI. Cost-minimization analysis in the treatment of spasticity in children with cerebral palsy with botulinum toxin type A: an observational, longitudinal, retrospective study. *Farm Hosp* 2016; 40: 412–416.
 97. Benecke R, Jost WH, Kanovsky P, Ruzicka E, Comes G, Grafe S. A new botulinum toxin type A free of complexing proteins for treatment of cervical dystonia. *Neurology* 2005; 64: 1949–1951.
 98. Roggenkämper P, Jost WH, Bihari K, Comes G, Grafe S. Efficacy and safety of a new botulinum toxin type A free of complexing proteins in the treatment of blepharospasm. *J Neural Transm* 2006; 113: 303–312.
 99. Jost WH, Kohl A, Brinkmann S, Comes G. Efficacy and tolerability of a botulinum toxin type a free of complexing proteins (NT 201) compared with commercially available botulinum toxin type a (BOTOX®) in healthy volunteers. *J Neural Transm* 2005; 112: 905–913.
 100. Park J, Lee MS, Harrison AR. Profile of Xeomin® (incobotulinumtoxinA) for the treatment of blepharospasm. *Clin Ophthalmol* 2011; 5: 725–732.
 101. Zoons E, Dijkgraaf MGW, Dijk JM, van Schaik IN, Tijssen MA. Botulinum toxin as treatment for focal dystonia: a systematic review of the pharmacotherapeutic and pharmacoeconomic value. *J Neurol* 2012; 259: 2519–2526.
 102. Dressler D. Routine use of Xeomin in patients previously treated with Botox: long term results. *Eur J Neurol* 2009; 16 (Suppl 2): 2–5.
 103. Scaglione F. Conversion ratio between Botox®, Dysport®, and Xeomin® in clinical practice. *Toxins (Basel)* 2016; 8: piiE65.
 104. Grosset DG, Tyrrell EG, Grosset KA. Switch from abobotulinumtoxinA (Dysport®) to incobotulinumtoxinA (Xeomin®) botulinum toxin formulation: a review of 257 cases. *J Rehabil Med* 2015; 47: 183–186.
 105. Lai JM, Francisco GE, Willis FB. Dynamic splinting after treatment with botulinum toxin type-A: a randomized controlled pilot study. *Adv Ther* 2009; 26: 241–248.
 106. Sun S-F, Hsu C-W, Sun H-P, Hwang C-W, Yang C-L, Wang J-L. Combined botulinum toxin type A with modified constraint-induced movement therapy for chronic stroke patients with upper extremity spasticity: a randomized controlled study. *Neurorehabil Neural Repair* 2010; 24: 34–41.
 107. Weber DJ, Skidmore ER, Niyonkuru C, Chang CL, Huber

- LM, Munin MC. Cyclic functional electrical stimulation does not enhance gains in hand grasp function when used as an adjunct to onabotulinumtoxin A and Task practice therapy: a single-blind, randomized controlled pilot study. *Arch Phys Med Rehabil* 2010; 91: 679–686.
108. Demetrios M, Khan F, Turner-Stokes L, Brand C, McSweeney S. Multidisciplinary rehabilitation following botulinum toxin and other focal intramuscular treatment for post-stroke spasticity. *Cochrane Database of Syst Rev* 2013 (6): CD009689.
 109. Mills PB, Finlayson H, Sudol M, O'Connor R. Systematic review of adjunct therapies to improve outcomes following botulinum toxin injection for treatment of limb spasticity. *Clin Rehabil* 2016; 30: 537–548.
 110. Kinnear BZ, Lannin NA, Cusick A, Harvey LA, Rawicki B. Rehabilitation therapies after botulinum toxin-a injection to manage limb spasticity: a systematic review. *Phys Ther* 2014; 94: 1569–1581.
 111. Farag J, Reebye R, Ganzert C, Mills P. Does casting after botulinum toxin injection improve outcomes in adults with limb spasticity? A systematic review. *J Rehabil Med* 2020; 52: jrm00005.
 112. PEDro Physiotherapy Evidence Database. Available from: <https://www.pedro.org.au/>.
 113. Carda S1, Invernizzi M, Baricich A, Cisari C. Casting, taping or stretching after botulinum toxin type A for spastic equinus foot: a single-blind randomized trial on adult stroke patients. *Clin Rehabil* 2011; 25: 1119–1927.
 114. Farina S, Migliorini S, Gandolfi M, Bertolasi L, Casarotto M, Manganotti P, Fiaschi A, Smania N. Combined effects of botulinum toxin and casting treatments on lower limb spasticity after stroke. *Funct Neurol* 2008; 23: 87–91.
 115. Baricich A, Carda S, Bertoni M, Maderna L, Cisari C. A single-blinded, randomized study of botulinum toxin type A combined with non-pharmacological treatment for spastic foot. *J Rehabil Med* 2008; 40: 870–872.
 116. Verplancke D, Snape S, Salisbury CF, Jones PW, Ward AB. A randomized controlled trial of botulinum toxin on lower limb spasticity following acute acquired severe brain injury. *Clin Rehabil* 2005; 19: 117–125.
 117. Karadag-Saygi E E, Cubukcu-Aydoseli K, Kablan N, Ofluglu D. Role of kinesiology taping combined with botulinum toxin to reduce plantar flexors spasticity after stroke. *Top Stroke Rehabil* 2010; 17: 318–320.
 118. Reiter F, Danni M, Lagalla G, Ceravolo G, Provinciali L. Low-dose botulinum toxin with ankle taping for treatment of spastic equinovarus foot after stroke. *Arch Phys Med Rehabil* 1998; 79: 532–535.
 119. Yasar E, Tok F, Safaz I, Balaban B, Yilmaz B, Alaca R. The efficacy of serial casting after botulinum toxin A injection in improving equinovarus deformity in patients with chronic stroke. *Brain Inj* 2010; 24: 736–739.
 120. Santamato A, Micello MF, Panza F, Fortunato F, Picelli A, Smania N, Logroscino G, Fiore P, Ranieri M. Adhesive taping vs. daily manual muscle stretching and splinting after botulinum toxin type A injection for wrist and fingers spastic overactivity in stroke patients: a randomized controlled trial. *J Neurol Sci* 2015; 350: 1–6.
 121. Lee J-Y, Kim S-N, Lee I-S, Jung H, Lee K-S, Koh S-E. Effects of extracorporeal shock wave therapy on spasticity in patients after brain injury: a meta-analysis. *J Phys Ther Sci* 2014; 26: 1641–1647.
 122. Wu Y-T, Yu H-K, Chen L-R, Chang C-N, Chen Y-M, Hu G-C. Extracorporeal shock waves versus botulinum toxin type A in the treatment of post-stroke upper limb spasticity – a randomized, noninferiority trial. *Arch Phys Med Rehabil* 2018; 99: 2143–2150.
 123. Santamato A, Notarnicola A, Panza F, Ranieri M, Micello MF, Manganotti P, et al. SBOTE study: extracorporeal shock wave therapy versus electrical stimulation after botulinum toxin type a injection for post-stroke spasticity-a prospective randomized trial. *Ultrasound Med Biol* 2013; 39: 283–291.
 124. Manganotti P, Amelio E. Long-term effect of shock wave therapy on upper limb hypertonia in patients affected by stroke. *Stroke* 2005; 36: 1967–1971.
 125. Li T-Y, Chang C-Y, Chou Y-C, Chen L-C, Chu H-Y, Chiang S-L, et al. Effect of radial shock wave therapy on spasticity of the upper limb in patients with chronic stroke. *Medicine (Baltimore)* 2016; 95: e3544.
 126. Moon S, Kim M, Jung S, Son S, JH L, H S, et al. The effect of extracorporeal shock wave therapy on lower limb spasticity in. *Ann Rehabil Med* 2013; 37: 461–470.
 127. Sohn MK, Cho KH, Kim Y-J, Hwang SL. Spasticity and electrophysiologic changes after extracorporeal shock wave therapy on gastrocnemius. *Ann Rehabil Med* 2011; 35: 599–604.
 128. Picelli A, La Marchina E, Gajofatto F, Pontillo A, Vangelista A, Filippini R, et al. Sonographic and clinical effects of botulinum toxin Type A combined with extracorporeal shock wave therapy on spastic muscles of children with cerebral palsy. *Dev Neurorehabil* 2017; 20: 160–164.
 129. Wilkenfeld AJL. Review of electrical stimulation, botulinum toxin, and their combination for spastic drop foot. *J Rehabil Res Dev* 2013; 50: 315.
 130. Leo A, Naro A, Molonia F, Tomasello P, Saccà I, Bramanti A, et al. Spasticity Management: the current state of transcranial neuromodulation. *PMR* 2017; 9: 1020–1029.
 131. Naro A, Leo A, Russo M, Casella C, Buda A, Crespantini A, et al. Breakthroughs in the spasticity management: are non-pharmacological treatments the future? *J Clin Neurosci* 2017; 39: 16–27.
 132. Crea S, Cempini M, Mazzoleni S, Carrozza MC, Posteraro F, Vitiello N. Phase-II clinical validation of a powered exoskeleton for the treatment of elbow spasticity. *Front Neurosci* 2017; 11: 261.
 133. Laver KE, Schoene D, Crotty M, George S, Lannin NA, Sherrington C. Telerehabilitation services for stroke. *Cochrane Database Syst Rev* 2013; (12) CD010255.
 134. Chen J, Jin W, Zhang X-X, Xu W, Liu X-N, Ren C-C. Tele-rehabilitation approaches for stroke patients: systematic review and meta-analysis of randomized controlled trials. *J Stroke Cerebrovasc Dis* 2015; 24: 2660–2668.
 135. Chen J, Jin W, Dong WS, Jin Y, Qiao FL, Zhou YF, et al. Effects of home-based telesupervising rehabilitation on physical function for stroke survivors with hemiplegia. *Am J Phys Med Rehabil* 2017; 96: 152–160.
 136. Roche N, Zory R, Sauthier A, Bonnyaud C, Pradon D, Bensmail D. Effect of rehabilitation and botulinum toxin injection on gait in chronic stroke patients: a randomized controlled study. *J Rehabil Med* 2015; 47: 31–37.
 137. Marsal C, Gracies JM, Dean C, Mesure S, Bayle N. Beliefs of rehabilitation professionals towards guided self-rehabilitation contracts for post stroke hemiparesis. *Top Stroke Rehabil* 2017; 24: 68–613.