

Improved Outcome in Children With Newly Diagnosed High-Risk Neuroblastoma Treated With Chemoimmunotherapy: Updated Results of a Phase II Study Using hu14.18K322A

Wayne L. Furman, MD¹; Beth McCarville, MD²; Barry L. Shulkin, MD²; Andrew Davidoff, MD²; Matthew Krasin, MD²; Chia-Wei Hsu, MS²; Haitao Pan, PhD²; Jianrong Wu, PhD³; Rachel Brennan, MD¹; Michael W. Bishop, MD¹; Sara Helmig, MD¹; Elizabeth Stewart, MD¹; Fariba Navid, MD⁴; Brandon Triplett, MD²; Victor Santana, MD²; Teresa Santiago, MD²; Jacquelyn A. Hank, PhD⁵; Stephen D. Gillies, PhD⁶; Alice Yu, MD, PhD^{7,8}; Paul M. Sondel, MD, PhD⁹; Wing H. Leung, MD, PhD⁹; Alberto Pappo, MD¹; and Sara M. Federico, MD¹

PURPOSE We evaluated whether combining a humanized antidisialoganglioside monoclonal antibody (hu14.18K322A) throughout therapy improves early response and outcomes in children with newly diagnosed high-risk neuroblastoma.

PATIENTS AND METHODS We conducted a prospective, single-arm, three-stage, phase II clinical trial. Six cycles of induction chemotherapy were coadministered with hu14.18K322A, granulocyte-macrophage colony-stimulating factor (GM-CSF), and low-dose interleukin-2 (IL-2). The consolidation regimen included busulfan and melphalan. When available, an additional cycle of parent-derived natural killer cells with hu14.18K322A was administered during consolidation (n = 31). Radiation therapy was administered at the end of consolidation. Postconsolidation treatment included hu14.18K322A, GM-CSF, IL-2, and isotretinoin. Early response was assessed after the first two cycles of induction therapy. End-of-induction response, event-free survival (EFS), and overall survival (OS) were evaluated.

RESULTS Sixty-four patients received hu14.18K322A with induction chemotherapy. This regimen was well tolerated, with continuous infusion narcotics. Partial responses (PRs) or better after the first two chemoimmunotherapy cycles occurred in 42 of 63 evaluable patients (66.7%; 95% CI, 55.0 to 78.3). Primary tumor volume decreased by a median of 75% (range, 100% [complete disappearance]-5% growth). Median peak hu14.18K322A serum levels in cycle one correlated with early response to therapy ($P = .0154$, one-sided t -test). Sixty of 62 patients (97%) had an end-of-induction partial response or better. No patients experienced progressive disease during induction. The 3-year EFS was 73.7% (95% CI, 60.0 to 83.4), and the OS was 86.0% (95% CI, 73.8 to 92.8), respectively.

CONCLUSION Adding hu14.18K322A to induction chemotherapy improved early objective responses, significantly reduced tumor volumes in most patients, improved end-of-induction response rates, and yielded an encouraging 3-year EFS. These results, if validated in a larger study, may be practice changing.

J Clin Oncol 40:335-344. © 2021 by American Society of Clinical Oncology

INTRODUCTION

Treatment for high-risk (HR) neuroblastoma includes induction therapy (chemotherapy and surgery), consolidation with high-dose chemotherapy followed by autologous hematopoietic stem-cell transplant (ASCT), radiotherapy, and postconsolidation treatment with isotretinoin and a monoclonal antibody (mAb) that targets the disialoganglioside GD2. Administration of a chimeric anti-GD2 mAb (dinutuximab) with granulocyte-macrophage colony-stimulating factor (GM-CSF), interleukin-2 (IL-2), and isotretinoin in the postconsolidation setting significantly improved 2- and 5-year event-free survival (EFS; 66% v 46%, $P = .01$ ¹;

$56.6 \pm 4.7\%$ v $46.1 \pm 5.1\%$, $P = .042$,² respectively). However, nearly half of all patients experienced relapse and succumb to disease.

Dinutuximab has been administered in postconsolidation to avoid chemotherapy-induced immunosuppression since it can adversely affect antibody-dependent cell-mediated cytotoxicity (ADCC). However, preclinical studies in neuroblastoma models and adult clinical studies demonstrated that concurrent chemotherapy with various monoclonal antibodies provides additive or synergistic benefits.³⁻¹⁰ We postulated that incorporating an anti-GD2 mAb throughout therapy for the treatment of HR neuroblastoma would improve clinical responses

ASSOCIATED CONTENT

Appendix Protocol

Author affiliations and support information (if applicable) appear at the end of this article.

Accepted on November 1, 2021 and published at ascopubs.org/journal/jco on December 6, 2021; DOI <https://doi.org/10.1200/JCO.21.01375>

CONTEXT

Key Objective

Despite aggressive multimodal treatment, nearly half of all patients with newly diagnosed high-risk (HR) neuroblastoma still succumb to their disease. Will the combination of a humanized antidiisialoganglioside monoclonal antibody (hu14.18K322A) with induction chemotherapy improve early response rates after two cycles and improve outcomes in children with newly diagnosed HR neuroblastoma?

Knowledge Generated

Early responses were significantly improved when compared with a historical group of children treated with identical chemotherapy, without hu14.18K322A. None of the patients in our trial experienced tumor progression during induction chemotherapy, and the overall outcomes are very encouraging.

Relevance

These results, if validated in a larger study, may change the standard of care for children with HR neuroblastoma.

and outcomes. We used a unique humanized anti-GD2 mAb, hu14.18K322A, that retains the binding specificity of dinutuximab, is 98% human, has a single point mutation to reduce complement-associated pain, and is produced in a YB2/O rat myeloma cell line to reduce fucosylation and enhance ADCC.¹¹ We demonstrated that combining hu14.18K322A with chemotherapy is tolerable and clinically active in patients with relapsed neuroblastoma.¹² Therefore, hu14.18K322A was administered concurrently with chemotherapy during induction in this phase II (NB2012), single-institution setting to assess efficacy and determine if chemoimmunotherapy warrants further evaluation in a randomized phase III study. The results of the first 42 patients, demonstrating significant improvement in early responses, were previously published.¹³ This report details the results of the entire cohort of 64 patients, their end-of-induction responses, estimates of event-free and overall survival (OS), and information on human antihuman antibody (HABA) development.

In ANBL02P1, the Children's Oncology Group (COG) investigators reported a 40% partial response (PR) rate after two cycles of induction therapy (cyclophosphamide and topotecan) in HR patients¹⁴; this response was similar to the response described in ANBL0532 in which patients received the same induction regimen.¹⁵ NB2012 used the identical chemotherapy induction regimen as ANBL02P1 and ANBL0532. The primary objectives of NB2012 were to assess response after two cycles of chemoimmunotherapy compared with a previous study with chemotherapy only and to estimate the EFS when hu14.18K322A was added to all phases of treatment. Changes in Curie scores (CSs) from diagnosis to end-of-induction chemotherapy were also assessed. Post-induction CSs have been previously validated as prognostic in two large independent cohorts of patients.¹⁶⁻¹⁸

PATIENTS AND METHODS

Study Design

This study was an open-label, single-arm phase II trial. The primary objective was to compare the response rates of the

first two cycles of chemoimmunotherapy (cyclophosphamide, topotecan, and hu14.18K322A, followed by GM-CSF and IL-2) with the response rates of two cycles of cyclophosphamide and topotecan alone reported in a previous study (ANBL02P1) in children with newly diagnosed HR neuroblastoma.¹⁴ At an interim analysis, the response rate exceeded the a priori benchmark for activity¹³; therefore, enrollment was expanded to estimate EFS as a primary objective.

Statistical Analysis Plan

In the COG phase III randomized trial ANBL0532,¹⁵ the chemotherapy components of the induction regimen were identical to NB2012 and the 95% CI of 3-year EFS was (46.5%, 55.3%). A Kaplan-Meier estimate-based design was used to test the null hypothesis of a 46% 3-year EFS rate, against the desired alternative 3-year EFS rate of 62%. A total of 61 patients provided 80% power to detect a 16% increase of 3-year EFS with a one-sided type I error rate of 5%. Analysis of EFS was performed in the intention-to-treat population, defined as all enrolled patients, and the efficacy-evaluable population, defined as all patients who had received at least one dose of study treatment and had at least one available postbaseline tumor assessment. EFS and OS were analyzed using the Kaplan-Meier method. ORR and 95% CIs were calculated using the Clopper-Pearson method. Summary statistics were provided for clinical and demographic characteristics and for AEs. The association between the EFS and the peak and trough level of hu14.18K322A were explored. We performed all statistical tests using SAS (version 9.4).

Patients

Children (< 19 years) with newly diagnosed HR neuroblastoma were eligible for enrollment. Diagnosis, staging, and response assessments were performed according to the 1993 International Criteria for Neuroblastoma Response,¹⁹ and HR neuroblastoma was defined by the criteria used by the COG.²⁰

This trial (NCT01857934) was approved by our institutional review board and opened in May 2013. The last patient




FIG 1. Study schema. BuMel, busulfan and melphalan; GM-CSF, granulocyte-macrophage colony-stimulating factor; IL-2, interleukin-2; NK, natural killer; PBSC, peripheral blood stem cell; XRT, radiation therapy.

completed therapy in July 2019. Written informed consent was obtained from participants in accordance with institutional guidelines (Protocol, online only).

Hu14.18K322A

The hu14.18K322A production cell line was provided by Merck Serono (Darmstadt, Germany) and manufactured for clinical use by the Children's GMP, LLC (Memphis, TN). With each cycle of therapy, hu14.18 K322A levels were measured by ELISA in serum obtained 1 hour after the first antibody infusion, as previously described.^{12,21} Patients were monitored for the development of HAMA before first dose of hu14.18K322A with every treatment cycle (induction, consolidation, and postconsolidation phase) with an ELISA as previously described.²²

Treatment

The schedule and dosages of induction chemotherapy agents cyclophosphamide, topotecan, cisplatin, etoposide, doxorubicin, and vincristine were identical to those used in ANBL02P1 and ANBL0532.^{14,15} Four daily doses of hu14.18K322A (days 2-5; administered over 4 hours, when possible) at 40 mg/m²/dose (once daily) were given with each cycle of induction chemotherapy. All patients received continuous infusions of opioids at standard dosages, approximately 30 minutes before antibody infusions. Systemic steroids were not allowed. Each cycle was followed by daily subcutaneous GM-CSF (250 μg/m² once per day) through the nadir until an absolute neutrophil count of ≥ 2,000/mm³ and subcutaneous IL-2 (10⁶ units/m²), once every other day for six doses (Fig 1).

Hematopoietic progenitor cells were collected after induction cycle 2 or 4. Primary tumors were resected at any time point after induction cycle 2, when the surgeon determined that a maximum resection could be safely achieved. Consolidation therapy included ASCT with a

Busulfan and Melphalan (BuMel)-conditioning regimen²³ and, for consenting patients, experimental therapy with an additional cycle of daily hu14.18K322A administration beginning 2 days (days 2-5) after ASCT (given on day 0) and parental natural killer (NK) cell infusions (when available), derived as previously described.²³ Tolerability of hu14.18K322A during recovery from consolidation therapy was evaluated in two dosage cohorts (25 mg/m² once per day for 4 days, n = 16 [with NK cells, n = 10]; 40 mg/m² once per day for 4 days, n = 26; with NK cells [n = 21]). The NK cells were infused 4 days after ASCT. When two of the first 42 patients developed a hemophagocytic lymphohistiocytosis-like syndrome,²⁴ the additional cycle of hu14.18K322A during consolidation was eliminated in the subsequent cohort (n = 17). After recovering from ASCT (typically within 43 days of stem-cell reinfusion), patients received intensity-modulated radiation therapy or scanned proton beam radiation therapy (2,340 cGy in 180 cGy fractions). Those with macroscopic residual disease after induction chemotherapy received an additional 720 cGy to those sites.

Radiation therapy was delivered to the presurgical gross tumor volume and sites of residual disease on the basis of imaging performed at the end of induction. Post-consolidation therapy with hu14.18K322A, GM-CSF, IL-2, and isotretinoin (doses and schedules identical to those reported by Yu et al¹) was started within 100 days from ASCT. Hu14.18K322A (40 mg/m² once per day for 4 days) was administered in place of dinutuximab.¹ Adverse events were assessed by the National Cancer Institute Common Terminology Criteria for Adverse Events (version 4.0). If narcotic infusions required increased dosages, pain was designated grade 3. Diagnosis, staging, and response assessments were performed according to the 1993 International Criteria for Neuroblastoma Response.¹⁹ Metastatic complete response was defined as no skeletal uptake

TABLE 1. Patient Characteristics at Diagnosis

Characteristic	NB2012 (N = 64), No. (%)
Age, months	
< 18	5 (8) ^a
≥ 18	59 (92)
Sex	
Female	27 (42)
Male	37 (58)
Race	
White	43 (67)
Black	18 (28)
Others	3 (5)
Unreported	—
INSS stage	
IIb	1 (2)
III	7 (11) ^b
IV	56 (87)
INRG stage	
L1	0
L2	8 (13) ^c
M	56 (87)
MYCN status	
Not amplified	43 (67)
Amplified	21 (33)
Shimada histology	
Favorable	6 (9)
Unfavorable	41 (64)
Not performed	17 (27)

Abbreviations: INRG, International Neuroblastoma Risk Group; INSS, International Neuroblastoma Staging System; UH, unfavorable histology.

^aThree with *MYCN* amplification and two without *MYCN* amplification but DNA index = 1.

^bFive with *MYCN* amplification and two without *MYCN* amplification but with UH.

^c6 of 8 *MYCN* amplified; 8 of 8 with unfavorable histology.

on nuclear imaging and complete response in bone marrow and other metastatic sites. Responses were evaluated after induction cycles 2 and 6, after consolidation, and at the end of therapy. Anatomic imaging was assessed by one study radiologist (M.B.M.). Semiquantitative MIBG scoring (ie, CS) was assessed by one nuclear medicine physician (B.S.) at diagnosis and after cycles 2 and 6 of induction chemotherapy as described by Yanik et al.¹⁸

RESULTS

Patient Characteristics

Sixty-three of the 64 patients had measurable or evaluable disease at enrollment. Patient characteristics are described in Table 1. Most patients were male (n = 37), White

(n = 43), and ≥ 18 months (n = 59), with a median age of 3.1 years (range, 0.5-15.2 years), and had International Neuroblastoma Staging System (INSS) stage IV or International Neuroblastoma Risk Group stage M disease (n = 56). These characteristics were similar to patients enrolled on COG studies ANBL02P1,¹⁴ ANBL0532,¹⁵ and ANBL12P1²⁵ (Appendix Table A1, online only). During consolidation, 25 patients received proton beam radiation therapy and 34 patients received intensity-modulated radiation therapy.

Safety and Toxicity

The addition of hu14.18K322A to induction chemotherapy was well tolerated, with continuous infusion narcotics adjusted to patient tolerance. Therapy-related grade 3 or 4 toxicities during induction chemotherapy-hu14.18K322A coadministration are described in Appendix Table A2 (online only). Toxicities attributed to antibody infusion included pain (10%; 38 patient episodes in 379 cycles), hypotension (3%; 12 of 379), cough (1%; 4 of 379), hypoxia (6%; 21 of 379), and infusion-related reactions (4%; 14 of 379).^{1,26} Events related to myelosuppression, liver function abnormalities, enterocolitis, and fever with

TABLE 2. Responses and Curie Scores During Induction Chemoimmunotherapy

Response	Responses			
	After Two Cycles (N = 64)	Metastatic Responses After Two Cycles	After Six Cycles (N = 62 evaluable ^c)	Metastatic Responses after Six Cycles
CR	2 (3%)	mCR 3/56	21 (34%)	mCR 40 (65%)
VGPR	8 (13%)	< mCR 53/56	21 (34%)	< mCR 22 (35%)
PR	32 (51%)		18 (29%)	
NR	21 (33%)		2 (3%)	
PD	0	0	0	0
NE	1 ^a	8/64 ^b	2 ^c	2 ^c
CSs				
CS at Diagnosis (N = 54^d)		CS After Six Cycles (N = 54^d)		
0-12	20 (37%)	0-2	47 (87%)	
13-25	23 (43%)	3-6	3 (5.6%)	
> 25	11 (20%)	≥ 13	4 (7.4%)	

Abbreviations: CR, complete response; CS, Curie score; INSS, International Neuroblastoma Staging System; mCR, metastatic complete response; NE, non evaluable; NR, no response; PD, progressive disease; PR, partial response; VGPR, very good partial response.

^aOne patient with *MYCN* amplification and localized tumor had resection of her primary tumor prior to beginning chemotherapy.

^bEight patients had no metastases at diagnosis.

^cOne patient was taken off study prior to completing induction chemotherapy because of severe diarrhea from a vasoactive intestinal peptide secreting tumor and one patient withdrew electively for parental preference.

^dFifty-six patients had INSS IV disease, two withdrew prior to completing induction chemotherapy.

or without infection were similar to those reported previously.^{13,14}

Tumor Response

Responses were evaluated using the International Neuroblastoma Response Criteria (INRC)¹⁹ after two cycles of therapy (early response) and at the end of induction (Table 2). Two patients had resection of their primary tumors before beginning therapy but had other sites of evaluable disease. Of the 63 children evaluable for early response, 42 (66.7%; 95% CI, 55.0 to 78.3%) experienced PR or better. Figure 2 demonstrates the INRC components of response for

individual patients after induction cycle two and the end of induction. All but one patient had a decrease in their primary tumor volume at cycle 2 with 42 of 63 achieving reductions of at least 50%. Fourteen had a metastatic very good partial response after induction cycle 2, and 3 had metastatic site complete responses. At the end of induction, 60 of 62 evaluable patients had PRs or better (96.8%; 95% CI, 88.8.0 to 99.6) including 40 with mCRs (65%).

MIBG scoring (CS) was performed at every evaluation time point. CSs at diagnosis for the 56 patients with INSS stage IV disease ranged from 1 to 28 (median, 15) and from 0 to 23




FIG 2. Disease response assessed by INRC for patients (A) after induction cycle 2 and (B) at the end of induction. Each bar represents an individual patient. The primary tumor response is demonstrated as a percentile change in volume of the primary tumor compared with the primary tumor volume at baseline. The color of the individual bars signifies the metastatic response including anatomic imaging, functional imaging (MIBG or PET), and bone marrow assessment. The blue bar represents a complete metastatic response. The red bar represents a metastatic VGPR. The teal bar represents a metastatic partial response. The orange bar represents no metastatic response. The purple bar signifies patients without metastatic disease at diagnosis. Bone marrow response is depicted by the following shapes: gray triangle represents the presence of bone marrow disease at the time of the assessment, blue oval represents bone marrow disease present at baseline that resolved at the time of the assessment, and the red star represents no bone marrow disease at baseline. Together, the primary tumor response, metastatic response, and bone marrow response are used to inform the INRC overall response. The overall response of the individual patient is presented by the letter below each bar. The letter C signifies a CR; V is a VGPR; P is a PR; N is a no response; and E means that the patient was not evaluable at that time point because of the primary tumor resection performed before therapy. X represents a stage IV patient who did not have a primary tumor at diagnosis. O represents two patients who discontinued protocol therapy before completing induction. CR, complete response; INRC, International Neuroblastoma Response Criteria; MIBG, [¹²⁵I]metaiodobenzylguanidine; PET, positron emission tomography; PR, partial response; VGPR, very good partial response.




FIG 3. Association of hu14.18K322A peak serum levels ($\mu\text{g/mL}$) and response after two cycles of chemoimmunotherapy. Relationship between response (CR or PR [$n = 42$] v SD or PD [$n = 17$]) and peak hu14.18K322A levels, measured on first day of mAb infusion, 1 hour after a 4-hour infusion. $P = .0154$ (one-tailed t -test). CR, complete response; mAb, monoclonal antibody; PD, progressive disease; PR, partial response; SD, stable disease.

(median, 0) at the end-of-induction chemotherapy. For the 54 patients with INSS stage IV disease who completed induction, 47 (87%) had end-of-induction CSs of two or less, as compared with a median CS of 15 at diagnosis (Table 2).

Quantitative and qualitative differences in NK cells on a subset of these children were also examined and previously reported.²⁷

Hu14.18K322A Levels and HAHA Response

Median peak hu14.18K322A serum levels on cycle 1 of induction correlated with the clinical responses that were performed after the first two cycles of induction chemoimmunotherapy (PR or greater; $n = 42$) versus those who had stable disease ($n = 17$; $P = .054$, one-sided Wilcoxon rank-sum test; Fig 3). There was no association of EFS with the peak level of hu14.18K322A in cycle 1 (peak levels were dichotomized, above median or below median, and compared using the log-rank test [$P = .682$], and there was no association of early response after two cycles [PR or greater] with the trough level of hu14.18K322 A at the start of cycle 2 [$P = .336$, two-sided Wilcoxon rank-sum test]).

Five of 64 evaluable patients (7.8%) developed significant HAHA responses to hu14.18K322A after chemoimmunotherapy. None had a HAHA response associated

with a drop in the peak hu14.18K322A level in cycles after the HAHA development (Appendix Table A3, online only). The five HAHA+ patients did not show any significant difference in EFS ($P = .75$, log-rank test) or noticeable differences in tolerance to antibody than the 59 HAHA-patients.

Effect of an Additional Cycle of hu14.18K322A With or Without Parentally Derived NK Cells During Consolidation

Fifty-nine of 64 patients received consolidation with BuMel (Fig 4) as previously described.²³ Forty-two of these received an additional cycle of hu14.18K322A (25 mg/m² once per day \times 4 [$n = 16$] or 40 mg/m² once per day \times 4 [$n = 26$]), and 31 also received parentally derived NK cells. Seventeen patients received BuMel without the additional cycle of hu14.18K322A with or without NK cells. There was no difference in EFS between patients who received ASCT with BuMel and those who received ASCT with BuMel or hu14.18K322A with or without NK cells (data not shown).

Survival Outcomes

As of November 4, 2020, 54 of the 64 patients were alive, with a median follow-up time for the 48 patients who did not experience an event of 4.3 years (range, 1.1-6.7 years). Two patients died of complications from therapy without




FIG 4. Flow diagram. ASCT, autologous hematopoietic stem-cell transplant; mAb, monoclonal antibody; MRD, minimal residual disease; PD, progressive disease.

evidence of disease (one because of transplant-related complications and the other because of non-neutropenic sepsis); one progressed after consolidation, four had persistent disease after completing all therapy; one patient experienced disease recurrence after postconsolidation minimal residual disease cycle 3; and eight patients experienced relapse at 4, 4, 6, 15, 15, 15, 18, and 31 months, respectively, after completing all therapy. Four of these patients, one with persistent disease and three with relapsed disease, also died of disease progression. The overall 3-year EFS and OS rates were 73.7% (95% CI, 60.0 to 83.4) and 86% (95% CI, 73.8 to 92.8), respectively (Fig 5).

DISCUSSION

We have demonstrated that the addition of hu14.18K322A to induction chemotherapy significantly improves early responses in patients with newly diagnosed HR neuroblastoma (66.7% v 39.1%; 95% CI 55.0 to 78.3 v 35.3 to 43.0; $P < .0001$) compared with ANBL0532.¹⁵ Additionally, our chemoimmunotherapy induction regimen resulted in CSs ≤ 2 in 47 of 54 patients with INSS IV disease (87%), a finding that has been correlated with improved outcomes in both COG and SIOPEN trials.^{17,18} The significant improvement in early responses and postinduction CSs

suggests that hu14.18K322A, GM-CSF, and IL-2 added to induction chemotherapy could significantly affect outcome. Although our sample size was small, our encouraging 3-year EFS of 73.7% (95% CI, 60 to 83.4) compares favorably with recent COG, GPOH, and SIOPEN trials (Appendix Table A4, online only).^{15,25,28,29} Park et al¹⁵ reported an end-of-induction (EoI) Objective Response (\geq PR) of 75.2% and a 3-year EFS from the time of random assignment for tandem transplant of 51.1% (95% CI, 47.1 to 55.0) for all 652 eligible patients.¹⁵ In our trial, the EoI Objective Response was 97%. More recently, Granger et al²⁵ evaluated BuMeI ASCT after a five-cycle COG induction regimen and reported an EoI Objective Response of 80% and a 3-year EFS of 55.6% \pm 4.2% for all 146 patients enrolled (Appendix Table A4).²⁵ Comparing NB2012 results with ANBL0032,^{1,15,25} ANBL0532¹⁵ and ANBL12P1²⁵ are complicated by the facts that in the study by Yu et al,¹ only patients who achieved a PR or better to induction therapy were included, Park et al¹⁵ only randomly assigned patients at end of induction and only 355 of 652 patients originally enrolled were randomly assigned, and in Granger's report, 39 of 146 patients (26.7%) discontinued treatment early.²⁵ The NB2012 EFS estimate begins at the time of enrollment and includes all patients. Importantly, we did not observe disease progression in any patients during induction,




FIG 5. EFS and OS for all patients enrolled. EFS, event-free survival; OS, overall survival.

whereas 7%-14% of patients enrolled in previous COG trials experienced disease progression before receiving ASCT.^{15,30}

Peak serum levels of hu14.18K322A during the induction and postconsolidation cycles generally exceeded the level of antibody expected to facilitate ADCC in vivo.³¹ Higher levels of serum hu14.18K322A in cycle 1 were associated with more rapid response to treatment (PR or better after two cycles), consistent with dinutuximab plus temozolomide and irinotecan in relapsed neuroblastoma.³² These findings may suggest that greater in vivo exposure to an anti-GD2 mAb might be associated with improved outcomes. Although there was no significant association of hu14.18K322A levels with EFS, this might reflect the overall favorable EFS for the patients in this trial, which limits the statistical power to associate differences in hu14.18K322A levels with favorable versus unfavorable outcomes. The incidence of HAHA was substantially less than that seen in our phase I trial (5 of 64 v 15 of 37 patients [40.5%; $P < .0001$, Fisher's exact test]).¹¹ Furthermore, the HAHA response here seemed weaker even in the five patients who developed HAHA (on the basis of the absence of any drop in peak hu14.18K322A levels for these five HAHA+ patients) compared with that observed in our previous phase I trial.¹¹ We think that this difference likely reflects the immunosuppressive effect of concurrent chemotherapy on the induction of an antidrug-antibody response (like HAHA or HACA), as also seen recently in a COG trial of dinutuximab plus temozolomide and irinotecan.³²

Although promising, these results must be interpreted within the context of a single-institution phase II study. Although the chemotherapy backbone was identical and the major prognostic factors of our patient populations were comparable with those of ANBL02P1,¹⁴ ANBL0532,¹⁵ and ANBL12P1²⁵ (Appendix Table A1), several other differences in therapy might have contributed to the excellent NB2012 response rates. For example, GM-CSF was used during induction chemoimmunotherapy, rather than filgrastim, both for its ability to enhance ADCC^{33,34} and for primary prophylaxis of febrile neutropenia.³⁵ Additionally, low-dose IL-2 was used for its ability to enhance ADCC.³⁶ The BuMel preparative regimen and additional cycle of hu14.18K322A ($n = 42$) with parental-derived NK cells ($n = 31$ of 42) during consolidation might have also affected EFS. Although BuMel was found to be a superior consolidation regimen within the context of rapid COJEC induction of SIOPEN,³⁷ an arm evaluating its efficacy within the COG induction on the current open COG HR trial ANBL1531 was recently closed because of inferior outcomes. NB2012 was not designed to evaluate the effects of these modifications on the overall outcome.

In conclusion, the addition of hu14.18K322A to induction chemotherapy resulted in a near doubling of early responses, major reductions in tumor volume, improved responses, and a very encouraging 3-year EFS of 73.7% (95% CI, 60 to 83.4). These results, if validated in a larger study, may change the standard of care for children with HR neuroblastoma.

AFFILIATIONS

- ¹Department of Oncology, St Jude Children's Research Hospital, Memphis, TN
²St Jude Children's Research Hospital, Memphis, TN
³University of Kentucky, Lexington, KY
⁴Department of Pediatrics, Children's Hospital Los Angeles, University of Southern California, Los Angeles, CA
⁵Departments of Pediatrics and Human Oncology, University of Wisconsin, Madison, WI
⁶Provenance Biopharmaceuticals, Carlisle, MA
⁷University of California San Diego, San Diego, CA
⁸Institute of Stem Cell and Translational Cancer Research, Chang Gung Memorial Hospital at Linkou and Chang Gung University, Taiwan
⁹Department of Pediatrics, University of Hong Kong, Hong Kong

CORRESPONDING AUTHOR

Wayne L. Furman, MD, St Jude Children's Research Hospital, 262 Danny Thomas Place, MS 260, Memphis, TN 38105; e-mail: wayne.furman@stjude.org.

PRIOR PRESENTATION

Presented in part at the ASCO Annual Meeting, June 3-7, 2016, Chicago, IL; ASCO Annual Meeting, June 1-5, 2017, Chicago, IL; and at Advances in Neuroblastoma Research virtual meeting, January 25-27, 2021.

SUPPORT

Supported by St Jude Children's Research Hospital Comprehensive Cancer Center Support Grant (2 P30 CA021765), American Lebanese Syrian Associated Charities, and Cookies for Kids' Cancer and Cure Childhood Cancer Foundation.

CLINICAL TRIAL INFORMATION

NCT01857934 (NB2012)

REFERENCES

1. Yu AL, Gilman AL, Ozkaynak MF, et al: Anti-GD2 antibody with GM-CSF, interleukin-2, and isotretinoin for neuroblastoma. *N Engl J Med* 363:1324-1334, 2010
2. Yu AL, Gilman AL, Ozkaynak MF, et al: Long-term follow-up of a phase III study of ch14.18 (dinutuximab) + cytokine immunotherapy in children with high-risk neuroblastoma: COG study ANBL0032. *Clin Cancer Res* 27:2179-2189, 2021
3. Nowak AK, Robinson BW, Lake RA: Synergy between chemotherapy and immunotherapy in the treatment of established murine solid tumors. *Cancer Res* 63:4490-4496, 2003
4. Lake RA, Robinson BW: Immunotherapy and chemotherapy—A practical partnership. *Nat Rev Cancer* 5:397-405, 2005
5. Hiddemann W, Kneba M, Dreyling M, et al: Frontline therapy with rituximab added to the combination of cyclophosphamide, doxorubicin, vincristine, and prednisone (CHOP) significantly improves the outcome for patients with advanced-stage follicular lymphoma compared with therapy with CHOP alone: Results of a prospective randomized study of the German Low-Grade Lymphoma Study Group. *Blood* 106:3725-3732, 2005
6. Coiffier B, Lepage E, Briere J, et al: CHOP chemotherapy plus rituximab compared with CHOP alone in elderly patients with diffuse large-B-cell lymphoma. *N Engl J Med* 346:235-242, 2002
7. Hurwitz H, Fehrenbacher L, Novotny W, et al: Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl J Med* 350:2335-2342, 2004
8. Slamon DJ, Leyland-Jones B, Shak S, et al: Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med* 344:783-792, 2001
9. Yoshida S, Kawaguchi H, Sato S, et al: An anti-GD2 monoclonal antibody enhances apoptotic effects of anti-cancer drugs against small cell lung cancer cells via JNK (c-Jun terminal kinase) activation. *Jpn J Cancer Res* 93:816-824, 2002
10. Kowalczyk A, Gil M, Horwacik I, et al: The GD2-specific 14G2a monoclonal antibody induces apoptosis and enhances cytotoxicity of chemotherapeutic drugs in IMR-32 human neuroblastoma cells. *Cancer Lett* 281:171-182, 2009
11. Navid F, Sondel PM, Barfield R, et al: Phase I trial of a novel anti-GD2 monoclonal antibody, Hu14.18K322A, designed to decrease toxicity in children with refractory or recurrent neuroblastoma. *J Clin Oncol* 32:1445-1452, 2014
12. Federico SM, McCarville MB, Shulkin BL, et al: A pilot trial of humanized anti-GD2 monoclonal antibody (hu14.18K322A) with chemotherapy and natural killer cells in children with recurrent/refractory neuroblastoma. *Clin Cancer Res* 23:6441-6449, 2017
13. Furman WL, Federico SM, McCarville MB, et al: A phase II trial of Hu14.18K322A in combination with induction chemotherapy in children with newly diagnosed high-risk neuroblastoma. *Clin Cancer Res* 25:6320-6328, 2019
14. Park JR, Scott JR, Stewart CF, et al: Pilot induction regimen incorporating pharmacokinetically guided topotecan for treatment of newly diagnosed high-risk neuroblastoma: A Children's Oncology Group study. *J Clin Oncol* 29:4351-4357, 2011

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/JCO.21.01375>.

AUTHOR CONTRIBUTIONS

Conception and design: Wayne L. Furman, Beth McCarville, Matthew Krasin, Jianrong Wu, Fariba Navid, Victor Santana, Stephen D. Gillies, Alice Yu, Paul M. Sondel, Wing H. Leung, Alberto Pappo, Sara M. Federico

Administrative support: Jacquelyn A. Hank

Provision of study materials or patients: Wayne L. Furman, Beth McCarville, Andrew Davidoff, Matthew Krasin, Rachel Brennan, Elizabeth Stewart, Brandon Triplett, Stephen D. Gillies, Alberto Pappo, Sara M. Federico

Collection and assembly of data: Wayne L. Furman, Beth McCarville, Barry L. Shulkin, Matthew Krasin, Rachel Brennan, Sara Helmig, Elizabeth Stewart, Jacquelyn A. Hank, Paul M. Sondel, Alberto Pappo, Sara M. Federico

Data analysis and interpretation: Wayne L. Furman, Beth McCarville, Barry L. Shulkin, Andrew Davidoff, Matthew Krasin, Chia-Wei Hsu, Haitao Pan, Jianrong Wu, Michael W. Bishop, Brandon Triplett, Victor Santana, Teresa Santiago, Jacquelyn A. Hank, Paul M. Sondel, Wing H. Leung, Alberto Pappo, Sara M. Federico

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

ACKNOWLEDGMENT

The authors thank Gwen Anthony, Deanna Welsh, clinical and laboratory personnel, operations staff, and all the patients and their families who participated in this study.

15. Park JR, Kreissman SG, London WB, et al: Effect of tandem autologous stem cell transplant vs single transplant on event-free survival in patients with high-risk neuroblastoma: A randomized clinical trial. *JAMA* 322:746-755, 2019
16. Yanik G, Naranjo A, Parisi MT, et al: Impact of post-induction Curie scores in high-risk neuroblastoma. *Biol Blood Marrow Transplant* 21:S107, 2015
17. Yanik GA, Parisi MT, Naranjo A, et al: Validation of postinduction Curie scores in high-risk neuroblastoma: A Children's Oncology Group and SIOPEN Group report on SIOPEN/HR-NBL1. *J Nucl Med* 59:502-508, 2018
18. Yanik GA, Parisi MT, Shulkin BL, et al: Semiquantitative mIBG scoring as a prognostic indicator in patients with stage 4 neuroblastoma: A report from the Children's Oncology Group. *J Nucl Med* 54:541-548, 2013
19. Brodeur GM, Pritchard J, Berthold F, et al: Revisions of the international criteria for neuroblastoma diagnosis, staging, and response to treatment. *J Clin Oncol* 11:1466-1477, 1993
20. Maris JM: The biologic basis for neuroblastoma heterogeneity and risk stratification. *Curr Opin Pediatr* 17:7-13, 2005
21. Hank JA, Gan J, Ryu H, et al: Immunogenicity of the hu14.18-IL2 immunocytokine molecule in adults with melanoma and children with neuroblastoma. *Clin Cancer Res* 15:5923-5930, 2009
22. Albertini MR, Gan J, Jaeger P, et al: Systemic interleukin-2 modulates the anti-idiotypic response to chimeric anti-GD2 antibody in patients with melanoma. *J Immunother Emphasis Tumor Immunol* 19:278-295, 1996
23. Talleur AC, Triplett BM, Federico S, et al: Consolidation therapy for newly diagnosed pediatric patients with high-risk neuroblastoma using busulfan/melphalan, autologous hematopoietic cell transplantation, anti-GD2 antibody, granulocyte-macrophage colony-stimulating factor, interleukin-2, and haploidentical natural killer cells. *Biol Blood Marrow Transplant* 23:1910-1917, 2017
24. Epperly R, Furman W, Hines M, et al: Secondary hemophagocytic syndrome after autologous hematopoietic cell transplant and immune therapy for neuroblastoma. *Pediatr Blood Cancer* 66:e27964, 2019
25. Granger MM, Naranjo A, Bagatell R, et al: Myeloablative busulfan/melphalan consolidation following induction chemotherapy for patients with newly diagnosed high-risk neuroblastoma: Children's Oncology Group trial ANBL12P1. *Transpl Cell Ther* 27:490.e1-490.e8, 2021
26. Ozkaynak MF, Gilman AL, London WB, et al: A comprehensive safety trial of chimeric antibody 14.18 with GM-CSF, IL-2, and isotretinoin in high-risk neuroblastoma patients following myeloablative therapy: Children's Oncology Group study ANBL0931. *Front Immunol* 9:1355, 2018
27. Nguyen R, Sahr N, Sykes A, et al: Longitudinal NK cell kinetics and cytotoxicity in children with neuroblastoma enrolled in a clinical phase II trial. *J Immunother Cancer* 88:e000176, 2020
28. Garaventa A, Poetschger U, Valteau-Couanet D, et al: Randomized trial of two induction therapy regimens for high-risk neuroblastoma: HR-NBL1.5 International Society of Pediatric Oncology European Neuroblastoma Group study. *J Clin Oncol* 39:2552-2563, 2021
29. Berthold F, Faldum A, Ernst A, et al: Extended induction chemotherapy does not improve the outcome for high-risk neuroblastoma patients: Results of the randomized open-label GPOH trial NB2004-HR. *Ann Oncol* 31:422-429, 2020
30. Kreissman SG, Seeger RC, Matthay KK, et al: Purged versus non-purged peripheral blood stem-cell transplantation for high-risk neuroblastoma (COG A3973): A randomised phase 3 trial. *Lancet Oncol* 14:999-1008, 2013
31. Albertini MR, Hank JA, Schiller JH, et al: Phase IB trial of chimeric antidisialoganglioside antibody plus interleukin 2 for melanoma patients. *Clin Cancer Res* 3:1277-1288, 1997
32. Mody R, Yu AL, Naranjo A, et al: Irinotecan, temozolomide, and dinutuximab with GM-CSF in children with refractory or relapsed neuroblastoma: A report from the Children's Oncology Group. *J Clin Oncol* 38:2160-2169, 2020
33. Kushner BH, Cheung NK: GM-CSF enhances 3F8 monoclonal antibody-dependent cellular cytotoxicity against human melanoma and neuroblastoma. *Blood* 73:1936-1941, 1989
34. Mueller BM, Romerdahl CA, Gillies SD, et al: Enhancement of antibody-dependent cytotoxicity with a chimeric anti-GD2 antibody. *J Immunol* 144:1382-1386, 1990
35. Furman WL, Fairclough DL, Huhn RD, et al: Therapeutic effects and pharmacokinetics of recombinant human granulocyte-macrophage colony-stimulating factor in childhood cancer patients receiving myelosuppressive chemotherapy. *J Clin Oncol* 9:1022-1028, 1991
36. Meropol NJ, Porter M, Blumenson LE, et al: Daily subcutaneous injection of low-dose interleukin 2 expands natural killer cells in vivo without significant toxicity. *Clin Cancer Res* 2:669-677, 1996
37. Ladenstein R, Potschger U, Pearson AD, et al: Busulfan and melphalan versus carboplatin, etoposide, and melphalan as high-dose chemotherapy for high-risk neuroblastoma (HR-NBL1/SIOPEN): An international, randomised, multi-arm, open-label, phase 3 trial. *Lancet Oncol* 18:500-514, 2017



AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Improved Outcome in Children With Newly Diagnosed High-Risk Neuroblastoma Treated With Chemoimmunotherapy: Updated Results of a Phase II Study Using hu14.18K322A

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/authors/author-center.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](#)).

Barry L. Shulkin

Consulting or Advisory Role: Navidea

Matthew Krasin

Consulting or Advisory Role: Debiopharm Group

Michael W. Bishop

Consulting or Advisory Role: Fennec Pharma

Research Funding: Pfizer

Fariba Navid

Research Funding: Bayer (Inst)

Brandon Triplett

Travel, Accommodations, Expenses: Miltenyi Biotec

Stephen D. Gillies

Employment: LinkedUp Bioscience

Leadership: LinkedUp Bioscience

Stock and Other Ownership Interests: Provenance Biopharmaceuticals

Consulting or Advisory Role: Delos Capital Partners

Patents, Royalties, Other Intellectual Property: Patent owner not related to this paper

Alice Yu

Leadership: OPKO Health

Stock and Other Ownership Interests: OPKO Health/GeneDx

Honoraria: EUSA Pharma

Consulting or Advisory Role: OBI Pharma

Speakers' Bureau: EUSA Pharma

Research Funding: United Therapeutics (Inst)

Patents, Royalties, Other Intellectual Property: Globo H-Diphtheria toxoid vaccine for cancer therapy, NKT stimulatory phenyl-glycolipids for cancer therapy and vaccine adjuvant, Cancer targeting peptides, Methods for suppressing cancer by inhibition of TMCC3

Travel, Accommodations, Expenses: EUSA Pharma

Paul M. Sondel

Research Funding: Invenra Inc (Inst)

Patents, Royalties, Other Intellectual Property: I have partial interest in patents related to my work at the University of Wisconsin-Madison WI, which are held by and managed by the University of Wisconsin Foundation. I am an unpaid medical advisor to Invenra Inc, a monoclonal antibody biotech firm in Madison WI. My UW laboratory is collaborating with Invenra and receiving research reagents from them for mutual research

Uncompensated Relationships: Invenra

Wing H. Leung

Employment: Miltenyi Biotec

Alberto Pappo

Honoraria: Bayer, Roche

Consulting or Advisory Role: Merck, Loxo/Bayer, EUSA Pharma, Debbio

No other potential conflicts of interest were reported.

APPENDIX

TABLE A1. Comparison of Patient Characteristics

Characteristic	NB2012 (n = 64) (Reference level), No. (%)	ANBL02P1 (n = 31), No. (%)	<i>P</i>	ANBL0532 (n = 652), No. (%)	<i>P</i>	ANBL12P1 (n = 146), No. (%)	<i>P</i>
Age, months							
< 18	5 (8)	5 (16)	NS	Not reported		22 (15)	NS
≥ 18	59 (92)	26 (84)				124 (85)	
Median, months	37.5	30		37.2		37.3	
Sex							
Female	27 (42)	6 (19)	.028	286 (44)	NS	Not reported	
Male	37 (58)	25 (81)		366 (56)			
INSS stage							
IIb	1 (2)	Non stage IV: 4 (13)	NS	7 (1)	NS	Non stage IV: 20 (14)	NS
III	7 (11)			68 (10)			
IV	56 (87)	27 (87)		577 (89)		126 (86)	
MYCN status							
Not amplified	43 (67)	15 (60)	NS	327 (57)	NS	75 (56)	NS
Amplified	21 (33)	10 (40)		249 (43)		58 (44)	
Unreported	0	6		76		13	
Shimada histology							
Favorable	6 (9)	1 (3)	NS	23 (3)	.013	7 (5)	NS
Unfavorable	41 (64)	21 (68)		588 (86)		119 (82)	
Not performed	17 (27)	9 (29)		71 (10)		20 (14)	

Abbreviations: INSS, International Neuroblastoma Staging System; NS, not significant.

TABLE A2. Treatment-Related Grade 3/4 Toxicities Reported During the First Six Cycles of Induction Therapy in at Least 5% of Patients Plus Toxicities of Interest

Toxic Effect	Cycle 1 (n = 64)	Cycle 2 (n = 64)	Cycle 3 (n = 64)	Cycle 4 (n = 63)	Cycle 5 (n = 62)	Cycle 6 (n = 62)
Allergic						
Allergic reaction	—	—	—	—	1	—
Anaphylaxis	1	—	—	—	—	—
Cardiac						
Hypertension	5	2	1	—	2	2
Hypotension	1	—	2	3	2	4
Febrile neutropenia	30	12	4	35	12	30
Fever	15	2	—	5	4	6
GI						
Abdominal pain	3	2	2	2	3	7
Abnormal alkaline aminotransferase	3	4	5	1	1	4
Abnormal AST	4	1	1	1	1	1
Abnormal bilirubin	4	1	—	2	1	4
Diarrhea	2	2	4	1	3	1
GGT elevated	5	5	1	1	2	6
Nausea	—	—	4	3	4	3
Vomiting	—	1	7	8	2	6
Infection						
Catheter-related infection	2	4	2	6	4	9
Enterocolitis infectious	9	4	5	13	7	8
Sepsis	—	—	1	1	1	4
Upper respiratory infection	13	10	3	11	8	19
Urinary tract infection	3	—	6	5	2	8
Metabolic						
Acidosis	5	7	9	5	4	2
Hyperglycemia	3	4	9	7	9	3
Hypocalcemia	5	1	5	4	3	5
Hypokalemia	16	5	24	11	19	12
Hyponatremia	7	1	13	7	9	5
Hypophosphatemia	7	—	22	1	24	6
Mucosal						
Esophagitis	1	—	—	2	—	7
Mucositis	1	—	—	21	1	21
Pain	18	8	5	3	2	2
Respiratory						
Apnea	—	1 ^a	—	1	—	—
Bronchospasm	2	—	—	—	1	—
Cough	1	—	2	1	—	—
Hypoxia	14	3	1	—	1	2
Pleural effusion	6	1	1	—	—	—
Respiratory failure ^b	1	—	—	—	—	—
Others						
Hearing impaired	—	—	2	4	2	9
Infusion-related reaction	6	1	4	2	1	—

Abbreviation: GGT, Gamma Glutamyl Transferase.

^aEpisode of apnea was related to opioid administration.

^bRespiratory failure was related to significant thoracic disease burden.

TABLE A3. Serum Antibody Levels in Three Representative HAHA– Patients and All Five Patients Who Developed HAHA During Therapy

Pt-ID	HAHA ^a	Antibody	C1	C2	C3	C4	C5	C6	Int	M1	M2	M3	M4	M5	M6
9	–	K322A ^b	15.6	16.7	17.0	15.7	14.5	14.2	NA	12.9	10.5	11.8	13.7	13.9	NA
		HAHA	0	0	0	0	0	0	0	0	0	0	0	0	0
12	–	K322A	27.7	16.1	17.9	26.7	15.3	14.4	3.5	9.4	12.0	11.8	8.8	9.8	NA
		HAHA	0	0	0	0	0	0	0	0	0	0	0	0	0
13	–	K322A	12.2	23.2	21.0	23.2	20.6	17.9	NA	11.4	10.7	13.3	9.0	13.9	NA
		HAHA	0	0	0	0	0	0	NA	0	0	0	0	0	0
23	+	K322A	1.2	13.9	17.7	22.2	15.5	15.5	8.0	7.4	NA	9.5	NA	12	NA
		HAHA	0	0	0.15	0.71	NA	0.40	0	0	0	NA	0	0	0
66	+	K322A	12.4	NA	7.4	15.1	13.1	13.3	12.5	12.4	11.0	14.1	11.2	18.6	NA
		HAHA	0	1.3	0	0	0	0	0	0	0	0	0	0.89	0
69	+	K322A	5.9	3.8	17.0	16.2	18.3	15.4	10.3	12.7	11.5	12.1	15.8	16.8	NA
		HAHA	0	0.78	0	0	0	0	0	0	0	0	0	0	0
112	+	K322A	12.4	13.9	12.8	10.1	10.8	7.6	10.5	9.3	13.3	12.1	8.4	6.2	NA
		HAHA	0.72	3.2	1.7	0.84	0.46	0.93	0.82	0	0	0	0	0	0
145	+	K322A	5	16.8	10.6	9.5	12.6	15	NA	13	20	21.1	33.1	19.5	NA
		HAHA	0	0	0	0	0	0	0.82	0	0	0	0	0	0

Abbreviations: C, induction cycle; HAHA, human antihuman antibody; Int, intensification phase; K322A, hu14.18K322A; M, maintenance cycle; Pt-ID, patient identification number.

^aHAHA levels, measured as optical density (OD) units, detected in a bridging ELISA assay with serum obtained at the initiation of the indicated cycle, before therapy began in that cycle. HAHA+ indicates patients with ≥ 0.7 OD unit increase at any time, as compared with the baseline C1 sample. The five patients who developed evidence of a HAHA response at any time are shown. Three representative patients (of 40) without a HAHA response are also shown.

^bK322A concentration ($\mu\text{g/mL}$) for each indicated cycle, obtained 1 hour after completion of the K322A infusion on day 1 of that cycle.

TABLE A4. Comparison of End of Induction Objective Responses, EFS and OS of ANBL12P1, ANBL0532, HR-NBL1.5, NB2004-HR and NB2012

Response	ANBL12P1 ²⁵ (N = 146; 125 evaluable for response)		ANBL0532 ¹⁵ (N = 652)		HR-NBL1.5 ²⁹ (N = 566) ^b		NB2004-HR ²⁹ (N = 389) ^c		NB2012 (N = 64)		
		<i>P</i> ^a		<i>P</i> ^a		<i>P</i> ^a		<i>P</i> ^a		<i>P</i> ^a	
CR	28/125 (22.4%)	.0896	277 (42.5%)	< .0001 ^d	16 (3%)	< .0001 ^d	109 (28%)	.4010	21 (34%)		
VGPR	30/125 (24%)	.1485			99 (17%)	.0011 ^d	59 (15%)	≤ .0001 ^d	21 (34%)		
PR	42/125 (33.6%)	.5258	213 (32.7%)	.5519	295 (52%)	.0006 ^d	177 (46%)	.0105 ^d	18 (29%)		
≥PR	80%	.0018 ^d	75.2%	< .0001 ^d	72%	< .0001 ^d	89%	.0483	97%		
MR	10/125 (8%)	NA	79 (12.1%)	.0510 ^d	87 (15%)	NA			0		
NR	9/125 (7.2%)	.41			50 (9%)	.1691	3 (1%)	.4475	2 (3%)		
PD	6/146 (4.1%)	NA	46 (7.1%)	NA	12 (2%)	NA	40 (10%)	NA	0		
D/C early	39/146 (26.7%)	< .0001 ^d	NE 37 (5.7%)	.7788	Death before response evaluation 7 (1%)	.1546			3 (4%)		
					rCOJEC (N = 313) ^e	N5 (N = 313) ^e	STD (N = 211)	EXP (N = 211)			
Mean 3-year EFS	55.6% ± 4.2%	.0131 ^d	51.1% ± 3.95%	.0005 ^d	44% (38%-50%)	47% (41%-53%)	< .0001 ^d	34% (28%-40%)	32% (26%-38%)	< .0001 ^d	73.7% ± 11.7%
							< .0001 ^d			< .0001 ^d	
Mean 3-year OS	74.5% ± 3.7%	.0640	Reported in graph only		60% (54%-66%)	65% (58%-70%)	< .0001 ^d	54% (46%-62%)	48% (40%-56%)	< .0001 ^d	86.0% ± 9.5%
							.0010 ^d			< .0001 ^d	

Abbreviations: CR, complete response; EFS, event free survival; MR, mixed response; NA, not applicable; NR, no response; OS, overall survival; PD, progressive disease; PR, partial response; VGPR, very good partial response.

^aCompared to NB2012.

^bResponses in HRNBL1.5 were assessed prior to surgical resection of primary tumor.²⁸

^cNB2004-HR was a randomized comparison between six courses of standard (STD) induction chemotherapy versus an experimental (EXP) induction regimen with two additional courses of topotecan, cyclophosphamide and etoposide.²⁹

^d*P* values from two proportions Z-test with significance level ≤ 0.05.

^eHR-NBL1.5 was a randomized comparison of two different induction regimens: rCOJEC (comprised eight cycles within 70 days alternating vincristine and carboplatin [course A], vincristine and cisplatin [course B], and vincristine, etoposide, and cyclophosphamide [course C] in the sequence ABCBACB. The interval between courses was 10 days irrespective of hematologic recovery.) and MSKCC N5 (comprised high-dose cyclophosphamide, doxorubicin with vincristine [CAV] alternating with high-dose cisplatin, etoposide [PE] for a total of five courses [CAV, PE, CAV, PE, and CAV] within 110 days.)²⁸