



HHS Public Access

Author manuscript

Vet Clin North Am Small Anim Pract. Author manuscript; available in PMC 2022 January 30.

Published in final edited form as:

Vet Clin North Am Small Anim Pract. 2020 July ; 50(4): 653–662. doi:10.1016/j.cvsm.2020.03.001.

Stress and Feline Health

C.A. Tony Buffington, DVM, PhD [Clinical Professor],

Department of Medicine & Epidemiology, UC Davis School of Veterinary Medicine, Davis, CA 95616, 212 Cedar Lane, Woodland, CA 95695

Melissa Bain, DVM, MS [Professor]

Clinical Animal Behavior, Department of Medicine & Epidemiology, UC Davis, School of Veterinary Medicine, Davis, CA 95616, 5219 El Cerrito Ave., Davis, CA 95618

Keywords

coping; early life stress; epigenetic modulation of gene expression; perception of control; perception of threat; resilience; threat response system

Introduction

Veterinarians strive to improve the health of their patients, and recently there has been a more concentrated effort to understand the relationship between stress, physical health, and emotional health. Recent research has investigated how the effects of stressful environments manifest in cats, expanding from and connecting to research in other species.^{1,2}

What is stress?

The term stress can be defined in various ways. From the perspectives of health and disease, stress is often defined in terms of *stressors*, which are events in the internal and external environment that result in a *stress response*. We think of stressors as events that are perceived as threats to one's perception of control. From this perspective, a stressor is anything that activates the central threat response system (CTRS).³ Stressors can vary along a continuum: positive to negative, duration, acute to chronic, and intensity, (mild to toxic). For the purposes of this chapter, we have adapted the "ecobiodevelopmental" framework proposed by the American Academy of Pediatrics, "to stimulate fresh thinking about the promotion of health and prevention of disease across the lifespan."⁴ which provides a helpful way to begin to think about stressors and stress responses. This framework acknowledges that all events that activate the CTRS are not equally threatening, and proposes a conceptual taxonomy comprising three distinct types of CTRS responses: mild

drbuffington@ucdavis.edu .

The Authors have nothing to disclose. Dr. Buffington will serve as the corresponding author.

*"Everyone knows what stress is, but nobody really knows."*Hans Selye

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

(positive), moderate (tolerable), and severe (toxic). The taxonomy is based on the differences in their potential to cause enduring physiologic disruptions as a result of the intensity and duration of the response.

Mild stress responses, defined as brief in duration and mild-to-moderate in magnitude, can help develop stress coping skills. Mild stress responses are part of normal development when they occur in the safe, predictable environments of stable and supportive relationships. Examples of events resulting in positive stress responses in young animals include non-threatening veterinary visits and exposure to novel environments and foods.

Moderate stress responses result from exposure to experiences that present greater threats, such as lack of stimulation/boredom or household instability, illness or injury, or exposure to a natural disaster. As with mild stress responses, when the event occurs in an otherwise safe environment, recovery to normal is likely. Thus, the essential feature that makes moderate stress response tolerable is the extent to which one's protective surroundings permit retention of sufficient perception of control to permit adaptive coping to occur.

In the most threatening circumstances, severe stress responses can result. Toxic stress responses, strong, frequent, or prolonged, are the most dangerous to long-term health and welfare. Examples of events that can result in severe stress responses include chronic abuse, severe or chronic disease, and adverse early life events like nutritional deprivation,⁵ maternal separation, or significant maternal threat during pregnancy (more on this later).⁶

Stress and the Lower Urinary Tract (LUT)

Abnormal signs referable to the lower urinary tract (LUTS) of domestic cats include variable combinations of increased voiding urgency and frequency, decreased volume, blood in the urine, and urination outside the litter box. Although there are many possible causes of these signs, the most common causes include: "idiopathic cystitis"; a stone, usually calcium oxalate or magnesium ammonium phosphate; or a urinary tract infection (usually in older cats with upper tract disease). In addition to infection and inflammation, stress responses also can activate the LUT, although some of the fundamental neural mechanisms and pathways linking psychosocial stress to altered behaviors and physiological disorders are still unclear. Holstege⁷ has pointed out that micturition plays a much more important role in the context of the survival of individuals and species in most mammals than it does in humans. For most mammals, urine also signals important messages such as the demarcation of territory of a specific individual⁸ and estrus.⁹ These urine functions demonstrate that voiding can occur for reasons other than to empty a full bladder, including the contexts of perception of threat and reproduction. Micturition responses in these situations require, and imply, supraspinal control of micturition.⁷

Part of the interest in feline stress biology arose from studies of cats with severe, recurrent chronic LUTS. Lower urinary tract signs have long been recognized in domestic cats, being described as "very common" in Kirk's 1925 veterinary textbook.¹⁰ Clinicians historically interpreted signs referable to the LUT to suggest an external (e.g., infectious agent, urinary stone) or structural disorder of the LUT (e.g., stricture or incontinence), although

environmental conditions were recognized as a risk factor by Kirk in 1925, and subsequently confirmed to be relevant by other investigators.^{11,12}

In 1996, Osborne et al., summarized the literature on feline LUTS, identifying some 30 distinct causes of LUTS.¹³ That so many potential causes result in similar LUTS demonstrates that the signs in themselves only represent the limited repertoire available to the LUT to respond to any insult. During this time, a syndrome described as “feline interstitial cystitis” (FIC) also was being studied.¹⁴ These studies found that additional problems outside the lower urinary tract were commonly present in these cats, and could be mitigated by multimodal environmental modification (MEMO).¹⁵ Subsequent research has revealed the complex nature of this condition. For example, some cats with severe, chronic LUTS seem to have a functional, rather than a structural, lower urinary tract disorder,¹ and that LUTS can occur even in presumably healthy cats exposed to stressful circumstances.^{12,16}

These findings suggest that terms such as “feline urological syndrome”,¹⁷ “feline lower urinary tract disease”,¹⁸ and “feline interstitial cystitis”¹⁴ historically used to describe patients with this syndrome do not describe the extent of the problems occurring in many afflicted cats. Conversely, the narrow focus on the LUT may have precluded thorough evaluation of the entire patient, which might have revealed variable combinations of clinical signs referable to other organ systems such as the gastrointestinal tract, skin, lung, cardiovascular, central nervous, endocrine, and immune systems.^{15,19} These comorbidities include some of the most common problems in feline medicine. They also lack a predictable pattern of onset across patients, and often precede the appearance of LUTS. This situation is similar in human beings with “central sensitivity syndromes”²⁰, suggesting that individual patterns of comorbidities may represent an amplification of underlying familial sensitivities.²¹

Perception of threat—The inconsistency of comorbidities observed in cats with chronic LUTS suggests that they may be variable manifestations of a common underlying problem.^{22,23} The underlying problem in these patients appears to be a sensitized CTRS,¹ which results in a relative predominance of sympathetic to parasympathetic nervous system responses,^{25,26} increased hypothalamic-pituitary – but not adrenal – activity,²⁴ as well as variable alterations in immune²⁵ and endocrine function.²⁶ Chronic activation of the CTRS seems to result in chronic “wear and tear” on body and brain systems, which can eventually progress to clinical signs referable to those organs most vulnerable in a particular individual.²⁷

In addition to alterations in the components of the CTRS distal to the hypothalamus, brain structures that modulate activity of the CTRS also appear to be affected.²⁸ For example, cognitive factors such as classical and operant conditioning, attention bias, and memory have been shown to play roles in the onset, development, and maintenance of chronic health problems.²⁸ Moreover, studies have shown that the individual’s expectations and ability to cope with potentially traumatic events are more important than the physical events *per se* for determining the immediate and long term consequences for health.²⁹

The sensitization of the CTRS appears to result from genetic disorders, developmental events, environmental threats, or variable combinations of these.³⁰ A variety of polymorphisms in genes that affect stress responsiveness, including catechol-O-methyltransferase,³¹ serotonin transporter,³² and alpha-2 adrenergic receptor, have been identified in humans.^{33,34} In cats, a genetic polymorphism of the alpha-2 adrenergic receptor that appears to enhance stress susceptibility has been reported.³⁵

During the last two decades, the science of Developmental Origins of Health and Disease³⁶ has emerged to investigate the role of early life experiences on the sensitivity of the CTRS. Evidence from clinical, epidemiological, and experimental observations has shown how evolutionarily conserved developmental processes can interact with environmental cues, often transmitted from the mother via the placenta to the offspring, to attempt to match the physiology of the developing organism to its post-natal environment. The sequence of events that has emerged from this research proposes that when a pregnant female is exposed to a sufficiently severe stressor, the neuro-endocrine products of the ensuing stress response may cross the placenta and affect the course of fetal development.³⁷ The biological "purpose" of transmitting such environmental cues to the fetus may be to guide the development of the fetal CTRS and associated behaviors to increase the probability of survival. The fetus may "use" information in its *in utero* environment to make predictive adaptive response "decisions". If a threatening or nutrient-limiting environment is perceived, the developmental trajectory of the fetus may change in response to the available information to enhance reproductive fitness in the predicted *ex utero* environment.

Studies of the enduring effects of stressful developmental experiences on health have now been published in a wide variety of mammalian species, from rodents to primates.⁶ Problems can arise when the actual *ex utero* environment does not match the predicted one. Studies in multiple species have found that cardiovascular disease, type 2 diabetes mellitus, metabolic syndrome, and respiratory, gastrointestinal, lower urinary tract, dental, and mood disorders all can result from a mismatch between the predicted and actual environment the individual inhabits.³⁸ Cognitive function, too, is affected by both genetic and developmental influences. Impaired coping to stressful situations, increased fear and anxiety-related behaviors, and dysregulation of the CTRS all have been found in adults exposed to adverse early life experiences.⁶

Recent research also has shown that one mechanism underlying the sensitization of the CTRS involves a process called epigenetic modulation of gene expression.³⁹ This general biological process results in such commonplace outcomes as sex- and organ-specific patterns of gene expression that lead to the final phenotype of the organism by silencing genes not appropriate to the particular tissue environment. The molecular mechanisms of epigenetics are beyond the scope of this review, but detailed explanations are freely available.⁴⁰

The effects of stressors on sensitization of the CTRS seem to depend both on the timing and magnitude of exposure to products of the maternal stress response in relation to the activity of the developmental "programs" that determine the maturation of the various body systems during gestation and early postnatal development. For example, the small adrenal cortices found in some cats with FIC suggest that the event occurred during a time when

adrenocortical maturation was occurring.⁴¹ If the developing fetus had been exposed to a stressor before adrenocortical maturation, the effect may not have been observed, whereas if it had occurred later in development adrenal size and subsequent adrenocortical responses to stress might have been increased. Given the number of orphaned or abandoned kittens obtained from shelters, the risk for sensitization of the CTRS in client-owned cats may be high.

While sensitization of the CTRS is more likely to occur during growth and maturation of the neural, endocrine, and immune systems, it is not restricted to the developmental period. Moreover, sensitization of the CTRS may be unmasked by another adverse experience later in life, possibly associated with another round of epigenetic modulation of gene expression.²⁴ Subsequent rounds of alterations in gene expression may be quite stable and resistant to current medical interventions. Additionally, sensitization of the CTRS may be part of a more general “survival phenotype” that includes smaller (or larger) size at birth.⁴²

Perception of control—Activation of the CTRS appears to depend on the balance between perceptions of threat and perceptions of control. Other recent research has focused on the roles of resilience and choice on stress responsiveness. The American Psychological Association’s website “The Road to Resilience” (<https://www.apa.org/helpcenter/road-resilience>) defines resilience as “the process of adapting well in the face of adversity, trauma, tragedy, threats, or significant sources of stress. It means “bouncing back” from difficult experiences.” A recent (September 15, 2019) special issue of *Biological Psychiatry* addressed the topic of the neurobiology of resilience from basic, clinical, and translational perspectives to summarize current knowledge of this important and rapidly expanding topic.⁴³ For example, Cathomas, et al.,²⁹ reviewed the physiological and transcriptional adaptations of specific brain circuits, the role of cellular and humoral factors of the immune system, the gut microbiota, and changes at the interface between the brain and the periphery on the neurobiological mechanisms of stress resilience. They proposed viewing resilience as a process that integrates multiple central and peripheral systems.

Belief in one’s ability to exert control over the environment and to produce desired results is essential for wellbeing, and probably a psychological and biological necessity. Converging evidence from animal research, clinical studies and neuroimaging suggests that the need for control is a biological imperative for survival. The ability to choose permits animals to increase their perception of control over their surroundings, which helps reduce activation of the CTRS.^{44,45} In fact, one of the Five Freedoms, freedom from fear, recently has been operationalized as the opportunity for choice and control.⁴⁶ And Rochlitz has declared that, “Provided extremes are avoided, if the cat has a variety of behavioural choices and is able to exert some control over its physical and social environment, it will develop more flexible and effective strategies for coping with stimuli.”⁴⁷

The negative effects of adverse early life events mediated by severe stress responses on the long-term health and welfare of cats demonstrate the importance of identification of risk factors and provision of effective education of clients by clinicians about appropriate environments for indoor or otherwise confined cats throughout their lives.⁴⁸ From this perspective, initial vaccination appointments are anything but “routine”, and may in fact be

the most important appointments of the animal's life. This is because such visits present opportunities to teach husbandry appropriate for the animal based on the environment it is confined to at a time when owners are likely to be most motivated and responsive to recommendations. (See also Environment and Feline Health: At Home and in the Clinic by Heath). These needs remain but modifications may be required as cats age. (See Feline Ageing: Promoting Physiological and Emotional Wellbeing by Gunn-Moore)

Shonkoff et al.,⁴ described three foundational domains to promote long-term health and welfare that can be adapted to preventive veterinary care for confined cats. These include:

1. A stable and responsive environment that provides consistent, nurturing, and protective interactions that enhance learning and support development of adaptive capacities (resilience and coping) that promote a well-regulated CTRS;
2. A safe and supportive physical, chemical, and built environment that is free from toxins and fear, allows active exploration without significant risk of harm, and offers support for families caring for the cat; and
3. Satisfactory nutrition for the cat's age, including feeding management practices that maintain an appropriate body condition (beginning with the future mother prior to conception if possible) while providing safety, choice, and mental stimulation to the extent possible.⁴⁹ (See also Feeding Cats for Optimal Mental and Behavioral Well-Being Part 2 by Dantas and Delgado)

Consideration of the role of a sensitized CTRS also has important therapeutic implications. For example, drugs to modulate gene expression are under active investigation in oncology⁵⁰ and psychiatry⁵¹ that may become available in veterinary medicine. Given that these drugs also may modulate expression of other genes in potentially unpredictable ways, the availability of a naturally occurring animal model of a disease likely influenced by these mechanisms offers the opportunity to test these compounds before they are introduced to human medicine.²¹ Recent research also suggests that drugs like some of the psychedelics⁵² and low doses of ketamine⁵³ might prove therapeutic for patients with central sensitivity syndromes.

Environmental modification also can modulate gene expression,^{54,55} which may explain the effectiveness of behavior-based approaches for treatment in humans⁵⁶ and multimodal environmental modification for therapy of cats with FIC.^{12,15} Effective environmental modification has been found to result in significant resolution of signs of comorbid disorders as well as LUTS in treated cats suffering from FIC,^{12,15} and to reverse the effects of maternal separation on stress reactivity in rodents.⁵⁷

The roles of environmental modification on feline health and welfare can be visualized in the illustration (Figure 1). Cats' perceptions of threat may be reduced by increasing safety through provision of separate sets of resources, opportunities to express species-typical behaviors, and elimination of conflict,⁵⁸ and their perception of control can be enhanced by offering resources as choices, and providing only those the cat chooses.⁵⁹ Changing the environment changes the context and the cat's expectations, which over time changes its history, permitting it to cope with its environment and benefitting its health and welfare.

Summary/Discussion

Changing our view of some chronic diseases from that of isolated organ-originating diseases in otherwise healthy individuals to one that considers the role of a sensitized CTRS (and the potential roles of early adverse experiences) has at least two important implications. First, it provides a more parsimonious explanation for many findings that previously were quite difficult to account for, including the unfortunate lack of beneficial effect of therapies directed at the peripheral organ of interest by a particular medical subspecialty, the presence of multiple comorbid disorders in many patients with chronic disorders, but not in patients with other individual organ diseases, the unpredictable order of appearance of the comorbidities, and the altered functioning of the CTRS. Second, and more importantly, it opens whole new areas of therapy that may escape consideration from the individual medical specialty perspective. The currently available data only suggest this possibility, however, but they do permit generation of the hypothesis.

References

1. Buffington CA. Idiopathic cystitis in domestic cats-beyond the lower urinary tract. *Journal of Veterinary Internal Medicine*. 2011;25(4):784–796. [PubMed: 21564297]
2. Buffington CAT. Pandora Syndrome in cats: diagnosis and treatment. *Today's Veterinary Practice*. 2018;8(5):31–41.
3. Elman I, Borsook D. Threat Response System: parallel brain processes in pain vis-à-vis fear and anxiety. *Frontiers in psychiatry*. 2018;9:29. [PubMed: 29515465]
4. Shonkoff JP. Building a new biodevelopmental framework to guide the future of early childhood policy. *Child Development*. 2010;81(1):357–367. [PubMed: 20331672]
5. Bouret S, Levin BE, Ozanne SE. Gene-Environment Interactions Controlling Energy and Glucose Homeostasis and the Developmental Origins of Obesity. *Physiological Reviews*. 2015;95(1):47–82. [PubMed: 25540138]
6. Guzman DB, Howell B, Sanchez M. Early life stress and development: preclinical science. *Posttraumatic Stress Disorder: From Neurobiology to Treatment*, ed Bremner JD (Hoboken, NJ: Wiley-Blackwell). 2016:61–80.
7. Holstege G. Micturition and the soul. *Journal of Comparative Neurology*. 2005;493(1):15–20.
8. Pryor PA, Hart BL, Bain MJ, Cliff KD. Causes of urine marking in cats and effects of environmental management on frequency of marking. *Journal of the American Veterinary Medical Association*. 2001;219(12):1709–1713. [PubMed: 11767919]
9. Hart BL. Facilitation by estrogen of sexual reflexes in female cats. *Physiology & behavior*. 1971;7(5):675–678. [PubMed: 5164360]
10. Kirk H. Retention of urine and urine deposits. In: Kirk H, ed. *The Diseases of the Cat and its General Management*. London: Bailliere, Tindall and Cox; 1925:261–267.
11. Caston HT. Stress and the feline urological syndrome. *Feline Practice*. 1973;3(6):14–22.
12. Stella JL, Lord LK, Buffington CA. Sickness behaviors in response to unusual external events in healthy cats and cats with feline interstitial cystitis. *Journal of the American Veterinary Medical Association*. 2011;238(1):67–73. [PubMed: 21194324]
13. Osborne CA, Kruger JM, Lulich JP. Feline lower urinary tract disorders. Definition of terms and concepts. *Veterinary Clinics of North America - Small Animal Practice*. 1996;26(2):169–179.
14. Buffington CAT, Chew DJ, Woodworth BE. Feline interstitial cystitis. *Journal of the American Veterinary Medical Association*. 1999;215(5):682–687. [PubMed: 10476717]
15. Buffington CAT, Westropp JL, Chew DJ, Bolus RR. Clinical evaluation of multimodal environmental modification in the management of cats with lower urinary tract signs. *Journal of Feline Medicine and Surgery*. 2006;8:261–268. [PubMed: 16616567]

16. Westropp JL, Kass PH, Buffington CA. Evaluation of the effects of stress in cats with idiopathic cystitis. *Am J Vet Res.* 2006;67(4):731–736. [PubMed: 16579769]
17. Osbaldiston GW, Taussig RA. Clinical report on 46 cases of feline urological syndrome. *Vet Med/Small Anim Clin.* 1970;65:461–468.
18. Osborne CA, Johnston GR, Polzin DJ, et al. Redefinition of the feline urologic syndrome: feline lower urinary tract disease with heterogeneous causes. *Vet Clin North Am Small Anim Pract.* 1984;14(3):409–438. [PubMed: 6429924]
19. Buffington CA, Westropp JL, Chew DJ, Bolus RR. Risk factors associated with clinical signs of lower urinary tract disease in indoor-housed cats. *J Am Vet Med Assoc.* 2006;228(5):722–725. [PubMed: 16506933]
20. Yunus MB. Editorial Review: An Update on Central Sensitivity Syndromes and the Issues of Nosology and Psychobiology. *Curr Rheumatol Rev.* 2015;11(2):70–85. [PubMed: 26138918]
21. Buffington CA. Developmental influences on medically unexplained symptoms. *Psychotherapy and Psychosomatics.* 2009;78(3):139–144. [PubMed: 19270468]
22. Yunus MB. Central Sensitivity Syndromes: Stress System Failure May Explain the Whole Picture Reply. *Seminars in Arthritis and Rheumatism.* 2009;39(3):220–221.
23. Neblett R, Cohen H, Choi Y, et al. The Central Sensitization Inventory (CSI): establishing clinically significant values for identifying central sensitivity syndromes in an outpatient chronic pain sample. *J Pain.* 2013;14(5):438–445. [PubMed: 23490634]
24. Daskalakis NP, Bagot RC, Parker KJ, Vinkers CH, de Kloet ER. The three-hit concept of vulnerability and resilience: toward understanding adaptation to early-life adversity outcome. *Psychoneuroendocrinology.* 2013;38(9):1858–1873. [PubMed: 23838101]
25. Fagundes CP, Glaser R, Kiecolt-Glaser JK. Stressful early life experiences and immune dysregulation across the lifespan. *Brain, behavior, and immunity.* 2013;27:8–12.
26. Neeck G. Pathogenic mechanisms of fibromyalgia. *Ageing Res Rev.* 2002;1(2):243–255. [PubMed: 12039441]
27. Danese A, McEwen BS. Adverse childhood experiences, allostasis, allostatic load, and age-related disease. *Physiology & behavior.* 2012;106(1):29–39. [PubMed: 21888923]
28. Ursin H. Brain sensitization to external and internal stimuli. *Psychoneuroendocrinology.* 2014;42:134–145. [PubMed: 24636510]
29. Cathomas F, Murrough JW, Nestler EJ, Han M-H, Russo SJ. Neurobiology of resilience: Interface between mind and body. *Biological psychiatry.* 2019.
30. Buffington CAT. Comorbidity of interstitial cystitis with other unexplained clinical conditions. *Journal of Urology.* 2004;172:1242–1248.
31. Stein DJ, Newman TK, Savitz J, Ramesar R. Warriors versus worriers. The role of COMT gene variants. *CNS Spectrums.* 2006;11(10):745–748. [PubMed: 17008817]
32. Jacobs N, Kenis G, Peeters F, Derom C, Vlietinck R, van Os J. Stress-related negative affectivity and genetically altered serotonin transporter function - Evidence of synergism in shaping risk of depression. *Archives of General Psychiatry.* 2006;63(9):989–996. [PubMed: 16953001]
33. Wust S, Federenko IS, Van Rossum EFC, et al. A psychobiological perspective on genetic determinants of hypothalamus-pituitary-adrenal axis activity. In: *Biobehavioral Stress Response: Protective and Damaging Effects.* Vol 1032.2004:52–62.
34. Rosmond R, Bouchard C, Bjorntorp P. A C-1291G polymorphism in the alpha(2A)-adrenergic receptor gene (ADRA2A) promoter is associated with cortisol escape from dexamethasone and elevated glucose levels. *Journal of Internal Medicine.* 2002;251(3):252–257. [PubMed: 11886485]
35. Lu P, Owens T, Buffington CAT. Association Between Alpha-2a adrenergic Receptor Gene Polymorphisms and Feline Idiopathic Cystitis Paper presented at: NIDDK International Symposium: Frontiers in Painful Bladder Syndrome and Interstitial Cystitis 2006; Bethesda, MD.
36. Penkler M, Hanson M, Biesma R, Müller R. DOHaD in science and society: emergent opportunities and novel responsibilities. *Journal of developmental origins of health and disease.* 2019;10(3):268–273. [PubMed: 30466503]
37. Miranda A, Sousa N. Maternal hormonal milieu influence on fetal brain development. *Brain and behavior.* 2018;8(2):e00920.

38. Gluckman PD, Buklijas T, Hanson MA. The developmental origins of health and disease (DOHaD) concept: past, present, and future. In: *The epigenome and developmental origins of health and disease*. Elsevier; 2016:1–15.
39. Aristizabal MJ, Anreiter I, Halldorsdottir T, et al. Biological embedding of experience: A primer on epigenetics. *Proceedings of the National Academy of Sciences*. 2019;10.1073/pnas.1820838116.
40. Wikipedia. Epigenetics. <http://en.wikipedia.org/wiki/Epigenetics>. Accessed December 20, 2019.
41. Westropp JL, Welk KA, Buffington CAT. Small adrenal glands in cats with feline interstitial cystitis. *J Urol*. 2003;170(6 (Pt 1)):2494–2497. [PubMed: 14634458]
42. Parlee SD, MacDougald OA. Maternal nutrition and risk of obesity in offspring: The Trojan horse of developmental plasticity. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*. 2014;1842(3):495–506. [PubMed: 23871838]
43. Murrrough JW, Russo SJ. The Neurobiology of Resilience: Complexity and Hope. *Biological psychiatry*. 2019;86(6):406–409. [PubMed: 31466560]
44. Leotti LA, Iyengar SS, Ochsner KN. Born to choose: The origins and value of the need for control. *Trends in cognitive sciences*. 2010;14(10):457–463. [PubMed: 20817592]
45. Falk JH. *Born to choose: Evolution, self, and well-being*. New York, NY: Routledge; 2017.
46. Greggor A, Vicino GA, Swaisgood RR, et al. Animal welfare in conservation breeding: applications and challenges. *Frontiers in veterinary science*. 2018;5:323. [PubMed: 30631770]
47. Rochlitz I. A review of the housing requirements of domestic cats (*Felis silvestris catus*) kept in the home. *Applied Animal Behaviour Science*. 2005;93:97–109.
48. Ellis SL, Rodan I, Carney HC, et al. AAFP and ISFM feline environmental needs guidelines. *J Feline Med Surg*. 2013;15(3):219–230. [PubMed: 23422366]
49. Dantas LM, Delgado MM, Johnson I, Buffington CT. Food puzzles for cats: feeding for physical and emotional wellbeing. *J Feline Med Surg*. 2016;18(9):723–732 [PubMed: 27102691]
50. Mummaneni P, Shord SS. Epigenetics and Oncology. *Pharmacotherapy*. 2014;34(5):495–505. [PubMed: 24619798]
51. Mahgoub M, Monteggia LM. Epigenetics and Psychiatry. *Neurotherapeutics*. 2013;10(4):734–741. 2013;10(4):734–741. [PubMed: 24092614]
52. Murnane K. The renaissance in psychedelic research: What do preclinical models have to offer. *Progress in brain research*. 2018;242:25. [PubMed: 30471682]
53. Nowacka A, Borczyk M. Ketamine applications beyond anesthesia—A literature review. *European journal of pharmacology*. 2019;860(172547):10.1016/j.ejphar.2019.172547.
54. Sale A, Berardi N, Maffei L. Environment And Brain Plasticity: Towards An Endogenous Pharmacotherapy. *Physiological Reviews*. 2014;94(1):189–234. [PubMed: 24382886]
55. Sale A. A Systematic Look at Environmental Modulation and Its Impact in Brain Development. *Trends in neurosciences*. 2018;41(1):4–17. [PubMed: 29128107]
56. Kohrt BA, Jordans MJ, Koirala S, Worthman CM. Designing mental health interventions informed by child development and human biology theory: a social ecology intervention for child soldiers in Nepal. *American journal of human biology : the official journal of the Human Biology Council*. 2015;27(1):27–40. [PubMed: 25380194]
57. Francis DD, Diorio J, Plotsky PM, Meaney MJ. Environmental enrichment reverses the effects of maternal separation on stress reactivity. *Journal of Neuroscience*. 2002;22(18):7840–7843. [PubMed: 12223535]
58. Herron ME, Buffington CAT. Environmental enrichment for indoor cats. *Compend Contin Educ Pract Vet*. 2010;32(12):E1–E5.
59. Herron ME, Buffington CA. Environmental enrichment for indoor cats: implementing enrichment. *Compend Contin Educ Vet*. 2012;34(1):E1–5.

KEY POINTS

1. Stress is often defined in terms of *stressors* – events in the internal and external environment that result in a *stress response*. Stressors increase the ratio of one’s perception of threat/perception of control.
2. The central stress response system can be sensitized by early adverse experiences, which can leave the individual more susceptible to perceive events in the environment as threats.
3. Environmental modifications to reduce perceptions of threat and enhance perceptions of control can promote resilience, which can improve health and welfare.

SYNOPSIS

The term stress can be defined in various ways. In the health sciences stress often is defined in terms of *stressors*; events that are perceived as threats to one’s perception of control. From this perspective, a stressor is anything that activates the central threat response system (CTRS). Recent research has found that the CTRS can be sensitized to environmental events through a process called epigenetic modulation of gene expression. Sensitization may result from adverse early life experiences, or by sufficiently severe events at any time of life. When CTRS activation is chronic, as can occur in confined individuals, health and welfare may be harmed. Environmental modification can mitigate the harmful effects of chronic CTRS activation by reducing the individual’s perception of threat and increasing its perception of control, which improves health and welfare.




Figure 1.

The “circle of behavior” illustrating how animals sense and respond to events in the environment. Starting at the top, when an animal acts (e.g., a cat jumps up onto a counter), the environment may respond (e.g., a shout or swat from a person to remove the cat from the counter). The sensations arising in the cat’s nervous system enter the cat’s brain to form a perception of the response, which is then compared to events in the cat’s history (genetic, epigenetic and environmental), the context in which the response was received, and its expectation of future events. These result in subsequent acts, depending on the threat or reward potential of the response. The circle is completed on a time constant of milliseconds throughout the life of the animal.