



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Although pivotal aflibercept trials used a fixed-dose regimen,¹² most retinal specialists favour a variable-dose method,⁹ suggesting that trials reflecting the real-world use of aflibercept are needed. Despite many unanswered questions, these trials present a promising step forward for the millions of patients reliant on vision-preserving anti-VEGF medications.

MAC declares no competing interests. PAK declares fees from DeepMind for consultancy in automated analysis of retinal imaging with deep learning, for Roche in personalised health care, for Novartis in personalised health care, for Apellis in automated analysis of geographic atrophy, and for BitFount in privacy protecting machine learning. PAK has received speaker fees from Heidelberg Engineering, Topcon, Allergan, and Bayer, outside the area of work commented on here. PAK declares equity ownership in Big Picture Medical, a telemedicine enterprise, unrelated to the topic commented on here.

Mark A Chia, *Pearse A Keane
pearse.keane1@nhs.net

Institute of Ophthalmology, University College London, London, UK; NIHR Biomedical Research Centre for Ophthalmology, Moorfields Eye Hospital NHS Foundation Trust, London EC1V 2PD, UK

- 1 Keane PA, Patel PJ, Liakopoulos S, Heussen FM, Sadda SR, Tufail A. Evaluation of age-related macular degeneration with optical coherence tomography. *Surv Ophthalmol* 2012; **57**: 389–414.
- 2 Adamis AP, Brittain CJ, Dandekar A, Hopkins JJ. Building on the success of anti-vascular endothelial growth factor therapy: a vision for the next decade. *Eye* 2020; **34**: 1966–72.

- 3 Arpa C, Khalid H, Chandra S, et al. Ten-year survival trends of neovascular age-related macular degeneration at first presentation. *Br J Ophthalmol* 2021; **105**: 1688–95.
- 4 Talks SJ, Stratton I, Peto T, et al. Aflibercept in clinical practice; visual acuity, injection numbers and adherence to treatment, for diabetic macular oedema in 21 UK hospitals over 3 years. *Eye* 2021; published online July 9. <https://doi.org/10.1038/s41433-021-01625-8>.
- 5 Stitt AW, Curtis TM, Chen M, et al. The progress in understanding and treatment of diabetic retinopathy. *Prog Retin Eye Res* 2016; **51**: 156–86.
- 6 Wong WL, Su X, Li X, et al. Global prevalence of age-related macular degeneration and disease burden projection for 2020 and 2040: a systematic review and meta-analysis. *Lancet Glob Health* 2014; **2**: e106–16.
- 7 Heier JS, Khanani AM, Quezada-Ruiz C, et al. Efficacy, durability, and safety of intravitreal faricimab up to every 16 weeks for neovascular age-related macular degeneration (TENAYA and LUCERNE): two randomised, double-masked, phase 3, non-inferiority trials. *Lancet* 2022; published online Jan 24. [https://doi.org/10.1016/S0140-6736\(22\)00010-1](https://doi.org/10.1016/S0140-6736(22)00010-1).
- 8 Wykoff CC, Abreu F, Adamis AP, et al. Efficacy, durability, and safety of intravitreal faricimab with extended dosing up to every 16 weeks in patients with diabetic macular oedema (YOSEMITE and RHINE): two randomised, double-masked, phase 3 trials. *Lancet* 2022; published online Jan 24. [https://doi.org/10.1016/S0140-6736\(22\)00018-6](https://doi.org/10.1016/S0140-6736(22)00018-6).
- 9 Rezaei KA, Stone TW. Global trends in retina survey. Chicago, IL: American Society of Retina Specialists, 2015.
- 10 Monés J, Srivastava SK, Jaffe GJ, et al. Risk of inflammation, retinal vasculitis, and retinal occlusion-related events with brodalumab: post hoc review of HAWK and HARRIER. *Ophthalmology* 2021; **128**: 1050–59.
- 11 Dugel PU, Koh A, Ogura Y, et al. HAWK and HARRIER: phase 3, multicenter, randomized, double-masked trials of brodalumab for neovascular age-related macular degeneration. *Ophthalmology* 2020; **127**: 72–84.
- 12 Heier JS, Brown DM, Chong V, et al. Intravitreal aflibercept (VEGF trap-eye) in wet age-related macular degeneration. *Ophthalmology* 2012; **119**: 2537–48.

COVID-19 in Europe: new challenges for addressing vaccine hesitancy



The COVID-19 pandemic and the roll-out of multiple new COVID-19 vaccines worldwide has brought global attention to the challenge of vaccine hesitancy, and it has also exposed new dynamics around routine vaccine confidence. European populations were recognised as being among the least vaccine confident in the world in 2016, with France being identified as the most vaccine sceptical.¹ Although trends over the past 5 years have shown that some countries have become more vaccine confident over time, other countries, such as Hungary and Romania, have become less confident over time.² Although some of these trends can explain the varying levels of confidence in COVID-19 vaccines,³ the issues around the acceptance of COVID-19 vaccines present new challenges.

There was already a volatile landscape of public sentiment around vaccines before COVID-19,⁴ but public sentiment around COVID-19 vaccines has been even more dynamic and rapidly changing than they have been around other vaccines,⁵ reflecting the

evolving epidemiological context and the multiple waves of infection, anxieties about the newness of the COVID-19 vaccines, the reported risks of side-effects, misinformation circulating on (social) media,

Published Online
February 3, 2022
[https://doi.org/10.1016/S0140-6736\(22\)00150-7](https://doi.org/10.1016/S0140-6736(22)00150-7)



and distrust of government and pharmaceutical industries. Against this backdrop, there have been protests in several European countries against not only COVID-19 vaccination, but broader pandemic control measures.

Although some strategies can successfully improve vaccine confidence, the context of the COVID-19 global pandemic is a challenging environment. The emergence of new variants such as omicron, for instance, has highlighted the unpredictability of the situation, leading both vaccinated and unvaccinated individuals to question the effectiveness of vaccination, creating a communication challenge for decision makers, the media, experts, and front-line health professionals.⁶ The pandemic and the multiple control measures—ranging from masking, physical distancing, SARS-CoV-2 testing, lockdowns, and quarantines, to the various vaccine requirements and mandates—have contributed to increasing societal polarisation around the limits of acceptable government controls and have had a role in widening the divide between those with differing ideologies, political opinions, or levels of compliance with control measures.^{7,8} This polarisation is particularly visible in relation to political influence on vaccine confidence. In some countries, such as Poland and the UK, political discourse has damaged public trust and reinforced division between those who vaccinate, wear masks, or respect physical distancing and those who do not, with political interests influencing policies and recommendations.^{9–11} The uncertain context of the global pandemic makes collaborative and consistent communication especially important across the political spectrum, with a united national and scientific voice providing clarity, guidance, and reassurance to populations.

Countries that worked to build confidence and address vaccine hesitancy before the pandemic were better equipped to respond to the challenges that arose during the COVID-19 pandemic. For example, the implementation of mandatory childhood vaccination in France in 2018 (pentavalent or hexavalent; measles, mumps, and rubella; meningococcal conjugate type-C; and 13-valent pneumococcal conjugate vaccines), combined with public engagement efforts, increased both uptake and public confidence in vaccination.^{12,13} Furthermore, the process generated consensus among medical societies in France, providing much-needed

endorsement to health professionals recommending vaccination to hesitant patients.

The use of top-down approaches and national vaccination or communication campaigns have done little to address specific concerns raised by specific communities, such as the mistrust of national health-care systems and authorities reported among Roma populations across Europe,¹⁴ lower health literacy in migrant populations,¹⁵ and concerns about fertility raised among some communities.¹⁶ Reaching remaining unvaccinated populations needs personal engagement and strengthening of the COVID-19 information environment by sharing positive information directly relevant to populations adapted to the social and cultural norms.¹⁷ Health professionals have a central role in providing tailored and targeted support to address vaccine hesitancy at a community level. In the UK, an organisation of general practitioners from the Black Women in Health group successfully addressed COVID-19 vaccine hesitancy and increased vaccine uptake among racially minoritised communities by organising webinars, group discussions, podcasts, and videos in different languages and dialects.¹⁸ Additionally, it has been shown in large cities in Europe that bringing the vaccine closer to specific communities, such as offering vaccines in shopping malls and easing the access to local vaccination centres, can help overcome obstacles and increase trust.¹⁹

The coming months will be crucial as we navigate a pandemic in the context of omicron and any future SARS-CoV-2 variants. Politicians across the region are making different choices in terms of policies around vaccine mandates, health passes, and other COVID-19 control measures—each weighing the limits of their health system to cope, the socioeconomic impacts of the high number of people infected with omicron who, although not hospitalised, are still unable to go to work or school, and the implications for their already worn trust relationships with the public. Ward and colleagues' study on the French health pass found it "encouraged vaccination of many who were hesitant or reluctant, but it has not reduced hesitancy itself", and highlights important considerations, particularly relating to longer-term public trust and vaccine confidence.²⁰ A UK study showed that vaccine passports were supported by those who were already positive about vaccination but hardened vaccine hesitancy and resistance among those for whom confidence-building

was especially important.²¹ Concerted efforts are needed to encourage vaccine uptake and reach those who are hesitant. Vaccine mandates should be handled with care to avoid alienating those who most need to be reached.

It is important to acknowledge that immunisation is only one element of a wider package of control measures to address COVID-19. Communication with the general public should avoid overemphasising the role of vaccination in ending the pandemic without acknowledging that other measures such as use of masks, ventilation, or physical distancing are also needed.²² In the early stages of the pandemic, optimism around rapid vaccine delivery in some European countries turned to distrust among some people when vaccine delivery was delayed, highlighting the importance of managing expectations,²³ which is particularly relevant as new variants are identified and campaigns for vaccine booster doses are implemented.

Efforts to increase vaccine confidence should be transnational. COVID-19 can easily spread between countries, and regular travel and the movement of returning diaspora can further contribute to the spread of the virus. Rumours, misinformation, and concerns about COVID-19 vaccination can travel across borders even faster than the virus, especially when facilitated by online social and digital media platforms. Coordinated approaches across European countries such as the [European Joint Action on Vaccination](#) are essential in providing consistent and harmonious communication. Increased efforts should also be placed on developing mechanisms for sharing lessons learnt about strategies that have successfully increased confidence in vaccination in Europe and globally, both for the uptake of routine vaccines and for future pandemic preparedness.

The University of Antwerp receives grants from GlaxoSmithKline Biologicals, Sanofi, Merck, MSD, Osivax, and Johnson & Johnson and from the Bill & Melinda Gates Foundation, PATH, Trasher Foundation, Flemish and Federal Government, and the European Commission. The Vaccine Confidence Project at the London School of Hygiene & Tropical Medicine receives grants from AstraZeneca, GlaxoSmithKline, Merck, and Johnson & Johnson. HL has received honoraria for developing guidance on risk communication from AstraZeneca and for leading roundtables on addressing vaccine hesitancy from GlaxoSmithKline.

Emilie Karafillakis, Pierre Van Damme, Greet Hendrickx,
*Heidi J Larson

heidi.larson@lshstm.ac.uk

Vaccine Confidence Project, London School of Hygiene & Tropical Medicine, London WC1E 7HT, UK (EK, HJL); Vaccine Confidence Project (European Regional Office), University of Antwerp, Antwerp, Belgium (PVD, GH, HJL)

- 1 Larson HJ, de Figueiredo A, Xiahong Z, et al. The state of vaccine confidence 2016: global insights through a 67-country survey. *EBioMedicine* 2016; **12**: 295–301.
- 2 European Union. The state of vaccine confidence in the EU & UK. 2020. https://ec.europa.eu/health/sites/default/files/vaccination/docs/2020_confidence_rep_en.pdf (accessed Dec 2, 2021).
- 3 de Figueiredo A, Larson HJ. Exploratory study of the global intent to accept COVID-19 vaccinations. *Commun Med* 2021; **1**: 30.
- 4 Larson HJ, Broniatowski DA. Volatility of vaccine confidence. *Science* 2021; **371**: 1289.
- 5 Valckx S, Crevecoeur J, Verelst F, et al. Individual factors influencing COVID-19 vaccine acceptance in between and during pandemic waves (July–December 2020). *Vaccine* 2022; **40**: 151–61.
- 6 Glampson B, Brittain J, Kaura A, et al. Assessing COVID-19 vaccine uptake and effectiveness through the north west London vaccination program: retrospective cohort study. *JMIR Public Health Surveill* 2021; **7**: e30010.
- 7 Sharfstein JM, Callaghan T, Carpiano RM, et al. Uncoupling vaccination from politics: a call to action. *Lancet* 2021; **398**: 1211–12.
- 8 Fridman A, Gershon R, Gneezy A. COVID-19 and vaccine hesitancy: a longitudinal study. *PLoS One* 2021; published online April 16, 2021. <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0250123>.
- 9 Pennycook G, McPhetres J, Bago B, Rand DG. Beliefs about COVID-19 in Canada, the United Kingdom, and the United States: a novel test of political polarization and motivated reasoning. *Pers Soc Psychol Bull* 2021; published online June 28. <https://doi.org/10.1177%2F01461672211023652>.
- 10 Walkowiak MP, Walkowiak D. Predictors of COVID-19 vaccination campaign success: lessons learnt from the pandemic so far. A case study from Poland. *Vaccines (Basel)* 2021; **9**: 1153.
- 11 Sowa P, Kiszkiel L, Laskowski PP, et al. COVID-19 vaccine hesitancy in Poland—multifactorial impact trajectories. *Vaccines (Basel)* 2021; **9**: 876.
- 12 Sante publique France. Semaine europeenne de la vaccination du 24 au 30 avril 2019: une campagne nationale inedite et des premiers resultats positifs de l'obligation vaccinale. 2019. <http://www.santepubliquefrance.fr/les-actualites/2019/semaine-europeenne-de-la-vaccination-du-24-au-30-avril-2019> (accessed Dec 20, 2021).
- 13 Weil-Olivier C. Mandatory vaccination in France. Presentation at HPV Prevention Board. December, 2021. <https://medialibrary.uantwerpen.be/files/5700/270b1c31-caf4-449c-8c09-027317abeaaf.pdf> (accessed Dec 20, 2021).
- 14 Holt E. Covid-19 vaccination among Roma populations in Europe. *Lancet Microbe* 2021; **2**: e289.
- 15 Crawshaw AF, Deal A, Rustage K, et al. What must be done to tackle vaccine hesitancy and barriers to COVID-19 vaccination in migrants? *J Travel Med* 2021; **28**: taab048.
- 16 Carmody ER, Zander D, Klein EJ, Mulligan MJ, Caplan AL. Knowledge and attitudes toward COVID-19 and vaccines among a New York Haredi-Orthodox Jewish community. *J Community Health* 2021; **46**: 1161–69.
- 17 Petousis-Harris H, Chan A. Research review. Countering vaccine misinformation: a practical guide for healthcare providers. 2021. <https://lnkd.in/g/WN6UbqX> (accessed Dec 20, 2021).
- 18 Imohi O, Momoh C, Ologun A. COVID-19 vaccine hesitancy—debunking the myths using a community engagement approach underpinned by NICE guidance. 2021. <https://www.nice.org.uk/sharedlearning/covid-19-vaccine-hesitancy-debunking-the-myths-using-a-community-engagement-approach-underpinned-by-nice-guidance> (accessed Dec 20, 2021).
- 19 Deal A, Hayward SE, Huda M, et al. Strategies and action points to ensure equitable uptake of COVID-19 vaccinations: a national qualitative interview study to explore the views of undocumented migrants, asylum seekers, and refugees. *J Migr Health* 2021; **4**: 100050.
- 20 Ward JK, Gauna F, Gagneux-Brunon A, et al. The French health pass holds lessons for mandatory COVID-19 vaccination. *Nat Med* 2022; published online Jan 12. <https://doi.org/10.1038/s41591-021-01661-7>.
- 21 de Figueiredo A, Larson HJ, Reicher SD. The potential impact of vaccine passports on inclination to accept COVID-19 vaccinations in the United Kingdom: evidence from a large cross-sectional survey and modelling study. *EclinicalMed* 2021; **40**: 10110.
- 22 Su Z, Wen J, McDonnell D, et al. Vaccines are not yet a silver bullet: the imperative of continued communication about the importance of COVID-19 safety measures. *Brain Behav Immun* 2021; **12**: 100204.
- 23 Warren GW, Lofstedt R. COVID-19 vaccine rollout risk communication strategies in Europe: a rapid response. *J Risk Res* 2021; **24**: 369–79.

For more on the **European Joint Action on Vaccination** see <https://eu-jav.com/>