

CORRECTION

Open Access



# Correction to: Statistical analysis plan for better evidence for selecting transplant fluids (BEST-fluids): a randomised controlled trial of the effect of intravenous fluid therapy with balanced crystalloid versus saline on the incidence of delayed graft function in deceased donor kidney transplantation

Elaine M. Pascoe<sup>1</sup>, Steven J. Chadban<sup>2,3</sup>, Magid A. Fahim<sup>1,4</sup>, Carmel M. Hawley<sup>1,4,5</sup>, David W. Johnson<sup>1,4,5</sup>, Michael G. Collins<sup>1,6,7\*</sup>, for the BEST-fluids Investigators and the Australasian Kidney Trials Network

## Correction to: Pascoe (2022)

<http://orcid.org/10.1186/s13063-021-05989-w>

Following the publication of the original article [1], we were notified that an incorrect additional file was published alongside the paper.

The original article has been corrected.

Published online: 07 February 2022

## Reference

1. Pascoe EM, et al. Statistical analysis plan for Better Evidence for Selecting Transplant Fluids (BEST-Fluids): a randomised controlled trial of the effect of intravenous fluid therapy with balanced crystalloid versus saline on the incidence of delayed graft function in deceased donor kidney transplantation. 2022, 2022;23:52. <https://doi.org/10.1186/s13063-021-05989-w>.

## Author details

<sup>1</sup>Australasian Kidney Trials Network, The University of Queensland, Brisbane, Australia. <sup>2</sup>Department of Renal Medicine, Royal Prince Alfred Hospital, Sydney, Australia. <sup>3</sup>Charles Perkins Centre, The University of Sydney, Sydney, Australia. <sup>4</sup>Department of Nephrology, Princess Alexandra Hospital, Brisbane, Australia. <sup>5</sup>The Translational Research Institute, Brisbane, Australia. <sup>6</sup>Department of Renal Medicine, Auckland District Health Board, Auckland City Hospital, Level 15, Park Road, Grafton, Auckland, New Zealand. <sup>7</sup>Department of Medicine, Faculty of Medical and Health Sciences, University of Auckland, Auckland, New Zealand.

The original article can be found online at <https://doi.org/10.1186/s13063-021-05989-w>.

\* Correspondence: [Michael.Collins@adhb.govt.nz](mailto:Michael.Collins@adhb.govt.nz)

<sup>1</sup>Australasian Kidney Trials Network, The University of Queensland, Brisbane, Australia

<sup>6</sup>Department of Renal Medicine, Auckland District Health Board, Auckland City Hospital, Level 15, Park Road, Grafton, Auckland, New Zealand

Full list of author information is available at the end of the article



© The Author(s). 2022 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.