




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Letter to the Editor

Clinical characteristics of SARS-CoV-2 fully-vaccinated patients dying with COVID-19 in Italy

Luigi Palmieri¹, Marco Florida^{2,*}, Marina Giuliano², Dorina Tiple³, Cinzia Lo Noce¹, Paola Meli⁴, Graziano Onder¹, on behalf of the Italian National Institute of Health COVID-19 mortality group

¹) Department of Cardiovascular, Endocrine-metabolic Diseases and Aging, Istituto Superiore di Sanità, Rome, Italy

²) National Center for Global Health, Istituto Superiore di Sanità, Rome, Italy

³) Department of Neuroscience, Istituto Superiore di Sanità, Rome, Italy

⁴) National Center for Innovative Technologies in Public Health, Istituto Superiore di Sanità, Rome, Italy

ARTICLE INFO

Article history:

Received 12 November 2021

Received in revised form

28 December 2021

Accepted 27 January 2022

Available online 15 February 2022

Editor: R. Chemaly

To the Editor,

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) vaccination campaign started in Italy on December 27, 2020. As of October 2021, more than 82% of the Italian population age ≥ 12 years was fully vaccinated with one of the four vaccines approved by the European Medicines Agency (BNT162b2, mRNA-1273, ChAdOx1, Ad26.COV2–S) [1]. As in other countries, vaccine access was progressively extended from high-risk and older individuals to younger populations. Concurrently with the increasing number of vaccinated people, deaths related to coronavirus disease 2019 (COVID-19) in vaccinated persons also increased [2]. The aim of this study was to evaluate the characteristics of persons who died with COVID-19 after completing the vaccine course and to compare them with those who died with COVID-19 but were not vaccinated or had a partial vaccination course.

Data were obtained from the Italian National Institute of Health (Istituto Superiore di Sanità) Integrated Surveillance System [3]. COVID-19–related deaths were defined as those occurring in patients who tested positive for SARS-CoV-2, independent of pre-existing comorbidities possibly contributing to death. Clinical

data were obtained from a random sample of hospital charts [4]. The sample analyzed was not selected on the basis of sample size calculations, but included all clinical records available in the database. Individual data were linked with those of SARS-CoV-2 vaccination registries.

Three groups of COVID-19–related deaths were defined according to vaccination status were defined:

1. Unvaccinated: deaths in patients with COVID-19 who had not received any vaccine dose;
2. Partially vaccinated with early infection: deaths of patients with COVID-19 who tested positive for SARS-CoV-2 within 14 days from the beginning of the vaccination course; and
3. Fully vaccinated: deaths of patients with COVID-19 who tested positive for SARS-CoV-2 14 days after completion of the vaccination course [5].

The partially vaccinated with early infection group identified patients who acquired SARS-CoV-2 infection in the first 14 days after their first vaccination, when the risk of infection is comparable to that of the unvaccinated population. The comparison of this group with the fully vaccinated group reduces the possible bias related to the fact that vaccination priority was initially given to the very old and frail.

Deaths of patients who tested positive for SARS-CoV-2 between February 1 and October 5, 2021 were evaluated. February 1 was chosen as the index date because it included the 5-week interval needed to complete the vaccination course after the introduction of SARS-CoV-2 vaccination in Italy. To compare patient characteristics, we used *t* tests for normally distributed variables, nonparametric Mann–Whitney U-test for skewed variables, and Fisher's exact test for categorical variables.

At a national level, during the study period, 38,096 COVID-19–related deaths occurred. Of these, 1440 (3.8%) were in fully vaccinated persons and 2130 (5.6%) in partially vaccinated individuals. The present analysis is based on a sample of 760

* Corresponding author. Marco Florida, National Centre for Global Health, Istituto Superiore di Sanità, Rome, Italy.

E-mail address: marco.florida@iss.it (M. Florida).

Table 1
Clinical characteristics of COVID-19 deaths by study group

	Unvaccinated <i>n</i> = 760 (A)	Partially vaccinated with early infection <i>n</i> = 379 (B)	Fully vaccinated <i>n</i> = 236 (C)	p-value (A) vs (C)	p-value (B) vs (C)
Demographics					
Age (y), mean (standard deviation)	77.1 (14.0)	83.5 (8.6)	85.5 (9.3)	<0.001	0.007
Women, <i>n</i> (%)	322 (42.4)	150 (39.6)	98 (41.5)	0.880	0.673
Comorbidities, <i>n</i> (%)					
Ischemic heart disease	199 (26.5)	110 (29.2)	85 (36.2)	0.005	0.075
Atrial fibrillation	190 (25.3)	127 (33.7)	81 (34.5)	0.007	0.861
Heart failure	103 (13.7)	60 (15.9)	59 (25.1)	<0.001	0.006
Stroke	71 (9.5)	35 (9.3)	27 (11.5)	0.382	0.410
Hypertension	484 (64.5)	265 (70.3)	164 (69.8)	0.156	0.928
Type 2 diabetes	218 (29.1)	106 (28.1)	66 (28.1)	0.805	1.000
Dementia	129 (17.2)	89 (23.6)	87 (37.0)	<0.001	<0.001
Chronic obstructive pulmonary disease	119 (15.9)	75 (19.9)	57 (24.3)	0.005	0.225
Cancer	99 (13.2)	47 (12.5)	43 (18.3)	0.056	0.060
Chronic liver disease	39 (5.2)	20 (5.3)	14 (6.0)	0.622	0.721
Autoimmune diseases	52 (6.9)	25 (6.6)	16 (6.8)	1.000	1.000
Obesity	120 (16.0)	54 (14.3)	19 (8.1)	0.002	0.021
Number of comorbidities, <i>n</i> (%)				<0.001	0.020
0	27 (3.6)	3 (0.8)	0 (0)		
1	77 (10.3)	37 (9.8)	14 (6.0)		
2	129 (17.2)	58 (15.4)	23 (9.8)		
≥3	517 (68.9)	279 (74.0)	198 (84.3)		
Number of comorbidities, mean (standard deviation)	3.8 (2.2)	4.1 (2.3)	5.0 (2.5)	<0.001	<0.001
Nonrespiratory complications, <i>n</i> (%)					
Acute renal injury	212 (28.2)	113 (30.1)	75 (32.2)	0.249	0.652
Acute cardiac injury	63 (8.4)	24 (6.4)	26 (11.2)	0.193	0.048
Coinfection	204 (27.1)	91 (24.3)	64 (27.5)	0.933	0.390

Unvaccinated indicates deaths occurring in patients with COVID-19 who had not received any vaccine dose (*n* with missing data for comorbidities = 10); partially vaccinated with early infection indicates deaths occurring in patients with COVID-19 who tested positive for severe acute respiratory syndrome coronavirus 2 within 14 days from the beginning of the vaccination course (*n* with missing data for comorbidities = 2); fully vaccinated indicates deaths occurring in patients infected with COVID-19 who tested positive for SARS coronavirus type 2 14 days after completion of the vaccination course (*n* with missing data for comorbidities = 1). COVID-19, coronavirus disease 2019.

deaths in the unvaccinated group (2.3% of deaths in the same group), 379 deaths in the partially vaccinated group (17.8%), and 236 deaths in the fully vaccinated group (16.4%). The representativeness of this sample is presented in the Appendix (Table S1), together with the distribution of vaccine types in the sample analyzed and in the whole population of deaths during the study period (Table S2) and with calendar time of the first dose administration in fully and partially vaccinated individuals (Fig. S1).

The main study results are shown in Table 1. Fully vaccinated individuals, compared to unvaccinated persons, were significantly older and had a higher prevalence of ischemic heart disease, atrial fibrillation, heart failure, dementia, and chronic obstructive pulmonary disease, with a lower prevalence of obesity. Compared with partially vaccinated individuals, fully vaccinated persons were older, had a higher prevalence of heart failure and dementia, and a lower prevalence of obesity. The mean number of comorbidities was significantly higher in fully vaccinated persons compared with the other two groups. Among nonrespiratory COVID-19 complications, acute cardiac injury was significantly more frequent in fully vaccinated compared to partially vaccinated individuals.

These findings show that people who died with COVID-19 after completing the vaccination course have a significantly higher burden of comorbidities compared with those who died unvaccinated or with a partial vaccination course. This more severe clinical complexity appears to be multidimensional, with a significantly higher prevalence of heart disease, dementia, and chronic obstructive pulmonary disease. The significantly lower prevalence of obesity in the fully vaccinated group may indicate a possible protective effect of vaccination against mortality in obese persons that should be investigated further.

In terms of study limitations, we were unable to evaluate additional groups of unvaccinated/partially vaccinated and fully vaccinated patients with COVID-19 who did not die, because the clinical records of hospitalized patients who survive are not collected by the mortality surveillance system. The fully vaccinated group received the first dose of the vaccine earlier than the partially vaccinated group (Fig. S1), and we cannot exclude that the differences observed between these two groups could be explained by the vaccination priority given to older and more complex patients. Selection bias from the general population was not likely to have occurred: the sample appeared to be representative of the population of patients deceased with COVID-19 and of the vaccine types used in Italy. All cases included were homogeneous in terms of health care facilities, being all in-hospital deaths, and the clinical records database used represents a random sample of all COVID-19 deaths in which all Italian regions are represented. Access to vaccination was uniform in the entire country, with large availability of vaccine doses in all regions. The timeframe of our analysis also excluded the first wave of the pandemic, when the burden on health care facilities in terms of hospitalizations and admissions to intensive care units was higher and the system suffered major stress.

In conclusion, we believe that the present findings may contribute to the definition of a population at higher risk of mortality after full vaccination. This population with more severe clinical complexity should be considered for possible interventions aimed at reducing the risks of infection, such as the use of additional vaccine doses.

Transparency declaration

Conflicts of interest/competing interests: None to declare. None of the authors has a commercial or other association, financial

interest, activity, relationship or association that might pose a conflict of interest.

No funding was received for this work.

Authors' contribution

GO designed the study and drafted the manuscript. LP was responsible for statistical analysis; LP, MF, MG, DT, CLN and PM substantially contributed to acquisition of data and critical revision of the manuscript. All the authors approved the final version to be published.

Members of the Italian National Institute of Health COVID-19 mortality group

Luigi Palmieri, Elvira Agazio, Pierfrancesco Barbariol, Antonino Bella, Eva Benelli, Luigi Bertinato, Matilde Bocci, Stefano Boros, Marco Bressi, Giovanni Calcagnini, Marco Canevelli, Federica Censi, Alessandra Ciervo, Elisa Colaizzo, Roberto Da Cas, Martina Del Manso, Corrado Di Benedetto, Chiara Donfrancesco, Massimo Fabiani, Francesco Facchiano, Marco Florida, Fabio Galati, Marina Giuliano, Tiziana Grisetti, Cecilia Guastadisegni, Ilaria Lega, Cinzia Lo Noce, Pietro Maiozzi, Valerio Manno, Margherita Martini, Marco Massari, Alberto Mateo Urdiales, Eugenio Mattei, Claudia Meduri, Paola Meli, Francesca Menniti Ippolito, Giada Minelli, Graziano Onder, Daniele Petrone, Patrizio Pezzotti, Flavia Pricci, Ornella

Punzo, Federica Quarata, Valeria Raparelli, Flavia Riccardo, Simone Rocchetto, Chiara Sacco, Paolo Salerno, Giulia Sarti, Debora Serra, Stefania Spila Alegiani, Matteo Spuri, Marco Tallon, Manuela Tamburo De Bella, Dorina Tiple, Marco Toccaceli Blasi, Federica Trentin, Brigid Unim, Luana Vaianella, Nicola Vanacore, Maria Fenicia Vescio, Emanuele Rocco Villani, Liliana Elena Weimer, Silvio Brusaferrò.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.cmi.2022.01.024>.

References

- [1] Government of Italy. Vaccination report. Available at: <https://www.governo.it/it/cscovid19/report-vaccini/>. [Accessed 27 October 2021].
- [2] Istituto Superiore di Sanità. Epidemia COVID-19. Available at: https://www.epicentro.iss.it/coronavirus/bollettino/Bollettino-sorveglianza-integrata-COVID-19_6-ottobre-2021.pdf. [Accessed 22 October 2021].
- [3] Onder G, Rezza G, Brusaferrò S. Case-fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. *JAMA* 2020;323:1775–6.
- [4] Palmieri L, Vanacore N, Donfrancesco C, Lo Noce C, Canevelli M, Punzo O, et al. Clinical characteristics of hospitalized individuals dying with COVID-19 by age group in Italy. *J Gerontol A Biol Sci Med Sci* 2020;75:1796–800.
- [5] US Centers for Disease Control and Prevention. Stay up to date with your vaccines. Available at: <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/fully-vaccinated.html>. [Accessed 22 October 2021].