

## Article

# Radiographic Bone Loss and Its Relation to Patient-Specific Risk Factors, LDL Cholesterol, and Vitamin D: A Cross-Sectional Study

Teresa Thim <sup>1,2</sup>, Konstantin Johannes Scholz <sup>1</sup>, Karl-Anton Hiller <sup>1</sup>, Wolfgang Buchalla <sup>1</sup>,  
Christian Kirschneck <sup>3</sup>, Jonathan Fleiner <sup>4</sup>, Johan Peter Woelber <sup>5,†</sup> and Fabian Cieplik <sup>1,\*,†</sup>

<sup>1</sup> Department of Conservative Dentistry and Periodontology, University Hospital Regensburg, 93053 Regensburg, Germany; teresa.thim@web.de (T.T.); konstantin.scholz@ukr.de (K.J.S.); karl-anton.hiller@ukr.de (K.-A.H.); wolfgang.buchalla@ukr.de (W.B.)

<sup>2</sup> Private Practice, 63110 Rodgau, Germany

<sup>3</sup> Department of Orthodontics, University Hospital Regensburg, 93053 Regensburg, Germany; christian.kirschneck@ukr.de

<sup>4</sup> Center of Dental Implantology, Periodontology and 3D-Imaging, 78462 Konstanz, Germany; fleiner@stricker-fleiner.de

<sup>5</sup> Department of Operative Dentistry and Periodontology, Faculty of Medicine, University of Freiburg, 79085 Freiburg, Germany; johan.woelber@uniklinik-freiburg.de

\* Correspondence: fabian.cieplik@ukr.de

† These authors contributed equally to this work.



**Citation:** Thim, T.; Scholz, K.J.; Hiller, K.-A.; Buchalla, W.; Kirschneck, C.; Fleiner, J.; Woelber, J.P.; Cieplik, F. Radiographic Bone Loss and Its Relation to Patient-Specific Risk Factors, LDL Cholesterol, and Vitamin D: A Cross-Sectional Study. *Nutrients* **2022**, *14*, 864. <https://doi.org/10.3390/nu14040864>

Academic Editor: Maria Luz Fernandez

Received: 14 January 2022

Accepted: 16 February 2022

Published: 18 February 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** The influence of patient-specific factors such as medical conditions, low-density lipoprotein cholesterol (LDL-C) or levels of 25-hydroxyvitamin D (25OHD) on periodontal diseases is frequently discussed in the literature. Therefore, the aim of this retrospective cross-sectional study was to evaluate potential associations between radiographic bone loss (RBL) and patient-specific risk factors, particularly LDL-C and 25OHD levels. Patients from a dental practice, who received full-mouth cone beam CTs (CBCTs) and blood-sampling in the course of implant treatment planning, were included in this study. RBL was determined at six sites per tooth from CBCT data. LDL-C and 25OHD levels were measured from venous blood samples. Other patient-specific risk factors were assessed based on anamnesis and dental charts. Statistical analysis was performed applying non-parametric procedures (Mann–Whitney U tests, error rates method). Data from 163 patients could be included in the analysis. RBL was significantly higher in male patients, older age groups, smokers, patients with high DMFT (decayed/missing/filled teeth) score, lower number of teeth, and high LDL-C levels ( $\geq 160$  mg/dL). Furthermore, patients with high 25OHD levels ( $\geq 40$  ng/mL) exhibited significantly less RBL. In summary, RBL was found to be associated with known patient-specific markers, particularly with age and high LDL-C levels.

**Keywords:** periodontal; bone loss; LDL; vitamin D; CBCT; radiographic bone loss

## 1. Introduction

Periodontal diseases are among the most prevalent non-communicable diseases in mankind with 1.1 billion prevalent cases of severe periodontitis worldwide according to the Global Burden of Disease 2019 study [1,2]. In Germany, 51.6% of younger adults (35–44 years) and 64.6% of younger seniors (65–74 years) are affected by moderate to severe periodontal disease, as reported in the 5th German Oral Health study (DMS V) [3].

The pathogenesis of periodontal diseases is associated with the presence of subgingival biofilms and considered to be based on a host-mediated dysbiosis of the oral microbiota due to an exaggerated response of the host immune system resulting in a loss of periodontal supporting tissues [4–6]. General medical conditions such as diabetes mellitus or habits like smoking are associated with periodontal disease [7,8]. Recently, also metabolic disorders

as well as systemic inflammation were discussed in literature as a risk of causing or accelerating periodontal bone loss [9].

One of these recently discussed conditions is hypercholesterolemia, and in particular high levels of low-density lipoprotein cholesterol (LDL-C) [10]. Hypercholesterolemia is a common phenomenon, and epidemiological data show that increased LDL-C levels ( $\geq 130$  mg/dL) were prevalent in 29.4% of American adults [11]. High serum levels of LDL-C are often accompanied with a diet high in saturated or trans fats and sugar, physical inactivity, smoking, obesity, type 2 diabetes mellitus, and high blood pressure [12,13]. High LDL-C levels are also a main risk factor for atherosclerotic cardiovascular diseases [14], which represented the leading cause of death among US-Americans in 2017 [11].

Another condition, which was lately brought into focus in context with periodontitis and radiographic bone loss, is vitamin D deficiency [15,16]. Serum levels of 25-hydroxy vitamin D (25OHD), which is an intermediate product in vitamin D metabolism, lower than 20 ng/mL are defined as deficiency [17]. 25OHD deficiency is a widespread condition around the world, especially in northern regions [18], since vitamin D is mostly synthesized in the skin if exposed to sunlight, while just a little part is supplied by nutrition, e.g., by consumption of oily fish [17]. There are several health benefits known which are associated with sufficient 25OHD levels as vitamin D is important for an adequate bone mineralization and for various functions of the immune system [19,20] as well as there is a reduced risk for cardiovascular diseases [21] or cancer [22].

There are some investigations about increased LDL-C and decreased 25OHD levels within periodontitis-patients compared to healthy controls [16,23–28]. However, to the best of the authors' knowledge, there is no study investigating on a "healthy" cross-section of daily dental patients and evaluating potential risk factors for periodontal bone loss using a highly validated method such as cone beam CTs.

Against this background, the aim of this retrospective cross-sectional study was to investigate associations between radiographic bone loss (RBL) and patient-specific general health parameters, particularly levels of LDL-C and 25OHD, in a cohort of patients treated in a private dental practice. The null-hypothesis tested was that levels of LDL-C or 25OHD, respectively, were not associated with RBL.

## 2. Materials and Methods

### 2.1. Study Design

The present study was designed as a retrospective cross-sectional study. The objective was to evaluate RBL according to patient-specific parameters including sex, age, smoking history, DMFT (decayed/missing/filled teeth) score, number of teeth, LDL-C, and 25OHD levels. Data were collected from a cohort of patients who received treatment planning for dental implants in a private practice.

The study design was approved by the internal review board of the University of Regensburg, Germany (reference: 21-2431-104; issued on 23 June 2021) in accordance with the 1964 Helsinki Declaration and its later amendments and comparable ethical standards. The study was registered at the German Clinical Trials Register (ref: DRKS00025827).

### 2.2. Patient Population

All patients from the patient pool of a private practice in Rodgau (Hessen, Germany), who had received a cone beam CT (CBCT) as well as analysis of LDL-C and 25OHD levels in the course of treatment planning for dental implants between February 2017 and October 2020 were screened for inclusion in this study. Patients were excluded if they had less than 16 teeth [29] or in case of insufficient CBCT quality. No other exclusion criteria were applied.

### 2.3. Medical and Dental History


A detailed anamnesis of the medical history and intake of medications was obtained from the dental charts and anamnesis forms, and if necessary, complemented by telephone interviews. Smoking history was recorded as pack-years (PY), which were calculated by

multiplying the number of packs of cigarettes smoked per day by the number of years the person has smoked [30]. Additionally, the intake of statins or vitamin D supplementation was checked. Dental charts were checked for numbers of teeth and teeth were charted as decayed, missing, or filled (DMFT index) according to the clinical oral examination prior to the implant treatment planning, which was double-checked with the CBCT radiographs.

#### 2.4. Radiographic Examination

All CBCTs were conducted with the medical indication of treatment planning for dental implants as full-mouth CBCTs (Orthophos XG 3D, Dentsply-Sirona, Bensheim, Germany). The field of view was set to  $8 \times 8$  cm, voxel size was  $160 \mu\text{m}$ , scan time was 5.1 s, the voltage was 85 kV, and the current was 7 mA.

For evaluation of the radiographic bone loss (RBL), the software package CoPeriodontiX 9.9 (Dental Wings, Chemnitz, Germany) was used [31]. For each patient, all teeth except the third molars were measured. RBL was defined as distance between alveolar crest (AC) and cemento-enamel junction (CEJ) or restoration margin (RM) in cases of teeth with restorations (e.g., crowns) on six sites per tooth (mesio-buccal, buccal, disto-buccal, mesio-oral, oral, disto-oral). For measurement of RBL, every single tooth was manually positioned three-dimensionally according to its longitudinal axis and its CEJ or RM. After adjusting, CEJ or RM and AC had to be marked on six aspects of the tooth so that RBL could be calculated by the program. Figure 1 shows the workflow for determining RBL in the program package.



**Figure 1.** (A–C) Tooth selection and positioning. Tooth 15 in a distal-mesial (A), vestibular-oral (B), and axial (C) plane. The crosshair is positioned according to the longitudinal axis and CEJ or RM (A,B) and parallel to the cross-axis of the tooth (C). Every single tooth was selected and adjusted manually. (D–F) Setting the reference points. The yellow line marks the longitudinal axis of tooth 15. It is presented in three different cross-sections, oral/vestibular (D), vestibular-distal/oral-mesial (E), and vestibular-mesial/oral-distal (F). Twelve reference points per tooth (as depicted in green and red color) were set on the CEJ or RM and AC for each side. The red dot marks the AC oral (D), oral-mesial (E), and oral-distal (F). The software measures RBL by calculating the distance between the dots.

All CBCTs were examined by one examiner (TT), who had been extensively trained by an expert (JF). For validation of the accuracy of the RBL measurements, ten randomly chosen CBCTs were re-evaluated and differences between the first and second measurement were assessed for intra-examiner agreement.

### 2.5. LDL-C and 25OHD Levels

All evaluations of serum LDL-C and 25OHD were conducted in the course of treatment planning for dental implants. Two venous blood samples (2 mL each) were taken in a fasting state by a trained examiner from the basilic vein. The samples were sent to a specialized and accredited laboratory (Institut für medizinische Diagnostik, Berlin, Germany) for evaluation of serum levels of LDL-C and 25OHD. LDL-C levels were determined by enzymatic tests and the physical unit was mg/dL. For evaluation of 25OHD, electrochemiluminescence immunoassay (ECLIA) was conducted and 25OHD levels were measured in ng/mL. All LDL-C and 25OHD data were retrieved retrospectively from the dental charts.

### 2.6. Data Analysis

The maximum of the six RBL values per tooth was determined as the descriptive value for each tooth. The median of these maxima over all existing teeth of a patient except third molars was used as the RBL value of a patient for analysis. From all patients, medians including first and third quartiles from RBL values were calculated.

Patients were categorized, as follows: Three age groups were formed ( $\leq 44$  years; 45–59 years;  $\geq 60$  years). Smoking history was differentiated in “non-smokers”, “smokers with  $\leq 15$  PY”, and “smokers with  $\geq 16$  PY”. For DMFT score and number of teeth, the patients were divided in four groups ( $\leq 11$ ; 12–18; 19–23;  $\geq 24$ ) or two groups ( $\leq 24$ ;  $\geq 25$ ), respectively. LDL-C levels were subdivided as “optimal” LDL-C ( $\leq 99$  mg/dL), “near optimal” LDL-C (100–129 mg/dL), “borderline high” LDL-C (130–159 mg/dL) and “high” LDL-C ( $\geq 160$  mg/dL) according to the U.S. National Cholesterol Education Program [32]. 25OHD levels were categorized as “deficiency” ( $\leq 19$  ng/mL), “insufficiency” (20–29 ng/mL), “sufficiency” (30–39 ng/mL), and “optimal” ( $\geq 40$  ng/mL) [17,19].

For analysis of RBL categorized to the different LDL-C groups, patients reporting intake of statins were excluded ( $n = 5$ ). Accordingly, patients who reported supplementing vitamin D were excluded for RBL analyses categorized to the different 25OHD groups ( $n = 46$ ).

Differences among experimental groups for matching parameters were evaluated statistically using non-parametric Mann-Whitney U tests on a significance level of  $\alpha = 0.05$ . For evaluation of a general influence of a given parameter on all groups, the level of significance was adjusted to  $\alpha^*(k) = 1 - (1 - \alpha)^{1/k}$  ( $k =$  number of pairwise tests) according to the error rates method, yielding an  $\alpha^*(3) = 0.01695423$  and an  $\alpha^*(6) = 0.00851244$  for three or six pairwise tests (i.e., three or four categorized groups), respectively [33]. All statistical analyses were performed using SPSS for Windows, version 26 (SPSS Inc., Chicago, IL, USA).

RBL, age, and LDL-C or 25OHD levels, respectively, were put into a three-dimensional curve-fitting model and depicted accordingly. TableCurve 3D automated surface fitting analysis software (SYSTAT Software Inc., Systat Software Inc, San Jose, CA, USA; version 4.0) was used to find equations to describe the three-dimensional empirical data.

## 3. Results


### 3.1. Patient Population

CBCTs, LDL-C, and 25OHD levels were available from 178 patients. Twelve patients with less than 16 teeth were excluded. Three patients were excluded because quality of CBCT was too poor for further analysis (extensive artifacts due to metal-based restorations). A total of 163 patients could be included in this study.

The patient cohort comprised 100 females (61.3%) and 63 males (38.7%), the median (first; third quartile) age was 53 (44; 62) years and 80.4% were non-smokers. All patients exhibited in median (first; third quartile) 25 (22; 27) teeth, a DMFT score of 19 (14; 22), LDL-C level of 127 (107; 156) mg/dL, and 25OHD level of 29 (20; 43) ng/mL (Table 1). Median (first; third quartile) period of time between the CBCT and taking of the blood samples was 15 (1; 50) days. Figure 2 shows the flow of patients through the stages of this study.

**Table 1.** Patient characteristics (DMFT score, number of teeth, LDL-C, 25OHD). Depiction of medians (first; third quartiles) and statistically significant differences from pairwise comparisons (Mann–Whitney U tests;  $\alpha = 0.05$ ).

	All Patients	≤44 Years	45–59 Years	≥ 60 Years	≤44 vs. 45–59	≤44 vs. ≥60	45–59 vs. ≥60
DMFT score §	19 (14; 22)	14.5 (11; 19)	18 (14; 20)	21 (18.3; 23)	0.025	0.000	0.000
Number of teeth §	25 (22; 27)	27 (24.8; 28)	25 (22.5; 27)	23 (19.3; 25)	0.005	0.000	0.002
LDL-C § [mg/dL]	127 (107; 156)	109.5 (91.3; 127.5)	132 (113; 151.5)	146.5 (118.3; 174.3)	0.000	0.000	0.028
25OHD [ng/mL]	29 (20; 42)	27.5 (19.8; 40)	30 (19.5; 47)	30 (20.5; 39)	—	—	—




**Figure 2.** The chart depicts the flow of patients through the stages of this study.

For evaluation of the general influence of age on a given parameter,  $\alpha$  was adjusted according to the error rates method to  $\alpha^*(3) = 0.01695243$ . *p*-value: pairwise significant difference ( $p \leq 0.05$ ); —: no pairwise significant difference; § significant influence of age groups on the respective parameter according to the error rates method.

Table 1 shows DMFT score, number of teeth, LDL-C and 25OHD categorized according to the distinct age groups. According to the error rates method, all age groups showed statistically significant differences with regard to DMFT score, number of teeth, and LDL-C. DMFT score and LDL-C were significantly higher, and number of teeth was significantly lower in older age groups as compared to younger age groups. For 25OHD, there were no statistically significant differences among the age groups.

### 3.2. RBL

The RBL validation measurements revealed a median (first; third quartile) difference between the first and second measurements of  $-0.1$  ( $-0.3$ ;  $0.1$ ) mm, thus showing sufficient intra-examiner accuracy. Median (first; third quartile) RBL was  $3.6$  ( $3.2$ ;  $4.2$ ) mm for all patients (Figure 3A). Females had significantly smaller median RBL ( $3.5$  mm) than males ( $3.8$  mm;  $p = 0.026$ ; Figure 3A).



**Figure 3.** Results of RBL analysis for (A) all patients ( $n = 163$ ) and divided into male ( $n = 63$ ) and female ( $n = 100$ ) and according to (B) the different age-groups ( $\leq 44$  y,  $n = 42$ ;  $45\text{--}59$  y,  $n = 65$ ;  $\geq 60$  y,  $n = 56$ ) and (C) smoking groups (non-smokers,  $n = 131$ ;  $1\text{--}15$  py,  $n = 15$ ;  $\geq 16$  py,  $n = 17$ ). Results are depicted as medians, first and third quartiles and asterisks depict statistically significant differences between the groups. \* marks significant differences with  $p \leq 0.05$ ; \*\*\* marks significant differences with  $p \leq 0.001$ .


Patients  $\leq 44$  years showed significantly smaller median RBL ( $3.2$  mm) than patients between  $45$  and  $59$  years ( $3.6$  mm) and patients  $\geq 60$  years ( $4.1$  mm). The differences in RBL were statistically significant between all age groups ( $p = 0.000$  in all cases; Figure 3B) and accordingly, there was a general influence of the parameter age on RBL according to the error rates method.

Smokers reporting a smoking history of  $\geq 16$  PY had significantly higher median RBL values ( $4.1$  mm) than non-smokers ( $3.6$  mm;  $p = 0.029$ ; Figure 3C). There were no significant differences between the two smoking groups or between the non-smokers and the smoking group with  $1\text{--}15$  PY.


DMFT score analysis showed that RBL increased with DMFT (Figure 4A). Patients with DMFT score  $\leq 11$  showed significantly smaller median RBL ( $3.5$  mm) as compared to patients with DMFT score between  $19$  and  $23$  ( $3.8$  mm;  $p = 0.036$ ) and patients with DMFT score  $\geq 24$  ( $4.0$  mm;  $p = 0.019$ ). Likewise, patients with DMFT score between  $12$  and  $18$  showed significantly smaller median RBL ( $3.5$  mm) than those with DMFT score  $\geq 24$  ( $p = 0.020$ ). It was also found that patients with  $24$  teeth and less had significantly higher median RBL ( $3.9$  mm) than the ones with  $\geq 25$  teeth ( $3.5$  mm;  $p = 0.000$ ), as shown in Figure 4B.

Figure 5 shows RBL according to LDL-C (Figure 5A) and 25OHD groups (Figure 5B). Patients with high LDL-C ( $\geq 160$  mg/dL) showed significantly higher median RBL ( $3.9$  mm) than those with optimal ( $\leq 99$  mg/dL;  $3.4$  mm;  $p = 0.000$ ), near optimal ( $100\text{--}129$  mg/dL;  $3.5$  mm;  $p = 0.009$ ) and borderline high LDL-C ( $130\text{--}159$  mg/dL;  $3.7$  mm;  $p = 0.033$ ). Accordingly, the error rates method revealed a general influence of the parameter LDL-C on RBL. Patients with optimal 25OHD ( $\geq 40$  ng/mL) showed significantly lower median RBL ( $3.4$  mm) than those with deficient ( $\leq 19$  ng/mL;  $3.6$  mm;  $p = 0.029$ ) and sufficient 25OHD ( $30\text{--}39$  ng/mL;  $3.8$  mm;  $p = 0.031$ ). No general influence of the parameter 25OHD on RBL was detected by the error rates method.


RBL, age, and LDL-C or 25OHD levels, respectively, were put into a three-dimensional curve-fitting model and depicted accordingly. When depicting age and LDL-C, Figure 6A shows an irreducible influence of both parameters on RBL. When depicting age and 25OHD, Figure 6B shows that RBL is mainly influenced by age but not by 25OHD levels.



**Figure 4.** Results of RBL analysis according to (A) the different DMFT score groups ( $\leq 11$ ,  $n = 23$ ; 12–18,  $n = 57$ ; 19–23,  $n = 67$ ;  $\geq 24$ ,  $n = 16$ ) and (B) number of teeth ( $\leq 24$ ,  $n = 73$ ;  $\geq 25$ ,  $n = 90$ ). Results are depicted as medians, first and third quartiles and asterisks depict statistically significant differences between the groups. \* marks significant differences with  $p \leq 0.05$ ; \*\*\* marks significant differences with  $p \leq 0.001$ .



**Figure 5.** Results of RBL analysis according to the different (A) LDL-C groups ( $\leq 99$ ,  $n = 30$ ; 100–129,  $n = 51$ ; 130–159,  $n = 41$ ;  $\geq 160$ ,  $n = 36$ ) and (B) 25OHD groups ( $\leq 19$ ,  $n = 38$ ; 20–29,  $n = 38$ ; 30–39,  $n = 21$ ;  $\geq 40$ ,  $n = 20$ ). Results are depicted as medians, first and third quartiles and asterisks depict statistically significant differences between the groups. \* marks significant differences with  $p \leq 0.05$ ; \*\* marks significant differences with  $p \leq 0.01$ ; \*\*\* marks significant differences with  $p \leq 0.001$ .



**Figure 6.** Influence of age and LDL-C or 25OHD levels, respectively, on RBL as fitted three-dimensional curves. (A) Influence of age and LDL-C level on RBL as fitted three-dimensional curve (Cosine Series Bivariate Order 5;  $r^2 = 0.44$ ). (B) Influence of age and 25OHD level on RBL as fitted three-dimensional curve (Fourier Series Simple Order  $2 \times 4$ ;  $r^2 = 0.37$ ).

#### 4. Discussion

The aim of the present study was to investigate potential associations between RBL and patient-specific parameters like sex, age, smoking history, DMFT score, number of teeth, LDL-C, and 25OHD in a cohort of patients, who received treatment planning for dental implants. Focusing on RBL as reference value allowed evaluation of the accumulated history of periodontal destruction and reflected a longer period of time, regardless of the current state of clinical periodontal health.

The data of our study showed that there were significantly higher RBLs in males and in the older age groups, which is in line with other studies. Helmi et al. also evaluated radiographic alveolar bone loss in a cohort-study and revealed significant higher RBLs in men and older age groups [34]. Eke et al. found a higher prevalence of periodontal disease in men as well as an increasing prevalence of periodontitis in the older age groups [35]. Aging is accompanied with modifications of the host immune response, which leads to greater susceptibility to infections and autoimmunity [36]. The higher RBL in males may be explained since men may be less attentive to their (oral) health and consequently may exhibit worse oral hygiene levels, leading to higher RBLs [37]. Furthermore, the immune response is different in men and women, whereby men show higher levels of pro-inflammatory cytokines during infections [37,38].

Furthermore, RBL was also found to be significantly higher in smokers with  $\geq 16$  PY compared to non-smokers. A similar outcome could be found in other studies, which had measured radiographic alveolar bone loss [34,39]. Smoking is known as a risk factor for onset and progression of periodontitis [40,41] and has also been included as grade modifier in the 2018 classification of periodontal diseases [42]. Smoking is known to impair the host response to the dental plaque biofilm and to be linked to increased levels of potentially destructive inflammatory cytokines and enzymes [43]. Furthermore, smoking diminishes the reparative capacity of periodontal cells, including fibroblasts, osteoblasts and cementoblasts, thus potentially resulting in a higher RBL in smokers [43].

The patient cohort investigated in the present study exhibited a median DMFT score of 19 with a median number of 25 teeth per patient, which clearly outnumbers the results reported in the fifth German Oral Health study for the respective age groups (mean DMFT score of 11.2 or 17.7 for age groups 35–44 or 65–74, respectively) [3]. The reason might be the fact that the investigated cohort were seeking for treatment with dental implants. The older age groups showed significantly higher DMFT score, significantly lower number of teeth, and significantly higher RBL than patients from the other groups. To the best of the authors' knowledge, there has been no other study investigating RBL and DMFT score or numbers of teeth. Levin et al. evaluated the DMFT score and number of missing teeth in periodontitis patients compared to healthy controls [44]. They did not find statistically significant differences regarding DMFT scores but did find a significantly higher number of missing teeth in periodontitis patients. Strauss et al. and Mattila et al. investigated the co-occurrence of periodontitis and caries and found significantly higher numbers of decayed teeth in patients with periodontitis [45,46]. Tooth loss represents the end stage of oral diseases such as periodontitis or caries, thus representing an objective marker for the accumulated inflammatory burden of oral disease. Therefore, there may be common risk factors for caries, periodontitis, and tooth loss such as nutrition, limitations in oral hygiene, and not seeking dental treatment [45].

Pre-conditions such as high LDL-C levels or insufficient levels of 25OHD may be linked to higher RBL [10,15]. Median LDL-C was 127 mg/dL in the patient cohort. According to Virani et al. mean LDL-C among American adults was 112.1 mg/dL and prevalence of LDL-C levels  $\geq 130$  mg/dL was 29.4% [11]. In our study, 47.9% were found to have LDL-C levels  $\geq 130$  mg/dL. Thus, the investigated patient cohort had slightly higher values of LDL-C than the average population. As found by Waskiewicz et al., patients suffering from tooth loss and thus seeking for treatment with dental implants might have higher LDL-C values [47]. Significantly higher LDL-C levels were found in the older patient groups which is in line with the literature [48,49], and may be due to an age-associated loss of hepatic



LDL receptors, higher body-mass index, larger waist circumference and lower sex hormone levels [49].

LDL-C was found to have a significant influence on RBL according to the error rates method and the high LDL-C group ( $\geq 160$  mg/dL) exhibited significantly higher RBL than all other groups in pairwise comparisons. Due to the general influence of age groups on LDL-C as well as RBL according to the error rates method, it cannot be clarified entirely from the data of this study which of both parameters had the bigger influence on RBL (see Figure 6A). However, there are a few more studies, which reported significant associations between high LDL-C and periodontitis when investigating clinical parameters [23–26]. Furthermore, a meta-analysis and meta-regression concluded that periodontitis patients had significantly higher levels of LDL-C [10]. Conversely, Monteiro et al. and Saxlin et al. did not find any significant differences between periodontitis patients and healthy control patients regarding levels of LDL-C [50,51]. Potential associations between LDL-C and periodontal status can be discussed in two ways. The presence of a periodontal infection negatively affects serum lipid levels by an altered immune cell function which leads to a dysregulation of the lipid metabolism [23,52]. On the other hand, high LDL-C levels lead to an increase in periodontal destruction because of an activation of osteoclasts and inhibition of osteoblasts [53] and by the release of pro-inflammatory cytokines [8,54]. Furthermore, higher LDL-C values can be interpreted as a marker of disease-promoting lifestyles, which primarily lead to a higher periodontal inflammation [12,55]. There are also common gen polymorphisms, which are risk factors for both diseases, periodontal disease and hyperlipidemia [56].

The median 25OHD level was 29 ng/mL in the present study, and 23.9% of all patients showed vitamin D deficiency ( $\leq 19$  ng/mL). Another German investigation found median 25OHD of 44.9 nmol/L, which correspond to 18 ng/mL, and 57.3% of 3,917 subjects were found to be deficient of 25OHD [57]. We found no notable difference regarding 25OHD levels between the different age groups, which is in line with the literature [58,59].

RBL was found to be significantly higher in patients with “deficient” 25OHD levels found as compared to patients with “optimal” ( $\geq 40$  ng/mL) 25OHD levels. These results are in line with a recently published systematic review and meta-analysis concluding that periodontitis is associated with lower 25OHD levels [15]. This concurs with the known bone-protective effect of higher 25OHD levels, which have been shown to decrease the ratio of RANKL to OPG expression by periodontal ligament fibroblasts controlling osteoclastogenesis [60]. Ketharanathan et al. focused on radiographic bone loss in their cohort study to investigate the impact of 25OHD levels in periodontitis patients. They found that patients with periodontal disease comprised higher radiographic alveolar bone loss (as measured on bitewing radiographs) and lower 25OHD which corresponds to our findings [16]. In another study, clinical attachment loss was evaluated and compared to 25OHD levels [58]. There was significantly less attachment loss in patients with high 25OHD levels, but only in the older age group ( $\geq 60$  years) [58], which matches with the results from the present study, where age definitely had a higher influence on RBL than 25OHD levels, as depicted in Figure 6B. Similar results have also been shown by other studies [27,28]. Noteworthy, Perić et al. found a tendency for better healing outcomes following non-surgical periodontal therapy in patients who took vitamin D as a supplement than in patients without vitamin D supplementation [61]. There are two potential ways that vitamin D may affect the periodontal status. First, there are effects on bone mineral density especially in the elderly [62], and second, vitamin D may reduce gingival inflammation through anti-inflammatory effects on the general host immune response [15,63,64]. In addition, low salivary levels of 25OHD were found to be associated with higher levels of inflammatory biomarkers in periodontitis patients [65]. Furthermore, there is evidence that vitamin D supplementation reduces systemic inflammation and levels of pro-inflammatory salivary cytokines [66,67] and gingival bleeding [64], whereas vitamin D deficiency is supposed to be a risk factor for periodontal treatment failure [68]. It is also suggested that genetic variants of the vitamin D receptor are a biomarker for periodontitis [69].

As a potential limitation of the present study, it must be emphasized that RBL shows a history of periodontal destruction and aging, but gives no information on the current state of clinical periodontal health. Although aging is also strongly associated with RBL (as shown in Figure 6) [70], periodontal disease is considered to be the major cause for alveolar bone loss [71]. Accordingly, other studies showed that there is a reliable relationship between clinical and radiographic bone loss [72–74]. Clinical bone loss precedes radiographic findings six to eight months [75]. In addition, a high accuracy of CBCTs in periodontal diagnosis, especially in visualizing periodontal intra-bony and furcation defects, has been shown [76,77].

While DMFT score, number of teeth, and smoking history also reflect a longer period of time, measurement of LDL-C and 25OHD levels just reflects a current snapshot. Nevertheless, it may be assumed that the determined LDL-C and 25OHD levels are a marker for individual lifestyle and health constitution of the patients and are stable for longer periods of time, particularly due to exclusion of patients receiving “treatment” in the form of statins or vitamin D supplementation.

Although diabetes mellitus and rheumatoid arthritis are known to be associated with periodontal disease [78,79], no sub-analysis regarding those parameters was possible in the present cohort of patients since there were only two diabetes mellitus and four rheumatoid arthritis patients. The small number may be explained by the fact that only patients were included who were treated with dental implants, where diabetes and rheumatoid arthritis are known to be relative contraindications for treatment with dental implants [80,81].

## 5. Conclusions

The present study detected significant associations between RBL and patient-specific parameters like sex, smoking history, DMFT score, number of teeth, 25OHD levels, and particularly age and LDL-C. While RBL gives no information on the current state of clinical periodontal health, but reflects the cumulated burden of periodontal destruction, the outcomes of this study support similar findings of previous studies investigating clinical periodontal parameters. Future studies using RBL measurements should also include clinical periodontal parameters as well as further investigations of the association between lifestyle- and nutrition-linked conditions such as LDL-C and 25OHD levels and periodontal bone loss.

**Author Contributions:** Conceptualization, F.C., J.P.W., T.T., K.J.S., J.F.; methodology, T.T., F.C., J.P.W., J.F., K.J.S.; software, J.F., T.T.; validation, K.-A.H., C.K., J.F., W.B.; formal analysis, K.-A.H., T.T., K.J.S., F.C., J.P.W.; investigation, T.T., K.-A.H.; data curation, K.-A.H.; writing—original draft preparation, T.T.; writing—review and editing, F.C., J.P.W., K.J.S., K.-A.H., W.B., C.K., J.F.; visualization, T.T., F.C., K.-A.H.; supervision, F.C., J.P.W. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding and was solely based on institutional funding.

**Institutional Review Board Statement:** The study was conducted in accordance with the guidelines of the Declaration of Helsinki, and approved by the Institutional Review Board of the University of Regensburg (reference: 21-2431-104; issued on 23 June 2021).

**Informed Consent Statement:** Patient consent was not applicable for this type of retrospective study.

**Data Availability Statement:** All data supporting the reported results are available upon request from the corresponding author.

**Acknowledgments:** Dental Wings (Chemnitz, Germany) is gratefully acknowledged for providing the software package CoPeriodontiX 9.9.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Chen, M.X.; Zhong, Y.J.; Dong, Q.Q.; Wong, H.M.; Wen, Y.F. Global, regional, and national burden of severe periodontitis, 1990–2019: An analysis of the Global Burden of Disease Study 2019. *J. Clin. Periodontol.* **2021**, *48*, 1165–1188. [[CrossRef](#)] [[PubMed](#)]

2. James, S.L.; Abate, D.; Abate, K.H.; Abay, S.M.; Abbafati, C.; Abbasi, N.; Abbastabar, H.; Abd-Allah, F.; Abdela, J.; Abdelalim, A.; et al. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: A systematic analysis for the Global Burden of Disease Study 2017. *Lancet* **2018**, *392*, 1789–1858. [[CrossRef](#)]
3. Jordan, R.A.; Bodechtel, C.; Hertrampf, K.; Hoffmann, T.; Kocher, T.; Nitschke, I.; Schiffner, U.; Stark, H.; Zimmer, S.; Micheelis, W. The Fifth German Oral Health Study (Fünfte Deutsche Mundgesundheitsstudie, DMS V)—Rationale, design, and methods. *BMC Oral Health* **2014**, *14*, 161. [[CrossRef](#)]
4. Bartold, P.M.; van Dyke, T.E. Periodontitis: A host-mediated disruption of microbial homeostasis. Unlearning learned concepts. *Periodontology 2000* **2013**, *62*, 203–217. [[CrossRef](#)]
5. Hajishengallis, G.; Korostoff, J.M. Revisiting the Page & Schroeder model: The good, the bad and the unknowns in the periodontal host response 40 years later. *Periodontology 2000* **2017**, *75*, 116–151. [[CrossRef](#)] [[PubMed](#)]
6. Jakubovics, N.S.; Goodman, S.D.; Mashburn-Warren, L.; Stafford, G.P.; Cieplik, F. The dental plaque biofilm matrix. *Periodontology 2000* **2021**, *86*, 32–56. [[CrossRef](#)] [[PubMed](#)]
7. Grossi, S.G.; Zambon, J.J.; Ho, A.W.; Koch, G.; Dunford, R.G.; Machtei, E.E.; Norderyd, O.M.; Genco, R.J. Assessment of risk for periodontal disease. I. Risk indicators for attachment loss. *J. Periodontol.* **1994**, *65*, 260–267. [[CrossRef](#)] [[PubMed](#)]
8. Loos, B.G.; van Dyke, T.E. The role of inflammation and genetics in periodontal disease. *Periodontology 2000* **2020**, *83*, 26–39. [[CrossRef](#)]
9. Jain, P.; Hassan, N.; Khatoon, K.; Mirza, M.A.; Naseef, P.P.; Kuruniyan, M.S.; Iqbal, Z. Periodontitis and Systemic Disorder—An Overview of Relation and Novel Treatment Modalities. *Pharmaceutics* **2021**, *13*, 1175. [[CrossRef](#)]
10. Nepomuceno, R.; Pigossi, S.C.; Finoti, L.S.; Orrico, S.R.P.; Cirelli, J.A.; Barros, S.P.; Offenbacher, S.; Scarel-Caminaga, R.M. Serum lipid levels in patients with periodontal disease: A meta-analysis and meta-regression. *J. Clin. Periodontol.* **2017**, *44*, 1192–1207. [[CrossRef](#)]
11. Virani, S.S.; Alonso, A.; Benjamin, E.J.; Bittencourt, M.S.; Callaway, C.W.; Carson, A.P.; Chamberlain, A.M.; Chang, A.R.; Cheng, S.; Delling, F.N.; et al. Heart Disease and Stroke Statistics—2020 Update: A Report from the American Heart Association. *Circulation* **2020**, *141*, e139–e596. [[CrossRef](#)] [[PubMed](#)]
12. Karr, S. Epidemiology and management of hyperlipidemia. *Am. J. Manag. Care* **2017**, *23*, S139–S148. [[PubMed](#)]
13. Stanhope, K.L.; Schwarz, J.M.; Keim, N.L.; Griffen, S.C.; Bremer, A.A.; Graham, J.L.; Hatcher, B.; Cox, C.L.; Dyachenko, A.; Zhang, W.; et al. Consuming fructose-sweetened, not glucose-sweetened, beverages increases visceral adiposity and lipids and decreases insulin sensitivity in overweight/obese humans. *J. Clin. Investig.* **2009**, *119*, 1322–1334. [[CrossRef](#)] [[PubMed](#)]
14. Ference, B.A.; Ginsberg, H.N.; Graham, I.; Ray, K.K.; Packard, C.J.; Bruckert, E.; Hegele, R.A.; Krauss, R.M.; Raal, F.J.; Schunkert, H.; et al. Low-density lipoproteins cause atherosclerotic cardiovascular disease. 1. Evidence from genetic, epidemiologic, and clinical studies. A consensus statement from the European Atherosclerosis Society Consensus Panel. *Eur. Heart J.* **2017**, *38*, 2459–2472. [[CrossRef](#)] [[PubMed](#)]
15. Machado, V.; Lobo, S.; Proença, L.; Mendes, J.J.; Botelho, J. Vitamin D and Periodontitis: A Systematic Review and Meta-Analysis. *Nutrients* **2020**, *12*, 2177. [[CrossRef](#)] [[PubMed](#)]
16. Ketharanathan, V.; Torgersen, G.R.; Petrovski, B.É.; Preus, H.R. Radiographic alveolar bone level and levels of serum 25-OH-Vitamin D3 in ethnic Norwegian and Tamil periodontitis patients and their periodontally healthy controls. *BMC Oral Health* **2019**, *19*, 83. [[CrossRef](#)] [[PubMed](#)]
17. Holick, M.F.; Chen, T.C. Vitamin D deficiency: A worldwide problem with health consequences. *Am. J. Clin. Nutr.* **2008**, *87*, 1080S–1086S. [[CrossRef](#)]
18. Amrein, K.; Scherkl, M.; Hoffmann, M.; Neuwersch-Sommeregger, S.; Köstenberger, M.; Tmava Berisha, A.; Martucci, G.; Pilz, S.; Malle, O. Vitamin D deficiency 2.0: An update on the current status worldwide. *Eur. J. Clin. Nutr.* **2020**, *74*, 1498–1513. [[CrossRef](#)] [[PubMed](#)]
19. Charoengam, N.; Holick, M.F. Immunologic Effects of Vitamin D on Human Health and Disease. *Nutrients* **2020**, *12*, 2097. [[CrossRef](#)]
20. Prietl, B.; Treiber, G.; Pieber, T.R.; Amrein, K. Vitamin D and immune function. *Nutrients* **2013**, *5*, 2502–2521. [[CrossRef](#)] [[PubMed](#)]
21. Zhang, R.; Li, B.; Gao, X.; Tian, R.; Pan, Y.; Jiang, Y.; Gu, H.; Wang, Y.; Wang, Y.; Liu, G. Serum 25-hydroxyvitamin D and the risk of cardiovascular disease: Dose-response meta-analysis of prospective studies. *Am. J. Clin. Nutr.* **2017**, *105*, 810–819. [[CrossRef](#)] [[PubMed](#)]
22. Yin, L.; Ordóñez-Mena, J.M.; Chen, T.; Schöttker, B.; Arndt, V.; Brenner, H. Circulating 25-hydroxyvitamin D serum concentration and total cancer incidence and mortality: A systematic review and meta-analysis. *Prev. Med.* **2013**, *57*, 753–764. [[CrossRef](#)] [[PubMed](#)]
23. Awartani, F.; Atassi, F. Evaluation of periodontal status in subjects with hyperlipidemia. *J. Contemp. Dent. Pract.* **2010**, *11*, 33–40. [[CrossRef](#)]
24. Shivakumar, T.; Patil, V.A.; Desai, M.H. Periodontal status in subjects with hyperlipidemia and determination of association between hyperlipidemia and periodontal health: A clinicobiochemical study. *J. Contemp. Dent. Pract.* **2013**, *14*, 785–789. [[CrossRef](#)] [[PubMed](#)]
25. Fentoğlu, O.; Oz, G.; Taşdelen, P.; Uskun, E.; Aykaç, Y.; Bozkurt, F.Y. Periodontal status in subjects with hyperlipidemia. *J. Periodontol.* **2009**, *80*, 267–273. [[CrossRef](#)] [[PubMed](#)]

26. Lee, S.; Im, A.; Burm, E.; Ha, M. Association between periodontitis and blood lipid levels in a Korean population. *J. Periodontol.* **2018**, *89*, 28–35. [[CrossRef](#)] [[PubMed](#)]
27. Anbarcioglu, E.; Kirtiloglu, T.; Öztürk, A.; Kolbakir, F.; Acıkgöz, G.; Colak, R. Vitamin D deficiency in patients with aggressive periodontitis. *Oral Dis.* **2019**, *25*, 242–249. [[CrossRef](#)]
28. Kim, H.; Shin, M.H.; Yoon, S.J.; Kweon, S.S.; Lee, Y.H.; Choi, C.K.; Kim, O.; Kim, Y.J.; Chung, H.; Kim, O.S. Low serum 25-hydroxyvitamin D levels, tooth loss, and the prevalence of severe periodontitis in Koreans aged 50 years and older. *J. Periodontal Implant Sci.* **2020**, *50*, 368–378. [[CrossRef](#)] [[PubMed](#)]
29. Ebersole, J.L.; Lambert, J.; Bush, H.; Huja, P.E.; Basu, A. Serum Nutrient Levels and Aging Effects on Periodontitis. *Nutrients* **2018**, *10*, 1986. [[CrossRef](#)]
30. Lebowitz, M.D.; Burrows, B. Quantitative relationships between cigarette smoking and chronic productive cough. *Int. J. Epidemiol.* **1977**, *6*, 107–113. [[CrossRef](#)]
31. Fleiner, J.; Hannig, C.; Schulze, D.; Stricker, A.; Jacobs, R. Digital method for quantification of circumferential periodontal bone level using cone beam CT. *Clin. Oral Investig.* **2013**, *17*, 389–396. [[CrossRef](#)]
32. National Cholesterol Education Program. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) Final Report. *Circulation* **2002**, *106*, 3143–3421.
33. Miller, R.G., Jr. *Simultaneous Statistical Inference*, 2nd ed.; Springer: New York, NY, USA, 1981; ISBN 9781461381228.
34. Helmi, M.F.; Huang, H.; Goodson, J.M.; Hasturk, H.; Tavares, M.; Natto, Z.S. Prevalence of periodontitis and alveolar bone loss in a patient population at Harvard School of Dental Medicine. *BMC Oral Health* **2019**, *19*, 254. [[CrossRef](#)]
35. Eke, P.I.; Dye, B.A.; Wei, L.; Slade, G.D.; Thornton-Evans, G.O.; Borgnakke, W.S.; Taylor, G.W.; Page, R.C.; Beck, J.D.; Genco, R.J. Update on Prevalence of Periodontitis in Adults in the United States: NHANES 2009 to 2012. *J. Periodontol.* **2015**, *86*, 611–622. [[CrossRef](#)]
36. Ebersole, J.L.; Graves, C.L.; Gonzalez, O.A.; Dawson, D.; Morford, L.A.; Huja, P.E.; Hartsfield, J.K.; Huja, S.S.; Pandruvada, S.; Wallet, S.M. Aging, inflammation, immunity and periodontal disease. *Periodontology 2000* **2016**, *72*, 54–75. [[CrossRef](#)] [[PubMed](#)]
37. Lipsky, M.S.; Su, S.; Crespo, C.J.; Hung, M. Men and Oral Health: A Review of Sex and Gender Differences. *Am. J. Mens. Health* **2021**, *15*, 15579883211016361. [[CrossRef](#)]
38. Shiau, H.J.; Reynolds, M.A. Sex differences in destructive periodontal disease: Exploring the biologic basis. *J. Periodontol.* **2010**, *81*, 1505–1517. [[CrossRef](#)]
39. Rosa, G.M.; Lucas, G.Q.; Lucas, O.N. Cigarette smoking and alveolar bone in young adults: A study using digitized radiographs. *J. Periodontol.* **2008**, *79*, 232–244. [[CrossRef](#)]
40. Knight, E.T.; Liu, J.; Seymour, G.J.; Faggion, C.M.; Cullinan, M.P. Risk factors that may modify the innate and adaptive immune responses in periodontal diseases. *Periodontology 2000* **2016**, *71*, 22–51. [[CrossRef](#)]
41. Ryder, M.I.; Couch, E.T.; Chaffee, B.W. Personalized periodontal treatment for the tobacco- and alcohol-using patient. *Periodontology 2000* **2018**, *78*, 30–46. [[CrossRef](#)]
42. Papananou, P.N.; Sanz, M.; Buduneli, N.; Dietrich, T.; Feres, M.; Fine, D.H.; Flemmig, T.F.; Garcia, R.; Giannobile, W.V.; Graziani, F.; et al. Periodontitis: Consensus report of workgroup 2 of the 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions. *J. Clin. Periodontol.* **2018**, *45* (Suppl. S20), S162–S170. [[CrossRef](#)] [[PubMed](#)]
43. Chaffee, B.W.; Couch, E.T.; Vora, M.V.; Holliday, R.S. Oral and periodontal implications of tobacco and nicotine products. *Periodontology 2000* **2021**, *87*, 241–253. [[CrossRef](#)] [[PubMed](#)]
44. Levin, L.; Zini, A.; Levine, J.; Weiss, M.; Lev, R.; Chebath Taub, D.; Hai, A.; Almozni, G. Demographic profile, Oral Health Impact Profile and Dental Anxiety Scale in patients with chronic periodontitis: A case-control study. *Int. Dent. J.* **2018**, *68*, 269–278. [[CrossRef](#)]
45. Strauss, F.-J.; Espinoza, I.; Stähli, A.; Baeza, M.; Cortés, R.; Morales, A.; Gamonal, J. Dental caries is associated with severe periodontitis in Chilean adults: A cross-sectional study. *BMC Oral Health* **2019**, *19*, 278. [[CrossRef](#)] [[PubMed](#)]
46. Mattila, P.T.; Niskanen, M.C.; Vehkalahti, M.M.; Nordblad, A.; Knuutila, M.L.E. Prevalence and simultaneous occurrence of periodontitis and dental caries. *J. Clin. Periodontol.* **2010**, *37*, 962–967. [[CrossRef](#)]
47. Waskiewicz, K.; Oth, O.; Kochan, N.; Evrard, L. Des facteurs de risque généralement négligés en chirurgie orale et en implantologie: Le taux élevé de LDL-cholestérol et le taux insuffisant de la vitamine D. *Rev. Med. Brux.* **2018**, *39*, 70–77. [[CrossRef](#)] [[PubMed](#)]
48. Wang, M.; Hou, X.; Hu, W.; Chen, L.; Chen, S. Serum lipid and lipoprotein levels of middle-aged and elderly Chinese men and women in Shandong Province. *Lipids Health Dis.* **2019**, *18*, 58. [[CrossRef](#)]
49. Rosada, A.; Kassner, U.; Weidemann, F.; König, M.; Buchmann, N.; Steinhagen-Thiessen, E.; Spira, D. Hyperlipidemias in elderly patients: Results from the Berlin Aging Study II (BASEII), a cross-sectional study. *Lipids Health Dis.* **2020**, *19*, 92. [[CrossRef](#)]
50. Monteiro, A.M.; Jardini, M.A.N.; Alves, S.; Giampaoli, V.; Aubin, E.C.Q.; Figueiredo Neto, A.M.; Gidlund, M. Cardiovascular disease parameters in periodontitis. *J. Periodontol.* **2009**, *80*, 378–388. [[CrossRef](#)]
51. Saxlin, T.; Suominen-Taipale, L.; Kattainen, A.; Marniemi, J.; Knuutila, M.; Ylöstalo, P. Association between serum lipid levels and periodontal infection. *J. Clin. Periodontol.* **2008**, *35*, 1040–1047. [[CrossRef](#)]
52. Iacopino, A.M.; Cutler, C.W. Pathophysiological relationships between periodontitis and systemic disease: Recent concepts involving serum lipids. *J. Periodontol.* **2000**, *71*, 1375–1384. [[CrossRef](#)]

53. Choukroun, J.; Khoury, G.; Khoury, F.; Russe, P.; Testori, T.; Komiyama, Y.; Sammartino, G.; Palacci, P.; Tunali, M.; Choukroun, E. Two neglected biologic risk factors in bone grafting and implantology: High low-density lipoprotein cholesterol and low serum vitamin D. *J. Oral Implantol.* **2014**, *40*, 110–114. [[CrossRef](#)] [[PubMed](#)]
54. Noack, B.; Jachmann, I.; Roscher, S.; Sieber, L.; Kopprasch, S.; Lück, C.; Hanefeld, M.; Hoffmann, T. Metabolic diseases and their possible link to risk indicators of periodontitis. *J. Periodontol.* **2000**, *71*, 898–903. [[CrossRef](#)] [[PubMed](#)]
55. Woelber, J.P.; Tennert, C. Chapter 13: Diet and Periodontal Diseases. *Monogr. Oral Sci.* **2020**, *28*, 125–133. [[CrossRef](#)] [[PubMed](#)]
56. Wang, X.; Li, W.; Song, W.; Xu, L.; Zhang, L.; Feng, X.; Lu, R.; Meng, H. Association of CYP1A1 rs1048943 variant with aggressive periodontitis and its interaction with hyperlipidemia on the periodontal status. *J. Periodontol. Res.* **2019**, *54*, 546–554. [[CrossRef](#)] [[PubMed](#)]
57. Linseisen, J.; Bechthold, A.; Bischoff-Ferrari, H.A.; Hintzpetter, B.; Leschik-Bonnet, E.; Reichrath, J.; Stehle, P.; Volkert, D.; Wolfram, G.; Zittermann, A. Vitamin D und Prävention Ausgewählter Chronischer Krankheiten—Stellungnahme. Available online: <https://www.dge.de/wissenschaft/weitere-publikationen/stellungnahmen/?L=0> (accessed on 13 January 2022).
58. Dietrich, T.; Joshipura, K.J.; Dawson-Hughes, B.; Bischoff-Ferrari, H.A. Association between serum concentrations of 25-hydroxyvitamin D3 and periodontal disease in the US population. *Am. J. Clin. Nutr.* **2004**, *80*, 108–113. [[CrossRef](#)] [[PubMed](#)]
59. Bettencourt, A.; Boleixa, D.; Reis, J.; Oliveira, J.C.; Mendonça, D.; Costa, P.P.; Da Silva, B.M.; Marinho, A.; Da Silva, A.M. Serum 25-hydroxyvitamin D levels in a healthy population from the North of Portugal. *J. Steroid Biochem. Mol. Biol.* **2018**, *175*, 97–101. [[CrossRef](#)]
60. Küchler, E.C.; Schröder, A.; Teodoro, V.B.; Nazet, U.; Scariot, R.; Spanier, G.; Proff, P.; Kirschneck, C. The role of 25-hydroxyvitamin D3 and vitamin D receptor gene in human periodontal ligament fibroblasts as response to orthodontic compressive strain: An in vitro study. *BMC Oral Health* **2021**, *21*, 386. [[CrossRef](#)] [[PubMed](#)]
61. Perić, M.; Maiter, D.; Cavalier, E.; Lasserre, J.F.; Toma, S. The Effects of 6-Month Vitamin D Supplementation during the Non-Surgical Treatment of Periodontitis in Vitamin-D-Deficient Patients: A Randomized Double-Blind Placebo-Controlled Study. *Nutrients* **2020**, *12*, 2940. [[CrossRef](#)] [[PubMed](#)]
62. Dawson-Hughes, B.; Harris, S.S.; Krall, E.A.; Dallal, G.E. Effect of calcium and vitamin D supplementation on bone density in men and women 65 years of age or older. *N. Engl. J. Med.* **1997**, *337*, 670–676. [[CrossRef](#)] [[PubMed](#)]
63. Botelho, J.; Machado, V.; Proença, L.; Delgado, A.S.; Mendes, J.J. Vitamin D Deficiency and Oral Health: A Comprehensive Review. *Nutrients* **2020**, *12*, 1471. [[CrossRef](#)]
64. Dietrich, T.; Nunn, M.; Dawson-Hughes, B.; Bischoff-Ferrari, H.A. Association between serum concentrations of 25-hydroxyvitamin D and gingival inflammation. *Am. J. Clin. Nutr.* **2005**, *82*, 575–580. [[CrossRef](#)] [[PubMed](#)]
65. Costantini, E.; Sinjari, B.; Piscopo, F.; Porreca, A.; Reale, M.; Caputi, S.; Murmura, G. Evaluation of Salivary Cytokines and Vitamin D Levels in Periodontopathic Patients. *Int. J. Mol. Sci.* **2020**, *21*, 2669. [[CrossRef](#)] [[PubMed](#)]
66. Garcia, M.N.; Hildebolt, C.F.; Miley, D.D.; Dixon, D.A.; Couture, R.A.; Spearie, C.L.A.; Langenwaller, E.M.; Shannon, W.D.; Deych, E.; Mueller, C.; et al. One-year effects of vitamin D and calcium supplementation on chronic periodontitis. *J. Periodontol.* **2011**, *82*, 25–32. [[CrossRef](#)] [[PubMed](#)]
67. Meghil, M.M.; Hutchens, L.; Raed, A.; Multani, N.A.; Rajendran, M.; Zhu, H.; Looney, S.; Elashiry, M.; Arce, R.M.; Peacock, M.E.; et al. The influence of vitamin D supplementation on local and systemic inflammatory markers in periodontitis patients: A pilot study. *Oral Dis.* **2019**, *25*, 1403–1413. [[CrossRef](#)] [[PubMed](#)]
68. Bashutski, J.D.; Eber, R.M.; Kinney, J.S.; Benavides, E.; Maitra, S.; Braun, T.M.; Giannobile, W.V.; McCauley, L.K. The impact of vitamin D status on periodontal surgery outcomes. *J. Dent. Res.* **2011**, *90*, 1007–1012. [[CrossRef](#)]
69. Yu, X.; Zong, X.; Pan, Y. Associations between vitamin D receptor genetic variants and periodontitis: A meta-analysis. *Acta Odontol. Scand.* **2019**, *77*, 484–494. [[CrossRef](#)]
70. Streckfus, C.F.; Parsell, D.E.; Streckfus, J.E.; Pennington, W.; Johnson, R.B. Relationship between oral alveolar bone loss and aging among African-American and Caucasian individuals. *Gerontology* **1999**, *45*, 110–114. [[CrossRef](#)]
71. Jeffcoat, M.K. Bone loss in the oral cavity. *J. Bone Miner. Res.* **1993**, *8* (Suppl. S2), S467–S473. [[CrossRef](#)]
72. Machtei, E.E.; Hausmann, E.; Grossi, S.G.; Dunford, R.; Genco, R.J. The relationship between radiographic and clinical changes in the periodontium. *J. Periodontol. Res.* **1997**, *32*, 661–666. [[CrossRef](#)] [[PubMed](#)]
73. Farook, F.F.; Alodwene, H.; Alharbi, R.; Alyami, M.; Alshahrani, A.; Almohammadi, D.; Alnasyan, B.; Aboelmaaty, W. Reliability assessment between clinical attachment loss and alveolar bone level in dental radiographs. *Clin. Exp. Dent. Res.* **2020**, *6*, 596–601. [[CrossRef](#)] [[PubMed](#)]
74. Jeffcoat, M.K. Radiographic Methods for the Detection of Progressive Alveolar Bone Loss. *J. Periodontol.* **1992**, *63* (Suppl. S4), 367–372. [[CrossRef](#)] [[PubMed](#)]
75. Goodson, J.M.; Haffajee, A.D.; Socransky, S.S. The relationship between attachment level loss and alveolar bone loss. *J. Clin. Periodontol.* **1984**, *11*, 348–359. [[CrossRef](#)] [[PubMed](#)]
76. Walter, C.; Schmidt, J.C.; Dula, K.; Sculean, A. Cone beam computed tomography (CBCT) for diagnosis and treatment planning in periodontology: A systematic review. *Quintessence Int.* **2016**, *47*, 25–37. [[CrossRef](#)] [[PubMed](#)]
77. Woelber, J.P.; Fleiner, J.; Rau, J.; Ratka-Krüger, P.; Hannig, C. Accuracy and Usefulness of CBCT in Periodontology: A Systematic Review of the Literature. *Int. J. Periodontics Restor. Dent.* **2018**, *38*, 289–297. [[CrossRef](#)] [[PubMed](#)]
78. Ceccarelli, F.; Saccucci, M.; Di Carlo, G.; Lucchetti, R.; Piloni, A.; Pranno, N.; Luzzi, V.; Valesini, G.; Polimeni, A. Periodontitis and Rheumatoid Arthritis: The Same Inflammatory Mediators? *Mediat. Inflamm.* **2019**, *2019*, 6034546. [[CrossRef](#)]

79. Polak, D.; Shapira, L. An update on the evidence for pathogenic mechanisms that may link periodontitis and diabetes. *J. Clin. Periodontol.* **2018**, *45*, 150–166. [[CrossRef](#)]
80. Aghaloo, T.; Pi-Anfruns, J.; Moshaverinia, A.; Sim, D.; Grogan, T.; Hadaya, D. The Effects of Systemic Diseases and Medications on Implant Osseointegration: A Systematic Review. *Int. J. Oral Maxillofac. Implants* **2019**, *34*, s35–s49. [[CrossRef](#)] [[PubMed](#)]
81. Naujokat, H.; Kunzendorf, B.; Wiltfang, J. Dental implants and diabetes mellitus—a systematic review. *Int. J. Implant Dent.* **2016**, *2*, 5. [[CrossRef](#)] [[PubMed](#)]