

RESEARCH ARTICLE

Prognostic value of ^{11}C -methionine volume-based PET parameters in IDH wild type glioblastoma

Bart R. J. van Dijken^{1*}, Alfred O. Ankras², Gilles N. Stormezand², Rudi A. J. O. Dierckx^{1,2}, Peter Jan van Laar^{1,3}, Anouk van der Hoorn¹

1 Department of Radiology, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands, **2** Department of Nuclear Medicine and Molecular Imaging, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands, **3** Department of Radiology, Zorggroep Twente, Almelo and Hengelo, the Netherlands

* b.r.j.van.dijken@umcg.nl



Abstract

Purpose

^{11}C -Methionine (^{11}C -MET) PET prognostication of isocitrate dehydrogenase (IDH) wild type glioblastomas is inadequate as conventional parameters such as standardized uptake value (SUV) do not adequately reflect tumor heterogeneity. We retrospectively evaluated whether volume-based parameters such as metabolic tumor volume (MTV) and total lesion methionine metabolism (TLMM) outperformed SUV for survival correlation in patients with IDH wild type glioblastomas.

Methods

Thirteen IDH wild type glioblastoma patients underwent preoperative ^{11}C -MET PET. Both SUV-based parameters and volume-based parameters were calculated for each lesion. Kaplan-Meier curves with log-rank testing and Cox regression analysis were used for correlation between PET parameters and overall survival.

Results

Median overall survival for the entire cohort was 393 days. MTV (HR 1.136, $p = 0.007$) and TLMM (HR 1.022, $p = 0.030$) were inversely correlated with overall survival. SUV-based ^{11}C -MET PET parameters did not show a correlation with survival. In a paired analysis with other clinical parameters including age and radiotherapy dose, MTV and TLMM were found to be independent factors.

Conclusions

MTV and TLMM, and not SUV, significantly correlate with overall survival in patients with IDH wild type glioblastomas. The incorporation of volume-based ^{11}C -MET PET parameters may lead to a better outcome prediction for this heterogeneous patient population.

OPEN ACCESS

Citation: van Dijken BRJ, Ankras AO, Stormezand GN, Dierckx RAJO, Jan van Laar P, van der Hoorn A (2022) Prognostic value of ^{11}C -methionine volume-based PET parameters in IDH wild type glioblastoma. PLoS ONE 17(2): e0264387. <https://doi.org/10.1371/journal.pone.0264387>

Editor: Kevin Camphausen, National Institutes of Health, UNITED STATES

Received: September 7, 2021

Accepted: February 9, 2022

Published: February 25, 2022

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0264387>

Copyright: © 2022 van Dijken et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: We have uploaded the minimal (anonymized) data set as [supporting information](#).

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Introduction

The use of amino acid PET such as L-methyl- ^{11}C -methionine (^{11}C -MET) for prognostication of gliomas has recently been recommended [1]. ^{11}C -MET PET has an excellent tumor-to-background discrimination and provides supplementary metabolic information in addition to MRI [2].

Survival and treatment strategies differ significantly among glioma patients, necessitating reliable preoperative prognostication [3]. Gliomas without isocitrate dehydrogenase (IDH) mutation, including the WHO grade 4 glioblastoma, are now recognized as the most malignant subtype [4]. These IDH wild type gliomas are heterogeneous tumors characterized by a more aggressive and infiltrative nature than IDH mutant gliomas [5]. Glioblastomas IDH wild type WHO grade 4 are defined due to either aggressive histological features such as microvascular proliferation and necrosis, or by specific molecular markers such as TERT promoter mutation, EGFR amplification, and chromosome 7 gain/chromosome 10 loss [4]. Outcome in glioblastomas is generally poor.

The prognostic use of ^{11}C -MET PET in gliomas has previously been explored but showed inconclusive results [2]. Studies were usually limited to conventional parameters such as standardized uptake value (SUV), which do not reliably detect the heterogeneity of IDH wild type glioblastomas [6]. Recently, volume-based parameters have been introduced, possibly better reflecting the heterogeneity of IDH wild type glioblastomas [7, 8]. Volume-based parameters such as metabolic tumor volume (MTV) and total lesion methionine metabolism (TLMM), a combination of MTV and SUVmean, provide additional information about the tumor extent. The aim of this study was to investigate the value of conventional and volume-based ^{11}C -MET PET parameters for prognostication in IDH wild type glioblastomas.

Methods

We performed a retrospective search among all ^{11}C -MET PET examinations in our tertiary university hospital between 2011–2018. Inclusion criteria were glioblastoma, IDH wild type, WHO grade 4 according to the 2021 WHO guidelines, confirmed by biopsy or surgical resection ≤ 3 months after PET examination [4]. Exclusion criteria were previous cranial surgery or cerebral irradiation, known other primary tumor, and pediatric patients (< 18 years). The study was approved by the institutional review board and the need for written informed consent was waived.

The tracer was prepared as previously published [9] and imaging was performed in accordance with the 2006 procedure guidelines of the European Association of Nuclear Medicine for brain tumor imaging using labelled amino acid analogues [10]. Patients underwent static PET imaging 20 minutes after intravenous injection of 200 MBq (range: 180–220 MBq) ^{11}C -MET in one bed position of five minutes. Images were acquired on a Biograph mCT PET/CT system (Siemens/CTI, Knoxville, TN, USA) with 2 mm spatial resolution. The images were reconstructed using Truex + TOF with three iterations and 21 subsets in a 400 x 400 matrix size (zoom 1.0) and a 2 mm Gaussian filter.

Volumes of interest (VOI) were defined using Syngo.via (Siemens Medical Solutions Inc., Knoxville, TN, USA). A circle large enough to include all visual uptake was manually drawn by one researcher (BD) and checked by a nuclear physician with 3 years of experience (GS). The VOI were semi-automatically defined using a 40% threshold of SUVmax [6], which is standard value for delineation used in our research group and leads to good tumor definition when compared to visual inspection. Syngo.via allowed for extraction of the following PET parameters within the VOI: SUVmax, SUVpeak, SUVmean, MTV and TLMM (Fig 1). TLMM was




Fig 1. Schematic overview of metabolic and volume-based ^{11}C -MET PET parameters. (A) Snapshot of lesion with positive uptake of ^{11}C -MET. (B) SUV_{max} is defined as voxel with highest uptake within the lesion. (C) SUV_{mean} is an average of all SUV values from included voxels within the lesion. (D) SUV_{peak} is defined as the average value within a 1cm^3 sphere surrounding the SUV_{max} . Volume-based parameters are shown in E-F. (E) MTV is the volume of increased uptake within the lesion borders. (F) For TLMM the SUV_{mean} and MTV within lesion borders are combined and multiplied to provide both metabolic and volume-based information. TLMM is automatically calculated and extracted within the Syngo.via software. Abbreviations: MTV = metabolic tumor volume, SUV = standardized uptake value, TLMM = total lesion methionine metabolism.

<https://doi.org/10.1371/journal.pone.0264387.g001>

automatically calculated in Syngo.via by multiplying SUV_{mean} and MTV within the lesion border, in an equal manner as the Total Lesion Glycolysis (TLG) in FDG imaging.

Statistics were done in SPSS (IBM corp, version 25.0, Armonk, NY). Log-rank testing after dichotomization based on median values and Cox regression analysis were used for univariate survival analysis. The influence of various clinical variables (age, tumor grade, extent of resection and radiation dose) was also determined since the sample size was insufficient for multivariate analysis.

Results

A total of 13 patients with a median age of 57 years (range 28–70) and of whom 62% were male were included in the study. Adjuvant treatment consisted of radiotherapy plus or minus chemotherapy. One patient refused to undergo additional treatment after biopsy and one patient had to discontinue radiotherapy due to adverse events after a dose of 32.4 Gy. General characteristics are shown in Table 1.

Median survival was 393 days. The results of univariate analysis for the different PET parameters are shown in Table 2. MTV and TLMM were the only parameters that significantly correlated with overall survival (hazard ratio [HR] 1.136 (95%CI 1.035–1.246), $p = 0.007$ and HR 1.022 (95%CI 1.002–1.043), $p = 0.030$, respectively). Kaplan-Meier curves for MTV and TLMM are shown in Fig 2. Paired analysis with other clinical parameters demonstrated that MTV and TLMM were correlated with survival, independent of age, extent of resection or radiation dose. SUV based parameters were not correlated with survival (Fig 3).

Discussion

Volume-based ^{11}C -MET PET parameters were significantly correlated with survival in this small retrospective study among IDH wild type glioblastoma patients, while SUV based

Table 1. General characteristics of included patients.

Patient	Gender	Age	WHO Grade	EOR	Treatment	RT dose	Survival
1	F	70	GBM, gr 4*	GTR	RT	57.6	1325
2	F	63	GBM, gr 4	STR	C+RT	60	195 [‡]
3	M	28	GBM, gr 4*	B	C+RT	60	32 [‡]
4	F	64	GBM, gr 4*	B	C+RT	32.4 [†]	221
5	F	57	GBM, gr 4	B	C+RT	60	418
6	M	65	GBM, gr 4	B	None	-	266
7	M	59	GBM, gr 4	B	C+RT	60	200
8	M	44	GBM, gr 4	GTR	C+RT	60	277
10	M	47	GBM, gr 4	B	C+RT	60	171
11	M	61	GBM, gr 4	B	C+RT	60	618
12	M	52	GBM, gr 4	B	C+RT	60	469
13	M	67	GBM, gr 4*	B	RT	59.4	688
15	F	60	GBM, gr 4*	B	C+RT	59.4	237

* defined by EGFR amplification

[†]discontinued radiotherapy due to adverse events.

<https://doi.org/10.1371/journal.pone.0264387.t001>

parameters were not. TLMM, combining both volume-based and metabolic information, can thus be potentially employed for prognostication in IDH wild type glioblastomas.

Conventional SUV measures are the most commonly used quantitative PET parameters. Several studies have investigated the use of ¹¹C-MET SUV measures for glioma prognostication, with inconclusive results [2, 11–14]. SUV measures, however, do not provide information about tumor heterogeneity, nor do they visualize the extent of the lesion. In the current study we did not find a correlation between SUV measures and survival in IDH wild type glioblastomas, while volume-based parameters did demonstrate this correlation.

Our findings are in accordance with another ¹¹C-MET PET study which also demonstrated that a larger MTV and TLMM were correlated with lower survival [7]. However, important limitations of that study were the inclusion of oligodendrogliomas and lack of correcting for IDH mutation status, whilst it is known that both survival and ¹¹C-MET uptake values significantly differ between IDH wild type and IDH-1 gliomas [15, 16]. Furthermore, IDH wild type glioblastomas, are currently seen as separate entity and have a poorer prognosis than IDH-1 gliomas [4]. The prognostic role of volume-based parameters in gliomas have also been demonstrated for other amino acid PET tracers, such as O-(2-[¹⁸F]fluoroethyl)-l-tyrosine (18F-FET) [17–19]. for oncology prognostication has also been demonstrated in other populations [6, 20–23].

Table 2. Univariate analysis of ¹¹C-MET PET parameters and survival.

Parameter	HR (95%CI)	p value
SUVmax	0.818 (0.583–1.148)	0.246
SUVpeak	0.833 (0.549–1.263)	0.389
SUVmean	0.669 (0.366–1.221)	0.190
MTV	1.136 (1.035–1.246)	0.007
TLMM	1.022 (1.002–1.043)	0.030

Abbreviations: HR = hazard ratio, MTV = metabolic tumor volume, SUV = standard uptake value, TLMM = total lesion methionine metabolism.

<https://doi.org/10.1371/journal.pone.0264387.t002>




Fig 2. Kaplan-Meier curves for MTV and TLMM. Survival curves for MTV and TLMM. Dichotomisation of the data based on the median (≥ 14 for MTV and ≥ 51 for TLMM). The group with higher values is displayed with the dotted line while the group with lower values is displayed with the bold line. The survival difference between patients with higher and lower values was statistically significant for both MTV ($p = 0.014$) and TLMM ($p = 0.041$). Abbreviations: MTV = metabolic tumor volume, TLMM = total lesion methionine metabolism.

<https://doi.org/10.1371/journal.pone.0264387.g002>

Volume-based PET parameters are strongly dependent on the used delineation method [6]. This was recently demonstrated for ^{18}F -FDG PET in non-small cell lung cancer patients, where MTV significantly differed with chosen delineation method [6]. According to the authors of that study, the approach that reached best agreement was 40% of SUVmax, corresponding to the method used in the present study [6]. However, these results have not yet been validated for ^{11}C -MET PET. Moreover, it is not known if these results can be translated to gliomas. More research should therefore be aimed at establishing a robust delineation method for ^{11}C -MET PET in gliomas.




Fig 3. Differences in metabolic and volumetric parameters in two glioblastoma patients. Left image: 52-year-old male with right temporoparietal glioblastoma. Right image: 47-year-old male with left frontal glioblastoma. Despite relatively similar SUVmax values, there was a significant difference in survival between the two patients. The tumor of the patient on the right side demonstrated higher MTV and TLMM values than the tumor on the left side, corresponding to the lower survival of this patient.

<https://doi.org/10.1371/journal.pone.0264387.g003>

The most important limitation of this study is the small sample size. Due to this, a comprehensive multivariate analysis was not possible. Nevertheless, in paired analysis with other clinical parameters TLMM demonstrated to be independently correlated to survival. Another limitation is the retrospective nature of this study. Due to this, molecular markers, such as TERT promoter mutation, EGFR amplification, and gain of chromosome 7 with loss of chromosome 10 status was not always known, contributing to the smaller sample size. Our results should thus be seen as an encouragement for larger prospective studies focusing on the prognostic role of volume-based ^{11}C -MET PET parameters. At present, a prospective study on ^{11}C -MET PET for treatment evaluation in glioma patients is ongoing at our tertiary center (Netherlands Trial Register number NL6536) which will also address the utility of volume-based parameters for prognostication.

Conclusion

Volume-based ^{11}C -MET PET parameters MTV and TLMM were significantly correlated with overall survival in patients with IDH wild type glioblastomas. Volume-based parameters could play a future role in tailoring treatment decisions in patients with IDH wild type glioblastoma.

Supporting information

S1 File.
(PDF)

S2 File.
(PDF)

S1 Data.
(XLSX)

Author Contributions

Conceptualization: Bart R. J. van Dijken, Alfred O. Ankrah, Rudi A. J. O. Dierckx, Peter Jan van Laar, Anouk van der Hoorn.

Data curation: Bart R. J. van Dijken.

Formal analysis: Bart R. J. van Dijken, Alfred O. Ankrah, Gilles N. Stormezand, Anouk van der Hoorn.

Investigation: Bart R. J. van Dijken, Alfred O. Ankrah, Anouk van der Hoorn.

Methodology: Bart R. J. van Dijken, Alfred O. Ankrah, Gilles N. Stormezand, Rudi A. J. O. Dierckx, Peter Jan van Laar, Anouk van der Hoorn.

Project administration: Bart R. J. van Dijken, Anouk van der Hoorn.

Supervision: Gilles N. Stormezand, Rudi A. J. O. Dierckx, Peter Jan van Laar, Anouk van der Hoorn.

Writing – original draft: Bart R. J. van Dijken, Anouk van der Hoorn.

Writing – review & editing: Bart R. J. van Dijken, Alfred O. Ankrah, Gilles N. Stormezand, Rudi A. J. O. Dierckx, Peter Jan van Laar, Anouk van der Hoorn.

References

1. Albert NL, Weller M, Suchorska B, et al. Response Assessment in Neuro-Oncology working group and European Association for Neuro-Oncology recommendations for the clinical use of PET imaging in gliomas. *Neuro Oncol*. 2016; 18:1199–108. <https://doi.org/10.1093/neuonc/now058> PMID: 27106405
2. Glaudemans AWJM, Enting RH, Heesters MAAM, et al. Value of 11C-methionine PET in imaging brain tumours and metastases. *Eur J Nucl Med Mol Imaging*. 2013; 40:615–635. <https://doi.org/10.1007/s00259-012-2295-5> PMID: 23232505
3. van den Bent MJ, Baumert B, Erridge SC, et al. Interim results from the CATNON trial (EORTC study 26053–22054) of treatment with concurrent and adjuvant temozolomide for 1p/19q non-co-deleted anaplastic glioma: a phase 3, randomised, open-label intergroup study. *Lancet*. 2017; 390:1645–1653. [https://doi.org/10.1016/S0140-6736\(17\)31442-3](https://doi.org/10.1016/S0140-6736(17)31442-3) PMID: 28801186
4. Louis DN, Perry A, Reifenberger G, et al. The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol*. 2016; 131:803–820. <https://doi.org/10.1007/s00401-016-1545-1> PMID: 27157931
5. Eckel-Passow JE, Lachance DH, Molinaro AM, et al. Glioma Groups Based on 1p/19q, IDH, and TERT Promoter Mutations in Tumors. *N Engl J Med*. 2015; 372:2499–2508. <https://doi.org/10.1056/NEJMoa1407279> PMID: 26061753
6. Guilherme KD, Vázquez García D, Kramer GM, et al. Repeatability of [18 F]FDG PET/CT Total Metabolic Active Tumour Volume and Total Tumour Burden in NSCLC Patients. *EJNMMI Res*. 2019; 9:14. <https://doi.org/10.1186/s13550-019-0481-1> PMID: 30734113
7. Kobayashi K, Hirata K, Yamaguchi S, et al. Prognostic value of volume-based measurements on (11)C-methionine PET in glioma patients. *Eur J Nucl Med Mol Imaging*. 2015; 42:1071–1080. <https://doi.org/10.1007/s00259-015-3046-1> PMID: 25852010
8. Yoo MY, Paeng JC, Cheon GJ, et al. Prognostic Value of Metabolic Tumor Volume on (11)C-Methionine PET in Predicting Progression-Free Survival in High-Grade Glioma. 2015; 49:291–297.
9. Langstrom B, Antoni G, Gullberg P, et al. Synthesis of L- and D-[methyl-11C]methionine. *J Nucl Med*. 1987; 28:1037–1040. PMID: 3585494
10. Vander Borgh T, Asembaum S, Bartenstein P, et al. EANM procedure guidelines for brain tumour imaging using labelled amino acid analogues. *Eur J Nucl Med Mol Imaging*. 2006; 33:1374–1380. <https://doi.org/10.1007/s00259-006-0206-3> PMID: 16932934
11. Ceyskens S, van Laere K, de Groot T, et al. [11C]methionine PET, Histopathology, and Survival in Primary Brain Tumors and Recurrence. *AJNR Am J Neuroradiol*. 2006; 27:1432–1437. PMID: 16908552
12. Cicuendez M, Lorenzo-Bosquet C, Cuberas-Borrós G, et al. Role of [(11)C] Methionine Positron Emission Tomography in the Diagnosis and Prediction of Survival in Brain Tumours. *Clin Neurol Neurosurg*. 2015; 139:328–333. <https://doi.org/10.1016/j.clineuro.2015.10.035> PMID: 26588352
13. Lopci E, Riva M, Olivari L, et al. Prognostic Value of Molecular and Imaging Biomarkers in Patients With Supratentorial Glioma. *Eur J Nucl Med Mol Imaging*. 2017; 44:1155–1164. <https://doi.org/10.1007/s00259-017-3618-3> PMID: 28110346
14. Poetsch N, Woehrer A, Gesperger J, et al. Visual and Semiquantitative 11C-methionine PET: An Independent Prognostic Factor for Survival of Newly Diagnosed and Treatment-Naïve Gliomas. *Neuro Oncol*. 2018; 20:411–419. <https://doi.org/10.1093/neuonc/nox177> PMID: 29016947
15. Takei H, Shinoda J, Ikuta S, et al. Usefulness of Positron Emission Tomography for Differentiating Gliomas According to the 2016 World Health Organization Classification of Tumors of the Central Nervous System. *J Neurosurg*. 2019; 16:1–10.
16. Ogawa T, Kawai N, Miyake K, et al. Diagnostic Value of PET/CT With 11C-methionine (MET) and 18 F-fluorothymidine (FLT) in Newly Diagnosed Glioma Based on the 2016 WHO Classification. *EJNMMI Res*. 2020; 10:44. <https://doi.org/10.1186/s13550-020-00633-1> PMID: 32382870
17. Suchorska B, Jansen NL, Linn J, et al. Biological tumor volume in 18F-FET-PET before radiochemotherapy correlates with survival in GBM. *Neurology*. 2015; 84:710–719. <https://doi.org/10.1212/WNL.0000000000001262> PMID: 25609769
18. Bauer EK, Stoffels G, Blau T, et al. Prediction of survival in patients with IDH-wildtype astrocytic gliomas using dynamic O-(2- [18 F]-fluoroethyl)-L-tyrosine PET. *Eur J Nucl Med Mol Imaging*. 2020; 47:1486–1495. <https://doi.org/10.1007/s00259-020-04695-0> PMID: 32034446
19. Ceccon G, Lohman P, Werner JM, et al. Early Treatment Response Assessment Using 18 F-FET PET Compared with Contrast-Enhanced MRI in Glioma Patients After Adjuvant Temozolomide Chemotherapy. *J Nucl Med*. 2021; 62:918–925. <https://doi.org/10.2967/jnumed.120.254243> PMID: 33158907
20. Morales-Lozano MIM, Viering O, Samnick S, et al. 18F-FDG and 11C-Methionine PET/CT in Newly Diagnosed Multiple Myeloma Patients: Comparison of Volume-Based PET Biomarkers. *Cancers (Basel)*. 2020; 12:1042. <https://doi.org/10.3390/cancers12041042> PMID: 32340251

21. Cegla P, Burchardt E, Roszak A, et al. Influence of Biological Parameters Assessed in [18F]FDG PET/CT on Overall Survival in Cervical Cancer Patients. *Clin Nucl Med*. 2019; 44:860–863. <https://doi.org/10.1097/RLU.0000000000002733> PMID: 31348081
22. Carretta A, Bandiera A, Muriana P, et al. Prognostic Role of Positron Emission Tomography and Computed Tomography Parameters in Stage I Lung Adenocarcinoma. *Radiol Oncol*. 2020. Online ahead of print. <https://doi.org/10.2478/raon-2020-0034> PMID: 32463388
23. Creff G, Devillers A, Depeursinge A, et al. Evaluation of the Prognostic Value of FDG PET/CT Parameters for Patients With Surgically Treated Head and Neck Cancer: A Systematic Review. *JAMA Otolaryngol Head Neck Surg*. 2020; 146:471–479. <https://doi.org/10.1001/jamaoto.2020.0014> PMID: 32215611