

Article

Differential Uptake and Translocation of Cadmium and Lead by Quinoa: A Multivariate Comparison of Physiological and Oxidative Stress Responses

Atif A. Bamagoos¹, Hesham F. Alharby¹ and Ghulam Abbas^{2,*} 

¹ Department of Biological Sciences, Faculty of Science, King Abdulaziz University, Jeddah 21589, Saudi Arabia; abamagoos@kau.edu.sa (A.A.B.); halharby@kau.edu.sa (H.F.A.)

² Department of Environmental Sciences, COMSATS University Islamabad, Vehari Campus, Vehari 61100, Pakistan

* Correspondence: g.a92pk@gmail.com

Abstract: Contamination of soils with cadmium (Cd) and lead (Pb) has emerged as a serious environmental issue that reduces crop productivity. However, the metals tolerance and accumulation potential of quinoa (*Chenopodium Quinoa* Willd) under the combined stress of Cd and Pb has not yet been explored. In the present hydroponic study, the physiological and biochemical characteristics of quinoa exposed to Cd and Pb were explored. Four-week-old plants of quinoa genotype 'Puno' were grown under different concentrations of Cd (0, 50 and 100 μM), Pb (0, 250 and 500 μM) alone as well as in combinations. The results showed that with increasing Cd and Pb levels in the nutrient solution, the plant biomass, stomatal conductance and chlorophyll contents were decreased. However, the concurrent application of higher concentrations of Cd (100 μM) and Pb (500 μM) caused even more reduction in the plant biomass (more than 50% than the control) and physiological attributes. The combined application of Pb and Cd caused oxidative stress through an overproduction of H_2O_2 (10-fold) and TBARS (12.5-fold), leading to decrease in membrane stability (52%). The oxidative stress was alleviated by a 7-fold, 10-fold and 9-fold overactivation of superoxide dismutase (SOD), peroxidase (POD) and catalase (CAT), respectively. An excessive uptake of Cd resulted in a limited uptake of Pb and K in the roots and shoots of quinoa plants. The Cd and Pb tolerance and uptake potential of Puno showed its ability to stabilize Cd and Pb in co-contaminated soils.

Keywords: heavy metals; stomatal conductance; antioxidants; co-contamination; phytoremediation



Citation: Bamagoos, A.A.; Alharby, H.F.; Abbas, G. Differential Uptake and Translocation of Cadmium and Lead by Quinoa: A Multivariate Comparison of Physiological and Oxidative Stress Responses. *Toxics* **2022**, *10*, 68. <https://doi.org/10.3390/toxics10020068>

Academic Editors: Alexander S. Lukatkin, Naser A. Anjum and Przemyslaw Malec

Received: 22 December 2021

Accepted: 2 February 2022

Published: 4 February 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

In recent times, various human activities, such as mining, the release of effluents from industries, rapid urbanization and agricultural practices, have increased cadmium (Cd) and lead (Pb) contamination in soils [1,2]. Cd and Pb are not necessary for the normal metabolic functions of plants, and thus both are considered non-essential elements [2,3]. The average concentration of Cd in non-contaminated soils is 0.1–0.2 mg kg^{-1} [4]. Cadmium is toxic to plants' growth and development, and has been ranked fourth among the highly poisonous elements due to its high mobility in the environment [1,5].

Cadmium is very toxic even at very low concentrations, so, the uptake of Cd by plants affects their growth and normal metabolic activities [6]. Various abnormalities, such as stunted plant growth, damage to pigments and the obliteration of cell membranes and organelles, have been reported under Cd stress [3,7]. An indirect effect of Cd stress in plants is the generation of reactive oxygen species (ROS) [8]. The main ROS which are deleterious to plants include singlet oxygen ($\frac{1}{2}\text{O}_2$), superoxide ($\text{O}_2^{\bullet-}$), hydrogen peroxide (H_2O_2) and hydroxyl (HO^{\bullet}) radicals [9,10]. These ROS have ill effects on macromolecules such as lipids, proteins, carbohydrates and nucleic acids [8,11]. These ROS are detoxified in

plants by antioxidant enzymes, namely, catalase (CAT), peroxidase (POD) and superoxide dismutase (SOD) [3,8].

Lead is another hazardous environmental pollutant having deleterious effects on both humans and plants [12,13]. The natural background concentration of Pb in soils is approximately 50 mg kg^{-1} soil, which has been increased many times due to various anthropogenic activities [8,14]. An excessive uptake of Pb causes many anomalies in plants, such as a restricted uptake of nutrients, limitations to photosynthesis and stomatal conductance, disturbance of the water balance and an increased production of ROS, leading to membrane damage [13,15,16]. Antioxidant enzymes and non-enzymes are both used to mitigate the ROS toxicity caused by Pb [8,17,18].

In the present climate change scenario, crop production and the rehabilitation of contaminated soils should be done by targeting such plants species which have high tolerances against multiple environmental stresses. In recent times, the most promising example of such crops is quinoa (*Chenopodium quinoa* Willd). Quinoa is an exemplary crop with respect to food security, its multiple uses and its supreme nutritional quality [19,20]. Quinoa has a high tolerance to salinity and drought [20,21]. It has also shown good tolerance and metal stabilization potential to single-metal-contaminated soils [3,13]. Nevertheless, the combined effects Cd and Pb on the morpho-physiological and metal uptake responses of quinoa have not been investigated until now. Thus, the current experiment was conducted to (a) unravel the combined effects of Cd and Pb on the growth and physiological characteristics of quinoa, (b) determine the tolerance mechanisms of quinoa to the combined stress of Cd and Pb and (c) evaluate the Cd and Pb accumulation and translocation potential of quinoa.

2. Materials and Methods

2.1. Plant Cultivation and Treatment Application

A hydroponic experiment was conducted using healthy plants of the quinoa genotype 'Puno' during 2020–2021. The mean lowest and highest temperatures in the glasshouse varied from 10 to 26 °C and the relative humidity varied from 55 to 72% during the study period [3]. Seedlings were raised in a sand culture under the mentioned conditions. Uniform seedlings at the age of thirty days were transplanted into Hoagland's nutrient solution (half strength). The plants were acclimatized for one week, and afterwards they were exposed to various treatments of Cd and Pb. Calculated amounts of cadmium chloride and lead chloride salts were applied to develop various concentrations of Cd (0, 50, 100 μM Cd) and Pb (0, 250, 500 μM Pb) either alone or combined. These levels were selected on the basis of previous studies [3,13]. After each seven days, the nutrient solution was refreshed and its pH was maintained at 6.5 ± 0.2 with diluted hydrochloric acid (HCl) or sodium hydroxide (NaOH). A completely randomized design was used in this experiment with 9 treatments and 4 replications. There were two plants in each replication of each treatment.

2.2. Harvesting of Plants and Growth Measurement

After 28 days of Cd and/or Pb treatment exposure, plants were harvested, and roots were separated from the shoots. Although all the plants survived, there was a severe decline in the plants' growth under the combined highest levels of both metals. For the removal of adsorbed Cd and Pb from the root surface, roots were cautiously washed with dilute acid (0.01 M HCl) followed by thorough washing in distilled water [13]. Plant length (shoot/root) was measured by meter rod. The samples were then air dried and oven dried for 48 h at 70 °C to measure the dry weight. Roots' and shoots' dry weights were measured separately using a digital weighing balance.

2.3. Ionic Analysis of Plant Samples

Shoot and root samples were crushed into fine powder and digested in HNO_3 and HClO_4 taken in a 2:1 ratio. Briefly, 5 mL each of concentrated HClO_4 and HNO_3 were added in the digestion flasks containing plant samples and kept overnight. The next day, 5 mL additional HNO_3 was added and samples were digested on a hot plate until 1–2 mL clear

digestate was left. After digestion, cooling of samples was done by adding distilled water. Whatman filter paper No. 42 was used for filtration of digested samples, then samples were stored in air-tight bottles. The K concentrations in plant samples were determined by a flame photometer (BWB-XP5), whereas an atomic absorption spectrometer (AAS; PinAAcle 900F, PerkinElmer, Inc. Los Angeles, CA, USA) was used to measure the concentrations of Cd and Pb.

2.4. Metal Tolerance and Translocation

Metal tolerance and translocation potential of plants was calculated through BCF (bioconcentration factor), TF (transfer factor) and TI (tolerance index) for both Cd and Pb following Iftikhar et al. [13]. The concentrations of Cd and Pb in roots and shoots were divided by the respective metal concentration in the growth medium for the calculation of the shoot BCF and root BCF. The TF was indicated as the ratio of the metal concentration in shoot to root. The ratio of the dry weight of the plants under stress to the dry weight of the control plants was expressed as TI [3].

2.5. Pigment Contents and Stomatal Conductance

Fresh leaves were frozen in liquid nitrogen (LN_2) to conserve the metabolic status of leaves for pigment content determination (Chl-a, Chl-b and total Chl (a+b)). The samples (1.0 g) were then crushed in acetone (80%) extraction buffer in darkness. The samples were centrifuged for 10 min (3000 rpm), and the absorbance was noted at wavelengths 663.2, 646.8 and 470 nm with the help of a UV/vis spectrophotometer (Lambda 25, PerkinElmer Inc., Los Angeles, CA, USA). The equations given by Lichtenthaler [22] were used for pigment contents calculations. Stomatal conductance was determined using a leaf porometer (Decagon Devices, Pullman, WA, USA) on a full sunny day from the fully grown second leaf from the top.

2.6. Oxidative Stress Attributes

For the determination of ROS (H_2O_2), trichloroacetic acid (0.1%) was used for the homogenization of 0.5 g leaf sample (frozen in LN_2) according to Islam et al. [23] and centrifuged for 20 min ($3000 \times g$). The reaction mixture contained plant extract (1 mL), potassium phosphate buffer (10 mM, pH 7.0, 1 mL) and potassium iodide (2 M, 1 mL). The absorbance was recorded at 390 nm by UV/vis spectrophotometer and the values of H_2O_2 were expressed in nmol g^{-1} fresh weight. The TBARS content (thiobarbituric acid reactive substances) were determined for the estimation of lipid peroxidation. Grinding of leaf samples (0.5 g frozen in LN_2) was done in acetone buffer (80%) at 4 °C. The obtained material was mixed in butyl hydroxytoluene and thiobarbituric acid, heated in a water bath at 95 °C and centrifuged at $3000 \times g$ for 10 min. The absorbance of samples was noted at 532 nm wavelength, and the TBARS concentration was expressed in nmol g^{-1} . The methodology of Sairam [24] was adopted for the estimation of the membrane stability index (MSI).

2.7. Enzymatic Activities

Antioxidant enzymes were measured from the youngest fully expanded leaves (0.5 g) frozen in LN_2 . The leaves were homogenized in phosphate buffer (0.1 M, pH 7.0) and centrifuged at $10,000 \times g$ at 4 °C for 25–30 min. Superoxide dismutase (SOD) was assayed following Dhindsa et al. [25]. For the determination of peroxidase (POD), the method of Hemeda and Klein [26] was adopted and POD was represented in μM guaiacol oxidized $\text{min}^{-1} \text{mg}^{-1}$ protein. Catalase (CAT) was estimated as detailed by Aebi [27], and was described as μM of H_2O_2 degraded $\text{min}^{-1} \text{mg}^{-1}$ protein.

2.8. Statistical Analysis

Analysis of variance (ANOVA) was used to analyze the data at 5% significance level using statistical software Statistix 8.1. The pairwise treatment comparison was done by

least significant difference test [28]. Data were also analyzed through XLSTAT 2014 for PCA (principal component analysis) and Pearson correlations.

3. Results

3.1. Plant Growth

The results revealed that with increasing the concentrations of Cd and Pb in the nutrient solution, the growth of the shoots and roots of quinoa was reduced (Table 1). Nevertheless, the elevated levels of Cd (100 μM) and Pb (500 μM) jointly caused the highest alleviation in the plant biomass with respect to the control. The reductions in the shoot length were 33% and 28% under the higher concentrations of Cd (100 μM) and Pb (500 μM) alone treatments. There were 35% and 26% reductions in the root length of quinoa under the elevated levels of Cd and Pb alone treatments as compared to the control treatment. Under the combined treatment of Cd (100 μM) + Pb (500 μM), the respective reductions in shoot and root lengths were 60% and 61% as compared to the control. Root and shoot dry weights were reduced in the same way. The shoot dry weight decreased by 38% and 24% at 100 μM of Cd and 500 μM of Pb, respectively. The dry weight of the root decreased by 31% and 22% under the exposures of 100 μM of Cd alone and 500 μM of Pb alone, respectively, in comparison to the control. The dry weights of the root and shoot were decreased respectively by 62% and 65% in the combined treatment of 100 μM of Cd + 500 μM of Pb.

Table 1. Shoot and root growth of quinoa plants exposed to various concentrations of cadmium and lead under hydroponic conditions.

| Treatments (μM) | Shoot Length (cm) | Root Length (cm) | Shoot Dry Weight (g plant^{-1}) | Root Dry Weight (g plant^{-1}) |
|------------------------------|--------------------|------------------|--|---|
| Control | 18.0 \pm 0.60 a | 23 \pm 0.5 a | 0.34 \pm 0.01 a | 0.32 \pm 0.02 a |
| Cd 50 | 14.0 \pm 0.23 c | 18 \pm 0.8 c | 0.26 \pm 0.008 b | 0.26 \pm 0.008 b |
| Cd 100 | 12.0 \pm 0.50 d | 15 \pm 0.7 d | 0.21 \pm 0.01 c | 0.22 \pm 0.01 c |
| Pb 250 | 16.0 \pm 0.40 b | 20 \pm 0.5 b | 0.30 \pm 0.013 ab | 0.30 \pm 0.012 ab |
| Pb 500 | 13.0 \pm 0.30 cd | 17 \pm 0.4 c | 0.26 \pm 0.005 b | 0.25 \pm 0.008 b |
| Cd 50 + Pb 250 | 12.0 \pm 0.50 d | 17 \pm 0.8 c | 0.24 \pm 0.01 bc | 0.24 \pm 0.015 bc |
| Cd 100 + Pb 250 | 10.5 \pm 0.20 e | 13 \pm 0.5 de | 0.18 \pm 0.008 cd | 0.18 \pm 0.012 d |
| Cd 50 + Pb 500 | 9.0 \pm 0.30 f | 12 \pm 0.4 e | 0.16 \pm 0.01 d | 0.19 \pm 0.008 d |
| Cd 100 + Pb 500 | 7.0 \pm 0.40 g | 9 \pm 0.5 f | 0.13 \pm 0.009 e | 0.14 \pm 0.01 e |

Data presented are the mean values ($n = 4$) followed by the standard error. Different letters in each column represent the significant difference at a 5% probability level.

3.2. Pigment Contents and Stomatal Conductance

By increasing the Cd and Pb contamination of the nutrient solution, the leaf pigment contents (Chl a, Chl b and total Chl) and stomatal conductance of the quinoa plants declined (Table 2). Chl a, Chl b and total Chl contents declined by 31%, 32% and 31%, respectively, at 100 μM Cd and by 14%, 18% and 16%, respectively, at 500 μM of Pb with respect to the control plants. The combination of 100 μM of Cd and 500 μM of Pb resulted in 55%, 60% and 57% declines in Chl a, Chl b and total Chl, respectively, in comparison to the control treatment. The stomatal conductance was decreased by 20%, 14% and 47% under 100 μM of Cd, 500 μM of Pb and their combination, respectively.

3.3. Oxidative Stress Attributes

Cadmium and Pb stress increased the TBARS and H_2O_2 contents (Figure 1A,B). Cadmium and Pb in combination resulted in more oxidative stress as compared to the individual stresses. The higher level of Cd caused a 4-fold increase in H_2O_2 contents and a 5-fold increase in TBARS contents. Lead contamination at 500 μM caused a 3-fold increase in TBARS contents and 3.3-fold increase in H_2O_2 with respect to the control. The combination of 100 μM of Cd with 500 μM of Pb caused a 10-fold enhancement in H_2O_2 and 12.5-fold enhancement in TBARS contents in comparison to the control treatment. The MSI was decreased, with an increase in the oxidative stress under the Pb and Cd contamination

(Figure 1C). The higher levels of Cd, Pb and their combination resulted in 24%, 15% and 55% declines, respectively, in the MSI with respect to the control treatment.

Table 2. Pigment contents and stomatal conductance of quinoa plants exposed to various concentrations of cadmium and lead under hydroponic conditions.

| Treatments (μM) | Chl-a ($\mu\text{g g}^{-1}$ FW) | Chl-b ($\mu\text{g g}^{-1}$ FW) | Total Chl ($\mu\text{g g}^{-1}$ FW) | Stomatal Conductance ($\text{mmol m}^{-2}\text{s}^{-1}$) |
|------------------------------|----------------------------------|----------------------------------|--------------------------------------|--|
| Control | 245 \pm 8.0 a | 140 \pm 5.0 a | 385 \pm 10.0 a | 500 \pm 10.0 a |
| Cd 50 | 205 \pm 7.0 bc | 110 \pm 3.0 bc | 315 \pm 13.0 bc | 450 \pm 8.0 b |
| Cd 100 | 170 \pm 3.0 cd | 95 \pm 4.0 cd | 265 \pm 4.0 d | 400 \pm 15.0 c |
| Pb 250 | 210 \pm 8.0 b | 115 \pm 2.0 bc | 325 \pm 10.0 cd | 470 \pm 8.0 ab |
| Pb 500 | 190 \pm 5.0 bc | 100 \pm 2.0 cd | 290 \pm 5.0 c | 430 \pm 16.0 b |
| Cd 50 + Pb 250 | 180 \pm 4.0 d | 90 \pm 3.0 d | 270 \pm 10.0 cd | 410 \pm 18.0 bc |
| Cd 100 + Pb 250 | 160 \pm 8.0 e | 80 \pm 4.0 e | 240 \pm 5.0 e | 350 \pm 10.0 d |
| Cd 50 + Pb 500 | 140 \pm 4.0 f | 65 \pm 3.0 f | 205 \pm 10.0 f | 310 \pm 8.0 e |
| Cd 100 + Pb 500 | 110 \pm 5.0 g | 55 \pm 2.0 g | 165 \pm 5.0 g | 265 \pm 12.0 f |

Data presented are the mean values ($n = 4$) followed by the standard error. Different letters in each column represent the significant difference at a 5% probability level.

3.4. Antioxidant Enzymes

At increasing concentrations of Cd and Pb, the activities of antioxidant enzymes including SOD, CAT and POD showed differential responses (Figure 2A–C). The antioxidant enzyme activities were the highest under the combination of 100 μM of Cd + 500 μM of Pb with respect to control. The activities of SOD, CAT and POD were respectively 3-fold, 4-fold and 3-fold higher at 100 μM of Cd as compared to the control. At a Pb level of 500 μM , there were 2.6-fold, 3-fold and 2.6-fold enhancements in SOD, CAT and POD activities, respectively, than the control treatment. Under the combination of Cd (100 μM) with Pb (500 μM), there were 7-fold, 9-fold and 10-fold increases in SOD, CAT and POD activities with respect to the control.




Figure 1. Cont.




Figure 1. Effects of different levels of Cd, Pb and their combinations on H₂O₂ contents (A), TBARS contents (B) and MSI (C) of quinoa. Different letters represent the significant difference among the treatments at a 5% significance level.

3.5. Potassium and Heavy Metals Accumulation

The accumulation of K was decreased in quinoa tissues in the presence of Cd and Pb (Figure 3A,B). The combined treatments of both metals resulted in the lowest accumulation of K in plant tissues. With increasing levels of Cd in the growth medium, the Cd concentration was also increased in the roots and shoots of quinoa plants (Figure 4A,B). When Cd was applied in combination with Pb, root and shoot Cd contents were even more enhanced. Root and shoot Pb accumulation was increased as Pb contamination was increased in the growth medium (Figure 4C,D). However, when Pb was applied in combination with various Cd levels, root and shoot Pb contents declined. Root Cd and Pb contents were more than the shoot Cd and Pb contents for all the Cd and Pb treatments. The BCFs of the shoots and roots with both Cd and Pb were higher than one for all the treatments (Table 3). The translocation of both metals from the root to the shoot was expressed as the TF. The values of the TF were less than one for both Cd and Pb under all the treatments. The TI under Pb treatments was greater than the TI under Cd treatments. However, with increases in metal contamination the TI was decreased in the case of both metals (Table 3). Under

100 μM of Cd, 500 of μM Pb, and their combinations, the values of the TI were 62%, 76% and 38%, respectively.




Figure 2. Effect of different levels of Cd, Pb and their combinations on the activities of SOD (A), CAT (B) and POD (C) of quinoa. Different letters represent the significant difference among the treatments at a 5% significance level.




Figure 3. Effect of different levels of Cd, Pb and their combinations on shoot (A) and root (B) K concentrations. Different letters represent the significant difference among the treatments at a 5% significance level.




Figure 4. Cont.




Figure 4. Effect of different levels of Cd, Pb and their combinations on Cd concentration in shoot (A), root (B) and Pb concentration in shoot (C) and root (D) of quinoa. Different letters represent the significant difference among the treatments at a 5% significance level.

Table 3. Bioconcentration factor (BCF), translocation factor (TF) and tolerance index (TI) of quinoa plants exposed to various concentrations of cadmium and lead under hydroponic conditions.

| Treatments (μM) | Cd | | | Pb | | | TI |
|-----------------|--------------|---------------|---------------|---------------|---------------|----------------|--------------|
| | BCF Shoot | BCF Root | TF | BCF Shoot | BCF Root | TF | |
| Cd 50 | 8.83 ± 0.5 b | 9.8 ± 0.4 c | 0.9 ± 0.05 a | – | – | 0.18 ± 0.005 b | 76.5 ± 3 b |
| Cd 100 | 5.78 ± 0.2 c | 6.4 ± 0.8 d | 0.90 ± 0.04 a | – | – | 0.21 ± 0.008 a | 61.8 ± 2 c |
| Pb 250 | – | – | 0.5 ± 0.03 c | 4.30 ± 0.05 a | 24.4 ± 1.8 ab | 0.16 ± 0.01 b | 88.2 ± 3 a |
| Pb 500 | – | – | 0.8 ± 0.02 b | 2.90 ± 0.04 c | 17.6 ± 1.5 c | 0.16 ± 0.005 c | 76.5 ± 3.1 b |
| Cd 50 + Pb 250 | 9.79 ± 0.5 b | 12.5 ± 0.7 b | 0.79 ± 0.05 b | 3.67 ± 0.2 b | 22.8 ± 2.4 b | 0.16 ± 0.003 c | 70.6 ± 3 bc |
| Cd 100 + Pb 250 | 6.67 ± 0.4 c | 7.6 ± 0.5 d | 0.88 ± 0.02 a | 1.59 ± 0.1 d | 10.2 ± 1.2 d | 0.16 ± 0.005 c | 52.9 ± 1.5 d |
| Cd 50 + Pb 500 | 13.3 ± 0.3 a | 16.9 ± 0.6 a | 0.79 ± 0.03 b | 4.83 ± 0.2 a | 27.0 ± 2.5 a | 0.18 ± 0.006 b | 47.1 ± 2.5 e |
| Cd 100 + Pb 500 | 8.77 ± 0.2 b | 10.6 ± 1.2 bc | 0.82 ± 0.04 b | 1.93 ± 0.2 d | 11.6 ± 1.3 d | 0.17 ± 0.003 c | 38.2 ± 2 f |

Data presented are the mean values ($n = 4$) followed by the standard error. Different letters in each column represent the significant difference at a 5% probability level.

3.6. Multivariate Analyses

A principal component analysis (PCA) and a Pearson correlation matrix (Table 4; Figure 5A,B) were used to depict the relations among different observations and response variables. The PCA divided all the response variables into eight factors from F1 to F8. Nevertheless, only three factors had the main contribution. The contributions of these three factors were 88%, 10%, and 1%, respectively. All the variables were divided into four main groups: (a) growth and physiological attributes, (b) oxidative stress attributes, (c) Cd contents and (d) Pb contents. The oxidative stress indicators (H_2O_2 , MDA and MSI) and antioxidant enzymes (SOD, CAT and POD) were clustered closely. Likewise, plant growth and physiological attributes had a close association (Figure 5A). A strong inverse relation was noticed between, on the one hand, ionic and oxidative attributes, and, on the other, the rest of the parameters through Pearson correlation matrix (Table 4). The PCA also divided treatments across various axes (Figure 5B). Separate treatments of both Cd and Pb were grouped along the positive x -axis. On the other hand, the combined treatments of Cd and Pb were grouped along the negative x -axis.




Figure 5. Comparison of different response variables (A) and different treatments (B) of quinoa exposed to Cd and Pb treatments using principal component analysis.

Table 4. Correlation matrix of different variables of quinoa under cadmium and lead stress. Values in bold represent the significant correlations among the variables.

| Variables | RDW | SDW | RL | SL | Chla | Chlb | TChl | H ₂ O ₂ | MDA | MSI | SOD | CAT | POD | Cond | SK | RK | SCd | RCd | SPb | RPb |
|-------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|-------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|---------------|----------|-----|
| SDW | 0.9879 | 1 | | | | | | | | | | | | | | | | | | |
| RL | 0.9895 | 0.9938 | 1 | | | | | | | | | | | | | | | | | |
| SL | 0.9808 | 0.9865 | 0.9885 | 1 | | | | | | | | | | | | | | | | |
| Chla | 0.9704 | 0.9820 | 0.9897 | 0.9887 | 1 | | | | | | | | | | | | | | | |
| Chlb | 0.9590 | 0.9755 | 0.9778 | 0.9909 | 0.9881 | 1 | | | | | | | | | | | | | | |
| TChl | 0.9687 | 0.9822 | 0.9878 | 0.9924 | 0.9982 | 0.9956 | 1 | | | | | | | | | | | | | |
| H ₂ O ₂ | −0.9717 | −0.9783 | −0.9728 | −0.9713 | −0.9704 | −0.9665 | −0.9716 | 1 | | | | | | | | | | | | |
| MDA | −0.9382 | −0.9381 | −0.9353 | −0.9450 | −0.9434 | −0.9489 | −0.9483 | 0.9837 | 1 | | | | | | | | | | | |
| MSI | 0.9607 | 0.9684 | 0.9618 | 0.9809 | 0.9747 | 0.9793 | 0.9793 | −0.9868 | −0.9826 | 1 | | | | | | | | | | |
| SOD | −0.9311 | −0.9483 | −0.9509 | −0.9508 | −0.9558 | −0.9536 | −0.9576 | 0.9833 | 0.9753 | −0.9676 | 1 | | | | | | | | | |
| CAT | −0.9651 | −0.9659 | −0.9626 | −0.9682 | −0.9649 | −0.9657 | −0.9680 | 0.9968 | 0.9935 | −0.9910 | 0.9810 | 1 | | | | | | | | |
| POD | −0.9168 | −0.9249 | −0.9274 | −0.9363 | −0.9410 | −0.9441 | −0.9449 | 0.9765 | 0.9931 | −0.9729 | 0.9886 | 0.9849 | 1 | | | | | | | |
| Cond | 0.9708 | 0.9781 | 0.9809 | 0.9776 | 0.9824 | 0.9746 | 0.9821 | −0.9957 | −0.9810 | 0.9849 | −0.9893 | −0.9929 | −0.9807 | 1 | | | | | | |
| SK | 0.9655 | 0.9718 | 0.9742 | 0.9839 | 0.9820 | 0.9792 | 0.9837 | −0.9902 | −0.9727 | 0.9899 | −0.9824 | −0.9895 | −0.9725 | 0.9930 | 1 | | | | | |
| RK | 0.9656 | 0.9723 | 0.9724 | 0.9812 | 0.9830 | 0.9716 | 0.9813 | −0.9863 | −0.9749 | 0.9920 | −0.9760 | −0.9862 | −0.9728 | 0.9914 | 0.9914 | 1 | | | | |
| SCd | −0.9070 | −0.9170 | −0.8818 | −0.8823 | −0.8640 | −0.8499 | −0.8609 | 0.8924 | 0.8665 | −0.8953 | 0.8324 | 0.8813 | 0.8314 | −0.8749 | −0.8567 | −0.8974 | 1 | | | |
| RCd | −0.9099 | −0.9245 | −0.8901 | −0.8990 | −0.8817 | −0.8735 | −0.8810 | 0.9113 | 0.8946 | −0.9209 | 0.8612 | 0.9041 | 0.8651 | −0.8955 | −0.8812 | −0.9197 | 0.9959 | 1 | | |
| SPb | −0.3167 | −0.3263 | −0.3771 | −0.4155 | −0.4289 | −0.5004 | −0.4584 | 0.3698 | 0.4131 | −0.4019 | 0.4296 | 0.4003 | 0.4452 | −0.4025 | −0.4364 | −0.3578 | −0.0098 | 0.0476 | 1 | |

4. Discussion

In the present investigation, the tolerance potential of quinoa genotype 'Puno' to the co-contamination of its growth medium with Cd and Pb was explored. The growth of quinoa significantly declined as the contamination levels of Cd and Pb were augmented in the growth medium. This was particularly true with the combined application of both metals in the growth medium. Many researchers have reported that under Cd stress there is a considerable reduction in plants' growth [9,29]. Research has shown that Cd does not have any important role in the growth and development of plants [30]. In line with the present study, Abdal et al. [3] found that the plant growth of quinoa was decreased under Cd stress, but that the addition of NaCl to the nutrient solution mitigated the negative effects of Cd. Similarly, Amjad et al. [8] found that the plant growth of quinoa was correspondingly decreased when soil Cd levels were increased from 30 to 90 mg kg⁻¹. This cadmium-induced decline in the growth of quinoa is attributed to numerous anomalies such as a restricted uptake of nutrients; the toxicity of the Cd ion; and a limitation of plant–water relations, the photosynthetic process and enzymatic activities [3,9]. Similar to Cd, the presence of Pb in the growth medium also reduced the plant growth of quinoa. These findings are in accordance with our previous studies [8,13], in which the growth of quinoa was decreased under Pb stress in both soil and solution culture conditions. Moreover, it was observed that mild salt stress (150 mM NaCl) had positive effects on quinoa exposed to Pb stress [13]. The reduction in plants' growth under Pb stress is mainly due to the interference of Pb with crucial physiological processes such as the chlorophyll biosynthesis, uptake of essential nutrients, stomatal conductance of leaves, and overproduction of ROS [12]. The interaction of Pb and Cd was particularly harmful due to the additive effects of both heavy metals on the mentioned physiological processes of quinoa. The results of the current investigation also showed that the higher level of Pb (500 µM) is less toxic than the individual higher level of Cd (100 µM). The higher level of Cd (100 µM) caused more decline in pigments, stomatal conductance and K accumulation than the higher level of Pb (500 µM), resulting in more decline in the plant biomass. These observations reflect that even a very low concentration of Cd is toxic to plants [6]. This is the very first study regarding the combined effects of Cd and Pb on quinoa. Previously, Amjad et al. [8] explored the separate effects of Cd and Pb on quinoa in a pot experiment. Hence, these findings are more realistic, considering the co-occurrence of both Cd and Pb in contaminated soils.

Stomatal conductance and pigment contents were also decreased by Cd and Pb and more so under their joint application. Reduction in these attributes of quinoa under Cd and Pb stress is well documented [3,8,13]. Moreover, it was noticed that these attributes were improved when 150 mM NaCl was supplied to quinoa growing under various levels of Cd [3] and Pb [13] stress. The possible reasons for the decline in photosynthetic pigments under Cd and Pb stress include (a) oxidative stress, (b) the direct influence of metal ions on the pigment biosynthesis pathways and (c) the replacement of magnesium with metal ions within the chlorophyll molecule [6,8,9]. Non-stomatal limitations of photosynthesis are the main reasons for the limited stomatal conductance of quinoa leaves under Pb and Cd stress [13,31]. As for pigments, the stomatal conductance and growth was decreased more at the 100 µM of Cd than 500 µM of Pb level, indicating the greater toxicity of Cd than Pb under the same growth conditions.

In the present study, the contamination of the nutrient solution with Cd and Pb caused an over-generation of ROS such as H₂O₂ and resulted in oxidative stress in quinoa. The higher accumulation of H₂O₂ caused the oxidative stress. The overproduction of ROS, for example, H₂O₂, caused lipid peroxidation (higher TBARS contents) and decreased the stability of the cell membranes. The overproduction of H₂O₂ and resultant damage to cell membranes is a well observed phenomenon under Cd and Pb stress. For example, Amjad et al. [8] found that soil Cd levels higher than 30 mg kg⁻¹ caused severe oxidative stress in quinoa. In the case of hydroponic experiments, it was observed that increasing the Cd levels from 50 to 200 µM [3] and Pb levels from 250 to 500 µM [13] caused severe oxidative stress in quinoa plants. Many researchers have observed that Cd and Pb stress caused oxidative

stress in plants other than quinoa as well [9,15]. Metal-induced overproduction of ROS and resultant damage to cellular membranes was more detrimental under the joint application of higher levels of both Cd (100 μ M) and Pb (500 μ M). A higher accumulation and toxicity of both Cd and Pb ions in the joint treatments may be the reason for the greater oxidative stress and cellular damage. Among various ROS, H_2O_2 is regarded as the most significant, because it has the tendency to be converted into even more poisonous ROS, i.e., hydroxyl anions [3,15]. Thus, in order to safeguard the cell from the damaging effects of these ROS, the detoxification of H_2O_2 is inevitable. Plants have been naturally gifted with antioxidant enzyme systems for the mitigation of ROS within the cellular organelles [8,32]. Superoxide dismutase is one of the most important enzymes, because it has the ability to convert the superoxide radicals into H_2O_2 and oxygen [9,13]. Both Cd and Pb caused considerable increases in the activity of SOD. This overactivation of SOD may be due to its ability to convert the superoxide radicals into H_2O_2 or the direct interaction of Cd and Pb with SOD [9,15]. Similar to these findings, increased activation of SOD in quinoa plants under metals stress is well documented [3,13]. The toxicity of H_2O_2 is further decreased when it is converted to molecular oxygen and water. This important step in the detoxification of H_2O_2 is achieved through the overexpression of CAT and POD [8,11]. The present study showed that under the combination of Cd and Pb, the activities of both these enzymes were enhanced. Similar increases in the activities of these enzymes were also observed under Cd and Pb stress both in soil and solution culture experiments in quinoa [3,8,13] as well as other plants [9,11]. Thus, it can be inferred that both CAT and POD play crucial roles in the detoxification of H_2O_2 and resultant cellular damage in plants under Cd and Pb stress [3,13].

The contents of K were decreased in plant tissues in the presence of Cd and Pb. Potassium is a macro nutrient and has many indispensable roles in the life cycles of plants. It plays crucial roles in osmotic adjustment, enzyme activation, cell enlargement, chlorophyll synthesis, the maintenance of cytoplasmic pH and membrane potential [33,34]. Consequently, the plants which can maintain higher cellular K levels are usually more tolerant of various environmental stresses [34]. We noticed that the excessive Cd and Pb uptake caused a corresponding decrease in K accumulation in quinoa [8,9]. Elevated levels of Cd and Pb in a nutrient solution result in a reduced uptake and translocation of mineral nutrients such as zinc (Zn), iron (Fe), calcium (Ca), manganese (Mn), magnesium (Mg) and potassium (K) due to cation competition at root uptake sites. A number of transporters and channels have been identified that may be involved in the process of metal uptake by plants [31,35–37]. For example, Cd can enter root cells as a divalent cation through (a) ZIP transporters such as orthologues of AtIRT1 and TcZNT1/TcZIP4, (b) orthologues of the wheat TaLCT1 transporter or (c) cation channels. Cadmium also enters the root cells as Cd-chelates by YSL (yellow stripe 1-like) proteins, which are oligopeptide transporters. Cadmium and Pb uptake via the symplastic pathway, and their entry into the xylem occurs via heavy metal P1B-ATPases, which are also known as heavy-metal transporting ATPases (HMAs) [31,36].

The current investigation revealed that with increasing concentrations of Cd and Pb in the growth medium, shoot and root Cd and Pb concentrations were also enhanced. This is the well observed phenomenon that when the metal concentration is increased in the growth medium of plants, it results in a corresponding increase in the metal concentration in the plants' tissues [9]. Similar to our results, many reports showed the excessive uptake of Pb and Cd in plants in the presence of these metals in the rhizosphere [3,14]. The interaction of Cd and Pb was anticipated and quite interesting. When Cd was applied in combination with Pb, the root and shoot Cd contents were even more increased. However, the root and shoot Pb contents were decreased. The higher accumulation of Cd in the presence of Pb may be due to the greater mobility and uptake of Cd than Pb [38]. On the other hand, the reason for the lower uptake of Pb may be the lower absorption and movement of Pb in the normal range of the pH of the growth medium [13,39]. So, it is quite evident that the separate applications of Cd and Pb in the growth medium have different effects than their combined application on the accumulation and translocation of Cd and Pb by plants [40,41].

In line with our results, it was found that the Cd uptake and its translocation from the soil to aerial parts of wheat plants was increased in the presence of Pb [41]. In a nutrient solution, roots accumulated more Cd and Pb as compared to shoots for all the provided concentrations of Pb and Cd. The greater retention of both metals in roots may be due to the greater production of phytochelatins (PCs) in roots and the complex formation of metal ions with these PCs. These PCs are predominantly retained within root vacuoles; hence, the movement of Cd and Pb from roots to shoots is greatly decreased [6,9,42]. The BCFs for both roots and shoots were higher than one for all the Cd and Pb treatments; however, the TFs were less than one for both Cd and Pb for all the treatments. The shoot BCF for Cd was higher than that for Pb, indicating the greater potential of quinoa to accumulate Cd than Pb in its aboveground parts under the same growth conditions. The BCF is usually less in soil culture experiments as compared to hydroponics due to lower availability and mobility of metals [8]. Cadmium and Pb translocation from the root to the shoot was less than one for the sole or combined treatments of Cd and Pb. In line with these findings, under both the soil and hydroponic conditions a limited transport of Cd and Pb from root to shoot in quinoa has been reported [3,13]. Metals tolerance and their translocation from root to shoot demonstrated that quinoa is suitable for restricting the movement of Cd and Pb from the root to the shoot in Cd and Pb co-contaminated soils. Similarly, many other plant species have been found suitable for restricting the metals within the soil and impairing their mobility to the aerial parts of the plants [40,42]. In a previous study by Amjad et al. [8], the health risk assessments of separate treatments of Cd and Pb revealed that due to the very low translocation of Cd and Pb from the root to the shoot, the consumption of quinoa poses no health risks to humans.

A multivariate comparison of different treatments and observations was done using a PCA. This data analysis technique is very suitable for depicting the association among different variables [3]. The PCA and Pearson correlations of our study showed inverse associations of tissue Cd and Pb contents with the growth and physiological attributes of quinoa. Contrarily, oxidative stress indicators (MDA, H₂O₂ and MSI) and antioxidant enzymes (SOD, POD and CAT) had positive associations with Cd and Pb contents. The sole and combined applications of Cd and Pb also had differential effects. The clustering of the combined treatments of both Cd and Pb in the negative axis was an indication of their more damaging effects on the plants' growth and physiology than their separate application.

5. Conclusions

Our results demonstrated that the quinoa genotype 'Puno' has differential tolerance and uptake potentials regarding Cd and Pb under the co-contamination of both metals. On an individual basis, Cd resulted in a greater decrease in the plant growth and physiological attributes of quinoa as compared to Pb. Oxidative stress and an abundant uptake of toxic ions (particularly Cd) were the main causes of the limited growth under the combined stress of Cd and Pb. Enhancement of the activities of antioxidant enzymes ameliorated the metal-induced oxidative damage in quinoa plants. The translocation factors for both metals were less than one, indicating that the greater portion of both Cd and Pb was sequestered in the roots as compared to the shoots. The Cd and Pb tolerance and uptake potential of Puno showed its ability to grow in Cd and Pb co-contaminated soils. However, further research is needed to determine the metal accumulation in quinoa grains and expected health risks posed by growing quinoa in Cd and Pb co-contaminated soils.

Author Contributions: Conceptualization, A.A.B., H.F.A. and G.A.; methodology, A.A.B., and H.F.A.; software, H.F.A.; validation, A.A.B. and H.F.A.; formal analysis, G.A.; investigation, H.F.A. and G.A.; resources, A.A.B. and H.F.A.; data curation, A.A.B. and G.A.; writing—original draft preparation, G.A.; writing—review and editing, A.A.B. and H.F.A.; visualization, H.F.A. and G.A.; project administration, G.A. All authors have read and agreed to the published version of the manuscript.

Funding: This project was funded by the Deanship of Scientific Research (DSR) at King Abdulaziz University, Jeddah, under grant no. G: 278-130-1442.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data will be available as requested.

Acknowledgments: This project was funded by the Deanship of Scientific Research (DSR) at King Abdulaziz University, Jeddah, under grant no. G: 278-130-1442. The authors, therefore, acknowledge with thanks the DSR for its technical and financial support.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Qayyum, M.F.; Rehman, M.Z.U.; Ali, S.; Rizwan, M.; Naeem, A.; Maqsood, M.A.; Khalid, H.; Rinklebe, J.; Ok, Y.S. Residual effects of monoammonium phosphate, gypsum and elemental sulfur on cadmium phytoavailability and translocation from soil to wheat in an effluent irrigated field. *Chemosphere* **2017**, *174*, 515–523. [[CrossRef](#)] [[PubMed](#)]
- Anwar, H.; Shahid, M.; Niazi, N.K.; Khalid, S.; Tariq, T.Z.; Ahmad, S.; Nadeem, M.; Abbas, G. Risk assessment of potentially toxic metal (loid) in *Vigna radiata* L. under wastewater and freshwater irrigation. *Chemosphere* **2021**, *265*, 129124. [[CrossRef](#)]
- Abdal, N.; Abbas, G.; Asad, S.A.; Ghfar, A.A.; Shah, G.M.; Rizwan, M.; Ali, S.; Shahbaz, M. Salinity mitigates cadmium-induced phytotoxicity in quinoa (*Chenopodium quinoa* Willd.) by limiting the Cd uptake and improved responses to oxidative stress: Implications for phytoremediation. *Environ. Geochem. Health* **2021**, *1–15*. [[CrossRef](#)]
- Waseem, A.; Arshad, J.; Iqbal, F.; Sajjad, A.; Mehmood, Z.; Murtaza, G. Pollution Status of Pakistan: A Retrospective Review on Heavy Metal Contamination of Water, Soil, and Vegetables. *BioMed Res. Int.* **2014**, *2014*, 1–29. [[CrossRef](#)] [[PubMed](#)]
- Shahid, M.; Dumat, C.; Khalid, S.; Niazi, N.K.; Antunes, P.M.C. Cadmium Bioavailability, Uptake, Toxicity and Detoxification in Soil-Plant System. *Rev. Environ. Contam. Toxicol.* **2016**, *241*, 73–137. [[CrossRef](#)]
- Zhang, S.; Ni, X.; Arif, M.; Zheng, J.; Stubbs, A.; Li, C. NaCl improved Cd tolerance of the euhalophyte *Suaeda glauca* but not the recretohalophyte *Limonium aureum*. *Plant Soil* **2020**, *449*, 303–318. [[CrossRef](#)]
- Shanying, H.E.; Xiaoe, Y.A.N.G.; Zhenli, H.E.; Baligar, V.C. Morphological and physiological responses of plants to cadmium toxicity: A review. *Pedosphere* **2017**, *27*, 421–443.
- Amjad, M.; Iqbal, M.M.; Abbas, G.; Farooq, A.B.U.; Naeem, M.A.; Imran, M.; Murtaza, B.; Nadeem, M.; Jacobsen, S.-E. Assessment of cadmium and lead tolerance potential of quinoa (*Chenopodium quinoa* Willd.) and its implications for phytoremediation and human health. *Environ. Geochem. Health* **2021**, *1–14*. [[CrossRef](#)]
- Rehman, S.; Abbas, G.; Shahid, M.; Saqib, M.; Farooq, A.B.U.; Hussain, M.; Farooq, A. Effect of salinity on cadmium tolerance, ionic homeostasis and oxidative stress responses in conocarpus exposed to cadmium stress. *Ecotoxicol. Environ.* **2019**, *171*, 146–153. [[CrossRef](#)]
- Shabbir, A.; Saqib, M.; Murtaza, G.; Abbas, G.; Imran, M.; Rizwan, M.; Naeem, M.A.; Ali, S.; Javeed, H.M.R. Biochar mitigates arsenic-induced human health risks and phytotoxicity in quinoa under saline conditions by modulating ionic and oxidative stress responses. *Environ. Pollut.* **2021**, *287*, 117348. [[CrossRef](#)]
- Shahid, M.; Farooq, A.B.U.; Rabbani, F.; Khalid, S.; Dumat, C. Risk assessment and biophysicochemical responses of spinach to foliar application of lead oxide nanoparticles: A multivariate analysis. *Chemosphere* **2019**, *245*, 125605. [[CrossRef](#)]
- Kushwaha, A.; Hans, N.; Kumar, S.; Rani, R. A critical review on speciation, mobilization and toxicity of lead in soil-microbe-plant system and bioremediation strategies. *Ecotoxicol. Environ. Saf.* **2018**, *147*, 1035–1045. [[CrossRef](#)] [[PubMed](#)]
- Iftikhar, A.; Abbas, G.; Saqib, M.; Shabbir, A.; Amjad, M.; Shahid, M.; Ahmad, I.; Iqbal, S.; Qaisrani, S.A. Salinity modulates lead (Pb) tolerance and phytoremediation potential of quinoa: A multivariate comparison of physiological and biochemical attributes. *Environ. Geochem. Health* **2021**, *44*, 257–272. [[CrossRef](#)] [[PubMed](#)]
- Arias, J.A.; Peralta-Videa, J.R.; Ellzey, J.T.; Ren, M.; Viveros, M.N.; Gardea-Torresdey, J.L. Effects of *Glomus deserticola* inoculation on Prosopis: Enhancing chromium and lead uptake and translocation as confirmed by X-ray mapping, ICP-OES and TEM techniques. *Environ. Exp. Bot.* **2010**, *68*, 139–148. [[CrossRef](#)]
- Shahid, M.; Pinelli, E.; Pourrut, B.; Silvestre, J.; Dumat, C. Lead-induced genotoxicity to *Vicia faba* L. roots in relation with metal cell uptake and initial speciation. *Ecotoxicol. Environ. Saf.* **2011**, *74*, 78–84. [[CrossRef](#)]
- Ahmadi, H.; Corso, M.; Verbruggen, N.; Clemens, S. CAX1 suppresses Cd-induced generation of reactive oxygen species in *Arabidopsis halleri*. *Plant. Cell. Environ.* **2018**, *41*, 2435–2448. [[CrossRef](#)] [[PubMed](#)]
- Mishra, K. Lead exposure and its impact on immune system: A review. *Toxicol. In Vitro* **2009**, *23*, 969–972. [[CrossRef](#)]
- Murtaza, B.; Naeem, F.; Shahid, M.; Abbas, G.; Shah, N.S.; Amjad, M.; Bakhat, H.F.; Imran, M.; Niazi, N.K.; Murtaza, G. A multivariate analysis of physiological and antioxidant responses and health hazards of wheat under cadmium and lead stress. *Environ. Sci. Pollut. Res.* **2018**, *26*, 362–370. [[CrossRef](#)]
- Iqbal, S.; Basra, S.M.A.; Afzal, I.; Wahid, A.; Saddiq, M.S.; Hafeez, M.B.; Jacobsen, S.-E. Yield potential and salt tolerance of quinoa on salt-degraded soils of Pakistan. *J. Agron. Crop Sci.* **2019**, *205*, 13–21. [[CrossRef](#)]
- Abbas, G.; Amjad, M.; Saqib, M.; Murtaza, B.; Asif Naeem, M.; Shabbir, A.; Murtaza, G. Soil sodicity is more detrimental than salinity for quinoa (*Chenopodium quinoa* Willd.): A multivariate comparison of physiological, biochemical and nutritional quality attributes. *J. Agron. Crop Sci.* **2021**, *207*, 59–73. [[CrossRef](#)]

21. Shabala, S.; Hariadi, Y.; Jacobsen, S.-E. Genotypic difference in salinity tolerance in quinoa is determined by differential control of xylem Na⁺ loading and stomatal density. *J. Plant Physiol.* **2013**, *170*, 906–914. [[CrossRef](#)] [[PubMed](#)]
22. Lichtenthaler, H.K. Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. *Methods Enzymol.* **1987**, *148*, 350–382.
23. Islam, E.; Liu, D.; Li, T.; Yang, X.; Jin, X.; Mahmood, Q.; Tian, S.; Li, J. Effect of Pb toxicity on leaf growth, physiology and ultrastructure in the two ecotypes of *Elsholtzia argyi*. *J. Hazard. Mater.* **2008**, *154*, 914–926. [[CrossRef](#)] [[PubMed](#)]
24. Sairam, R.K.; Rao, K.V.; Srivastava, G.C. Differential response of wheat genotypes to long term salinity stress in relation to oxidative stress, anti-oxidant activity and osmolyte concentration. *Plant Sci.* **2002**, *163*, 1037–1046. [[CrossRef](#)]
25. Dhindsa, R.S.; Plumb-Dhindsa, P.; Thorpe, T.A. Leaf Senescence: Correlated with Increased Levels of Membrane Permeability and Lipid Peroxidation, and Decreased Levels of Superoxide Dismutase and Catalase. *J. Exp. Bot.* **1981**, *32*, 93–101. [[CrossRef](#)]
26. Hemeda, H.M.; Klein, B.P. Effects of Naturally Occurring Antioxidants on Peroxidase Activity of Vegetable Extracts. *J. Food Sci.* **1990**, *55*, 184–185. [[CrossRef](#)]
27. Aebi, H. Catalase in vitro. *Methods Enzymol.* **1984**, *105*, 121–126.
28. Steel, R.; Torrie, J.; Dickey, D. *Principles and Procedures of Statistics: A Biometrical Approach*, 3rd ed.; McGraw-Hill: New York, NY, USA, 1997.
29. Panda, A.; Rangani, J.; Kumari, A.; Parida, A. Efficient regulation of arsenic translocation to shoot tissue and modulation of phytochelatin levels and antioxidative defense system confers salinity and arsenic tolerance in the Halophyte *Suaeda maritima*. *Environ. Exp. Bot.* **2017**, *143*, 149–171. [[CrossRef](#)]
30. Shabir, R.; Abbas, G.; Saqib, M.; Shahid, M.; Shah, G.M.; Akram, M.; Niazi, N.K.; Naeem, M.A.; Hussain, M.; Ashraf, F. Cadmium tolerance and phytoremediation potential of acacia (*Acacia nilotica* L.) under salinity stress. *Int. J. Phytoremediation* **2018**, *20*, 739–746. [[CrossRef](#)]
31. Gallego, S.M.; Pena, L.B.; Barcia, R.A.; Azpilicueta, C.E.; Iannone, M.F.; Rosales, E.P.; Zawoznik, M.S.; Groppa, M.D.; Benavides, M.P. Unravelling cadmium toxicity and tolerance in plants: Insight into regulatory mechanisms. *Environ. Exp. Bot.* **2012**, *83*, 33–46. [[CrossRef](#)]
32. Shahid, M.; Dumat, C.; Khalid, S.; Schreck, E.; Xiong, T.; Niazi, N.K. Foliar heavy metal uptake, toxicity and detoxification in plants: A comparison of foliar and root metal uptake. *J. Hazard. Mater.* **2017**, *325*, 36–58. [[CrossRef](#)] [[PubMed](#)]
33. Munns, R.; Tester, M. Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol.* **2008**, *59*, 651–681. [[CrossRef](#)] [[PubMed](#)]
34. Amjad, M.; Akhtar, S.S.; Yang, A.; Akhtar, J.; Jacobsen, S.E. Antioxidative Response of Quinoa Exposed to Iso-Osmotic, Ionic and Non-Ionic Salt Stress. *J. Agron. Crop Sci.* **2015**, *201*, 452–460. [[CrossRef](#)]
35. Li, X.; Zhang, X.; Wang, X.; Yang, X.; Cui, Z. Bioaugmentation-assisted phytoremediation of lead and salinity co-contaminated soil by *Suaeda salsa* and *Trichoderma asperellum*. *Chemosphere* **2019**, *224*, 716–725. [[CrossRef](#)] [[PubMed](#)]
36. Sharma, P.; Dubey, R.S. Lead toxicity in plants. *Braz. J. Plant Physiol.* **2005**, *17*, 35–52. [[CrossRef](#)]
37. Pourrut, B.; Shahid, M.; Dumat, C.; Winterton, P.; Pinelli, E. Lead uptake, toxicity, and detoxification in plants. *Rev. Environ. Contam. Toxicol.* **2011**, *213*, 113–136. [[PubMed](#)]
38. Souri, Z.; Karimi, N.; De Oliveira, L.M. Antioxidant enzymes responses in shoots of arsenic hyperaccumulator, *Isatis cappadocica* Desv., under interaction of arsenate and phosphate. *Environ. Technol.* **2017**, *39*, 1316–1327. [[CrossRef](#)]
39. Abbaspour, A.; Kalbasi, M.; Hajrasuliha, S.; Fotovat, A. Effect of Organic Matter and Salinity on Ethylenediaminetetraacetic Acid-Extractable and Solution Species of Cadmium and Lead in Three Agricultural Soils. *Commun. Soil Sci. Plant Anal.* **2008**, *39*, 983–1005. [[CrossRef](#)]
40. Nawrot, N.; Wojciechowska, E.; Pazdro, K.; Szmagliński, J.; Pempkowiak, J. Uptake, accumulation, and translocation of Zn, Cu, Pb, Cd, Ni, and Cr by *P. australis* seedlings in an urban dredged sediment mesocosm: Impact of seedling origin and initial trace metal content. *Sci. Total Environ.* **2021**, *768*, 144983. [[CrossRef](#)]
41. Xu, J.; Hu, C.; Wang, M.; Zhao, Z.; Zhao, X.; Cao, L.; Lu, Y.; Cai, X. Changeable effects of coexisting heavy metals on transfer of cadmium from soils to wheat grains. *J. Hazard. Mater.* **2021**, *423*, 127182. [[CrossRef](#)]
42. Yan, X.; An, J.; Yin, Y.; Gao, C.; Wang, B.; Wei, S. Heavy metals uptake and translocation of typical wetland plants and their ecological effects on the coastal soil of a contaminated bay in Northeast China. *Sci. Total Environ.* **2021**, *803*, 149871. [[CrossRef](#)] [[PubMed](#)]