



AUTHOR CORRECTION OPEN



Author Correction: The 1-minute sit-to-stand test to detect desaturation during 6-minute walk test in interstitial lung disease

Keiji Oishi , Kazuto Matsunaga , Maki Asami-Noyama, Tasuku Yamamoto, Yukari Hisamoto, Tetsuya Fujii, Misa Harada, Junki Suizu, Keita Murakawa, Ayumi Chikumoto, Kazuki Matsuda, Haruka Kanesada, Yujiro Kikuchi, Kazuki Hamada, Sho Uehara, Ryo Suetake, Syuichiro Ohata, Yoriyuki Murata, Yoshikazu Yamaji, Kenji Sakamoto, Kosuke Ito, Hisayuki Osoreda, Nobutaka Edakuni, Tomoyuki Kakugawa, Tsunahiko Hirano and Masafumi Yano

npj Primary Care Respiratory Medicine (2022)32:9; <https://doi.org/10.1038/s41533-022-00274-y>

Correction to: *npj Primary Care Respiratory Medicine* <https://doi.org/10.1038/s41533-022-00268-w>, published online 27 January 2022

In Table 2 of this article, the heading for the column labelled “Nadir SpO₂ < 90% in the 6MWT” was mistakenly located over the column labelled “Nadir SpO₂ < 90% in the 1STST”. The original article has been corrected.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022