



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Contents lists available at ScienceDirect

Environmental Research

journal homepage: www.elsevier.com/locate/envres

Review article



The role of outdoor and indoor air quality in the spread of SARS-CoV-2: Overview and recommendations by the research group on COVID-19 and particulate matter (RESCOP commission)

Prisco Piscitelli^{a,r}, Alessandro Miani^{a,b,*}, Leonardo Setti^{a,c,**}, Gianluigi De Gennaro^{a,d}, Xavier Rodo^e, Begona Artinano^f, Elena Vara^g, Lisa Rancan^g, Javier Arias^h, Fabrizio Passariniⁱ, Pierluigi Barbieri^{a,j}, Alberto Pallavicini^k, Alessandro Parente^{l,x}, Edoardo Cavalieri D'Oro^m, Claudio De Maio^m, Francesco Saladino^m, Massimo Borelliⁿ, Elena Colicino^o, Luiz Marcos G. Gonçalves^p, Gianluca Di Tanna^q, Annamaria Colao^r, Giovanni S. Leonardi^s, Andrea Baccarelli^t, Francesca Dominici^u, John P.A. Ioannidis^v, José L. Domingo^w, on behalf of the RESCOP Commission established by Environmental Research (Elsevier)

^a Italian Society of Environmental Medicine (SIMA), Milan, Italy^b Department of Environmental Science and Policy, University of Milan, Milan, Italy^c Department of Industrial Chemistry, University of Bologna, Bologna, Italy^d Department of Biology, University of Bari "Aldo Moro", Bari, Italy^e ICREA and Climate & Health Program, ISGlobal, Barcelona, Spain^f Unit Atmospheric Pollution and POP Characterization, CIEMAT, Madrid, Spain^g Department of Biochemistry and Molecular Biology, School of Medicine, Complutense University, Madrid, Spain^h School of Medicine, Complutense University, Madrid, Spainⁱ Interdepartmental Centre for Industrial Research "Renewable Sources, Environment, Blue Growth, Energy", University of Bologna, Rimini, Italy^j Department of Chemical and Pharmaceutical Sciences, University of Trieste, Trieste, Italy^k Department of Life Sciences, University of Trieste, Trieste, Italy^l Université libre de Bruxelles (ULB), Ecole Polytechnique de Bruxelles, Département d'Aéro-Thermo-Mécanique, Brussels, Belgium^m Chemical, Biological, Radiological and Nuclear Unit (NBCRE), Italian National Fire and Rescue Service, Milan, Italyⁿ UMG School of PhD Programmes, University Magna Graecia of Catanzaro, Italy^o Department of Environmental Medicine and Public Health at the Icahn School of Medicine at Mount Sinai, New York, USA^p Natalnet Associate Labs, Universidade Federal do Rio Grande do Norte, Natal, Brazil^q BioStatistics & Data Science Division, Meta-Research and Evidence Synthesis Unit, The George Institute for Global Health, Faculty of Medicine, University of New South Wales, Sydney, Australia^r UNESCO Chair on Health Education and Sustainable Development, University of Naples Federico II, Naples, Italy^s Department of Public Health, Environments and Society, London School of Hygiene and Tropical Medicine (LSHTM), London, UK^t Chair of the Department of Environmental Health Sciences, Columbia University, New York, USA^u Harvard T.H. Chan School of Public Health, Boston, USA^v Departments of Medicine, of Epidemiology and Population Health, of Biomedical Data Science and of Statistics, Stanford University, Stanford, CA, USA^w Laboratory of Toxicology and Environmental Health, Universitat Rovira i Virgili, School of Medicine, Reus, Spain^x Brussels Institute for Thermal-fluid systems and clean Energy (BRITE), Brussels, Belgium

* Corresponding author. Department of Environmental Science and Policy, University of Milan, Via Celoria 2, 20133, Milan, Italy.

** Corresponding author.

E-mail addresses: piscitelli@unescochairnapoli.it (P. Piscitelli), alessandro.miani@unimi.it (A. Miani), leonardo.setti@unibo.it (L. Setti), gianluigi.degennaro@uniba.it (G. De Gennaro), xavier.rodo@isglobal.org (X. Rodo), b.artinano@ciemmat.es (B. Artinano), evarami@ucm.es (E. Vara), lisaranc@ucm.es (L. Rancan), javierar@ucm.es (J. Arias), fabrizio.passarini@unibo.it (F. Passarini), barbierp@units.it (P. Barbieri), pallavic@units.it (A. Pallavicini), alessandro.parente@ulb.be (A. Parente), edoardo.cavalieridoro@vigilifuoco.it (E.C. D'Oro), claudio.demaio@vigilifuoco.it (C. De Maio), francescosaladino72@gmail.com (F. Saladino), massimo.borelli@gmail.com (M. Borelli), elena.colicino@mssm.edu (E. Colicino), lmarcos@dca.ufm.br (L.M.G. Gonçalves), gditanna@georgeinstitute.org.au (G. Di Tanna), colao@unescochairnapoli.it (A. Colao), gleonardi@doctors.org.uk (G.S. Leonardi), ab4303@cumc.columbia.edu (A. Baccarelli), fdominic@hsp.harvard.edu (F. Dominici), jioannid@stanford.edu (J.P.A. Ioannidis), joseluis.domingo@urv.cat (J.L. Domingo).

<https://doi.org/10.1016/j.envres.2022.113038>

Received 20 January 2022; Received in revised form 24 February 2022; Accepted 24 February 2022

Available online 26 February 2022

0013-9351/© 2022 Elsevier Inc. All rights reserved.

ARTICLE INFO

Keywords:

COVID-19
Air quality
Air pollution
Indoor environments
Ventilation
CO2 monitoring

ABSTRACT

There are important questions surrounding the potential contribution of outdoor and indoor air quality in the transmission of SARS-CoV-2 and perpetuation of COVID-19 epidemic waves. Environmental health may be a critical component of COVID-19 prevention. The public health community and health agencies should consider the evolving evidence in their recommendations and statements, and work to issue occupational guidelines. Evidence coming from the current epidemiological and experimental research is expected to add knowledge about virus diffusion, COVID-19 severity in most polluted areas, inter-personal distance requirements and need for wearing face masks in indoor or outdoor environments. The COVID-19 pandemic has highlighted the need for maintaining particulate matter concentrations at low levels for multiple health-related reasons, which may also include the spread of SARS-CoV-2. Indoor environments represent even a more crucial challenge to cope with, as it is easier for the SARS-CoV-2 to spread, remain vital and infect other subjects in closed spaces in the presence of already infected asymptomatic or mildly symptomatic people. The potential merits of preventive measures, such as CO₂ monitoring associated with natural or controlled mechanical ventilation and air purification, for schools, indoor public places (restaurants, offices, hotels, museums, theatres/cinemas etc.) and transportations need to be carefully considered. Hospital settings and nursing/retirement homes as well as emergency rooms, infectious diseases divisions and ambulances represent higher risk indoor environments and may require additional monitoring and specific decontamination strategies based on mechanical ventilation or air purification.

1. Introduction

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is the etiologic agent of the COVID-19 pandemic. The incubation period of COVID-19 lasts from 2 to 12 days, with an average of 5.1 days and several possible symptoms (Lauer et al., 2020). Enormous international scientific efforts across a range of disciplines are underway to understand the factors determining the transmission and infectivity of the new coronavirus, with the aim of minimizing its spread, reducing the diffusion rate and developing new therapeutic treatments or vaccines. Healthy people probably most commonly get infected via inhalation of viral particles spread by patients during normal speaking, sneezing and coughing (Chatterjee, 2020). However, it is necessary to consider the possibility of SARS-CoV-2 spreading via other routes than by infected droplets (Morawska and Cao, 2020). Surfaces touched by infected persons, water and sewage, garbage, or soil may also represent routes of viral spreading, but their relative contribution remains contentious (Onakpoya et al., 2021). As resulting from the review performed by Marquès and Domingo (2021), SARS-CoV-2 can last several hours or even few days on different surfaces, but it can be inactivated by using the common available disinfectants and biocides. Consequently, it seems that the persistence of SARS-CoV-2 on inert surfaces should not be regarded as an issue of special concern for the transmission of the new coronavirus if compared to the main airborne contagion route through aerosols Marquès and Domingo (2021).

The main transmission route of SARS-CoV-2 seems to be the droplet route (particles >5 µm) (Jin et al., 2020). However, on May 2021 the CDC updated their COVID-19 prevention guidelines specifically referring to the SARS-CoV-2 transmission also from aerosolized particles (“inhalation of very fine respiratory droplets and aerosol particles”) that are much smaller than large respiratory droplets and can remain suspended in the environment – especially indoor – for minutes or possibly hours, thus “increasing the amount of viral particles to which a person is exposed”. Actually, CDC recognized that “once infectious droplets and particles are exhaled, they move outward from the source” and that the transmission of SARS-CoV-2 may occur also at distances greater than six feet (2 m) from the infectious source, despite it being less likely than at closer distances (CDC, 2020). There is debate about how long viral particles can remain infectious while suspended in air. Regardless, the risk for infection decreases with increasing distance from the source and increasing time after exhalation. “Heavier respiratory droplets containing the virus fall to the ground or other surfaces under the force of gravity and the very fine droplets and aerosol particles remain in the air stream” (CDC, 2020), being progressively diluted in relation to the growing volume and streams of air they encounter. The progressive loss of viral viability and infectiousness over time is also influenced by

environmental factors such as temperature, humidity, and ultraviolet radiation such as sunlight (CDC, 2020). Such factors may be key contributors to the emerging seasonal pattern of SARS-CoV-2 epidemic waves.

1.1. COVID-19 and air pollution: the research group on COVID-19 and particulate matter (RESCOP)

The potential relationship between COVID-19 epidemic waves and air pollution poses tantalizing questions. Domingo et al. (2020) suggested that the burden of COVID-19 is more severe in those areas characterized by higher levels of particulate matter (PM). Setti et al. (2020a) observed an association between the number of Italian provinces with daily averaged PM₁₀ concentrations exceeding the European limit values (set at 50 µg/m³) and the subsequent number of COVID-19 cases. Significant associations were found between mean concentrations of PM_{2.5} during the month of February 2020 and the total number of Italian COVID-19 cases on 31 March 2020. These ecological analyses need to be seen with great caution. However, Setti et al. (2020b) also evidenced that air samples collected from open spaces in the industrial zone in Bergamo province were positive for SARS-CoV-2 RNA. Researchers demonstrated viral RNA in 30% of PM₁₀ (particulate matter) of outdoor air samples (Setti et al., 2020b). Research carried out by the Harvard School of Public Health confirmed an association between increases in particulate matter concentrations and mortality rates due to COVID-19 (Wu et al., 2020a).

An international research group was initially fostered by the Italian Society of Environmental Medicine (SIMA) with the aim of investigating possible links between COVID-19 and particulate matter, as well as the topic of indoor air quality in relation to COVID-19 diffusion (RESCOP, *Research Group on COVID-19 and Particulate Matter*). Starting from this initiative, *Environmental Research* (Elsevier journal) established the RESCOP thematic Commission on COVID-19 and outdoor/indoor air quality, which has promoted a Special Issue devoted to this specific topic, resulting in the submission of about 150 papers and publication of 76 of them from all over the world.

The primary objective of the RESCOP Commission was to involve international experts already working on these specific topics by developing analytical protocols according to which it could be possible to confirm or exclude the presence of SARS-CoV-2 RNA in indoor environments or on outdoor particulate matter, as well as performing specific assessments of the potential persistent viability and virulence (infectious potential) of the virus load eventually detected on particulate matter and indoor environments. Obviously, detection of viral genomic material alone does not prove the presence of intact virus and infectivity.

The secondary objective of the Commission was to examine whether the viral detection on PM10/PM2.5 could be used as early indicator of COVID-19 epidemics recurrence, and whether such information could be used for preventive purposes. Searching for SARS-CoV-2 could be implemented also to assess contamination of indoor environments such as hospitals, ambulatories, restaurants, shopping malls, shops, factories, pubs, as well as airplanes, trains, cruise ships etc.

The third objective of the Commission activities was to deepen the virus diffusion and health aggression models in relation to pollution parameters. Information coming from the articles published by the Commission on the Special Issue are expected to add knowledge about virus diffusion, COVID-19 severity in most polluted areas, inter-personal distance requirements and need for wearing face masks in indoor or outdoor environments as well as to whether one should change recommendations for inter-personal distance in specific circumstances.

Moreover, the research outcomes might help to highlight the need for maintaining particulate matter concentrations at low levels for multiple health-related reasons, including potentially the spread of SARS-CoV-2 (Setti et al., 2020a). The levels of particulate matter and NO₂ should be as much as possible close to the new thresholds recently set by WHO in the updated Air Quality Guidelines presented in September 2021 (WHO, 2021).

1.2. Environmental determinants of SARS-CoV-2 transmission

The systematic review carried out by Maleki et al. (2021) reported that the main transmission route of SARS-CoV-2 is the droplet route (particles > 5 µm) (Jin et al., 2020). However, Mecnas et al. (2020) showed that low humidity and temperature may increase the vitality of SARS-CoV-2 in an aerosol. This aspect is particularly important in relation to indoor environments because – according to Morawska and Cao (2020) and Paules et al. (2020) – small aerosol particles with a higher viral load probably can be transferred up to 10 m from the emission source. On the other side, the persistence of SARS-CoV-2 in hospital wards (COVID-19 outbreak) in Wuhan was confirmed by the number of virus copies (from 1 to 42 copies/m³) in a study of Liu et al. (2020a).

1.3. COVID-19 increased risk of transmission in indoor environments: possible preventive measures

Specific preventive measures should be seriously considered for indoor environments and public transportations, given the potentially increased risk of SARS-CoV-2 transmission in case of inadequate ventilation or air handling in enclosed spaces, within which “the concentration of exhaled respiratory fluids, especially very fine droplets and aerosol particles, can build-up in the air space” (CDC, 2020). The risk is higher in those crowded indoor environments where prolonged exposures (>15 min) and exhalation of infected respiratory fluids may increase by some activities (e.g. from physical exercise, raised voice such as shouting, singing etc.).

Physical distancing and use of well-fitting masks may not be enough in crowded, high-risk activities. In this perspective, the added value of specific systems for Indoor Air Quality monitoring needs further research. Di Gilio et al. (2021), validated that CO₂ portable monitors (10 cm × 10cm) can be used to assess the potential risk of infected droplets/aerosols inhalation when CO₂ levels reach a defined threshold (fixed at 700 ppm in more recent studies) and foster the natural ventilation of the space by simply opening doors and windows for the time needed to reduce CO₂ levels. Of note, normal breathing of a child aged 7–9 years old generates 14 L of CO₂ per hour, which is 50% lower than the amount produced by a teenager, while in conditions of moderate physical activity, a 15 years old student can release up to 85 L of CO₂ per hour. In addition, air purifiers or controlled mechanical ventilation (CMV) devices (when certified by scientific reports concerning their safety and efficacy in air dilution and particles’ removal) could be

installed to allow a correct and constant air exchange of indoor environments without the need of opening the windows or where natural air exchange is not sufficient to remain below the threshold of 700 ppm.

Formal decision and cost-effectiveness analyses are needed to establish the utility of such monitors and devices and how widely they should be adopted, if at all. Obviously, the level of epidemic activity and the fatality rate in an area may affect the cost-effectiveness ratios. These ratios may also change as different viral strains of different infectivity become dominant and as fatality rate decreases sharply with some interventions, primarily vaccinations. Based on current knowledge, the minimum functional standard is represented by decentralized double flow CMV with pre-filtration of the incoming air (that can be heated in the winter months so as not to reduce the indoor gradient of temperature). The air flow per hour to be exchanged must be determined in m³ considering the volume of indoor space. For a room of 90 m³, 10–15 cycles per hour of complete air exchange are needed to mitigate the risk of breathing in infected material. This corresponds to 900–1400 m³ of air in entrance and the same extracted from the indoor environment each hour. However, taking into account the high cost of the above-mentioned technologies, a cost-effective system could be represented by the installation of continuous air extractors which can allow huge amount of air exchange by simply extracting indoor air from the enclosed spaces (rooms, offices, etc.) thus facilitating the maintenance of good air quality from natural ventilation through windows and door openings.

If choosing centralized systems, it should be highlighted the issue of expensive and continuous sanitizing of air ducts. Furthermore, air purification systems (able to kill bacteria and viruses) may be considered – especially for indoor public settings – making sure that they use high performance HEPA filters or ULPA filters DFS capable of nanometric disinfectant filtration (removal of nanoparticles).

Concerning special indoor environments such as hospital settings and nursing/retirement homes, tested by Zupin et al., (2021), further research should determine whether there are warning thresholds of indirect indicators of contagion risk (i.e. CO₂) and assess the viability and infectivity potential of SARS-CoV-2 present in the air of COVID hospitals, ambulances and emergency rooms.

Once cost-effectiveness issues are addressed by non-conflicted studies and further evidence accumulates, EU Member States’ Governments may foster and develop financial mechanisms to support the adoption of the most appropriate systems to reduce the risk of contagion on public transportations (trains, buses, airplanes, cruise ships etc.) and indoor environments (i.e. schools, banks, offices, workplaces, halls, hotels, theatres, cinemas, disco etc.). Specific pilots are under development by several organizations such as the United Nations International Center for Genetic Engineering and Biotechnology (ICGEB) for the treatment of indoor air inside cruise ships, and by the Italian Society of Environmental Medicine for museums.

A Reference Guide for Indoor Air Quality (IAQ) management and monitoring may be adopted in each indoor public setting - starting from schools and Universities - with a IAQ coordinator appointed to address the most common issues related to IAQ management.

1.4. HVAC and local exhaust ventilation systems

Borro et al. (2021) and Aghalari et al. (2021) have demonstrated that SARS-CoV-2 is mainly transmitted through exhalations from the airways of infected persons, so that Heating, Ventilation and Air Conditioning (HVAC) systems might play a role in spreading the infection in indoor environments. Borro and colleagues modeled the role of HVAC systems in the diffusion of the contagion through a Computational Fluid Dynamics (CFD) simulations of cough at Vatican children’s hospital in Rome. Both waiting rooms and hospital rooms were modeled as indoor scenarios. A specific Infection-Index (η) parameter was used to estimate the amount of contaminated air inhaled by each person present in the simulated indoor scenarios. The potential role of exhaust air ventilation

systems placed above the coughing patient's mouth was also assessed.

Their CFD-based simulations showed that HVAC air-flow remarkably enhances infected droplets diffusion in the whole indoor environment within 25 s from the cough event, despite the observed dilution of saliva particles containing the virus. In the waiting room simulation, the Infection-Index (η) parameter increases the faster the higher the HVAC airflow. Greater flows of air conditioning correspond to greater diffusion of the infected droplets.

The proper use of Local Exhaust Ventilation systems (LEV) simulated in the hospital room reduced infected droplets spreading from the patient's mouth in the first 0.5 s following the cough event. In the hospital room, the use of LEV system reduced the η index computed for the patient hospitalized at the bed next to the spreader, with a decreased possibility of contagion.

1.5. Other interventions with potential effect on COVID-19 spread

Grimalt et al. (2021) performed a systematic sampling and analysis of airborne SARS-CoV-2 RNA in different hospital areas including ICU rooms to assess viral spread, finding out the highest occurrence of RNA in the rooms with COVID-19 patients and the adjacent corridor. There was a significant transfer of viruses from the COVID-19 patients' rooms to the corridors, while patients intubated and connected to respirators that filtered all exhaled air resulted in significantly lower viral concentrations in adjacent corridors. Systematic air filtration was performed in rooms with COVID-19 infected and uninfected patients, in corridors adjacent to these rooms and intensive care units, and in open spaces, showing the opportunity to use ventilation systems in order to minimize possible contagion risk, as well as the importance of direct monitoring of SARS-CoV-2 in the air of indoor environments to prevent unexpected viral exposures.

Building design strategies may also be important to consider in mitigating threats to occupants. The review by Megahed et al. (2021) aims to draw architects' attention towards the high risk of airborne transmission of diseases by providing the latest updates and solutions to better understand the environmental and health issues associated with COVID-19. Based on the complexity of the problem and the need for interdisciplinary research, this study presents a conceptual model that addresses the integration of engineering controls, design strategies and air disinfection techniques.

Finally, some of the societal and individual-level measures used to curb COVID-19 transmission may have effects on air quality. For examples, lockdown measures had drastic impacts on social and economic situations, but they may have generated some positive (even if short-lived) impacts on the environment. Recent data released by National Aeronautics and Space Administration (NASA) and European Space Agency (ESA) indicate a reduction in the levels of environmental pollution particularly in dimensions of air pollution across the megacities of the globe (Khan et al., 2021). On the other side, Dargahi et al. (2021) have demonstrated the presence of SARS-CoV-2 virus in the internal and external surfaces of various masks used by patients and staff with SARS coronavirus, as well as the possibility of airborne transmission in Imam Khomeini Hospital of Ardabil (Iran).

1.6. Air pollution and COVID-19

Previous research has assessed the presence of SARS-CoV-2 in outdoor air particulate matter (PM) in urban areas of Northern Italy and USA (Setti et al., 2020a). Linillos-Pradillo et al. (2021) investigated the presence of SARS-CoV-2 RNA in outdoor air samples (on PM₁₀, PM_{2.5} and PM₁). Six samples of PM₁₀, PM_{2.5} and PM₁ were collected between May 4th and 22nd 2020 in Madrid, on quartz fiber filters by using MCV high volume samplers (30 m³ h⁻¹ flow) with three inlets (Digital DHA-80) for sampling PM₁₀, PM_{2.5} and PM₁. RNA extraction and amplification was performed according to the protocol recently set by Setti et al. (2020b) in Italy. No presence of SARS-CoV-2 in quartz fiber

filters samplers for PM₁₀, PM_{2.5} and PM₁ fractions was observed. The authors concluded that the absence of viral genomes could be due to different factors including: limited social interactions, masks, and economic activities resulting in reduced circulation of the coronavirus, lower daily PM concentration in outdoor air, and higher temperature that characterizes spring season.

Wu et al. (2020b) described the challenges and outline promising directions and opportunities concerning the assessment of long-term exposure to air pollution and possible increase in the severity of COVID-19 health outcomes, including death. The same authors had already highlighted that higher historical PM_{2.5} exposures in the U.S. were positively associated with higher county-level COVID-19 mortality rates after accounting for many area-level confounders (Wu et al., 2020a).

At this stage, research published is yet inconclusive regarding the role of air pollution on the geographic spread of the disease both regionally and globally. Further studies and more systematic research on the relationship of air pollution and COVID-19 should be encouraged and conducted. Of course, there are many other reasons to act decisively on reducing air pollution, besides any direct effect on COVID-19 transmission. Exposure to ambient air pollution is related to 4.2 million premature deaths per year worldwide (WHO 2021) and it is associated with a variety of adverse health outcomes, such as respiratory and cardiovascular morbidity. Furthermore, exposure to air pollution can increase human sensitivity to respiratory pathogens via damage to the respiratory track or via airborne transmission on the surface of particulate matter, and might represent an additional factor influencing COVID-19 morbidity and mortality rates.

Barnett-Itzhaki et al. (2021) examined the association between population-weighted long term exposure to PM_{2.5} and NO_x, and the morbidity and mortality over time following the detection of the first COVID-19 positive case in 36 OECD countries. They found that PM 2.5 concentrations in 2015–2017 were positively correlated with COVID-19 morbidity and mortality after 10, 20, 40 and 60 days since the first confirmed case in all countries. NO_x concentrations in 2015–2017 and country's density (population/km²) were also positively correlated with COVID-19 morbidity and mortality on the 60th day. All multivariate linear regressions of PM_{2.5} concentration models showed consistent signals. The authors concluded that long-term exposure to air pollutants concentrations exceeding WHO 2005 guidelines might exacerbate morbidity and mortality rates from COVID-19.

These results should be validated carefully given their ecological design. However, they could raise another red flag globally among decision makers about the need of reducing air pollution and its harmful effects.

The questioned link between air pollution and COVID-19 spreading or related mortality represents a hot topic that has immediately been regarded in the light of divergent views. Becchetti et al. (2021) analyzed available literature concerning the link between air quality, as measured by different pollutants and a number of COVID-19 outcomes, such as the number of positive cases, deaths, and excess mortality rates documenting the existence of substantial evidence produced worldwide concerning the role played by air pollution on health in general and on COVID-19 outcomes in particular. These results support both long-term exposure effects and short-term consequences (including the hypothesis of particulate matter acting as viral "carrier"). The authors concluded that the link between air pollution and COVID-19 outcomes is corroborated by several different research methodologies. It can be argued, therefore, that policy implications should be drawn from a rational assessment of these findings as "not taking any action" represents an action itself. In this perspective, interesting work has been performed by the Scientific Foresight Unit (STOA) of the European Parliament Research Service (EU Parliament, 2021).

De Angelis et al. (2021) carried out an ecological study to assess the association between long-term exposure to particulate matter (PM) and NO₂ on COVID-19 incidence and all-cause mortality taking into account

demographic, socioeconomic and meteorological variables. Several demographic and socioeconomic variables were associated with COVID-19 incidence and all-cause mortality. An increase in average winter temperature was associated with a non-linear decrease in COVID-19 incidence and all-cause mortality, while an opposite trend emerged for the absolute humidity parameters. An increase of $10 \mu\text{g}/\text{m}^3$ in the mean annual concentrations of PM_{2.5} and PM₁₀ over the previous years was associated with a 58% and 34% increase in COVID-19 incidence rate, respectively. Similarly, a $10 \mu\text{g}/\text{m}^3$ increase of annual mean PM_{2.5} concentrations was associated with a 23% increase in all-cause mortality. An inverse association was found between NO₂ levels and COVID-19 incidence and all-cause mortality. Similar evidence was reported by Mele et al. (2021) and Gujral et al. (2021) who monitored the incidence in three different neural networks for the cities of Paris, Lyon and Marseille as well as in Los Angeles and Ventura counties of California, respectively. The effect of the air quality on viral spread was demonstrated also by Sarawut-Sangkham et al. (2021) in the Bangkok Metropolitan region.

There is evidence that chronic and short-term exposure to air pollution exacerbates symptoms and increases mortality rates for similar respiratory diseases. This is consistent with early studies of COVID-19 mortality rates, but these results need to be confirmed and further consolidated by controlling for individual-level risk factors. Due to intense industrialization and urbanization, air pollution has become a serious global concern as a hazard to human health.

It is known that exposure to atmospheric particulate matter (PM) causes severe health problems in human and significant damage to the physiological systems. Hence, it is important to understand the adverse effects of PM in human health in their multidimensional nature. The review performed by Zhu et al. (2021) provided insights on the detrimental effects of PM in various human health problems including respiratory, circulatory, nervous, and immune systems along with their possible mechanisms of toxicity. Furthermore, the potential effects of short- and long-term exposure to atmospheric pollution on COVID-19 risk and fatality rates were well described in the analysis of the first epidemic wave in Northern Italy (Ho et al., 2021) or in the Catalan Tarragona Province in Spain (Marquès and Domingo, 2022). In the U.S., Zhou et al. (2021) found strong evidence that wildfires amplified the effect of short-term exposure to PM_{2.5} on COVID-19 cases and deaths, although with substantial heterogeneity across counties.

1.7. Standardization issues

Robotto et al. (2021) described a coherent preliminary approach to SARS-CoV-2 indoor and outdoor air sampling in order to overcome the evident lack of standardization. Three aspects are highlighted here in order to use this experience like a standard protocol adopted by the RESCOP Commission. First, quality and consistency to air sampling relies on the development of recovery tests using standard materials and investigating sampling materials, techniques, timing, sample preservation and pre-treatments. Second, in order to overcome the shortcomings of every single sampling technique, coupling different samplers in parallel sampling could be an efficient strategy to collect more information and make data more reliable. Third, with regards to airborne virus sampling, the results could be confirmed by simplified emission and dilution models.

2. Conclusions

Evidence coming from the current epidemiological and experimental research is expected to add knowledge about virus diffusion, COVID-19 severity in most polluted areas, inter-personal distance requirements and the effects of wearing face masks in different circumstances in indoor or outdoor environments. Moreover, the research outcomes might help to highlight the need for maintaining particulate matter concentrations at low levels, for many health-related reasons, besides whatever

COVID-19-related effects. Indoor environment represents a challenge to cope with, as it is easier for SARS-CoV-2 to spread, remain vital and infect other subjects in presence of already infected asymptomatic or mildly symptomatic people in enclosed spaces. Effective preventive measures should be adopted in indoor public places and transportations, factoring effectiveness, epidemic activity, and cost considerations. Concerning special indoor environments such as hospital settings and nursing/retirement homes, further research is needed to determine warning thresholds of indirect indicators of contagion risk (i.e. CO₂), as well as to assess the viability and infective potential of SARS-CoV-2 present in the air of COVID hospitals, ambulances and emergency rooms.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments:

Authors are grateful to all the members of the RESCOP study group who are not listed among the authors of this paper and particularly to: Prof. Frank Kelly, head of the Global Centre of Air pollution research at Imperial College in London; Prof. Antonio Alcami from Severo Ochoa Center for Molecular Biology in Madrid; Prof. Darby Jack and Prof. Steven Chillrud from Columbia University, New York; Prof. M. Cristina Tirado von der Pahlen, director of the UCLA Institute for the Environment and Sustainability (Los Angeles); Prof. Cyril Gueydan, Mr. Oliveir Berten, Ms. Anne Botteaux from Université libre de Bruxelles (ULB), and Dr. Sandrine Bladt from Bruxelles Environnement; Prof. Antonio Marco Pantaleo (Imperial College, London); Prof. Francesco Salustri (University of Oxford); Prof. Igor Pereira (University of Rio Grande Do Norte, Brazil); Prof. Nguyen Tien Huy (University of Nagasaki, Japan); Prof. Riccardo Pansini (University of Dali, Yunnan, China), and Prof. Davide Fornacca (University of Geneva). Authors are also grateful to Daniel Lovegrove (Elsevier) for his support and advice in setting up the RESCOP Commission on Environmental Research journal.

References

- Aghalari, Z., Dahms, H.U., Sosa-Hernandez, J.E., Oyervides-Muñoz, M.A., Parra-Saldívar, R., 2021 Feb 4. Evaluation of SARS-CoV-2 transmission through indoor air in hospitals and prevention methods: a systematic review. *Environ. Res.* 110841.
- Barnett-Itzhaki, Z., Levi, A., 2021 Apr 1. Effects of chronic exposure to ambient air pollutants on COVID-19 morbidity and mortality-A lesson from OECD countries. *Environ. Res.* 195, 110723.
- Becchetti, L., Beccari, G., Conzo, G., Conzo, P., De Santis, D., Salustri, F., 2021 Feb 1. Air quality and COVID-19 adverse outcomes: divergent views and experimental findings. *Environ. Res.* 193, 110556.
- Borro, L., Mazzei, L., Raponi, M., Piscitelli, P., Miani, A., Secinaro, A., 2021 Feb. The role of air conditioning in the diffusion of Sars-CoV-2 in indoor environments: a first computational fluid dynamic model, based on investigations performed at the Vatican State Children's hospital. *Environ. Res. Epub* 2020 Oct 15193:110343.
- CDC, 2020. National Center for Immunization and Respiratory Diseases (NCIRD), Division of Viral Diseases. CDC COVID-19 Science Briefs. Centers for Disease Control and Prevention (US, Atlanta (GA)). Scientific Brief: SARS-CoV-2 Transmission. [Updated 2021 May 7]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK570442/>.
- Chatterjee, P., Nagi, N., Agarwal, A., Das, B., Banerjee, S., Sarkar, S., Gupta, N., Gangakhedkar, R.R., 2020 Feb. The 2019 novel coronavirus disease (COVID-19) pandemic: a review of the current evidence. *Indian J. Med. Res.* 151 (2–3), 147.
- Dargahi, A., Jeddi, F., Ghobadi, H., Vosoughi, M., Karami, C., Sarailoo, M., Hadisi, A., Mokhtari, S.A., Haghghi, S.B., Sadeghi, H., Alighadri, M., 2021 May 1. Evaluation of masks' internal and external surfaces used by health care workers and patients in coronavirus-2 (SARS-CoV-2) wards. *Environ. Res.* 196, 110948.
- De Angelis, E., Renzetti, S., Volta, M., Donato, F., Calza, S., Placidi, D., Lucchini, R.G., Rota, M., 2021 Apr 1. COVID-19 incidence and mortality in Lombardy, Italy: an ecological study on the role of air pollution, meteorological factors, demographic and socioeconomic variables. *Environ. Res.* 195, 110777.
- Di Gilio, A., Palmisani, J., Pulimeno, M., Cerino, F., Cacace, M., Miani, A., de Gennaro, G., 2021 Nov. CO(2) concentration monitoring inside educational buildings as a strategic tool to reduce the risk of Sars-CoV-2 airborne transmission. *Environ. Res.* 202, 111560 doi: 10.1016/j.envres.2021.111560 Epub 2021 Jul 3.

- Domingo, J.L., Rovira, J., 2020 May 11. Effects of air pollutants on the transmission and severity of respiratory viral infections. *Environ. Res.* 109650.
- EU Parliament, 2021. **Pollution and the Spread of COVID-19.** EPRS_STU, p. 697192. [http://www.europarl.europa.eu/RegData/etudes/STUD/2021/697192/EPRS_STU\(2021\)697192_EN.pdf](http://www.europarl.europa.eu/RegData/etudes/STUD/2021/697192/EPRS_STU(2021)697192_EN.pdf).
- Grimalt, J.O., Vílchez, H., Fraile-Ribot, P.A., Marco, E., Campins, A., Orfila, J., Van Drooge, B.L., Fanjul, F., 2022 Mar 1. Spread of SARS-CoV-2 in hospital areas. *Environ. Res.* 204, 112074.
- Gujral, H., Sinha, A., 2021 Mar 1. Association between exposure to airborne pollutants and COVID-19 in Los Angeles, United States with ensemble-based dynamic emission model. *Environ. Res.* 194, 110704.
- Ho, C.C., Hung, S.C., Ho, W.C., 2021 Aug 1. Effects of short-and long-term exposure to atmospheric pollution on COVID-19 risk and fatality: analysis of the first epidemic wave in Northern Italy. *Environ. Res.* 199, 111293.
- Jin, Y., Yang, H., Ji, W., Wu, W., Chen, S., Zhang, W., Duan, G., 2020 Apr. Virology, epidemiology, pathogenesis, and control of COVID-19. *Viruses* 12 (4), 372.
- Khan, I., Shah, D., Shah, S.S., 2021 Feb. COVID-19 pandemic and its positive impacts on environment: an updated review. *Int. J. Environ. Sci. Technol.* 18 (2), 521–530.
- Lauer, S.A., Grantz, K.H., Bi, Q., Jones, F.K., Zheng, Q., Meredith, H.R., Azman, A.S., Reich, N.G., Lessler, J., 2020 May 5. The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Ann. Intern. Med.* 172 (9), 577–582.
- Linillos-Pradillo, B., Rancan, L., Ramiro, E.D., Vara, E., Artfñano, B., Arias, J., 2021 Apr 1. Determination of SARS-CoV-2 RNA in different particulate matter size fractions of outdoor air samples in Madrid during the lockdown. *Environ. Res.* 195, 110863.
- Liu, K., Fang, Y.Y., Deng, Y., Liu, W., Wang, M.F., Ma, J.P., Xiao, W., Wang, Y.N., Zhong, M.H., Li, C.H., Li, G.C., 2020 May 5. Clinical characteristics of novel coronavirus cases in tertiary hospitals in Hubei Province. *Chin. Med. J.* 133 (9), 1025.
- Maleki, M., Anvari, E., Hopke, P.K., Noorimotlagh, Z., Mirzaee, S.A., 2021 Feb 18. An updated systematic review on the association between atmospheric particulate matter pollution and prevalence of SARS-CoV-2. *Environ. Res.* 110898.
- Marquès, M., Domingo, J.L., 2021. Contamination of inert surfaces by SARS-CoV-2: persistence, stability and infectivity. A review. *Environ. Res.* 193, 110559.
- Marquès, M., Domingo, J.L., 2022. Positive association between outdoor air pollution and the incidence and severity of COVID-19. A review of the recent scientific evidence. *Environ. Res.* 203, 111930.
- Mecenas, P., Bastos, R.T., Vallinoto, A.C., Normando, D., 2020 Sep 18. Effects of temperature and humidity on the spread of COVID-19: a systematic review. *PLoS One* 15 (9), e0238339.
- Megahed, N.A., Ghoneim, E.M., 2021 Feb 1. Indoor Air Quality: rethinking rules of building design strategies in post-pandemic architecture. *Environ. Res.* 193, 110471.
- Mele, M., Magazzino, C., Schneider, N., Strezov, V., 2021 Mar 1. NO2 levels as a contributing factor to COVID-19 deaths: the first empirical estimate of threshold values. *Environ. Res.* 194, 110663.
- Morawska and Cao, Morawska, L., Johnson, G.R., Ristovski, Z.D., et al., 2020. Size distribution and sites of origin of droplets expelled from the human respiratory tract during expiratory activities. *Aerosol. Sci.* 40 (3), 256–269, 2009.
- Onakpoya, I.J., Heneghan, C.J., Spencer, E.A., Brassey, J., Plüddemann, A., Evans, D.H., Conly, J.M., Jefferson, T., 2021 Mar 24. SARS-CoV-2 and the role of fomite transmission: a systematic review. *F1000Res* 10, 233.
- Paules, C.I., Marston, H.D., Fauci, A.S., 2020 Feb 25. Coronavirus infections—more than just the common cold. *JAMA* 323 (8), 707–708.
- Robotto, A., Quaglino, P., Lembo, D., Morello, M., Brizio, E., Bardi, L., Civra, A., 2021. SARS-CoV-2 and indoor/outdoor air samples: a methodological approach to have consistent and comparable results. *Environ. Res.* 195, 110847.
- Sangkham, S., Thongtip, S., Vongruang, P., 2021 Jun 1. Influence of air pollution and meteorological factors on the spread of COVID-19 in the Bangkok Metropolitan Region and air quality during the outbreak. *Environ. Res.* 197, 111104.
- Setti, L., Passarini, F., De Gennaro, G., Barbieri, P., Licen, S., Perrone, M.G., Piazzalunga, A., Borelli, M., Palmisani, J., Di Gilio, A., Rizzo, E., 2020 Sep 1. Potential role of particulate matter in the spreading of COVID-19 in Northern Italy: first observational study based on initial epidemic diffusion. *BMJ Open* 10 (9), e039338.
- Setti, L., Passarini, F., De Gennaro, G., Barbieri, P., Perrone, M.G., Borelli, M., Palmisani, J., Di Gilio, A., Torboli, V., Fontana, F., Clemente, L., 2020 Sep 1. SARS-Cov-2RNA found on particulate matter of Bergamo in Northern Italy: first evidence. *Environ. Res.* 188, 109754.
- Wu, X., Nethery, R.C., Sabath, M.B., Braun, D., Dominici, F., 2020 Nov 1. Air pollution and COVID-19 mortality in the United States: strengths and limitations of an ecological regression analysis. *Sci. Adv.* 6 (45), eabd4049.
- Wu, X., Braun, D., Schwartz, J., Kioumourtzoglou, M.A., Dominici, F., 2020 Jul 1. Evaluating the impact of long-term exposure to fine particulate matter on mortality among the elderly. *Sci. Adv.* 6 (29), eaba5692.
- Zhou, X., Josey, K., Kamareddine, L., Caine, M.C., Liu, T., Mickley, L.J., Cooper, M., Dominici, F., 2021 Aug 1. Excess of COVID-19 cases and deaths due to fine particulate matter exposure during the 2020 wildfires in the United States. *Sci. Adv.* 7 (33), eabi8789.
- Zhu, C., Maharajan, K., Liu, K., Zhang, Y., 2021 May 5. Role of atmospheric particulate matter exposure in COVID-19 and other health risks in human: a review. *Environ. Res.* 111–281.
- Zupin, L., Licen, S., Milani, M., Clemente, L., Martello, L., Semeraro, S., Fontana, F., Ruscio, M., Miani, A., Crovella, S., Barbieri, P., 2021 Oct 24. Evaluation of residual infectivity after SARS-CoV-2 aerosol transmission in a controlled laboratory setting. *Int. J. Environ. Res. Publ. Health* 18 (21), 11172.