

Macrophage MerTK promotes profibrogenic cross-talk with hepatic stellate cells via soluble mediators


Authors

Mirella Pastore, Alessandra Caligiuri, Chiara Raggi, Nadia Navari, Benedetta Piombanti, Giovanni Di Maira, Elisabetta Roviola, Marie-Pierre Piccinni, Letizia Lombardelli, Federica Logiodice, Krista Rombouts, Salvatore Petta, Fabio Marra

Correspondence

fabio.marra@unifi.it (F. Marra).

Graphical abstract



Highlights

- MerTK, a member of the TAM family of proteins, is highly expressed in MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages.
- In these macrophages, activation of MerTK induces phosphorylation of Akt, STAT3, ERK1/2, p38 and increased expression of VEGF-A.
- MerTK activation in macrophages modulates the secretome to promote the profibrogenic phenotype of human HSCs.
- Profibrogenic effects of macrophages expressing high levels of MerTK were blocked by knockdown or inhibition of MerTK.

Lay summary

Fibrosis represents the process of scarring occurring in patients with chronic liver diseases. This process depends on production of scar tissue components by a specific cell type, named hepatic stellate cells, and is regulated by interaction with other cells. Herein, we show that activation of MerTK, a receptor present in a population of macrophages, causes the production of factors that act on hepatic stellate cells, increasing their ability to produce scar tissue.

Macrophage MerTK promotes profibrogenic cross-talk with hepatic stellate cells via soluble mediators



Mirella Pastore,¹ Alessandra Caligiuri,¹ Chiara Raggi,¹ Nadia Navari,¹ Benedetta Piombanti,¹ Giovanni Di Maira,¹ Elisabetta Rovida,² Marie-Pierre Piccinni,³ Letizia Lombardelli,³ Federica Logiodice,³ Krista Rombouts,⁴ Salvatore Petta,⁵ Fabio Marra^{1,*}

¹Department of Experimental and Clinical Medicine, University of Florence, Florence Italy; ²Department of Experimental and Clinical Biomedical Sciences "Mario Serio", University of Florence, Florence Italy; ³Department of Experimental and Clinical Medicine and Center of Excellence for Research, Transfer and High Education DENOTHE of the University of Florence, Florence Italy; ⁴Regenerative Medicine and Fibrosis Group, Institute for Liver and Digestive Health, University College London, Royal Free Hospital, United Kingdom; ⁵Section of Gastroenterology and Hepatology, PROMISE, University of Palermo, Palermo Italy

JHEP Reports 2022. <https://doi.org/10.1016/j.jhepr.2022.100444>

Background & Aims: Activation of Kupffer cells and recruitment of monocytes are key events in fibrogenesis. These cells release soluble mediators which induce the activation of hepatic stellate cells (HSCs), the main fibrogenic cell type within the liver. Mer tyrosine kinase (MerTK) signaling regulates multiple processes in macrophages and has been implicated in the pathogenesis of non-alcoholic steatohepatitis-related fibrosis. In this study, we explored if MerTK activation in macrophages influences the profibrogenic phenotype of HSCs.

Methods: Macrophages were derived from THP-1 cells or differentiated from peripheral blood monocytes towards MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages. The role of MerTK was assessed by pharmacologic and genetic inhibition. HSC migration was determined in Boyden chambers, viability was measured by the MTT assay, and proliferation was evaluated by the BrdU incorporation assay.

Results: Gas-6 induced MerTK phosphorylation and Akt activation in macrophages, and these effects were inhibited by UNC569. During polarization, MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages exhibited activation of STAT3, ERK1/2, p38 and increased expression of VEGF-A. Activation of MerTK in THP-1 macrophages induced a secretome which promoted a significant increase in migration, proliferation, viability and expression of profibrogenic factors in HSCs. Similarly, conditioned medium from MerTK⁺ macrophages induced a significant increase in cell migration, proliferation, STAT3 and p38 phosphorylation and upregulation of IL-8 expression in HSCs. Moreover, conditioned medium from Gas-6-stimulated Kupffer cells induced a significant increase in HSC proliferation. These effects were specifically related to MerTK expression and activity in macrophages, as indicated by pharmacologic inhibition and knockdown experiments.

Conclusions: MerTK activation in macrophages modifies the secretome to promote profibrogenic features in HSCs, implicating this receptor in the pathogenesis of hepatic fibrosis.

Lay summary: Fibrosis represents the process of scarring occurring in patients with chronic liver diseases. This process depends on production of scar tissue components by a specific cell type, named hepatic stellate cells, and is regulated by interaction with other cells. Herein, we show that activation of MerTK, a receptor present in a population of macrophages, causes the production of factors that act on hepatic stellate cells, increasing their ability to produce scar tissue.

© 2022 The Author(s). Published by Elsevier B.V. on behalf of European Association for the Study of the Liver (EASL). This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Liver fibrosis is a wound-healing response that occurs in the context of chronic liver injury, and is characterized by excessive extracellular matrix (ECM) accumulation and fibrous scar formation, leading to liver dysfunction.¹ Hepatic stellate cells (HSCs) are the main cellular effectors of hepatic fibrogenesis. Through a

process termed 'activation', HSCs acquire a myofibroblast-like phenotype, characterized by increased proliferation, contractility, chemotaxis and ECM production.² The development of fibrosis is the result of a complex cross-talk involving diverse cell types, either resident in the liver, or infiltrating the tissue as a response to injury. Hepatic macrophages comprise Kupffer cells (KCs) and infiltrating, bone marrow-derived monocytes/macrophages.³ Various populations of hepatic macrophages have been identified, expressing specific markers and exhibiting specialized functions.⁴ Environmental stimuli define macrophage polarization into classic M1, and alternative M2, which can be further differentiated into diverse subtypes, each induced by different molecules and eliciting different signals.⁵ During the early stages

Keywords: Gas-6; THP-1; M2c-like macrophages; NASH; liver fibrosis.

Received 1 February 2021; received in revised form 29 December 2021; accepted 4 January 2022; available online 1 February 2022

* Corresponding author. Address: Department of Experimental and Clinical Medicine, University of Florence, Largo Brambilla 3, I50134 Florence, Italy; Tel.: +39055 7945425, fax: +39055 7947104.

E-mail address: fabio.marra@unifi.it (F. Marra).






Fig. 1. MerTK inhibition or silencing affect fibrogenic pathways in THP-1 macrophages. (A) Total RNA was isolated from serum-deprived THP-1 cells in unstimulated conditions (Control) (THP-1 monocytes) or after treatment with PMA (10 ng/ml) (THP-1 macrophages). *MerTK* expression was evaluated by qRT-PCR and normalized to β -actin. (B) 30 μ g of cell lysate were analyzed by western blotting, using antibodies against MerTK; β -Actin was used as a control for equal loading. PMA-treated serum-deprived THP-1 macrophages for 6 days were used for following experiments. (C) Serum-deprived THP-1 macrophages were exposed to Gas-6 for the indicated time points. 30 μ g of cell lysates were analyzed by western blotting with the indicated antibodies. (D) Serum-deprived THP-1 macrophages were exposed to Gas-6 (200 ng/ml) or to PMA (100 ng/ml) for the indicated time points. 30 μ g of cell lysates were analyzed by western blotting with the indicated antibodies. (E) Serum-deprived THP-1 macrophages were exposed to UNC569 or DMSO before stimulation with Gas-6 for 15 min. 30 μ g of cell lysates were analyzed by western blotting, using the indicated antibodies. (F-H) THP-1 macrophages were transfected with MERTK-specific siRNA (light purple columns) or with NT siRNA (purple columns), (F) Total RNA was isolated 48 h and 72 h after transfection and *MERTK* expression was determined by qRT-PCR and normalized to β -actin. (G-H) 72 h after transfection, THP-1 macrophages were exposed to Gas-6 for 8 h. Total RNA was isolated and the expression of (G) *VEGF-A* and (H) *TGF β 1* was evaluated by qRT-PCR and normalized to β -actin. Densitometries of p-MerTK/MerTK and p-Akt/Akt expression (n = 3) were shown in the graphs. Data are mean \pm SEM. Statistical significance was assessed by Student's *t* test. **p* <0.05 vs. THP-1 monocytes; **p* <0.05 vs. Control; ***p* <0.05 vs. THP-1 macrophage stimulated with Gas-6 without inhibition for MerTK; Δ *p* <0.05 vs. NT siRNA; $\dagger\dagger$ *p* <0.05 vs. NT siRNA unstimulated; \ddagger *p* <0.05 vs. NT siRNA stimulated with Gas-6. NT, non-targeting; PMA, phorbol 12-myristate 13-acetate; qRT-PCR, quantitative real-time PCR; siRNA, small-interfering RNA.

of liver injury, bone marrow-derived monocytes are recruited to the liver and differentiate into M1 macrophages.⁶ M1 macrophages can rapidly switch to an M2 phenotype and mediate tissue repair in case of cessation of liver injury.⁷ However, when the lesion is persistent, M2 macrophages mediate an important pro-fibrotic function.⁸

Myeloid-epithelial-reproductive tyrosine kinase (MerTK) is a member of the Tyro-3, Axl, and Mer (TAM) receptor tyrosine kinase family⁹ and is expressed at high levels in macrophages displaying the M2c-like phenotype, characterized by a marked involvement in tissue repair, matrix remodeling and promotion of angiogenesis through secretion of pro-angiogenic factors.^{10,11}

Growth arrest-specific gene 6 (Gas-6), a glycoprotein considered a new potential fibrosis biomarker,¹²⁻¹⁴ is one of the best characterized ligands for MerTK. A single nucleotide polymorphism (SNP) associated with lower MerTK expression has been linked to the reduced progression of liver fibrosis in patients with non-alcoholic fatty liver disease (NAFLD) or HCV.¹⁵⁻¹⁷ In hepatic tissue from patients with NAFLD, MerTK was found to be mostly expressed in macrophages and partially in HSCs, and was loosely aggregated within inflammatory foci.¹⁶ Moreover, activation of MerTK in HSCs resulted in the induction of profibrogenic actions.

Cross-talk between HSCs and macrophages is crucial for fibrosis development,¹⁸ but the biological role played by




Fig. 2. Migration, proliferation, viability, and activated signaling pathways in HSCs exposed to CM from MerTK-inhibited THP-1 macrophages. (A-C-E) Serum-starved HSCs were exposed to SF medium (Control, white columns) or to CM from THP-1 macrophages left untreated (DMSO, purple columns), treated with UNC569 (light purple columns) or stimulated with Gas-6 in the presence (light purple columns) or absence (purple columns) of UNC569. (B-D-F) Serum-starved HSCs were exposed to SF medium (Control, white columns) or to CM of THP-1 macrophages transfected with NT siRNA (purple columns) or MERTK-siRNA (light purple columns) and left untreated or treated with Gas-6. (A-B) HSC migration was measured in modified Boyden chambers. (C-D) HSC proliferation was assayed by BrdU Cell Proliferation ELISA Kit. (E-F) HSC viability was measured by MTT assay. (G) Serum-starved HSCs were exposed to CM from THP-1 macrophages unstimulated or stimulated Gas-6 in the presence or absence of UNC569 or (H) to CM from THP-1 macrophages transfected with NT siRNA or MERTK-siRNA stimulated with or without Gas-6 for 24 h. 30 μg of total cell lysates were subjected to immunoblot analysis to detect different proteins or phosphoproteins. Densitometries of p-STAT3/STAT3, p-P38/P38 and IL8/β-Actin expression (n = 3) were shown in the graphs. Data are mean ± SEM. Statistical significance was assessed by Student's *t* test. **p* <0.05 vs. SF medium; ***p* <0.05 vs. CM of untreated THP-1 macrophages; §*p* <0.05 vs. CM of THP-1 macrophage stimulated with Gas-6 without inhibition for MerTK; †*p* <0.05 vs. SF medium; ††*p* <0.05 vs. CM of NT siRNA unstimulated THP-1 macrophages; ¥*p* <0.05 vs. CM of NT siRNA THP-1 macrophage stimulated with Gas-6. CM, conditioned media; HSC(s), hepatic stellate cells; NT, non-targeting; SF, serum free; siRNA, small-interfering RNA.

macrophage-derived MerTK is still unclear. Herein, we show that macrophages expressing high levels of MerTK promote a profibrogenic phenotype in HSCs (via paracrine signals), inducing a significant increase in cell migration, viability, proliferation and the expression of profibrogenic factors.

Materials and methods

Cell culture and treatments

The THP-1 human monocytic leukemic cell line, kindly provided by Dr. Elisabetta Rovida (University of Florence, Italy), was

differentiated into macrophages by 48 h or 144 h of incubation with phorbol 12-myristate 13-acetate (PMA) (10 ng/ml) at 37°C in 5% CO₂, as described by Kurihara *et al.*^{19,20} Human monocytes were isolated from buffy coats of healthy blood donors who provided written informed consent. A two-step procedure was employed, as described by De Almeida *et al.*,²¹ using Lympholyte-H (Cedarlane, Burlington, CA, USA) followed by slight hyperosmolar Percoll gradient (GE Healthcare Life Sciences, RD, St. Louis, MO, USA). Primary human HSCs were isolated from normal liver tissue unsuitable for transplantation as previously described in detail.²² All patients provided written, informed




Fig. 3. HSC expression of proinflammatory and profibrogenic genes in response to CM from MerTK-inhibited THP-1 macrophages. Serum-starved HSCs were exposed to SF medium (Control, white columns) or to CM from THP-1 macrophages left untreated (DMSO, purple columns), treated with UNC569 (light purple columns) or stimulated with Gas-6 in the presence (light purple columns) or absence (purple columns) of UNC569 for 48 h. (A–H) Total RNA was isolated and mRNA expression of different genes was evaluated by qRT-PCR and normalized for the expression of β -actin. Data are mean \pm SEM. Statistical significance was assessed by Student's *t* test. **p* < 0.05 vs. SF medium; ***p* < 0.05 vs. CM of untreated THP-1 macrophages; § *p* < 0.05 vs. CM of THP-1 macrophage stimulated with Gas-6 without inhibition for MerTK. CM, conditioned media; HSC(s), hepatic stellate cell(s); qRT-PCR, quantitative real-time PCR; SF, serum free.

consent for cell isolation, and the study was approved by the local institutional review board. KCs were purchased from Thermo Scientific (Waltham, MA USA).

Studies with conditioned medium

THP-1 monocytes were differentiated into macrophages by 144 h of incubation with 10 ng/ml PMA at 37°C in 5% CO₂, hereafter referred to as “THP-1 macrophages”. siRNA-transfected THP-1 macrophages were incubated for 24 h in serum-free medium and then stimulated with Gas-6 (200 ng/ml) for 8 h. When the MerTK inhibitor UNC569²³ was used, both THP-1-derived and circulating monocyte-derived MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages were incubated in serum-free medium for 24 h and then exposed to 5 μM UNC569 or to DMSO for 90 min before stimulation with Gas-6 (200 ng/ml) for 8 h. All macrophages were then rinsed 3 times with PBS and further incubated overnight in serum-free medium. Conditioned medium (CM) was collected, centrifuged at 2,000 x g for 10 min and stored at -80°C for further use. Although the traditional classification (M1, M2a, M2b, M2c), which is based on induction of *in vitro* polarization, does not accurately describe the phenotypic heterogeneity of *in vivo* macrophages, for ease of reading, the CM of MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages is defined as ‘M2c CM’.

Statistical analysis

Data in bar graphs represent mean \pm SD from at least 3 independent experiments. Luminograms are representative of at least 3 experiments with similar results.

Statistical analysis was performed using Student's *t* test.

For further details regarding the materials and methods used, please refer to the [CTAT table and supplementary information](#).

Results

MerTK expression and activation in THP-1 macrophages

To elucidate the involvement of MerTK in the interaction between macrophages and HSCs, we initially studied macrophages derived from THP-1 cells, an immortalized monocytic line which differentiates into macrophages when exposed to PMA.¹⁹ In PMA-differentiated THP-1 macrophages, the expression of MerTK, but not of Axl and Tyro3, was substantially increased (data not shown).²⁰ Accordingly, *MERTK* mRNA levels were several thousand-fold higher in THP-1 macrophages after 6 days of treatment with PMA (Fig. 1A). This increase in MerTK expression was confirmed by western blot analysis (Fig. 1B). Consistent with previous findings,^{24,25} exposure of THP-1 macrophages to Gas-6, a ligand of MerTK, increased MerTK




Fig. 4. HSC expression of proinflammatory and profibrogenic markers in response to CM from MERTK-silenced THP-1 macrophages. Serum-starved HSCs were exposed to SF medium (Control, white columns) or to CM from THP-1 macrophages transfected with NT siRNA (purple columns) or MERTK-siRNA (light purple columns) and left untreated or treated with Gas-6 for 48 h. (A-H) Total RNA was isolated and mRNA expression of different genes was evaluated by qRT-PCR and normalized to β -actin. Data are mean \pm SEM. Statistical significance was assessed by Student's *t* test. $\dagger p < 0.05$ vs. SF medium; $\dagger\dagger p < 0.05$ vs. CM of NT siRNA unstimulated THP-1 macrophages; $\forall p < 0.05$ vs. CM of NT siRNA THP-1 macrophage stimulated with Gas-6. CM, conditioned media; HSCs, hepatic stellate cells; NT, non-targeting; qRT-PCR, quantitative real-time PCR; SF, serum free; siRNA, small-interfering RNA.

phosphorylation and downstream activation of Akt, in a time-dependent manner (Fig. 1C). In contrast, exposure to PMA did not result in any increase in MerTK phosphorylation (Fig. 1D).

Pre-incubation with UNC569, an inhibitor specific to MerTK, reduced MerTK phosphorylation and downstream Akt activation²³ (Fig. 1E). To rule out that the effects of UNC569 were due to non-specific inhibition, THP-1 macrophages were transfected with MERTK-specific small-interfering RNA (siRNA) and knock-down efficiency was evaluated by real-time PCR (Fig. 1F). Exposure of THP-1 macrophages to Gas-6 induced a significant increase in gene expression of vascular endothelial growth factor-A (VEGF-A) and transforming growth factor- β 1 (TGF β 1) and these effects were completely abolished by MerTK silencing (Fig. 1G and H). Taken together, these data indicate that, in macrophages, MerTK regulates the expression of factors implicated in liver repair, in a MerTK-dependent fashion.

In THP-1 macrophages, MerTK activation induces the release of soluble factors which modulate the biology of HSCs

To examine the involvement of MerTK signaling in the cross-talk between macrophages and HSCs, we studied the effects of CM obtained from THP-1 macrophages on the biological features of HSCs. HSCs exposed to THP-1-CM showed a significant increase in migration, that was more evident when macrophages were treated with Gas-6 (Fig. 2A). To evaluate whether these effects

were mediated by MerTK activation in macrophages, we used CM from THP-1 macrophages with pharmacologic or genetic inhibition of MerTK. Using both these strategies, increased migration of HSCs in response to CM from Gas-6-stimulated THP-1 macrophages was significantly reduced (Fig. 2A,B). Similar results were obtained when we explored the effects of THP-1-CM on HSC proliferation (Fig. 2C,D) and viability (Fig. 2E,F). In HSCs exposed to CM from Gas-6-stimulated THP-1 macrophages, increased activation of STAT3 and p38 was observed, together with an increase in IL-8 protein expression. These effects were inhibited by pre-treatment of THP-1 macrophages with UNC569 or after MerTK silencing (Fig. 2G-H). To better elucidate the role of THP-1 macrophage MerTK in modulating the profibrogenic properties of HSCs, we analyzed the expression of genes upregulated upon HSC activation. In HSCs treated with CM from Gas-6-stimulated THP-1 macrophages, mRNA expression of tissue inhibitor of metalloproteinase 1 (TIMP1), TGF β 1, interleukin 8 (IL8), VEGF-A, actin alpha 2, smooth muscle (ACTA2), matrix metalloproteinase 2 (MMP2), collagen type I alpha 1 chain (COL1A1) and matrix metalloproteinase 9 (MMP9) was significantly increased compared to that in HSCs treated with CM from unstimulated macrophages (Fig. 3). These effects were blunted by macrophage pre-treatment with UNC569, indicating that the actions of Gas-6 are mediated by MerTK (Fig. 3). Next, we stimulated HSCs with CM from THP-1




Fig. 5. Expression and activation of MerTK and downstream signaling in monocytes differentiated towards M2c-like macrophages. (A) Morphologic analysis of culture of naïve (M0) macrophages derived from peripheral blood monocytes and of peripheral blood monocytes that were treated with M-CSF (50 ng/ml) for 5 days (M2-like macrophages), followed by cultivation with M-CSF (50 ng/ml) and IL-10 (50 ng/ml) for additional 3 days (M2c-like macrophages). (B-C) Cells were stained for CD206, CD163, CD209 or MerTK and analyzed by flow cytometry. (B) Representative Dot plots depicting the gating strategy for analyzing MerTK expression in (CD206⁺ CD163⁺ CD209⁻) macrophages. (C) Histograms show MFI. (D) CD206 CD163 and MERTK mRNA expression was analyzed by qRT-PCR and normalized to β -actin. (E-F) Macrophages were deprived of serum and lysed. (G-H) Serum-deprived M2c macrophages were exposed to UNC569 or DMSO for 90 min before stimulation with Gas-6 for 15 min. 30 μ g of cell lysates were analyzed by western blotting to detect different proteins or phosphoproteins. Densitometries of MerTK/ β -Actin, p-MerTK/MerTK, p-STAT3/STAT3, p-STAT5/STAT5, p-Akt/Akt p-ERK1/2/ERK1/2, p-P38/P38 and VEGF-A/ β -Actin expression (n = 3) were shown in the graphs. Data are mean \pm SEM. Statistical significance was assessed by Student's *t* test. **p* <0.05 vs. M0 macrophages; *p* <0.05 vs. M2 macrophages; ***p* <0.05 vs. Control M2c; §*p* <0.05 vs. M2c macrophage stimulated with Gas-6 without inhibition for MerTK. M-CSF, macrophage colony-stimulating factor; MFI, mean fluorescence intensity; qRT-PCR, quantitative real-time PCR.

macrophages after MERTK depletion, with or without exposure to Gas-6. Upon MerTK knockdown, CM from macrophages showed a reduced capability to induce expression of fibrogenic genes in HSCs (Fig. 4). Taken together, these data indicate that MerTK expressed by differentiated THP-1 cells mediates profibrogenic actions in HSCs via soluble mediators.

Profibrogenic effects of MerTK activation in MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages via cross-talk with HSCs

Macrophage polarization towards a M2c-like phenotype is associated with a marked increase in MerTK expression.¹¹ Because THP-1 cells only partially recapitulate the biologic effects of tissue-infiltrating macrophages, we next investigated

whether the effects of MerTK activation could be reproduced in a population of monocyte-derived MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages (Fig. 5A). Exposure to a culture medium rich in macrophage colony-stimulating factor (M-CSF) and IL-10 induced high expression of CD206 (a marker for alternatively activated macrophages²⁶), CD163 (a specific marker for M2c cells²⁷), and low expression of CD209 (a specific marker for M2a phenotype¹¹) (Fig. 5C). Flow cytometry analyses confirmed a high expression of MerTK in the CD206⁺/CD163⁺/CD209⁻ phenotype (Fig. 5B,C). Furthermore, MERTK expression was gradually increased along MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophage differentiation. CD206 expression significantly increased on the fifth day of exposure to M-CSF, while a




Fig. 7. HSC expression of proinflammatory and profibrogenic genes in response to CM from MerTK-inhibited M2c macrophages. Serum-starved HSCs were exposed to SF medium (Control, white columns) or to CM of M2c macrophages left untreated (DMSO, purple columns), treated with UNC569 (light purple columns) or stimulated with Gas-6 in the presence (light purple columns) or absence (purple columns) of UNC569 for 48 h. (A-H) Total RNA was isolated and levels of different genes were determined by qRT-PCR and normalized to β -actin. Data are mean \pm SEM. Statistical significance was assessed by Student's *t* test. **p* <0.05 vs. SF medium; ***p* <0.05 vs. CM of untreated M2c macrophages; § *p* <0.05 vs. CM of M2c macrophage stimulated with Gas-6 without inhibition for MerTK. CM, conditioned media; HSC(s), hepatic stellate cell(s); qRT-PCR, quantitative real-time PCR; SF, serum-free.

induction of a profibrogenic phenotype, we examined the activation of different signaling pathways. In HSCs exposed to CM from Gas-6-stimulated MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages, we observed increased activation of STAT3 and p38, together with an increase in IL-8 expression. These effects were inhibited by pre-treatment of MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages with UNC569 (Fig. 6D).

We next evaluated the regulation of profibrogenic genes in HSCs treated with CM from Gas-6-stimulated MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages. Expression of *TIMP1*, *TGFβ1*, *IL8*, *VEGF-A*, *ACTA2*, *MMP2*, *COL1A1* and *MMP9* was increased compared to that in HSCs treated with CM from unstimulated macrophages. These actions were significantly reduced when macrophages were pre-treated with UNC569, indicating that MerTK activation was required (Fig. 7). Taken together, these data indicate that activation of MerTK in alternatively activated macrophages results in the upregulation of profibrogenic signals in HSCs.

Gas-6 expression in differentiated macrophages and in KCs

Gas-6 was produced by THP-1, M2c-like macrophages and KCs, with higher levels in MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages and in KCs (Fig. S1A-D). However, pharmacologic inhibition or *MERTK* gene silencing did not result in significant changes in the levels of Gas-6 (Fig. S1A-D). Gas-6 mRNA levels were significantly increased in THP-1 macrophages after 2 and 6

days of treatment with PMA (Fig. S1E), as well as during macrophage polarization towards a M2c-like phenotype (Fig. 8A). Of note, Gas-6 expression was markedly elevated in KCs compared to naïve (M0) or M2c-like-polarized macrophages (Fig. 8A). KCs expressed transcripts for *MERTK* at levels comparable to those measured in differentiated macrophages (data not shown).

MerTK activation in KCs induces HSC proliferation

To investigate whether MerTK expression in KCs contributes to the fibrogenic cross-talk with HSCs, we obtained CM from cultured human KCs in different experimental conditions.

HSCs exposed to KC-CM showed a significant increase in cell proliferation (Fig. 8B). This effect was more evident when KCs were treated with Gas-6 and significantly reduced upon pre-treatment of KCs with UNC569.

Discussion

Hepatic fibrogenesis is the result of a complex cross-talk among different resident and infiltrating cell types.²⁹ Accumulating evidence points to the pivotal role of macrophages in the generation of profibrogenic signals as well as in the process of fibrosis resolution.³⁰ Evidence for different macrophage phenotypes adds complexity to the system³¹ and the relative role of differentially polarized cells has not been completely unraveled. In




Fig. 8. Expression of Gas-6 in KCs and effect of MerTK activation in KCs on HSC proliferation. Total RNA was isolated from serum-deprived peripheral monocyte-derived macrophages and from KCs. (A) *Gas-6* mRNA expression was analyzed by qRT-PCR and normalized to β -actin. (B) Serum-starved HSCs were exposed to SF medium (Control, white column) or to CM from KCs left untreated (DMSO, purple columns), treated with UNC569 (light purple column) or stimulated with Gas-6 in the presence (light purple column) or absence (purple column) of UNC569. HSC proliferation was assayed by BrdU Cell Proliferation ELISA Kit. Data are mean \pm SEM. Statistical significance was assessed by Student's *t* test. #*p* <0.05 vs. M0 macrophages. **p* <0.05 vs. SF medium; ***p* <0.05 vs. CM of untreated KCs; §*p* <0.05 vs. CM of KCs stimulated with Gas-6 without inhibition for MerTK. CM, conditioned media; HSC(s), hepatic stellate cell(s); KCs, Kupffer cells; qRT-PCR, quantitative real-time PCR; SF, serum-free.

this study, we provide evidence that activation of the tyrosine kinase receptor MerTK on the surface of M2c-like-polarized macrophages modifies the secretome to induce profibrogenic actions in HSCs.

MerTK has been under intense investigation over the last few years for its involvement in immune regulatory activities.³² A first indication of a possible link between MerTK and hepatic fibrosis has been provided by a genome-wide association study in patients with hepatitis C, where a SNP in MerTK was more prevalent in patients with advanced fibrosis.¹⁵ We recently reported a higher hepatic expression of MERTK in patients with NAFLD and in murine models of fibrogenesis.¹⁶ Moreover, 2 intronic SNPs in *MERTK* rs6726639 A³³ and rs4374383 A¹⁵⁻¹⁷ were associated with decreased hepatic MerTK expression and a reduced risk of liver fibrosis. We also previously provided evidence for a direct role of MerTK in the regulation of the profibrogenic actions of HSCs,¹⁶ because exposure of cultured human HSCs to Gas-6 resulted in induction of a profibrogenic phenotype. Moreover circulating levels of Gas-6 were increased in patients with cirrhosis (concentrations around 60-80 ng/ml^{13,14}) and in animal models of chronic liver damage (concentrations around 40 ng/ml¹⁴). Taken together, these data suggested a relevant role of MerTK activation in the fibrogenic process,

especially in the context of non-alcoholic steatohepatitis (NASH).¹⁶

Immunohistochemical analysis of livers with NASH fibrosis demonstrated that macrophages are major contributors to the expression of MerTK.¹⁶ Macrophages localize in close proximity to activated HSCs, and have been shown to play a key role in the progression of liver injury from inflammation to fibrosis, and eventually cirrhosis and hepatocellular carcinoma.³⁴ Based on this background, we differentiated cultured macrophages to an M2c-like phenotype, which is known to express MerTK at high levels, and evaluated the contribution of activation of this kinase in the regulation of the HSC phenotype. Specifically, we employed CM obtained from macrophage cultures, which is known to contain soluble factors, exosomes and microparticles that may modulate the phenotype of HSCs.³⁵ We first used THP-1 macrophages, which express high levels of *MerTK*, *Gas-6*, *VEGF-A* and *TGF β 1*, factors characterizing a wound-healing macrophagic phenotype which promotes cellular proliferation and blood vessel development,³⁶⁻³⁸ as a cellular model.^{20,38} Our findings indicate that Gas-6 induces MerTK phosphorylation and activation of Akt, a well-recognized downstream effector of MerTK,²³ confirming the functionality of this pathway.

More important, CM from Gas-6-treated THP-1 cells induced the increase in migration, proliferation and viability of HSCs, and these actions were blocked by exposure of macrophages to UNC569, a small-molecule inhibitor specific for MerTK.²³ The role of MerTK-expressing macrophages in a profibrogenic cross-talk with HSCs was further supported by experiments using siRNAs to downregulate MerTK expression, which provided similar results to those obtained with the chemical inhibitor. Interestingly, exposure of HSCs to medium conditioned by MerTK-expressing THP-1 cells also resulted in a significant upregulation of the expression of several genes known to be involved in fibrogenesis, such as *TIMP1*, *TGF β 1*, *ACTA2*, *MMP2*, *COL1A1* and *MMP9*. In addition, upregulated expression of *VEGF-A*, which participates in the regulation of angiogenesis during tissue repair,³⁹ and of *IL8*, a proinflammatory chemokine,⁴⁰ at the gene and protein levels, was observed. Also in this case, stimulation of macrophages with Gas-6 increased the effects, while pharmacologic inhibition or genetic knockdown of MerTK blocked these actions. Interestingly, Gas-6 also upregulated the expression of *TGF β 1* and *VEGF-A* in macrophages, indicating that paracrine and autocrine loops involving these factors are regulated by MerTK.

To provide additional data supporting our findings, we used MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages derived from peripheral blood monocytes. This subpopulation expresses high levels of MerTK and, consistently, we observed a marked increase in activation of signaling pathways such as STAT3, ERK1/2, p38,⁴¹ and in the expression of VEGF-A,⁴² all downstream targets of MerTK. Of note, no differences between M2-like and MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages were observed for ERK1/2 and p38 activation, possibly reflecting the fact that several agonists converge on these pathways. Also, in these differentiated and well-characterized cells, Gas-6 induced a substantial increase in phosphorylation of MerTK and Akt, and downstream targets of MerTK such as STAT5, STAT3, ERK1/2 and p38, which was reverted by pre-treatment with UNC569. Using these freshly prepared cells, we corroborated the findings indicating that CM from Gas-6-stimulated MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages promotes the profibrogenic actions of HSCs, inducing a

significant increase in HSC migration, cell viability, cell proliferation and expression of profibrotic and proinflammatory genes. Lastly, we found that CM from Gas-6-stimulated MerTK⁺/CD206⁺/CD163⁺/CD209⁻ macrophages induced, in HSCs, the phosphorylation of STAT3 and p38, which are fundamental in the profibrogenic phenotype of HSCs,⁴³ and increased the expression of IL8, an inflammatory chemokine that is closely associated with angiogenesis.⁴⁰ Importantly, the results obtained in THP-1 cells and in peripheral monocytes derived M2c-like macrophages could be extended to KCs, where transcripts for MerTK were detected, and Gas-6 was expressed at the gene and protein levels.

Functionally, CM from Gas-6-treated KCs increased HSC proliferation, and this effect was blocked by exposure of KCs to UNC569. Taken together, these data strongly suggest that macrophage MerTK could play an important role in the liver fibrogenic process by upregulating several functions associated with the activated phenotype of HSCs. These studies provide an additional mechanism to support previously reported data indicating that holo- or myeloid-specific MerTK targeting in NASH mice decreases liver fibrosis.³⁸

Data reported herein are in clear agreement with previous studies highlighting the reduced risk of liver fibrosis in carriers of the (rs4374383 G>A) variant of MerTK, both in hepatitis C and in NASH.^{15–17} Moreover, in a recent study, myeloid-specific targeting of MerTK reduced fibrosis development in NASH mice. Interestingly, shedding of MerTK from the surface of macrophages limited profibrogenic activities, which were dependent, at least in part, on TGFβ1-mediated signals.³⁸ These findings are

consistent with recent data reporting single-cell RNA sequence analysis of cirrhotic human livers, showing that MerTK is primarily expressed in liver macrophages.⁴⁴

These data should be put in the context of the growing interest in other members of the TAM family, such as Axl, in the pathophysiology of liver fibrosis.⁴⁵ Blocking activation of this receptor in different cell types was reported to blunt fibrogenesis in experimental models.^{13,14} Because tyrosine kinase receptors are generally considered druggable targets, different lines of evidence, including those reported in the present study, indicate the TAM family should be the topic of further investigation as a new target for pharmacologic treatment of hepatic fibrosis.^{12,13}

The identification of a role for MerTK in fibrogenesis expands the number of hepatic conditions where this receptor is implicated. In patients with acute-on-chronic liver failure, monocytes and macrophages expressing MerTK were increased, and MerTK expression was associated with a reduced ability to produce proinflammatory cytokines, resulting in suppression of the innate immune response to microbes.⁴⁶ On the other hand, in mice with acute liver failure, MerTK-deficient animals exhibited persistent liver injury and inflammation, identifying a protective role of MerTK⁺ macrophages in this setting.⁴⁷

In conclusion, MerTK expressed on macrophages induces the secretion of factors which modify the phenotype of stellate cells in a profibrogenic fashion, highlighting a novel cross-talk between intrahepatic cells. Together with other data, our findings warrant additional investigation to substantiate the role of this receptor as a therapeutic target for antifibrotic treatments.

Abbreviations

CM, conditioned medium; ECM, extracellular matrix; Gas-6, growth arrest-specific gene 6; HSC(s), hepatic stellate cells; KC(s), Kupffer cell(s); M-CSF, macrophage colony-stimulating factor; MerTK, Myeloid-epithelial-reproductive tyrosine kinase; NAFLD, non-alcoholic fatty liver disease; NASH, non-alcoholic steatohepatitis; PMA, phorbol 12-myristate 13-acetate; siRNA, small-interfering RNA; TGFβ1, transforming growth factor-β1; TIMP1, tissue inhibitor of metalloproteinase 1; VEGF-A, vascular endothelial growth factor-A.

Financial support

This work was supported by the Horizon 2020 project 'Elucidating Pathways of Steatohepatitis' (Epos) under Grant Agreement no. 634413 and the University of Florence. MP was supported in part by a "Mario Coppo" Hepatology Scholarship from the Italian Association for the Study of the Liver (AISF).

Conflict of interest

The authors declare no conflicts of interest to disclose related to this manuscript.

Please refer to the accompanying ICMJE disclosure forms for further details.

Authors' contributions

MP, FM designed the study and wrote the manuscript. MP, AC, CR, NN, BP, GD, ER, MPP, LL, FL, KB and SP provided materials, performed the experiments, collected the data, and analyzed the results. MP, AC, FM supervised the project and critically revised the manuscript.

Data availability statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jhepr.2022.100444>.

References

Author names in bold designate shared co-first authorship

- [1] Cordero-Espinoza L, Huch M. The balancing act of the liver: tissue regeneration versus fibrosis. *J Clin Invest* 2018;128:85–96.
- [2] Tsuchida T, Friedman SL. Mechanisms of hepatic stellate cell activation. *Nat Rev Gastroenterol Hepatol* 2017;14:397–411.
- [3] Guillot A, Tacke F. Liver macrophages: old dogmas and new insights. *Hepatol Commun* 2019;3:730–743.
- [4] Sica A, Invernizzi P, Mantovani A. Macrophage plasticity and polarization in liver homeostasis and pathology. *Hepatology* 2014;59:2034–2042.
- [5] **Spiller KL, Wrona EA, Romero-Torres S, Pallotta I, Graney PL, Witherel CE, et al.** Differential gene expression in human, murine, and cell line-derived macrophages upon polarization. *Exp Cell Res* 2016;347:1–13.
- [6] Schuster S, Cabrera D, Arrese M, Feldstein AE. Triggering and resolution of inflammation in NASH. *Nat Rev Gastroenterol Hepatol* 2018;15:349–364.
- [7] **Pradere JP, Kluwe J, De Minicis S, Jiao JJ, Gwak GY, Dapito DH, et al.** Hepatic macrophages but not dendritic cells contribute to liver fibrosis by promoting the survival of activated hepatic stellate cells in mice. *Hepatology* 2013;58:1461–1473.
- [8] Braga TT, Agudelo JS, Camara NO. Macrophages during the fibrotic process: M2 as friend and foe. *Front Immunol* 2015;6:602.
- [9] Pastore M, Grimaudo S, Pipitone RM, Lori G, Raggi C, Petta S, et al. Role of myeloid-epithelial-reproductive tyrosine kinase and macrophage polarization in the progression of atherosclerotic lesions associated with nonalcoholic fatty liver disease. *Front Pharmacol* 2019;10:604.
- [10] **Kim SY, Lim EJ, Yoon YS, Ahn YH, Park EM, Kim HS, et al.** Liver X receptor and STAT1 cooperate downstream of Gas6/Mer to induce anti-inflammatory arginase 2 expression in macrophages. *Sci Rep* 2016;6:29673.

- [11] Zizzo G, Hilliard BA, Monestier M, Cohen PL. Efficient clearance of early apoptotic cells by human macrophages requires M2c polarization and MerTK induction. *J Immunol* 2012;189:3508–3520.
- [12] Smirne C, Rigamonti C, De Benedittis C, Sainaghi PP, Bellan M, Burlone ME, et al. Gas6/TAM signaling components as novel biomarkers of liver fibrosis. *Dis Markers* 2019;2019:2304931.
- [13] Tutusaus A, de Gregorio E, Cucarull B, Cristóbal H, Aresté C, Graupera I, et al. A functional role of GAS6/TAM in nonalcoholic steatohepatitis progression implicates AXL as therapeutic target. *Cell Mol Gastroenterol Hepatol* 2020;9:349–368.
- [14] Bárcena C, Stefanovic M, Tutusaus A, Joannas L, Menéndez A, García-Ruiz C, et al. Gas6/Axl pathway is activated in chronic liver disease and its targeting reduces fibrosis via hepatic stellate cell inactivation. *J Hepatol* 2015;63:670–678.
- [15] Patin E, Kutalik Z, Guernon J, Bibert S, Nalpas B, Jouanguy E, et al. Genome-wide association study identifies variants associated with progression of liver fibrosis from HCV infection. *Gastroenterology* 2012;143:1244–1252.e1212.
- [16] Petta S, Valenti L, Marra F, Grimaudo S, Tripodo C, Bugianesi E, et al. MERTK rs4374383 polymorphism affects the severity of fibrosis in non-alcoholic fatty liver disease. *J Hepatol* 2016;64:682–690.
- [17] Jiménez-Sousa M, Gómez-Moreno AZ, Pineda-Tenor D, Brochado-Kith O, Sánchez-Ruano JJ, Artaza-Varasa T, et al. The myeloid-epithelial-reproductive tyrosine kinase (MERTK) rs4374383 polymorphism predicts progression of liver fibrosis in hepatitis C virus-infected patients: a longitudinal study. *J Clin Med* 2018;7.
- [18] Pakshir P, Hinz B. The big five in fibrosis: macrophages, myofibroblasts, matrix, mechanics, and miscommunication. *Matrix Biol* 2018;68–69:81–93.
- [19] Kurihara Y, Furue M. Interferon- γ enhances phorbol myristate acetate-induced cell attachment and tumor necrosis factor production via the NF- κ B pathway in THP-1 human monocytic cells. *Mol Med Rep* 2013;7:1739–1744.
- [20] Liao D, Wang X, Li M, Lin PH, Yao Q, Chen C. Human protein S inhibits the uptake of AcLDL and expression of SR-A through Mer receptor tyrosine kinase in human macrophages. *Blood* 2009;113:165–174.
- [21] de Almeida MC, Silva AC, Barral A, Barral Netto M. A simple method for human peripheral blood monocyte isolation. *Mem Inst Oswaldo Cruz* 2000;95:221–223.
- [22] Casini A, Pinzani M, Milani S, Grappone C, Galli G, Jezequel AM, et al. Regulation of extracellular matrix synthesis by transforming growth factor beta 1 in human fat-storing cells. *Gastroenterology* 1993;105:245–253.
- [23] Christoph S, Deryckere D, Schlegel J, Frazer JK, Batchelor LA, Trakhimets AY, et al. UNC569, a novel small-molecule mer inhibitor with efficacy against acute lymphoblastic leukemia in vitro and in vivo. *Mol Cancer Ther* 2013;12:2367–2377.
- [24] Cai B, Kasikara C, Doran AC, Ramakrishnan R, Birge RB, Tabas I. MerTK signaling in macrophages promotes the synthesis of inflammation resolution mediators by suppressing CaMKII activity. *Sci Signal* 2018;11.
- [25] Alciato F, Sainaghi PP, Sola D, Castello L, Avanzi GC. TNF- α , IL-6, and IL-1 expression is inhibited by GAS6 in monocytes/macrophages. *J Leukoc Biol* 2010;87:869–875.
- [26] Suzuki Y, Shirai M, Asada K, Yasui H, Karayama M, Hozumi H, et al. Macrophage mannose receptor, CD206, predict prognosis in patients with pulmonary tuberculosis. *Sci Rep* 2018;8:13129.
- [27] Ambarus CA, Krausz S, van Eijk M, Hamann J, Radstake TR, Reedquist KA, et al. Systematic validation of specific phenotypic markers for in vitro polarized human macrophages. *J Immunol Methods* 2012;375:196–206.
- [28] Zizzo G, Cohen PL. Antibody cross-linking of CD14 activates MerTK and promotes human macrophage clearance of apoptotic neutrophils: the dual role of CD14 at the crossroads between M1 and M2c polarization. *Inflammation* 2018;41:2206–2221.
- [29] Li H, Zhou Y, Wang H, Zhang M, Qiu P, Zhang R, et al. Crosstalk between liver macrophages and surrounding cells in nonalcoholic steatohepatitis. *Front Immunol* 2020;11:1169.
- [30] Tacke F, Zimmermann HW. Macrophage heterogeneity in liver injury and fibrosis. *J Hepatol* 2014;60:1090–1096.
- [31] Zigmund E, Samia-Grinberg S, Pasmanik-Chor M, Brazowski E, Shibolet O, Halpern Z, et al. Infiltrating monocyte-derived macrophages and resident kupffer cells display different ontogeny and functions in acute liver injury. *J Immunol* 2014;193:344–353.
- [32] Zhang L, DeBerge M, Wang J, Dangi A, Zhang X, Schroth S, et al. Receptor tyrosine kinase MerTK suppresses an allogenic type I IFN response to promote transplant tolerance. *Am J Transpl* 2019;19:674–685.
- [33] Cavalli M, Pan G, Nord H, Wallén Arzt E, Wallerman O, Wadelius C. Genetic prevention of hepatitis C virus-induced liver fibrosis by allele-specific downregulation of MERTK. *Hepatol Res* 2017;47:826–830.
- [34] Ambade A, Sathishchandran A, Saha B, Gyongyosi B, Lowe P, Kodys K, et al. Hepatocellular carcinoma is accelerated by NASH involving M2 macrophage polarization mediated by hif-1. *Oncoimmunology* 2016;5:e1221557.
- [35] Friedman SL, Arthur MJ. Activation of cultured rat hepatic lipocytes by Kupffer cell conditioned medium. Direct enhancement of matrix synthesis and stimulation of cell proliferation via induction of platelet-derived growth factor receptors. *J Clin Invest* 1989;84:1780–1785.
- [36] Wynn TA, Vannella KM. Macrophages in tissue repair, regeneration, and fibrosis. *Immunity* 2016;44:450–462.
- [37] Tanaka M, Siemann DW. Gas6/Axl signaling pathway in the tumor immune microenvironment. *Cancers (Basel)* 2020;12.
- [38] Cai B, Dongiovanni P, Corey KE, Wang X, Shmarakov IO, Zheng Z, et al. Macrophage MerTK promotes liver fibrosis in nonalcoholic steatohepatitis. *Cell Metab* 2020;31:406–421.e407.
- [39] Ferrari G, Cook BD, Terushkin V, Pintucci G, Mignatti P. Transforming growth factor-beta 1 (TGF-beta1) induces angiogenesis through vascular endothelial growth factor (VEGF)-mediated apoptosis. *J Cell Physiol* 2009;219:449–458.
- [40] Zhu B, Lin N, Zhang M, Zhu Y, Cheng H, Chen S, et al. Activated hepatic stellate cells promote angiogenesis via interleukin-8 in hepatocellular carcinoma. *J Transl Med* 2015;13:365.
- [41] Cummings CT, Deryckere D, Earp HS, Graham DK. Molecular pathways: MERTK signaling in cancer. *Clin Cancer Res* 2013;19:5275–5280.
- [42] Howangyin KY, Zlatanova I, Pinto C, Ngkelo A, Cochain C, Rouanet M, et al. Myeloid-epithelial-reproductive receptor tyrosine kinase and milk fat globule epidermal growth factor 8 coordinately improve remodeling after myocardial infarction via local delivery of vascular endothelial growth factor. *Circulation* 2016;133:826–839.
- [43] Kagan P, Sultan M, Tachlytski I, Safran M, Ben-Ari Z. Both MAPK and STAT3 signal transduction pathways are necessary for IL-6-dependent hepatic stellate cells activation. *PLoS One* 2017;12:e0176173.
- [44] Ramachandran P, Dobie R, Wilson-Kanamori JR, Dora EF, Henderson BEP, Luu NT, et al. Resolving the fibrotic niche of human liver cirrhosis at single-cell level. *Nature* 2019;575:512–518.
- [45] Linger RM, Keating AK, Earp HS, Graham DK. TAM receptor tyrosine kinases: biologic functions, signaling, and potential therapeutic targeting in human cancer. *Adv Cancer Res* 2008;100:35–83.
- [46] Bernsmeier C, Pop OT, Singanayagam A, Triantafyllou E, Patel VC, Weston CJ, et al. Patients with acute-on-chronic liver failure have increased numbers of regulatory immune cells expressing the receptor tyrosine kinase MERTK. *Gastroenterology* 2015;148:603–615.e614.
- [47] Triantafyllou E, Pop OT, Possamai LA, Wilhelm A, Liaskou E, Singanayagam A, et al. MerTK expressing hepatic macrophages promote the resolution of inflammation in acute liver failure. *Gut* 2018;67:333–347.