

# Mesenchymal stem cell treatment for COVID-19

Ruonan Xu,<sup>a,1\*</sup> Zhiqian Feng,<sup>b,1</sup> and Fu-Sheng Wang<sup>a\*</sup>

<sup>a</sup>Senior Department of Infectious Diseases, The Fifth Medical Center of PLA General Hospital, National Clinical Research Center for Infectious Diseases, Beijing, China

<sup>b</sup>The Second School of Clinical Medicine, Southern Medical University, Guangzhou, China



## Summary

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection has caused a global pandemic since late 2019 that resulted in more than 360 million population infection. Among them, less than 7% of infected individuals develop severe or critical illness. Mass vaccination has been carried out, but reinfection and vaccine breakthrough cases still occur. Besides supportive and antiviral medications, much attention has been paid in immunotherapies that aim at reducing pathological changes in the lungs. Mesenchymal stem cells (MSCs) is used as an option because of their immunomodulatory, anti-inflammatory, and regenerative properties. As of January 16, 2022, when ClinicalTrials.gov was searched for “Mesenchymal stem cells and COVID-19,” over 80 clinical trials were registered. MSC therapy was found to be safe and some effective in preclinical and clinical studies. Here, we summarize the major pathological characteristics of COVID-19 and provide scientific and rational evidence for the safety and possible effectiveness of MSCs in COVID-19 treatment.

eBioMedicine 2022;77:  
103920  
Published online xxx  
<https://doi.org/10.1016/j.ebiom.2022.103920>

**Copyright** © 2022 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

**Keywords:** COVID-19; Mesenchymal stem cell; Immune therapy; Lung injury

## Introduction

Coronavirus disease 2019 (COVID-19), caused by the novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has developed into a global pandemic that began in late 2019. As of January 16, 2022, a total of 5,551,970 deaths have occurred out of a total of 328,675,785 COVID-19 cases.<sup>1</sup> Owing to the emergence of novel viral variants, including omicron, the risks of new infections, reinfections, and vaccine breakthrough infections has increased considerably worldwide, leading to a situation where the current epidemic is not yet well controlled. Similar to SARS-CoV-1 and MERS-CoV, SARS-CoV-2 belongs to the beta coronavirus family.<sup>2</sup> Individuals infected by the virus are frequently classified based on the clinical severity of their cases and designated as asymptomatic, mild, moderate, severe, or critical.<sup>3</sup> Severe and critical patients, particularly older adults and those with co-existing illnesses, have a high risk of mortality<sup>4</sup> and characterized by over-activated immune disorders. These include inflammatory hyper-responses and cytokine storms, which contribute to pulmonary tissue damage, air exchange dysfunction, acute respiratory distress syndrome (ARDS), multiple organ failure, and even death. To resolve the immune


dysfunction arising from COVID-19, besides symptomatic treatment and supplemental oxygen therapy, certain immunotherapy-related clinical trials have been conducted, including treatments based on the use of glucocorticoids,<sup>5</sup> interleukin (IL)-1 family blockers,<sup>6,7</sup> anti-IL-6 antibodies,<sup>8</sup> convalescent plasma,<sup>9</sup> and antiviral specific neutralizing antibodies.<sup>10</sup> Published data have demonstrated that immune-modulation regimens are well tolerated and show some effectiveness in clinical trial; however, further investigation is needed to confirm their efficacy in large-cohort randomized, controlled studies.

Mesenchymal stem cells (MSCs) possess immunomodulatory, anti-inflammatory, and regenerative properties.<sup>11</sup> The safety and some effectiveness of MSCs have been investigated in various clinical trials for the treatment of several disorders, including graft-versus-host disease, inflammatory bowel disease, osteoarthritis, rheumatoid arthritis, and multiple sclerosis.<sup>12</sup> The therapeutic roles of MSCs in preclinical models and clinical trials of acute lung injury, ARDS, and lung fibrosis suggest the feasibility of MSC treatment for COVID-19. For these reasons, MSC therapy can be considered as an optional interventional method for improving clinical outcomes in severe and critical patients. By January 16, 2022, when ClinicalTrials.gov was searched using the terms “Mesenchymal stem cells and COVID-19,” there were over 80 clinical trials registered (Figure 1). This review summarizes the pathological features of COVID-19 patients, along with the current progress and

\*Corresponding authors.

E-mail addresses: [xuruonan2004@aliyun.com](mailto:xuruonan2004@aliyun.com) (R. Xu), [fswang302@163.com](mailto:fswang302@163.com) (F.-S. Wang).

<sup>1</sup> These authors contributed equally to this work.



**Figure 1.** A summary of clinical trials registered in ClinicalTrials.gov by searching “Mesenchymal stem cells and COVID-19”. (a) A map for all registered clinical trials. (b) The chart for the recruitment status of clinical trials.

challenges of using MSCs to treat COVID-19, and also establishes the rationality of using MSCs in the treatment of the disease.

### Pathological features of COVID-19

During acute infection with SARS-CoV-2, viral replication and attack may directly result in injury to the respiratory tract and pulmonary tissues, which can trigger a series of over-active immune responses that not only play an antiviral role but also induce tissue inflammation and damage. Autopsies of dead patients with critical illness showed that the histological characteristics of lesions in the lungs were virus-induced and led to diffuse alveolar epithelial damage and inflammatory cell infiltration. This resulted in alveolar destruction with alveolar lining cell necrosis, protein-rich exudates, hyaline membrane formation, and endothelial cell membrane destruction. All of these eventually led to an increase in alveolar-capillary membrane permeability and loss of aerated lung tissue. Moreover, certain patients had fibrous deposition, mural fibrosis, and microcystic honeycombing in the lungs.<sup>13–16</sup> In contrast to other types of pneumonia, SARS-CoV-2 appears more likely to cause severe thrombotic disorders and widespread thrombosis with microangiopathy.<sup>14,17,18</sup> In mild and moderate cases, innate and adaptive immune responses are well coordinated, which contributes to viral symptom resolution and convalescence.<sup>19</sup> However, in severe cases, over-activation and massive infiltration of immunocytes such as macrophages, natural killer cells, and CD4+ and CD8+ T cells into the lungs result in high levels of cytokine and chemokine production.<sup>20</sup> Uncontrolled pro-inflammatory feedback triggers a cytokine storm and accelerates local tissue damage and systemic injuries.<sup>21</sup> Severe patients showed a significant increase in plasma levels of cytokines, including IL-2, IL-6, IL-7, IL-10, granulocyte-colony stimulating factor, interferon-inducible protein 10, monocyte

chemoattractant protein-1, macrophage inflammatory protein (MIP)-1 $\alpha$ , and tumor necrosis factor (TNF)- $\alpha$ .<sup>22–25</sup> The antiviral immune-related interferon (IFN) type I response was also highly impaired in the majority of patients with severe symptoms. Furthermore, the total numbers of peripheral CD4+ T cells and CD8+ T cells were significantly lower in severe and critical cases,<sup>26,27</sup> particularly in patients with lymphocytopenia, impaired type I IFN production, and excessive activation of Th1 and inflammatory monocytes.<sup>28</sup> A high viral load may further aggravate the ongoing inflammatory cascades.<sup>29</sup>

Generally, pulmonary inflammatory lesions were primarily resolved in the convalescence stage, while > 80% of severe/critical patients and 60% of mild/moderate patients still exhibited residual computed tomography (CT) abnormalities three months after discharge. These predominantly included ground-glass opacity followed by strip-like fibrosis.<sup>30</sup> A recent result of a one-year follow-up study showed that 25% of individuals discharged from hospitalization still had chest CT abnormalities, including residual linear opacities and multifocal reticular/cystic lesions, and older patients with severe pneumonia, ARDS, and lymphopenia were more likely to develop CT abnormalities for over a year.<sup>31</sup> Pulmonary fibrosis is seen as an important factor affecting long-term clinical outcomes.<sup>32,33</sup> In an extended study of SARS-CoV-2-infected patients, lung fibrosis persisted for over seven years, suggesting that lung fibrosis could be a persistent sequela of pulmonary inflammatory injury.<sup>34</sup> The increased expression of angiotensin-converting enzyme 2, transforming growth factor beta, connective tissue growth factor, and fibronectin<sup>35</sup> and endothelial/epithelial-to-mesenchymal transition have all been believed to account for pulmonary fibrosis and vascular destruction post-COVID-19.<sup>36</sup>


Extrapulmonary manifestations, including thrombotic complications, myocardial dysfunction and arrhythmia, acute coronary syndromes, hepatocellular

injury, gastrointestinal symptoms, acute kidney injury, and hyperglycemia, may further complicate the natural course of COVID-19 in severe and critical cases.<sup>37–39</sup> A broad distribution of SARS-CoV-2 in the respiratory, cardiovascular, genitourinary, digestive, and immune systems and in endo/exocrine glands and the skin was identified in autopsy specimens.<sup>38,40</sup> Renal autopsy revealed coronavirus-like particles in the cytoplasm of the renal proximal tubular epithelium, accompanied by prominent acute renal tubular injury.<sup>41</sup> Human islet autopsy showed selective SARS-CoV-2 infection in  $\beta$  cells, followed by attenuated pancreatic insulin secretion and increased  $\beta$  cell apoptosis.<sup>42</sup> In the heart, both compositional and gene expression changes were recorded during acute SARS-CoV-2 infection.<sup>43</sup>

### Rationality of MSC treatment for COVID-19

MSCs may exert immunomodulation via autocrine, paracrine, and endocrine pathways (Figure 2).<sup>44–46</sup> MSCs can modulate host innate and adaptive immune responses, thus reducing proinflammatory cytokine production.<sup>45,46</sup> Given the previously described

pathological characteristics of COVID-19,<sup>47,48</sup> MSCs are considered an optional candidate for the treatment of COVID-19. In preclinical models of acute lung injury, the benefit of MSCs was associated with the alleviation of local and systemic inflammation,<sup>49</sup> the amelioration of immune cell activation,<sup>50</sup> and mitigated lung injury.<sup>45</sup> MSCs have been shown to reduce the infiltration of macrophages, neutrophils, and DCs into the lungs and to decrease the levels of MIP-2, TNF- $\alpha$ , IL-6, IL-1 $\beta$ , and IL-12p70 in bronchoalveolar lavage fluid.<sup>45,46,49,51,52</sup> Stanniocalcin-2 was found to mediate the effect of MSCs through modulating macrophage polarization from the M1 to M2 phenotype.<sup>53</sup> In addition, MSCs induce mature DCs to become tolerogenic DCs, maintain immune tolerance, and negatively regulate the immune response.<sup>54</sup> Furthermore, MSCs can attenuate pulmonary inflammation by regulating different T cell subsets; for example, MSCs can inhibit the activation and proliferation of T cells and induce the polarization of CCR2+CD4+ T cell subsets enriched with forkhead box P3 and IL-10.<sup>55</sup> They can also inhibit the production of proinflammatory IL-17 and IFN- $\gamma$  in T cells<sup>49</sup> and alleviate the infiltration of proinflammatory



**Figure 2.** A schematic overview of the pathological characteristics and potential of MSCs for treating COVID-19. (1) During the acute phase, SARS-CoV-2 infection induces an inflammatory hyper-response and pulmonary tissue damage, which may lead to the partial fibrosis of lung tissues. (2) Immunomodulatory effects of MSCs contribute to reducing inflammation. (3) MSCs promote proliferation of alveolar epithelial cells and vascular endothelial cells and reduce their apoptosis. (4) MSCs contribute to decreasing collagen deposition and inhibiting epithelial–mesenchymal transformation.

CD8+ T cells<sup>56</sup> and B cells expressing chemokines for neutrophil recruitment and immunoglobulin production.<sup>57</sup> Additionally, MSC-derived extracellular vesicles could also participate in immunoregulation by decreasing the absolute count of neutrophils and TNF- $\alpha$  levels in an *ex vivo* lung injury model<sup>58</sup> and by transferring functional mitochondria to macrophages to reduce proinflammatory cytokine production and increase their phagocytic capacity.<sup>59</sup>

MSCs possess regenerative and differentiation properties that contribute to the repair of injured tissues. MSCs have been shown to improve alveolar fluid clearance and lung function by secreting paracrine factors that regulate membrane transport, restore the alveolar epithelial and pulmonary microvascular endothelial lining, reduce pulmonary edema, and ultimately promote the restoration of lung structure. Different cytokines are released through paracrine pathways, including keratinocyte growth factor (KGF), angiopoietin (Ang)-1, hepatocyte growth factor (HGF), and vascular endothelial growth factor (VEGF).<sup>60–62</sup> KGF produced by MSCs may rapidly and specifically contribute to the proliferation and differentiation of alveolar epithelial type II cells.<sup>60</sup> KGF is also related to surfactant production,<sup>61</sup> alveolar epithelial cell apoptosis relief,<sup>63</sup> and sodium-dependent alveolar fluid transport recovery.<sup>64</sup> Furthermore, HGF and VEGF may have synergistic effects on the restoration of endothelial intercellular junctions, reduction of caveolin-1 protein expression, and endothelial apoptosis.<sup>65</sup> Additionally, Ang-1 produced by MSCs has the capacity to increase endothelial cell proliferation during vascular remodeling, promote vascular stabilization during inflammation, and improve human alveolar epithelial type II cell permeability through directly affecting cytoskeletal remodeling.<sup>62,66,67</sup>

MSCs have the capacity to reshape the lung cell microenvironment by reducing the levels of profibrogenic factors. MSC infusion contributes to lung structure recovery by restoring lung epithelial cell proliferation and decreasing collagen deposition in a bleomycin-induced idiopathic pulmonary fibrosis model.<sup>68,69</sup> The efficacy of MSCs is also associated with the correction of inappropriate epithelial–mesenchymal transformation.<sup>70</sup>

### Clinical trials of MSCs for COVID-19 treatment

The safety of MSC treatment for patients with ARDS has been demonstrated in the studies listed in Table 1.<sup>71–76</sup> In addition, MSCs simultaneously promote the restoration of pathological lung tissues.<sup>77</sup> Therefore, MSC therapy is considered an alternative option for patients with severe COVID-19.

To date, several clinical trials have utilized different sources of MSCs (Table 2).<sup>78</sup> Compared with conventional treatment, MSC add-on therapy may improve clinical symptoms, which are characterized by the relief

Numbers	Trial ID	Sources of MSCs	Phase	Numbers of patients	Doses and administration routes	Primary outcomes	Refs.
1	NCT01902082	AD-MSCs	1	12	A single dose of $1 \times 10^6$ cells/ kg; IV	No adverse events or serious adverse events. The clinical effect at this dose is weak.	71
2	Not reported	BM-MSCs	Case report	2	A single dose of $2 \times 10^6$ cells/ kg; IV	Oxygenation and pulmonary compliance were improved. Levels of inflammation markers were reduced.	72
3	Not reported	UC-MSCs	1	9	A single dose of $1.0 \times 10^6$ cells/kg or $5.0 \times 10^6$ cells/kg or $1.0 \times 10^7$ cells/kg; IV	Tolerated without serious adverse events.	73
4	NCT01775774	BM-MSCs	1	9	A single dose of $1 \times 10^6$ cells/kg or $5 \times 10^6$ cells/kg; IV	Well tolerated and safe.	74
5	NCT02097641	BM-MSCs	2a	60	A single dose of $10 \times 10^6$ cells/kg; IV	No MSC-related predefined hemodynamic or respiratory adverse events.	75
6	NCT03042143.	UC-MSCs	1	9	A single dose of $1$ or $4 \times 10^8$ ; IV	Well tolerated. Adverse events included apyrexia, non-sustained ventricular tachycardia, and deranged liver function.	76

**Table 1: Completed clinical studies of MSCs in the treatment of patients with ARDS.**

MSCs, mesenchymal stem cells; AD-MSCs, adipose-derived MSCs; BM-MSCs, bone marrow autologous MSCs; UC-MSCs, umbilical cord-derived MSCs; IV, intravenous injection.

Numbers	Trial ID	Source of MSCs	Phase	Numbers of patients	Doses and administration routes	Primary outcomes	Refs.
1	NCT04252118	UC-MSCs	1	18	Three doses of $3 \times 10^7$ cells; IV	Safe and well tolerated.	79
2	ChiCTR2000029990	MSC	1	10	A single dose of $1 \times 10^6$ cells/ kg; IV	No adverse effects. Pulmonary function and symptoms significantly improved.	80
3	Not reported	MB-MSCs	Case report	2	Three doses of $1 \times 10^6$ cells/ kg; IV	Values of immune indicators increased, and levels of inflammation indicators decreased. Lung exudate lesions were absorbed.	81
4	ChiCTR2000029606	MB-MSCs	1	44	Three doses of $3 \times 10^7$ cells; IV	Safe and well tolerated. SpO <sub>2</sub> was significantly improved, and chest imaging results were improved.	82
5	ChiCTR2000031494	UC-MSCs	1	41	A single dose of $2 \times 10^6$ cells/ kg; IV	Clinical symptoms were relieved and alleviated in a shorter time. Levels of inflammatory factors rapidly decreased, and lymphocyte count returned to normal levels in less time.	83
6	Not reported	UC-MSCs	1	31	$1 \times 10^6$ cells/ kg were suspended in 100 ml normal saline. The volume of infused UC-MSCs was 100–300 mL; IV	No adverse events were observed. Laboratory parameters tended to improve.	84
7	Not reported	Convalescent plasma combination with UC-MSCs	Case report	1	Three doses of $1 \times 10^6$ cells/kg; IV	Absolute lymphocyte count increased, bilateral infiltrates were absorbed, and pulmonary function was significantly improved.	85
8	Not reported	UC-MSCs or PL-MSCs	Case series	11	Three doses of $2 \times 10^8$ cells; IV	No serious adverse events. Levels of inflammatory biomarkers significantly decreased.	86
9	Not reported	hESC-IMRCs	1	27	One to three doses of $3 \times 10^6$ cells/ kg; IV	No adverse events or abnormal responses related to cell therapy occurred. The area of lung fibrosis lesions decreased.	87
10	NCT04269525	UC-MSCs	A pilot study	16	Four doses of $1 \times 10^8$ cells; IV	No infusion-related or allergic reactions. Oxygenation index was improved, and CT scan improved. Restoration of lymphocyte count and subsets.	88
11	NCT04416139	UC-MSCs	1	5	A single dose of $1 \times 10^6$ cells /kg; IV	PaO <sub>2</sub> /FIO <sub>2</sub> was improved.	89
12	Not reported	hESC-IMRCs	Case report	1	Two doses of $3 \times 10^6$ cells/ kg; IV	Levels of proinflammatory cytokines were decreased. SpO <sub>2</sub> improved.	90
13	NCT04288102	UC-MSCs	2	100	Three doses of $4 \times 10^7$ cells; IV	Safe and potentially effective. Lung lesion volume, especially solid component lesion volume, was reduced. The 6-minute walk test result improved.	91
14	NCT04457609	UC-MSCs	1	40	A single dose of $1 \times 10^6$ cells/ kg; IV	Safe and well tolerated. Survival rate increased.	92

Table 2 (Continued)

Numbers	Trial ID	Source of MSCs	Phase	Numbers of patients	Doses and administration routes	Primary outcomes	Refs.
15	NCT04392778	WJ-MSCs	1	30	Three doses of $3 \times 10^6$ cells/kg; IV	All the indicators of anti-inflammation, anti-fibrosis signs in the lungs, and levels of immunodulatory markers were dramatically improved.	93
16	NCT04355728	UC-MSCs	1/2a	24	Two doses of $100 \pm 20 \times 10^6$ cells; IV	No serious adverse events were observed. Mortality and time to recovery decreased significantly.	94
17	Not reported	MSCs	1	23	Two or three doses of $1 \times 10^6$ cells/kg; IV	Safe and well tolerated. Pulmonary function and overall outcomes were improved.	95
18	Not reported	UC-MSCs	1	210	A single dose of $1-2 \times 10^6$ cells/kg; IV	No adverse effects. SaO <sub>2</sub> improved.	96
19	Not reported	AD-MSCs	1	13	One to three doses of $0.98 \times 10^6$ cells/kg; IV	No adverse events. Clinical symptoms and values of inflammatory parameters improved.	97
20	NCT04339660	UC-MSCs	2	58	A single dose of $1 \times 10^6$ cells /kg; IV	MSC-treated patients had fewer adverse events. Clinical symptoms, values of inflammatory parameters, and CT scan improved. MSCs promoted the production of SARS-CoV-2-specific antibodies.	98
21	IRCT20190717044241N2	WJ-MSCs	1	5	Three doses of $150 \times 10^6$ cells; IV	Safe and well tolerated. Levels of inflammatory biomarkers improved.	99
22	Not reported	UC-MSCs	Case report	1	Three doses of $5 \times 10^7$ cells/ kg; IV	No obvious side effects. Values of clinical indices and clinical symptoms were improved.	100
23	Not reported	UC-MSCs	Case report	1	Five doses of $1.5 \times 10^6$ cells/ kg; IV	The number of lymphocytes was increased. Respiratory and renal functions were improved.	101
24	Not reported	WJ-MSCs	Case report	1	A single dose of $1 \times 10^6$ cells /kg; IV	The percentage and counts of T cells were increased. CT scan was improved.	102
25	Not reported	MSCs	A retrospective study	25	One to three doses of $1 \times 10^6$ cells/kg; IV	Three cases experienced treatment-related side effects including liver dysfunction, heart failure, and allergic rash.	103
26	Not reported	ExoFlo	1	27	15 ml of ExoFlo was added to 100 mL of normal saline; IV	No adverse events. Clinical status, oxygenation, and laboratory values improved, levels of acute phase reactants declined.	105

**Table 2: Completed clinical studies of MSCs in the treatment of patients with COVID-19.**

MSCs, mesenchymal stem cells; MB-MSCs, menstrual blood-derived MSCs; PL-MSCs, placenta-derived MSCs; hESC-IMRCs, human embryonic stem cell–derived immunity– and matrix–regulatory cells; WJ-MSCs, Wharton's jelly-derived MSCs; ExoFlo, exosomes derived from allogeneic bone marrow MSCs; SaO<sub>2</sub>, arterial oxygen saturation; SpO<sub>2</sub>, oxygen saturation; CT, computed tomography.

of dyspnea and a shorter recovery time.<sup>79–90</sup> CT images of the lungs also showed notable improvements in pulmonary lesions,<sup>81–83,86,91</sup> including a decrease in pathological lobes, ground-glass opacity, and consolidations.<sup>83</sup> Other studies have shown that MSC add-on therapy enhanced the survival of severe/critical patients<sup>82,92</sup> and even those who developed ARDS.<sup>86,93–95</sup> The choice of a time window for MSC-based therapies was considered crucial; MSC transfusion for severe patients prior to intubation could reduce the risk of disease progression and 28-day mortality.<sup>83,96</sup> Furthermore, early initiation of MSC treatment might contribute to a higher extubation rate in patients who received mechanical ventilation.<sup>97</sup>

The underlying mechanisms of MSC-based remedies for the maintenance of homeostasis, immune reconstitution, and tissue repair have been investigated in detail, and the effect of MSCs on hematopoiesis was found to correlate with the activation of a novel subpopulation of VNN2+ hematopoietic stem/progenitor like (HSPC-like) cells, which were mobilized following MSC infusion.<sup>98</sup> Notably, the changes in inflammatory indices following MSC treatment were inconsistent. Partial data from clinical trials showed that MSCs exerted anti-inflammatory and immunoregulatory effects through reducing the levels of inflammatory biomarkers (C-reactive protein, IL-6, IL-2, IL-12, IL-8, TNF- $\alpha$ , and IFN- $\gamma$ ) and increasing the level of anti-inflammatory IL-10 and T cell counts.<sup>80,81,83,84,94,95,98–102</sup> In other trials, significant changes in inflammatory factor levels were not observed.<sup>82,91,103</sup> Therefore, the effect of MSC treatment on inflammation should be further studied in large-cohort randomized controlled trials.

In our double-blind, placebo-controlled phase 2 trial, MSC administration significantly reduced solid component lesion volume within 28 days in patients with severe COVID-19.<sup>91</sup> During the subsequent one-year follow-up period, the patients who received MSC treatment showed reduced solid component lesion volume and improved pulmonary function, indicating that MSC therapy might have long-term benefits.<sup>104</sup> Bone marrow MSC-derived exosomes have also been reported to be safe and effective in reducing acute phase reactant and neutrophil counts, increasing T cell counts, and improving oxygenation in severe cases.<sup>105</sup> These data support the therapeutic potential of MSCs for severe cases with progressive lung damage.

Although most clinical trials showed that MSC treatment is generally safe and well tolerated, some MSC-related adverse events have still been recorded. The major clinical symptoms included facial flushing, fever,<sup>79</sup> shivering,<sup>86</sup> headache,<sup>99</sup> liver dysfunction, heart failure, and allergic rash.<sup>95</sup> However, adverse events were recorded in both MSC-treated patients and controls, indicating that the side effects might be infusion-related.<sup>94</sup> Furthermore, the speed of transfusion and the baseline status of patients with co-existing

illnesses may be associated with the occurrence of adverse events.

### Challenges and perspective

Large-scale vaccination has provided essential protection by establishing herd immunity, which will further contribute to the control of the SARS-CoV-2 pandemic. However, there are still re-infection and post-vaccination breakthrough cases in COVID-19. Many infected individuals are asymptomatic or show mild/moderate symptoms; however, it is difficult to ensure the full recovery of severe and critical patients using current medications, including symptomatic treatment, antiviral therapy, and respiratory support treatment. MSCs may serve as an alternative immunotherapeutic option for severely affected individuals and contribute to the improvement of COVID-19 outcomes. Notably, clinical application protocols for MSCs are diverse, and optimized standard protocols will be required to maximize their therapeutic effect. Moreover, the choice of a time window for treatment, the identification of COVID-19 patients at suitable phases, and the regimen (including dosage, interval time, and round) all need to be comprehensively optimized.

Most importantly, adverse reactions related to MSC treatment and infusion also require caution. Low-grade fever, facial flushing, headache, and allergic rash after MSC infusion were reported, although these side effects were transient and generally disappeared within 24 h without any treatment. However, treatment of these symptoms may be required in severe cases. In addition, the risks of thromboembolic events associated with intravenous infusions of high doses of MSCs should also be noted. To improve safety, it is best to monitor the electrocardiogram and percutaneous oxygen saturation during MSC infusion, and to test for laboratory parameters, including liver and kidney function, blood routine, CRP, inflammatory cytokines, coagulation, procalcitonin, blood gases, and heart function, before MSC infusion.<sup>106</sup>

Follow-up studies extending over a year in length have reported that severe COVID-19 patients still experienced symptoms and persistent physiological and radiographic abnormalities.<sup>31</sup> These data indicate that some severe COVID-19 patients will still require suitable interventions to improve their long-term prognoses. MSCs have been suggested for post-COVID-19 treatment for their antifibrotic and regeneration/differentiation properties. To determine the fate of the infused cells and to interpret the interactions between the colonized MSCs and immunocytes, MSC tracking *in vivo* is required. Furthermore, specific airspace and plasma biomarkers are required to predict and verify the efficacy of MSCs against COVID-19.

Both fresh and frozen MSCs with proven efficacy in clinical applications are readily available, and it is

noteworthy that dead or apoptotic MSCs found in MSC preparations exert the same immunomodulatory properties as the "living MSCs" by releasing phosphatidylserine.<sup>107</sup> In addition, MSCs derived from different donors and tissues lead to individual heterogeneity, which complicates the progress of consistent and standardized stem cell production.<sup>108,109</sup> Therefore, it is necessary to screen qualified MSCs to avoid uncertain effects.

In summary, MSC therapy appear to serve as an alternative candidate for ameliorating inflammation, repairing lung tissue injury, and preventing long-term pulmonary disability in patients with COVID-19. However, the different groups of enrolled patients, therapeutic regimen and the time window for treatment of MSCs may lead to the disparities of data in the previous clinical trials. Prospective, multicenter, randomized and controlled trials with large sample size are still necessary to further confirm therapeutic efficacy of MSCs for the patients with severe and critical COVID-19.

### Outstanding questions

Although available data from preliminary clinical trials have proved the safety and some efficiency of MSC treatment for COVID-19 patients, there are certain disparities caused by the inconsistency of inclusion and exclusion criteria. Therefore, determining the characteristics of patients who may gain more clinical benefits from MSC treatment will require more multicenter, randomized, controlled trials and long-term follow-up studies. Moreover, concomitant treatments may exert a synergistic or antagonistic role with MSCs, and treatment regimens need to be optimized. Further mechanistic studies are also needed to verify the efficacy of MSCs in COVID-19 treatment.

### Search strategy and selection criteria

Data for this review were identified by searches of PubMed and Google Scholar and references from relevant articles using the search terms "SARS-CoV-2," "COVID-19," "immunotherapy," "Mesenchymal stem cells," "ARDS," "Clinical trial," "Lung injury," "pulmonary fibrosis," "Cytokine storm," "immune response," "clinical features," "follow-up," and related terms. Only articles published in English between 1994 and 2021 were included in the study. Articles were selected based on their relevance and the authors' best judgments.

### Contributors

Ruonan Xu and Fu-Sheng Wang contributed to the conception and design of the review. The draft was written by Ruonan Xu and Zhiqian Feng. Fu-Sheng Wang, Ruonan Xu critically and systemically revised the manuscript. All authors contributed to the article and approved the submitted version.

### Declaration of interests

The authors declare no competing interests.

### Acknowledgments

This study was supported by the funds from Innovative Research Group Project of the National Natural Science Foundation of China (81721002), the National Key Research and Development Program of China (2017YFA0105700). We also declare that the funders had no role in paper design, data collection, data analysis, interpretation and writing of the paper. Figure 1 was created using BioRender software.

### References

- 1 WHO Coronavirus (COVID-19) Dashboard: <https://covid19.who.int/> January 16, 2022.
- 2 Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med*. 2020;382(8):727–733. <https://doi.org/10.1056/NEJMoa2001017>.
- 3 National Institutes of Health. *Coronavirus Disease 2019 (COVID-19) Treatment Guidelines*. Bethesda (MD): National Institutes of Health (US); 2021.
- 4 Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA*. 2020;323(11):1061–1069. <https://doi.org/10.1001/jama.2020.1585>.
- 5 Sterne J, Murthy S, Diaz JV, et al. Association between administration of systemic corticosteroids and mortality among critically ill patients with COVID-19: a meta-analysis. *JAMA*. 2020;324(13):1330–1341. <https://doi.org/10.1001/jama.2020.17023>.
- 6 Cavalli G, Dagna L. The right place for IL-1 inhibition in COVID-19. *Lancet Respir Med*. 2021;9(3):223–224. [https://doi.org/10.1016/S2213-2600\(21\)00035-7](https://doi.org/10.1016/S2213-2600(21)00035-7).
- 7 Cavalli G, De Luca G, Campochiaro C, et al. Interleukin-1 blockade with high-dose anakinra in patients with COVID-19, acute respiratory distress syndrome, and hyperinflammation: a retrospective cohort study. *Lancet Rheumatol*. 2020;2(6):e325–e331. [https://doi.org/10.1016/S2665-9913\(20\)30127-2](https://doi.org/10.1016/S2665-9913(20)30127-2).
- 8 Michot JM, Albiges L, Chaput N, et al. Tocilizumab, an anti-IL-6 receptor antibody, to treat COVID-19-related respiratory failure: a case report. *Ann Oncol*. 2020;31(7):961–964. <https://doi.org/10.1016/j.annonc.2020.03.300>.
- 9 Duan K, Liu B, Li C, et al. Effectiveness of convalescent plasma therapy in severe COVID-19 patients. *Proc Natl Acad Sci U S A*. 2020;117(17):9490–9496. <https://doi.org/10.1073/pnas.2004168117>.
- 10 Gupta A, Gonzalez-Rojas Y, Juarez E, et al. Early treatment for COVID-19 with SARS-CoV-2 neutralizing antibody sotrovimab. *N Engl J Med*. 2021;385(21):1941–1950. <https://doi.org/10.1056/NEJMoa2107934>.
- 11 Lopes-Pacheco M, Robba C, Rocco P, Pelosi P. Current understanding of the therapeutic benefits of mesenchymal stem cells in acute respiratory distress syndrome. *Cell Biol Toxicol*. 2020;36(1):83–102. <https://doi.org/10.1007/s10565-019-09493-5>.
- 12 Wang LT, Ting CH, Yen ML, et al. Human mesenchymal stem cells (MSCs) for treatment towards immune- and inflammation-mediated diseases: review of current clinical trials. *J Biomed Sci*. 2016;23(1):76. <https://doi.org/10.1186/s12929-016-0289-5>.
- 13 Edler C, Schröder AS, Aepfelbacher M, et al. Dying with SARS-CoV-2 infection—an autopsy study of the first consecutive 80 cases in Hamburg, Germany. *Int J Legal Med*. 2020;134(4):1275–1284. <https://doi.org/10.1007/s00414-020-02317-w>.
- 14 Ackermann M, Verleden SE, Kuehnel M, et al. Pulmonary vascular endothelialitis, thrombosis, and angiogenesis in COVID-19. *N Engl J Med*. 2020;383(2):120–128. <https://doi.org/10.1056/NEJMoa2015432>.
- 15 Xu Z, Shi L, Wang Y, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med*. 2020;8(4):420–422. [https://doi.org/10.1016/S2213-2600\(20\)30076-X](https://doi.org/10.1016/S2213-2600(20)30076-X).



- 16 Carsana L, Sonzogni A, Nasr A, et al. Pulmonary post-mortem findings in a series of COVID-19 cases from northern Italy: a two-centre descriptive study. *Lancet Infect Dis*. 2020;20(10):1135–1140. [https://doi.org/10.1016/S1473-3099\(20\)30434-5](https://doi.org/10.1016/S1473-3099(20)30434-5).
- 17 Magro C, Mulvey JJ, Berlin D, et al. Complement associated microvascular injury and thrombosis in the pathogenesis of severe COVID-19 infection: a report of five cases. *Transl Res*. 2020;220:1–13. <https://doi.org/10.1016/j.trsl.2020.04.007>.
- 18 Varga Z, Flammer AJ, Steiger P, et al. Endothelial cell infection and endothelitis in COVID-19. *Lancet*. 2020;395(10234):1417–1418. [https://doi.org/10.1016/S0140-6736\(20\)30937-5](https://doi.org/10.1016/S0140-6736(20)30937-5).
- 19 Sette A, Crotty S. Adaptive immunity to SARS-CoV-2 and COVID-19. *Cell*. 2021;184(4):861–880. <https://doi.org/10.1016/j.cell.2021.01.007>.
- 20 Szabo PA, Dogra P, Gray JJ, et al. Longitudinal profiling of respiratory and systemic immune responses reveals myeloid cell-driven lung inflammation in severe COVID-19. *Immunity*. 2021;54(4):797–814. <https://doi.org/10.1016/j.immuni.2021.03.005>.
- 21 Tay MZ, Poh CM, Rénia L, Macary PA, Ng L. The trinity of COVID-19: immunity, inflammation and intervention. *Nat Rev Immunol*. 2020;20(6):363–374. <https://doi.org/10.1038/s41577-020-0311-8>.
- 22 Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
- 23 Song JW, Zhang C, Fan X, et al. Immunological and inflammatory profiles in mild and severe cases of COVID-19. *Nat Commun*. 2020;11(1):3410. <https://doi.org/10.1038/s41467-020-17240-2>.
- 24 Chen G, Wu D, Guo W, et al. Clinical and immunological features of severe and moderate coronavirus disease 2019. *J Clin Invest*. 2020;130(5):2620–2629. <https://doi.org/10.1172/JCI137244>.
- 25 Qin C, Zhou L, Hu Z, et al. Dysregulation of immune response in patients with coronavirus 2019 (COVID-19) in Wuhan, China. *Clin Infect Dis*. 2020;71(15):762–768. <https://doi.org/10.1093/cid/ciaa248>.
- 26 Sun Y, Dong Y, Wang L, et al. Characteristics and prognostic factors of disease severity in patients with COVID-19: the Beijing experience. *J Autoimmun*. 2020;112: 102473. <https://doi.org/10.1016/j.jaut.2020.102473>.
- 27 Chen G, Wu D, Guo W, et al. Clinical and immunological features of severe and moderate coronavirus disease 2019. *J Clin Invest*. 2020;130(5):2620–2629. <https://doi.org/10.1172/JCI137244>.
- 28 Zhou Y, Fu B, Zheng X, et al. Pathogenic T-cells and inflammatory monocytes incite inflammatory storms in severe COVID-19 patients. *Natl Sci Rev*. 2020;7(6):998–1002. <https://doi.org/10.1093/nsr/nwaa041>.
- 29 Hadjadj J, Yatim N, Barnabei L, et al. Impaired type I interferon activity and inflammatory responses in severe COVID-19 patients. *Science*. 2020;369(6504):718–724. <https://doi.org/10.1126/science.abc6027>.
- 30 Zhou M, Xu J, Liao T, et al. Comparison of residual pulmonary abnormalities 3 months after discharge in patients who recovered from COVID-19 of different severity. *Front Med*. 2021;8: 682087. <https://doi.org/10.3389/fmed.2021.682087>. (Lausanne).
- 31 Pan F, Yang L, Liang B, et al. Chest CT patterns from diagnosis to 1 year of follow-up in COVID-19. *Radiology*. 2021 211199. <https://doi.org/10.1148/radiol.2021211199>.
- 32 Spagnolo P, Balestro E, Aliberti S, et al. Pulmonary fibrosis secondary to COVID-19: a call to arms? *Lancet Respir Med*. 2020;8(8):750–752. [https://doi.org/10.1016/S2213-2600\(20\)30222-8](https://doi.org/10.1016/S2213-2600(20)30222-8).
- 33 Grillo F, Barisione E, Ball L, Mastracci L, Fiocca R. Lung fibrosis: an undervalued finding in COVID-19 pathological series. *Lancet Infect Dis*. 2021;21(4):e72. [https://doi.org/10.1016/S1473-3099\(20\)30582-X](https://doi.org/10.1016/S1473-3099(20)30582-X).
- 34 Wu X, Dong D, Ma D. Thin-Section computed tomography manifestations during convalescence and long-term follow-up of patients with severe acute respiratory syndrome (SARS). *Med Sci Monit*. 2016;22:2793–2799. <https://doi.org/10.12659/msm.896985>.
- 35 Xu J, Xu X, Jiang L, Dua K, Hansbro PM, Liu G. SARS-CoV-2 induces transcriptional signatures in human lung epithelial cells that promote lung fibrosis. *Respir Res*. 2020;21(1):182. <https://doi.org/10.1186/s12931-020-01445-6>.
- 36 Eapen MS, Lu W, Gaikwad AV, et al. Endothelial to mesenchymal transition: a precursor to post-COVID-19 interstitial pulmonary fibrosis and vascular obliteration? *Eur Respir J*. 2020;56(4):2003167. <https://doi.org/10.1183/13993003.03167-2020>.
- 37 Gupta A, Madhavan MV, Sehgal K, et al. Extrapulmonary manifestations of COVID-19. *Nat Med*. 2020;26(7):1017–1032. <https://doi.org/10.1038/s41591-020-0968-3>.
- 38 Yao XH, Luo T, Shi Y, et al. A cohort autopsy study defines COVID-19 systemic pathogenesis. *Cell Res*. 2021;31(8):836–846. <https://doi.org/10.1038/s41422-021-00523-8>.
- 39 Lopes-Pacheco M, Silva PL, Cruz FF, et al. Pathogenesis of multiple organ injury in COVID-19 and potential therapeutic strategies. *Front Physiol*. 2021;12: 593223. <https://doi.org/10.3389/fphys.2021.593223>.
- 40 Puelles VG, Lütgehetmann M, Lindenmeyer MT, et al. Multiorgan and renal tropism of SARS-CoV-2. *N Engl J Med*. 2020;383(6):590–592. <https://doi.org/10.1056/NEJMc2011400>.
- 41 Su H, Yang M, Wan C, et al. Renal histopathological analysis of 26 post-mortem findings of patients with COVID-19 in China. *Kidney Int*. 2020;98(1):219–227. <https://doi.org/10.1016/j.kint.2020.04.003>.
- 42 Wu CT, Lidsky PV, Xiao Y, et al. SARS-CoV-2 infects human pancreatic  $\beta$  cells and elicits  $\beta$  cell impairment. *Cell Metab*. 2021;33(8):1565–1576. <https://doi.org/10.1016/j.cmet.2021.05.013>.
- 43 Delorey TM, Ziegler C, Heimberg G, et al. COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular targets. *Nature*. 2021;595(7865):107–113. <https://doi.org/10.1038/s41586-021-03570-8>.
- 44 Wang Y, Chen X, Cao W, Shi Y. Plasticity of mesenchymal stem cells in immunomodulation: pathological and therapeutic implications. *Nat Immunol*. 2014;15(11):1009–1016. <https://doi.org/10.1038/ni.3002>.
- 45 Ionescu L, Byrne RN, van Haaften T, et al. Stem cell conditioned medium improves acute lung injury in mice: *in vivo* evidence for stem cell paracrine action. *Am J Physiol Lung Cell Mol Physiol*. 2012;303(11):L967–L977. <https://doi.org/10.1152/ajplung.00144.2011>.
- 46 Gupta N, Su X, Popov B, Lee JW, Serikov V, Matthay MA. Intrapulmonary delivery of bone marrow-derived mesenchymal stem cells improves survival and attenuates endotoxin-induced acute lung injury in mice. *J Immunol*. 2007;179(3):1855–1863. <https://doi.org/10.4049/jimmunol.179.3.1855>.
- 47 Da SK, Gobatto A, Costa-Ferro Z, et al. Is there a place for mesenchymal stromal cell-based therapies in the therapeutic armamentarium against COVID-19? *Stem Cell Res Ther*. 2021;12(1):425. <https://doi.org/10.1186/s13287-021-02502-7>.
- 48 Khoury M, Cuenca J, Cruz FF, Figueroa FE, Rocco P, Weiss DJ. Current status of cell-based therapies for respiratory virus infections: applicability to COVID-19. *Eur Respir J*. 2020;55(6). <https://doi.org/10.1183/13993003.00858-2020>.
- 49 Hackstein H, Lippitsch A, Krug P, et al. Prospectively defined murine mesenchymal stem cells inhibit Klebsiella pneumoniae-induced acute lung injury and improve pneumonia survival. *Respir Res*. 2015;16:123. <https://doi.org/10.1186/s12931-015-0288-1>.
- 50 Phinney DG, Di Giuseppe M, Njah J, et al. Mesenchymal stem cells use extracellular vesicles to outsource mitophagy and shuttle microRNAs. *Nat Commun*. 2015;6:8472. <https://doi.org/10.1038/ncomms9472>.
- 51 Hayes M, Curley GF, Masterson C, Devaney J, O'Toole D, Laffey JG. Mesenchymal stromal cells are more effective than the MSC secretome in diminishing injury and enhancing recovery following ventilator-induced lung injury. *Intensive Care Med Exp*. 2015;3(1):29. <https://doi.org/10.1186/s40635-015-0065-y>.
- 52 Curley GF, Ansari B, Hayes M, et al. Effects of intratracheal mesenchymal stromal cell therapy during recovery and resolution after ventilator-induced lung injury. *Anesthesiology*. 2013;118(4):924–932. <https://doi.org/10.1097/ALN.0b013e318287ba08>.
- 53 Lv H, Liu Q, Sun Y, et al. Mesenchymal stromal cells ameliorate acute lung injury induced by LPS mainly through stanniocalcin-2 mediating macrophage polarization. *Ann Transl Med*. 2020;8(6):334. <https://doi.org/10.21037/atm.2020.02.105>.
- 54 Lu Z, Chang W, Meng S, et al. Mesenchymal stem cells induce dendritic cell immune tolerance via paracrine hepatocyte growth factor to alleviate acute lung injury. *Stem Cell Res Ther*. 2019;10(1):372. <https://doi.org/10.1186/s13287-019-1488-2>.
- 55 Cao M, Liu H, Dong Y, et al. Mesenchymal stem cells alleviate idiopathic pneumonia syndrome by modulating T cell function through CCR2-CCL2 axis. *Stem Cell Res Ther*. 2021;12(1):378. <https://doi.org/10.1186/s13287-021-02459-7>.
- 56 Zhu J, Feng B, Xu Y, et al. Mesenchymal stem cells alleviate LPS-induced acute lung injury by inhibiting the proinflammatory function of Ly6C(+) CD8(+) T cells. *Cell Death Dis*. 2020;11(10):829. <https://doi.org/10.1038/s41419-020-03036-1>.
- 57 Feng B, Zhu J, Xu Y, et al. Immunosuppressive effects of mesenchymal stem cells on lung B cell gene expression in LPS-induced acute lung injury. *Stem Cell Res Ther*. 2020;11(1):418. <https://doi.org/10.1186/s13287-020-01934-x>.

- 58 Park J, Kim S, Lim H, et al. Therapeutic effects of human mesenchymal stem cell microvesicles in an *ex vivo* perfused human lung injured with severe *E. Coli* pneumonia. *Thorax*. 2019;74(1):43–50. <https://doi.org/10.1136/thoraxjnl-2018-211576>.
- 59 Morrison TJ, Jackson MV, Cunningham EK, et al. Mesenchymal stromal cells modulate macrophages in clinically relevant lung injury models by extracellular vesicle mitochondrial transfer. *Am J Respir Crit Care Med*. 2017;196(10):1275–1286. <https://doi.org/10.1164/rccm.201701-0170OC>.
- 60 Ulich TR, Yi ES, Longmuir K, et al. Keratinocyte growth factor is a growth factor for type II pneumocytes *in vivo*. *J Clin Invest*. 1994;93(3):1298–1306. <https://doi.org/10.1172/JCI117086>.
- 61 Yano T, Mason RJ, Pan T, Deterding RR, Nielsen LD, Shannon JM. KGF regulates pulmonary epithelial proliferation and surfactant protein gene expression in adult rat lung. *Am J Physiol Lung Cell Mol Physiol*. 2000;279(6):L1146–L1158. <https://doi.org/10.1152/ajplung.2000.279.6.L1146>.
- 62 Fang X, Neyrinck AP, Matthay MA, Lee JW. Allogeneic human mesenchymal stem cells restore epithelial protein permeability in cultured human alveolar type II cells by secretion of angiopoietin-1. *J Biol Chem*. 2010;285(34):26211–26222. <https://doi.org/10.1074/jbc.M110.119917>.
- 63 Oswari J, Matthay MA, Margulies SS. Keratinocyte growth factor reduces alveolar epithelial susceptibility to *in vitro* mechanical deformation. *Am J Physiol Lung Cell Mol Physiol*. 2001;281(5):L1068–L1077. <https://doi.org/10.1152/ajplung.2001.281.5.L1068>.
- 64 Guery BP, Mason CM, Dobard EP, Beaucaire G, Summer WR, Nelson S. Keratinocyte growth factor increases transalveolar sodium reabsorption in normal and injured rat lungs. *Am J Respir Crit Care Med*. 1997;155(5):1777–1784. <https://doi.org/10.1164/ajrccm.155.5.9154891>.
- 65 Yang Y, Chen QH, Liu AR, Xu XP, Han JB, Qiu HB. Synergism of MSC-secreted HGF and VEGF in stabilising endothelial barrier function upon lipopolysaccharide stimulation via the Rac1 pathway. *Stem Cell Res Ther*. 2015;6:250. <https://doi.org/10.1186/s13287-015-0257-0>.
- 66 Cho CH, Kim KE, Byun J, et al. Long-term and sustained COMP-ANG1 induces long-lasting vascular enlargement and enhanced blood flow. *Circ Res*. 2005;97(1):86–94. <https://doi.org/10.1161/01.RES.0000174093.64855.a6>.
- 67 Korhonen EA, Lampinen A, Giri H, et al. Tier controls angiopoietin function in vascular remodeling and inflammation. *J Clin Invest*. 2016;126(9):3495–3510. <https://doi.org/10.1172/JCI84923>.
- 68 Zakaria DM, Zahran NM, Arafa S, Mehanna RA, Abdel-Moneim RA. Histological and physiological studies of the effect of bone marrow-derived mesenchymal stem cells on bleomycin induced lung fibrosis in adult albino rats. *Tissue Eng Regen Med*. 2021;18(1):127–141. <https://doi.org/10.1007/s13770-020-00294-0>.
- 69 Ortiz LA, Gambelli F, McBride C, et al. Mesenchymal stem cell engraftment in lung is enhanced in response to bleomycin exposure and ameliorates its fibrotic effects. *Proc Natl Acad Sci U S A*. 2003;100(14):8407–8411. <https://doi.org/10.1073/pnas.1432929100>.
- 70 Ono M, Ohkouchi S, Kanehira M, et al. Mesenchymal stem cells correct inappropriate epithelial-mesenchyme relation in pulmonary fibrosis using stanniocalcin-1. *Mol Ther*. 2015;23(3):549–560. <https://doi.org/10.1038/mt.2014.217>.
- 71 Zheng G, Huang L, Tong H, et al. Treatment of acute respiratory distress syndrome with allogeneic adipose-derived mesenchymal stem cells: a randomized, placebo-controlled pilot study. *Respir Res*. 2014;15(1):39. <https://doi.org/10.1186/1465-9921-15-39>.
- 72 Simonson OE, Mougiakakos D, Heldring N, et al. *In vivo* effects of mesenchymal stromal cells in two patients with severe acute respiratory distress syndrome. *Stem Cells Transl Med*. 2015;4(10):1199–1213. <https://doi.org/10.5966/sctm.2015-0021>.
- 73 Yip HK, Fang WF, Li YC, et al. Human umbilical cord-derived mesenchymal stem cells for acute respiratory distress syndrome. *Crit Care Med*. 2020;48(5):e391–e399. <https://doi.org/10.1097/CCM.0000000000004285>.
- 74 Wilson JG, Liu KD, Zhuo H, et al. Mesenchymal stem (stromal) cells for treatment of ARDS: a phase 1 clinical trial. *Lancet Respir Med*. 2015;3(1):24–32. [https://doi.org/10.1016/S2213-2600\(14\)70291-7](https://doi.org/10.1016/S2213-2600(14)70291-7).
- 75 Matthay MA, Calfee CS, Zhuo H, et al. Treatment with allogeneic mesenchymal stromal cells for moderate to severe acute respiratory distress syndrome (START study): a randomised phase 2a safety trial. *Lancet Respir Med*. 2019;7(2):154–162. [https://doi.org/10.1016/S2213-2600\(18\)30418-1](https://doi.org/10.1016/S2213-2600(18)30418-1).
- 76 Gorman E, Shankar-Hari M, Hopkins P, et al. Repair of acute respiratory distress syndrome by stromal cell administration (REALIST) trial: a phase 1 trial. *EClinicalMedicine*. 2021;4(1):101167. <https://doi.org/10.1016/j.eclinm.2021.101167>.
- 77 Wick KD, Leligdowicz A, Zhuo H, Ware LB, Matthay MA. Mesenchymal stromal cells reduce evidence of lung injury in patients with ARDS. *JCI Insight*. 2021;6(12):e148983. <https://doi.org/10.1172/jci.insight.148983>.
- 78 Zanirati G, Provenzi L, Libermann LL, et al. Stem cell-based therapy for COVID-19 and ARDS: a systematic review. *NPJ Regen Med*. 2021;6(1):73. <https://doi.org/10.1038/s41536-021-00181-9>.
- 79 Meng F, Xu R, Wang S, et al. Human umbilical cord-derived mesenchymal stem cell therapy in patients with COVID-19: a phase 1 clinical trial. *Signal Transduct Target Ther*. 2020;5(1):172. <https://doi.org/10.1038/s41392-020-00286-5>.
- 80 Leng Z, Zhu R, Hou W, et al. Transplantation of ACE2(-) mesenchymal stem cells improves the outcome of patients with COVID-19 pneumonia. *Aging Dis*. 2020;11(2):216–228. <https://doi.org/10.14336/AD.2020.0228>.
- 81 Tang L, Jiang Y, Zhu M, et al. Clinical study using mesenchymal stem cells for the treatment of patients with severe COVID-19. *Front Med*. 2020;14(5):664–673. <https://doi.org/10.1007/s11684-020-0810-9>.
- 82 Xu X, Jiang W, Chen L, et al. Evaluation of the safety and efficacy of using human menstrual blood-derived mesenchymal stromal cells in treating severe and critically ill COVID-19 patients: an exploratory clinical trial. *Clin Transl Med*. 2021;11(2):e297. <https://doi.org/10.1002/ctm2.297>.
- 83 Shu L, Niu C, Li R, et al. Treatment of severe COVID-19 with human umbilical cord mesenchymal stem cells. *Stem Cell Res Ther*. 2020;11(1):361. <https://doi.org/10.1186/s13287-020-01875-5>.
- 84 Guo Z, Chen Y, Luo X, He X, Zhang Y, Wang J. Administration of umbilical cord mesenchymal stem cells in patients with severe COVID-19 pneumonia. *Crit Care*. 2020;24(1):420. <https://doi.org/10.1186/s13054-020-03142-8>.
- 85 Peng H, Gong T, Huang X, et al. A synergistic role of convalescent plasma and mesenchymal stem cells in the treatment of severely ill COVID-19 patients: a clinical case report. *Stem Cell Res Ther*. 2020;11(1):291. <https://doi.org/10.1186/s13287-020-01802-8>.
- 86 Hashemian SR, Aliannejad R, Zarrabi M, et al. Mesenchymal stem cells derived from perinatal tissues for treatment of critically ill COVID-19-induced ARDS patients: a case series. *Stem Cell Res Ther*. 2021;12(1):91. <https://doi.org/10.1186/s13287-021-02165-4>.
- 87 Wu J, Zhou X, Tan Y, et al. Phase 1 trial for treatment of COVID-19 patients with pulmonary fibrosis using hESC-IMRCs. *Cell Prolif*. 2020;53(12):e12944. <https://doi.org/10.1111/cpr.12944>.
- 88 Feng Y, Huang J, Wu J, et al. Safety and feasibility of umbilical cord mesenchymal stem cells in patients with COVID-19 pneumonia: a pilot study. *Cell Prolif*. 2020;53(12):e12947. <https://doi.org/10.1111/cpr.12947>.
- 89 Iglesias M, Butrón P, Torre-Villalvazo I, et al. Mesenchymal stem cells for the compassionate treatment of severe acute respiratory distress syndrome due to COVID 19. *Aging Dis*. 2021;12(2):360–370. <https://doi.org/10.14336/AD.2020.1218>.
- 90 Wu J, Hu Z, Wang L, et al. First case of COVID-19 infused with hESC derived immunity- and matrix-regulatory cells. *Cell Prolif*. 2020;53(12):e12943. <https://doi.org/10.1111/cpr.12943>.
- 91 Shi L, Huang H, Lu X, et al. Effect of human umbilical cord-derived mesenchymal stem cells on lung damage in severe COVID-19 patients: a randomized, double-blind, placebo-controlled phase 2 trial. *Signal Transduct Target Ther*. 2021;6(1):58. <https://doi.org/10.1038/s41392-021-00488-5>.
- 92 Dilogo IH, Aditianihsih D, Sugiarto A, et al. Umbilical cord mesenchymal stromal cells as critical COVID-19 adjuvant therapy: a randomized controlled trial. *Stem Cells Transl Med*. 2021;10(9):1279–1287. <https://doi.org/10.1002/sctm.21-0046>.
- 93 Adas G, Cukurova Z, Yasar KK, et al. The systematic effect of mesenchymal stem cell therapy in critical COVID-19 patients: a prospective double controlled trial. *Cell Transplant*. 2021;30. <https://doi.org/10.1177/09636897211024942>.
- 94 Lanzoni G, Linetsky E, Correa D, et al. Umbilical cord mesenchymal stem cells for COVID-19 acute respiratory distress syndrome: a double-blind, phase 1/2a, randomized controlled trial. *Stem Cells Transl Med*. 2021;10(5):660–673. <https://doi.org/10.1002/sctm.20-0472>.
- 95 Häberle H, Magunia H, Lang P, et al. Mesenchymal stem cell therapy for severe COVID-19 ARDS. *J Intensive Care Med*. 2021;36(6):681–688. <https://doi.org/10.1177/0885066621997365>.

- 96 EN O, Pekkoc-Uyanik KC, Alpaydin N, Gulay GR, Simsek M. Clinical experience on umbilical cord mesenchymal stem cell treatment in 210 severe and critical COVID-19 cases in Turkey. *Stem Cell Rev Rep*. 2021;17(5):1917–1925. <https://doi.org/10.1007/s12015-021-10214-x>.
- 97 Sánchez-Guijo F, García-Arranz M, López-Parra M, et al. Adipose-derived mesenchymal stromal cells for the treatment of patients with severe SARS-CoV-2 pneumonia requiring mechanical ventilation. A proof of concept study. *EClinicalMedicine*. 2020;25: 100454. <https://doi.org/10.1016/j.eclinm.2020.100454>.
- 98 Zhu R, Yan T, Feng Y, et al. Mesenchymal stem cell treatment improves outcome of COVID-19 patients via multiple immunomodulatory mechanisms. *Cell Res*. 2021;1–19. <https://doi.org/10.1038/s41422-021-00573-y>.
- 99 Saleh M, Vaezi AA, Aliannejad R, et al. Cell therapy in patients with COVID-19 using Wharton's jelly mesenchymal stem cells: a phase I clinical trial. *Stem Cell Res Ther*. 2021;12(1):410. <https://doi.org/10.1186/s13287-021-02483-7>.
- 100 Liang B, Chen J, Li T, et al. Clinical remission of a critically ill COVID-19 patient treated by human umbilical cord mesenchymal stem cells: a case report. *Medicine*. 2020;99(31):e21429. <https://doi.org/10.1097/MD.00000000000021429>. (Baltimore).
- 101 Tao J, Nie Y, Wu H, et al. Umbilical cord blood-derived mesenchymal stem cells in treating a critically ill COVID-19 patient. *J Infect Dev Ctries*. 2020;14(10):1138–1145. <https://doi.org/10.3855/jidc.13081>.
- 102 Zhang Y, Ding J, Ren S, et al. Intravenous infusion of human umbilical cord Wharton's jelly-derived mesenchymal stem cells as a potential treatment for patients with COVID-19 pneumonia. *Stem Cell Res Ther*. 2020;11(1):207. <https://doi.org/10.1186/s13287-020-01725-4>.
- 103 Chen X, Shan Y, Wen Y, Sun J, Du H. Mesenchymal stem cell therapy in severe COVID-19: a retrospective study of short-term treatment efficacy and side effects. *J Infect*. 2020;81(4):647–679. <https://doi.org/10.1016/j.jinf.2020.05.020>.
- 104 Shi L, Yuan X, Yao W, et al. Human mesenchymal stem cells treatment for severe COVID-19: 1-Year follow-up results of a randomized, double-blind, placebo-controlled trial. *Ebiomedicine*. 2021;75: 103789. <https://doi.org/10.1016/j.ebiom.2021.103789>.
- 105 Sengupta V, Sengupta S, Lazo A, Woods P, Nolan A, Bremer N. Exosomes derived from bone marrow mesenchymal stem cells as treatment for severe COVID-19. *Stem Cells Dev*. 2020;29(12):747–754. <https://doi.org/10.1089/scd.2020.0080>.
- 106 Xu R, Shi L, Xie W, et al. Diagnosis and treatment guidelines for mesenchymal stem cell therapy for coronavirus disease 2019 (Beijing, 2021). *Infect Dis Immun*. 2021;1(2).
- 107 He X, Hong W, Yang J, et al. Spontaneous apoptosis of cells in therapeutic stem cell preparation exert immunomodulatory effects through release of phosphatidylserine. *Signal Transduct Target Ther*. 2021;6(1):270. <https://doi.org/10.1038/s41392-021-00688-z>.
- 108 Xie Y, Liu S, Wang L, et al. Individual heterogeneity screened umbilical cord-derived mesenchymal stromal cells with high Treg promotion demonstrate improved recovery of mouse liver fibrosis. *Stem Cell Res Ther*. 2021;12(1):359. <https://doi.org/10.1186/s13287-021-02430-6>.
- 109 Tai C, Wang L, Xie Y, Gao T, Huang F, Wang B. Analysis of key distinct biological characteristics of human Placenta-Derived mesenchymal stromal cells and individual heterogeneity attributing to donors. *Cells Tissues Organs*. 2021;210(1):45–57. <https://doi.org/10.1159/000513038>.