



Published in final edited form as:

Arch Immunol Ther Exp (Warsz). ; 70(1): 9. doi:10.1007/s00005-022-00648-7.

Recombinant Human Lactoferrin Reduces Inflammation and Increases Fluoroquinolone Penetration to Primary Granulomas during Mycobacterial Infection of C57Bl/6 Mice

Thao K.T. Nguyen^{1,2}, Zainab Niaz¹, Marian L. Kruzel¹, Jeffrey K. Actor¹

¹Department of Pathology and Laboratory Medicine, UTHealth McGovern Medical School, Houston, TX, USA

²The University of Texas MD Anderson Cancer Center – UTHealth Graduate School of Biomedical Sciences, Houston, TX 77030, USA

Abstract

Infection with *Mycobacterium tuberculosis* (*Mtb*) results in the primary formation of a densely packed inflammatory foci that limits entry of therapeutic agents into pulmonary sites where organisms reside. No current therapeutic regimens exist that modulate host immune responses to permit increased drug penetration to regions of pathological damage during tuberculosis disease. Lactoferrin is a natural iron-binding protein previously demonstrated to modulate inflammation and granuloma cohesiveness, while maintaining control of pathogenic burden. Studies were designed to examine recombinant human lactoferrin (rHLF) to modulate histological progression of *Mtb*-induced pathology in a non-necrotic model using C57Bl/6 mice. The rHLF was oral administered at times corresponding to initiation of primary granulomatous response, or during granuloma maintenance. Treatment with rHLF demonstrated significant reduction in size of primary inflammatory foci following *Mtb* challenge, and permitted penetration of ofloxacin fluoroquinolone therapeutic to sites of pathological disruption where activated (foamy) macrophages reside. Increased drug penetration was accompanied by retention of endothelial cell integrity. Immunohistochemistry revealed altered patterns of M1-like and M2-like phenotypic cell localization post infectious challenge, with increased presence of M2-like markers found evenly distributed throughout regions of pulmonary inflammatory foci in rHLF treated mice.

Keywords

Lactoferrin; Mycobacterium tuberculosis; *Mtb*; Granuloma; Immunopathology; M1/M2 Phenotype; Tuberculosis

Use of this AM is subject to the publisher's embargo period and AM terms of use. Under no circumstances may this AM be shared or distributed under a Creative Commons or other form of open access license, nor may it be reformatted or enhanced, whether by the Author or third parties. See here for Springer Nature's terms of use for AM versions of subscription articles: <https://www.springernature.com/gp/open-research/policies/accepted-manuscript-terms>

Corresponding Author: Jeffrey K. Actor, Ph.D., Professor, Department of Pathology and Laboratory Medicine, MSB 2.214, McGovern Medical School, 6431 Fannin, Houston, TX 77030, USA, Jeffrey.K.Actor@uth.tmc.edu.

Publisher's Disclaimer: This AM is a PDF file of the manuscript accepted for publication after peer review, when applicable, but does not reflect post-acceptance improvements, or any corrections.

Declaration of Competing Interests/Disclosures: None declared.

INTRODUCTION

Despite world-wide extensive research and eradicating efforts, *Mycobacterium tuberculosis* (*Mtb*) remains a major infectious pathogen to the human population with approximately 10.0 million infected cases and 1.5 million deaths reported in 2020 globally (World Health Organization 2020). The initial host-pathogen interaction and complex immunological responses culminate in an inflammatory pathology within pulmonary tissue, characterized as a primary granulomatous disease (Adams 1976; Hunter et al. 2006a). These granulomas and associated lesions obstruct normal pulmonary functions. Active research investigates how *Mtb* directly induces the granulomatous response, focusing on areas of pathological damage and bacterial burden that are most severe during primary infection.

Mtb associated factors are involved in the recognition and activation of host cells within the bronchial regions, leading to uptake by alveolar macrophages (Akira et al. 2006). This interaction triggers a series of immune responses via the production and release of cytokines by infected and responding cells (Hossain and Norazmi 2013). For example, trehalose 6'6-dimycolate (TDM), an abundant mycobacterial mycolic acid, directly triggers a granulomatous response (Hunter et al. 2006a; Nguyen et al. 2020) that resembles early (primary) infection pathology. TDM-activated macrophages release pro-inflammatory mediators, such as TNF- α and IL-1- β , along with additional chemotactic factors, to further recruit immune cells to areas of infection (Khader et al. 2009; Monin and Khader 2014; Welsh et al. 2008). Over time, additional recruited immune cells participate in formation of organized, sphere-shaped primary inflammatory structures (Martin et al. 2016). There exists a balance between host and organism; the granulomas formed during active *Mtb* infection contain and limit bacterial dissemination (Russell 2007), yet organisms have adopted mechanisms to survive and grow inside macrophage host cells. Indeed, recruited naïve macrophages can become new potential sites for *Mtb* to shelter and replicate (Davis and Ramakrishnan 2009).

Clinically, the physical nature of the primary granuloma also limits penetration of antimycobacterial therapeutics; as granulomas mature, reduction in vascularization limits drug delivery to within granulomas where large populations of *Mtb* may reside (Dartois 2014; Driver et al. 2012; Paige and Bishai 2010). This is not dissimilar to constraints seen in drug delivery to solid tumors (Minchinton and Tannock 2006). The physical nature of the host immune response therefore represents a challenge in treating *Mtb*-infected patients – it contributes to prolongation of treatment while permitting a small population of *Mtb* to escape elimination. The lack of complete penetration of antimycobacterial agents also indirectly increases risk of developing antibiotic-resistance (Kaplan et al. 2003, Schito et al. 2015, Sotgiu et al. 2015). Therefore, multiple lines of investigation currently are aimed at improving drug delivery through focused targeting that either manipulates the cohesiveness of the granuloma structure, or alters immune responses to *Mtb* during granuloma development (Kaplan 2020). These studies typically use models to mimic innate (primary) infection, or models with development of lesions with liquefactive necrosis (Driver et al. 2012).

Novel approaches to *Mtb* treatment include host-directed therapies (Ndlovu and Marakalala 2016) which are focused in two major fronts; the first augments immune response using immune-based treatments (Vilaplana et al. 2013), while the second alters the resultant immunopathology (Nguyen et al. 2021). Examples of immune-based treatment methods include agents that target the macro-autophagic compartment, such as vitamin D and retinoic acid, which increase phagocytosis to enhance the lysosomal degradative processes inside macrophages (Estrella et al. 2011). Ibuprofen, an anti-inflammatory drug, functions as an adjunct treatment by facilitating pyrazinamide to alleviate pathological damage in the lungs (Byrne et al. 2007; Eisen et al. 2013; Vilaplana et al. 2013). These immune-based approaches have significant potential, demonstrating superior disease outcomes in clinical trial results (Hayford et al. 2020; Zhang et al. 2018). While promising, the obstacle in drug delivery to macrophages inside established granulomas persists (Gupta et al. 2016). Therefore, therapies that target pathologies, in addition to antimycobacterial function, may subsequently be more useful as clinical tools. A well-known example of the immunopathological alteration approach is blocking excess TNF- α , a key pro-inflammatory cytokine required for granuloma formation (Flynn et al. 1995; Plessner et al. 2007). TNF- α inhibitors have been used to treat other inflammatory diseases effectively (Lv et al. 2014; Skerry et al. 2012; Zhang et al. 2021). However, its current downside is that TNF- α inhibitors can lead to increased bacterial dissemination; abolishing the granuloma containment modality runs the risk of *Mtb* reactivation (Chakravarty et al. 2008; Fernandez-Ruiz and Aguado 2018; Keane et al. 2001; Wallis et al. 2004).

Lactoferrin, a glycoprotein known for its ability to bind iron, has been extensively studied for its role as an immune modulator in host defense in disease models (Kruzel et al. 2017; Rosa et al. 2017). Lactoferrin falls into the category of “immune modulators”, and has been identified to reduce pathology of the tuberculoid granuloma (Hwang et al. 2017). As a modulator, lactoferrin demonstrates efficacy to boost immune memory responses in vaccine models, while reducing pro-inflammatory response in lipopolysaccharide exposed mouse models (Crouch et al. 1992; Hwang et al. 2009a, 2011; Kruzel et al. 2002; Legrand 2012). Relative to the study described in this report, bovine lactoferrin significantly reduced inflammatory pathology in *Mtb*-infected mice (Welsh et al. 2011). Both human and bovine lactoferrins were also shown effective to limit inflammation in a non-infectious TDM-induced granuloma mouse model (Nguyen et al. 2021; Welsh et al. 2010).

Lactoferrin treatment reduced the M1 phenotypic response to TDM, and limits pathological damage in murine lungs (Nguyen et al. 2020, 2021). The lactoferrin treatment also significantly increased penetration of a second-line anti-mycobacterial agent fluoroquinolone into TDM-induced granulomas (Nguyen et al. 2021). Such anti-inflammatory response in the lactoferrin-treated mice, along with other data showing lactoferrin reduces pro-inflammatory phenotypes in macrophages (Hwang et al. 2016; Wisgrill et al. 2018), suggests that lactoferrin had a modulating pro-inflammatory response on granulomas. The studies presented here extend these observations to examine if lactoferrin can modulate primary granuloma permeability and drug distribution during active *Mtb* infection. Here, the effect of lactoferrin on the permeability of primary granulomas (non-necrotic) to a fluoroquinolone is examined in the *Mtb*-infected C57Bl/6 mouse model, when lactoferrin is administered as a prophylactic (defined as when given prior to

granuloma formation) or therapeutic (defined as when given after granuloma establishment) intervention. These experiments shed light on mechanisms underlying changes to the early *Mtb* pathological events due to lactoferrin adjunct treatment.

MATERIALS AND METHODS

Mice

Five-week-old female C57BL/6 mice were purchased from Envigo (Houston, TX) with 18–20 g initial body mass. Eight to ten mice were used per group, per time each point indicated. All *Mtb* infections occurred in biosafety level 3 facilities under the University of Texas Health Science Center at Houston institutional guidelines (IBC-18-014), with approval from the animal ethics committee (AWC-17-0089).

Recombinant Human Lactoferrin and Ofloxacin, and Delivery to Mice

CHO-expressed recombinant human lactoferrin (rHLF; >98% purity; <10% iron saturated; <0.5 EU·mg⁻¹; Cat# LFH-101) was kindly provided as lyophilized powder by PharmaReview Corporation (Houston, TX) (Choi et al. 2008; Kruzel et al. 2021). The rHLF was reconstituted in dH₂O to a concentration of 10 mg·mL⁻¹. From day 14 to day 28 after *Mtb* infection, mice were given 1 mg·(100 μL)⁻¹·mouse⁻¹ of rHLF by oral gavage every other day as prophylactic treatment, similar to reported use in models of mycobacterial granulomatous responses (Hwang et al. 2017; Welsh et al. 2010, 2011). One mg/ml was found to be more productive than a 100 μg/ml dose (Fig. S.1). From day 21 to day 28 after *Mtb* infection, another group of mice were given 1 mg·(100 μL)⁻¹·mouse⁻¹ of recombinant human lactoferrin by oral gavage every other day as therapeutic treatment. Ofloxacin (Sigma Life Science; O8757-1G) was given at 100 μL of 30 mg·mL⁻¹·mouse⁻¹, solubilized in DMSO and diluted 1:10 with phosphate buffered saline (PBS), was intraperitoneal administered 30 min prior to sacrifice (Batard et al. 2011; Hwang et al. 2008).

Mtb Infection

Aerosol infections were done as previously reported (Hwang et al. 2009c, 2011), using *Mycobacterium tuberculosis*, strain Erdman (TMC 107, American Type Cell Culture). Organisms were cultured in Middlebrook 7H9 broth with 10% supplement (5% bovine serum albumin, 2% dextrose, and 0.5% Tween 20 in distilled water) to log phase. Pelleted bacteria were resuspended in PBS and diluted to 3 × 10⁸ colony forming units (CFU) per ml using McFarland standards. Bacteria were sonicated to disperse aggregates. The bacterial CFUs for each time point, including day 1 post infection, were confirmed by plating serial dilutions on Middlebrook 7H11 agar plates (Hardy Diagnostics, Santa Maria, CA) using the large right lobe of the mouse lung that was weighed and homogenized into 2 mL PBS, which were incubated at 37°C for 3 – 4 weeks. Mice were infected for four weeks, with treatments described above, using the protocol shown in Fig. 1. Bovine lactoferrin, shown to have biological equivalency to the rHLF (Hwang et al. 2017) was used for analysis of CFU dissemination to other organs and to reconfirm bioequivalency in dose response treatments (Fig. S.2).

Histological Assessment

The small left lobe of the mouse lung was collected and fixed in 10% buffered formalin. For histologic analysis, the lung was sectioned (5 μm thick) and stained with hematoxylin and eosin (H&E) and acid-fast staining as per standard procedures (Hwang et al. 2011). The histological assessment of the lung tissue following aerosol infection was done as previously reported (Hwang et al. 2009c). Multiple sections from at least 6 mice per group were analyzed using Motic DSAssistant digital software (version 1.0.7.44; Kowloon Bay, Kowloon, HK) (Nguyen et al. 2021). H&E stained and acid-fast stained slides were viewed by a trained pathologist, with descriptive results obtained in an experimentally blinded manner.

Immunohistochemistry

Fixed lung was embedded in paraffin, sectioned, and stained for immunohistochemical examination, similar to methods described (Hwang et al. 2019), diluted at 1:2000, was performed according to manufacturer's instructions with a modification of 20 min at low pH for antigen retrieval, and visualized using standard HRP techniques and DAB chromogen using Dako reagents (Dako, Agilent, Santa Clara, CA). In a similar manner, M1-like marker CD38 (Invitrogen, ThermoFisher, Cat# 14-0381-02), diluted at 1:1000, M2-like marker CD 206 (Bioss, Cat# bs-4727R), diluted at 1:1000, and endothelial cell marker CD31/PECAM-1 (Cell Signaling, Cat# 77699T) diluted at 1:200 were used for visualization on serial slide sections. Hematoxylin counterstained slides were viewed by a trained pathologist, with descriptive results obtained in an experimentally blinded manner.

Quantitative Assessment of Pulmonary Inflammation

High resolution scanned images of H&E-stained slides were assessed for lung inflammation and granulomas using Motic DSAssistant software (Kowloon Bay, Kowloon, HK), in a two step process using Fiji ImageJ (version 1.52o 23 April 2019, National Institutes of Health, Bethesda, MD) with plugin MorphoLibJ (McQuin et al. 2018), described in part in (Nguyen et al. 2021). Minimum and maximum values for hue, saturation, and brightness were set at: 120, 255; 0, 255; and 0, 255, respectively. Total cell area measurement used a modified equation detailed elsewhere where peak threshold was set at 164 (Schneider et al. 2012) (Fig. S.3). Values were averaged within treatment groups and normalized to non-treated controls.

H&E-stained slides were also used to capture photos of granulomas under the Olympus BX51 microscope using the Nuance Cri Multispectral Imaging System FX (PerkinElmer). The granuloma section was first identified and captured under H&E brightfield scope, then ofloxacin's fluorescent signals were captured under 40 \times lens with FITC filter (emission restriction set between 540–560 nm) after 120 ms of exposure. All microscopic settings and factors were maintained throughout the photo taking process with image files having the same dimensions of 1392 \times 1040 with 72 dpi resolution. For each granuloma, the total fluorescent area or ofloxacin absorption area and the total granuloma area were measured in pixel units using CellProfiler (software version 3.1.5) pipeline algorithm (McQuin et al. 2018), with described modifications (Diem et al. 2015). The measurement on the selected lung section is reported as percent area that represents the ratio of ofloxacin absorption over

the total granuloma area, based on original observations of (Batard et al. 2011). All data was graphed and statistical analyzed in GraphPad Prism (version 5.03).

Statistical Analysis

Data obtained was compared across groups then analyzed using a paired Student *t* test or one-way ANOVA with a Tukey post-hoc test; differences between means were considered statistically significant at a value of $p < 0.05$. Data are presented are a combination of 2–3 experimental repeats. Each experiment incorporated 8–10 mice per group.

RESULTS

Oral Administration of Recombinant Human Lactoferrin Alleviates Pulmonary Inflammation Post *Mtb* Infection

Previous studies using bovine derived lactoferrin given in drinking water demonstrated that continuous-access oral delivery during *Mycobacterium tuberculosis* infection could reduce inflammation-related primary granulomatous pathology in mice (Welsh et al. 2011). A defined protocol was adopted to determine if rHlf would initiate a similar protective response (Fig. 1). Mice were aerosol infected with *Mtb*, strain Erdman, and rHLF was given by oral gavage beginning either prior to (prophylactic), or post (therapeutic) expected initiation of granulomatous responses. Histological assessment (Fig. 2 and Fig. S.1) revealed marked reduction of pulmonary inflammation in both the prophylactic and therapeutic rHLF treated groups, in a similar manner to that previously reported with the bovine LF treatment. Specifically, the rHLF treatments demonstrated reduced complexity of granulomatous responses, smaller foci of inflammation, with less cellular accumulation and density in areas of inflammation. Quantitative measurement of inflammation confirmed histological reduction in pathology due to the rHLF treatments.

Whole right lobes of mouse lung were collected at four weeks post infections and processed for quantitative assessment of primary granulomatous response. Serial sections of H&E stained tissue sections were high resolution scanned to assess area occupied by inflammation (Fig. 3). The *Mtb* infected group had the largest occluded regions, with $24.79\% \pm 4.2\%$ relative area occupied by inflammatory response. Both lactoferrin treatment modalities resulted in reduction of pathology. The prophylactic treatment resulted in reduction of granuloma area to $21.42\% \pm 4.1\%$ of lung sections observed, while the therapeutic treatment significantly reduced occupied space to $17.36\% \pm 4.9\%$ ($p < 0.001$). Of interest, there was no change in CFU in the treated group in lung, liver or spleen tissue at four weeks post infection (Fig. S.2), suggesting that (1) the short term administration of lactoferrin did not alter pathogenic burden, and (2) the alteration due to treatments did not result in significant net dissemination to other tissues.

Increased Penetration of Fluoroquinolone to Inflammatory Foci after Treatment with rHLF

The altered pathology of less dense granulomatous structures raised the hypothesis that vascular structure would be maintained in the rHLF treated groups, which could subsequently result in enhanced penetration of mycobacterial therapeutic agents to within regions of inflammation. To test this, the naturally fluorescent fluoroquinolone, ofloxacin,

was intravenously administered to mice 30 min prior to sacrifice at the 28 days post infection time point. Figure 4 reveals the penetration patterns of ofloxacin to within regions of pulmonary granulomatous response. The histologically dense inflammatory response in the *Mtb* alone group did not permit penetration of the fluoroquinolone, with little to no signal entering granulomatous foci. In contrast, both the prophylactic and the therapeutic rHLF treatment protocols resulted in granulomas that were permissible to ofloxacin penetration. Assessment of serial sections by high resolution scanning revealed significant differences between groups (Fig. 5). The *Mtb* alone infected group had a relative fluorescent distribution of signal $13.76\% \pm 7.5\%$, confirming the relative difficulty in penetration of ofloxacin to within inflammatory regions. For comparison, the relative fluorescence of normal mouse lung had an average of $90.25\% \pm 3.5\%$ penetration, reflecting antibiotic distribution to the lung at 30 min post delivery. Of interest, in alignment with visual observations described above, both rHLF treated groups demonstrated elevated ofloxacin penetration within granulomas. The prophylactic treatment permitted relative increases to $21.68\% \pm 14.1\%$, and the therapeutic treatment showed significant increase at $47.15\% \pm 14.9\%$ presence of fluorescent signal. Furthermore, high power observation of signal revealed accumulation of ofloxacin correlating with presence of activated foamy macrophages (Fig. 6).

rHLF Treatment Correlates with Retention of Vascular Structure (Endothelial Cell Integrity) in Regions of Pulmonary Inflammation

The *Mtb* alone infected group demonstrated responses consistent with pulmonary disruption to alveolar structure and associated vascular tissue. To further examine the effect of the rHLF treatment on vascular structure, histological sections were stained with PECAM-1. Granulomas in the *Mtb* alone group demonstrated collapsed alveoli with central accumulation of immune cells in regions that disrupt blood distribution (Fig. 7). Central foci in these mice were devoid of staining, representing reduced vascularization to regions where organisms are expected to reside. In contrast, the rHLF treated mice demonstrated retention of vascularized structures within areas of inflammation, which also corresponded in matched sections to regions demonstrating ofloxacin penetration.

Altered Localization of M1/M2-Like Macrophages in Primary Granulomas Following rHLF Treatment

Recent reports demonstrated altered distribution of macrophage populations within primary granulomas, which may dictate pathological outcomes (Pisu et al. 2020a, 2020b). As a preliminary investigation into this topic, sections were immunohistochemically stained for general M1-like and M2-like antigens. Figure 8 depicts immunohistochemical staining for the M1-like marker CD38 and for the M2-like marker CD206. The non-treated *Mtb* infected lungs revealed a concentrated pattern of staining demonstrating high presence of both M1-like and M2-like macrophages, primarily cuffing vascular regions surrounding regions of inflammation. Limited numbers of both the M1-like and the M2-like cells were visible within the granuloma itself in the *Mtb* alone infected group. In contrast, the rHLF treated mice exhibited a only small number of M1-like cells that were lightly stained, residing inside the granulomatous response. Of note, the M2-like staining pattern in the lactoferrin treated groups showed M2-like macrophages which were evenly and diffusely distributed throughout the primary granulomatous pathology.

DISCUSSION

This is the first report that recombinant human lactoferrin can modulate the early granulomatous response during primary *Mtb* infection in mice, in a manner nearly identical to that reported for bovine lactoferrin (Welsh et al. 2011). The results presented indicate utility for human lactoferrin administered orally as a therapeutic approach to limit pathological damage during primary granuloma development. The intervention led to increased penetration of ofloxacin to regions where *Mtb* typically reside. Our observations in this model are in line with studies examining induced granulomas using purified mycobacterial mycolic acid TDM (Nguyen et al. 2021), where administration of rHLF achieved inflammation reduction in the lungs and greater ofloxacin distribution throughout granulomatous structures after treatment.

Lactoferrin-induced modulation in inflammatory response within mycobacterial-induced granulomas is likely the result of two concurrent events; that of significantly less proinflammatory cytokine production and reduction in recruited M1-like macrophages (Nguyen et al. 2021). Alveolar macrophages uptake *Mtb* during primary infection and become essentially the key cell phenotype to recruit and activate additional naïve macrophages to sites of infection (Akira et al. 2006; McClean and Tobin 2016). Classically activated macrophages polarize to the M1-like phenotype and exhibit functional phagocytosis and killing of bacteria. As a pathological by-product, they initiate immune cell recruitment via secreted proinflammatory cytokines to aid in the establishment of granulomatous tissue structures. Temporally, the subsequent recruitment and introduction of M2-like macrophages allows immune modification of the aggressive inflammatory response, permitting an immunological “brake” to an overwhelming pathological response (Hunter et al. 2006a; Huang et al. 2015; Marino et al. 2015; Nguyen et al. 2020; Thiriote et al. 2020). While further investigation is required, it is theorized that a balance in the presence of macrophage phenotypes within the granulomatous structures is essential to control pathological mediators by innate immune cells (Pisu et al. 2020a, 2020b), which is critical to regulation of IL-1 β (T cell activation and migration) (Schmitz et al. 2005) and TNF- α (vasodilatation and leukocyte adhesion to epithelium) (Page et al. 2018) necessary as a response to control bacterial growth.

A major observation of this study was that the second-line mycobacterial therapeutic fluorquinolone was able to penetrate within inflammatory foci. Similar to reported results using the TDM non-infectious model of pathology, significant amount of ofloxacin was found within granulomas following treatment with human lactoferrin, especially, but not limited to, regions of reduced inflammation. Concurrently, the lactoferrin-treated mice exhibited granulomatous responses with maintained vascular structures and open alveolar spaces. Acute inflammation and reactivity during *Mtb* infection occurs primarily peripheral to vascular regions, coinciding with destruction in continuity of endothelial lined blood vessels (Hunter et al. 2018; Hwang et al. 2019). In the experiments presented in this report, lactoferrin treatment allowed greater regions of lung tissue, and alveolar spaces, to remain unobstructed, as evident using the PECAM-1 endothelial surface marker. This coincided with observational retention of blood vessels within regions of pathology. The lessened pathological damage in the lactoferrin treated animals, along with significant maintenance of

vascular architecture, likely permitted ofloxacin transport inside of granulomas, as evident by the drug's fluorescent signal observed within and around the endothelial-lined vessels. Therefore, we hypothesize the promising use of lactoferrin as a safe adjuvant molecule to increase delivery of standard *Mtb* therapeutics, especially in newly exposed cohorts of post-primary status individuals. This may also have the potential to reduce overall treatment times, limit drug sensitivity development, and reduce antibiotic side effects in patients undergoing treatment. At this time, it is certainly recognized that the line of research using fluoroquinolone may not extend to other classes of antimycobacterial agents.

A major concern with current host-directed therapy is the increase of bacterial dissemination during treatment, such as seen when TNF- α blockers are used (Chakravarty et al. 2008; Fernandez-Ruiz and Aguado 2018; Keane et al. 2001; Wallis et al. 2004). It is crucial that the role of established granulomas is maintained throughout the anti-mycobacterial treatment; specifically, the activation of recruited immune cells must be maintained to limit organism spread to other tissues. Our observations (supplemental data) demonstrated no increase in lung CFUs in the treated group and no change in levels of detected organisms in liver and spleen. Such observation aligns with previous studies using bovine lactoferrin in a similar *Mtb* infectious model (Welsh et al. 2011). This shows potential therapeutic application of recombinant human lactoferrin to reduce pathological damage due to infection. And it does so without the major side effect of other host-directed therapies, including the compromising of *Mtb* confinement at the site of initial inflammation.

Lactoferrin immune-modulating effects in our *Mtb* infectious model are consistent with other infectious models reported (Actor et al. 2009; Drago-Serrano et al. 2017; Sienkiewicz et al. 2021). Its utility as a host-directed therapeutic has been demonstrated in other diseases where host inflammation plays a key role in pathology development (Actor et al. 2009; Doursout et al. 2013). Lactoferrin was used successfully in clinical trials a prophylactic to prevent development of enterocolitis and sepsis (Manzoni et al. 2009; Ochoa et al. 2012; Turin et al. 2014). In a similar manner, the prophylactic administration presented here was done prior to the establishment of granulomatous pathology. However, significant results were surprisingly identified when lactoferrin was given as a therapeutic at day 21 post infection at a time after granulomas have initiated in the lungs. Of clinical importance, when lactoferrin was used therapeutically it both decreased overall lung inflammation and increased ofloxacin distribution in granulomas.

Mechanistically, lactoferrin can induce dendritic cell activation and maturation, specifically supporting functions of antigen-presenting cell populations that act as to bridge innate and adaptive immunity (de la Rosa et al. 2008; Hwang and Actor 2009; Hwang et al. 2015; Rascon-Cruz et al. 2021; Spadaro et al. 2014). Lactoferrin can directly enhance the antigen presenting activity and T cell stimulation function in mycobacterial-infected macrophages (Hwang et al. 2009b). Lactoferrin can also modulate T cell activities in multiple ways (Actor et al. 2009). *In vitro*, T cell maturation and expression of T cell ζ -chain, a component of T cell receptor complex involved in receptor signaling pathways, can be increased upon treatment with lactoferrin (Dhennin-Duthille et al. 2000; Frydecka et al. 2002; Zimecki et al. 1991); T cell adhesive molecules involved in cell-to-cell contact are also increased in the presence of lactoferrin (Zimecki et al. 1999). Clearly, lactoferrin can influence T helper cell

polarization (Fischer et al. 2006). Bovine lactoferrin administered orally functions to induce both systemic and mucosal responses (Debbabi et al. 1998), with specific increases in IFN- γ Th1 T cell responses (Takakura et al. 2006) and NK cell activity in mice (Kuhara et al. 2006) likely by increasing IL-12 and related cytokines (Hwang et al. 2007; Kuhara et al. 2006). Together, these suggest possible modulations of T cell activities, combined with enhanced antigen-presenting cell maturation and macrophage recruitment effects, which theoretically shift the overall outcome of granulomas to successful containment of pathogens

Previous observations using the TDM-induced granuloma model demonstrated that lactoferrin could lessen the presence of inflammatory mediators post initiation of pathology (Nguyen et al. 2021), as well as preferentially recruit M2-like cells to granulomas (Nguyen et al. 2021). It is reasonable to infer that similar reductions using human lactoferrin at four weeks post *Mtb* infection would occur, since TDM is significantly released from mycobacteria (Hunter et al. 2006a, 2006b). This also suggests an important mechanistic link between macrophage recruitment and lactoferrin treatment during *Mtb* infection, perhaps via the alteration of surface adhesion markers on monocytes that would permit increased interaction with endothelial cells (Baveye et al. 2000; Yeom et al. 2011). However, it remains unknown if lactoferrin can differentially affect M1-like versus M2-like activities; such experiments have only been explored on classical M1-like activated monocytes (Hwang et al. 2009b; Nguyen et al. 2020). This suggests a premise that macrophage polarization could occur in the presence of lactoferrin, perhaps via the reduction of locally produced proinflammatory cytokines (Kruzel et al. 2002; Mantovani et al. 2004; Nguyen et al. 2021; Thiriot et al. 2020; Wang et al. 2014). And it is unknown at this time if mouse lactoferrin present in the lungs would act in a synergistic manner with the endogenously delivered molecule.

The studies presented here set a foundation to further investigate usage of human lactoferrin in combination with standard treatments for *Mtb* infection. Bacterial dissemination in later time points after infection, and prolonged treatments, should be studied to understand the full potential of its use as a host-directed therapeutic. In reality, it is not clinically feasible to identify individuals prior to development of pathology and granuloma formation. Therefore, more experiments are needed to determine the extent of human lactoferrin on the adaptive immune response when treatments begin after exposure at the primary granuloma establishment and maintenance stages. Furthermore, these studies were done with a focus on the early granulomatous response; experiments should include effects on caseous granulomatous, such as that seen when using the pathologically aggressive C3HeB/FeJ mouse model (Driver et al. 2012). Additional avenues of research should include lactoferrin delivery routes (Sfeir et al. 2004); it may be advantageous for directed aerosol delivery to regions most affected during tuberculosis infection. Overall, a role for clinical utility of human lactoferrin to modify the aggressive immune function during primary *Mtb* infection may exist, which would allow greater efficacy of treatments. In turn, this would potentially reduce standard treatment duration, antibiotic side effects, and overall pathological damage in patients.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

This work was given in part at the 14th International Conference on Lactoferrin Structure, Function and Applications, held in Lima, Peru (2019). This research was performed in part to fulfill requirements for the MS degree from The University of Texas MD Anderson Cancer Center, UTHealth Graduate School of Biomedical Sciences; located within the University of Texas Medical Center in Houston, Texas 77030. We thank Gustavo Ayala, MD, for use of the Nuance Cri Multispectral Imaging System FX. This project was supported in part by NIH Grant 1R42-AI117990.

REFERENCES

- Actor JK, Hwang SA, Kruzel ML (2009) Lactoferrin as a natural immune modulator. *Curr Pharm Des* 15: 956–1973
- Adams DO (1976) The granulomatous inflammatory response. A review. *Am J Pathol* 84:164–192 [PubMed: 937513]
- Akira S, Uematsu S, Takeuchi O (2006) Pathogen recognition and innate immunity. *Cell* 124:783–801 [PubMed: 16497588]
- Batard E, Jamme F, Villette S et al. (2011) Diffusion of ofloxacin in the endocarditis vegetation assessed with synchrotron radiation UV fluorescence microspectroscopy. *PLoS One* 6:e19440
- Baveye S, Elass E, Fernig DG et al. (2000) Human lactoferrin interacts with soluble CD14 and inhibits expression of endothelial adhesion molecules, E-selectin and ICAM-1, induced by the CD14-lipopolysaccharide complex. *Infect Immun* 68:6519–6525 [PubMed: 11083760]
- Byrne ST, Denkin SM, Zhang Y (2007) Aspirin and ibuprofen enhance pyrazinamide treatment of murine tuberculosis. *J Antimicrob Chemother* 59:313–316 [PubMed: 17185297]
- Chakravarty SD, Zhu G, Tsai MC et al. (2008) Tumor necrosis factor blockade in chronic murine tuberculosis enhances granulomatous inflammation and disorganizes granulomas in the lungs. *Infect Immun* 76:916–926 [PubMed: 18212087]
- Choi BK, Actor JK, Rios S et al. (2008) Recombinant human lactoferrin expressed in glycoengineered *Pichia pastoris*: effect of terminal N-acetylneuraminic acid on in vitro secondary humoral immune response. *Glycoconj J* 25:581–593 [PubMed: 18365311]
- Crouch SP, Slater KJ, Fletcher J (1992) Regulation of cytokine release from mononuclear cells by the iron-binding protein lactoferrin. *Blood* 80:235–240 [PubMed: 1535239]
- Dartois V (2014) The path of anti-tuberculosis drugs: from blood to lesions to mycobacterial cells. *Nat Rev Microbiol* 12:159–167 [PubMed: 24487820]
- Davis JM, Ramakrishnan L (2009) The role of the granuloma in expansion and dissemination of early tuberculous infection. *Cell* 136:37–49 [PubMed: 19135887]
- de la Rosa G, Yang D, Tewary P et al. (2008) Lactoferrin acts as an alarmin to promote the recruitment and activation of APCs and antigen-specific immune responses. *J Immunol* 180:6868–6876 [PubMed: 18453607]
- Debbabi H, Dubarry M, Rautureau M et al. (1998) Bovine lactoferrin induces both mucosal and systemic immune response in mice. *J Dairy Res* 65:283–293 [PubMed: 9627847]
- Dhennin-Duthille I, Masson M, Damiens E et al. (2000) Lactoferrin upregulates the expression of CD4 antigen through the stimulation of the mitogen-activated protein kinase in the human lymphoblastic T Jurkat cell line. *J Cell Biochem* 79:583–593 [PubMed: 10996849]
- Diem K, Magaret A, Klock A et al. (2015) Image analysis for accurately counting CD4+ and CD8+ T cells in human tissue. *J Virol Methods* 222:117–121 [PubMed: 26073660]
- Doursout MF, Horton H, Hoang L et al. (2013) Lactoferrin moderates LPS-induced hypotensive response and gut injury in rats. *Int Immunopharmacol* 15:227–231 [PubMed: 23267765]
- Drago-Serrano ME, Campos-Rodriguez R, Carrero JC et al. (2017) Lactoferrin: Balancing Ups and downs of inflammation due to microbial infections. *Int J Mol Sci* 18:501

- Driver ER, Ryan GJ, Hoff DR et al. (2012) Evaluation of a mouse model of necrotic granuloma formation using C3HeB/FeJ mice for testing of drugs against *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother* 56:3181–3195 [PubMed: 22470120]
- Eisen DP, McBryde ES, Walduck A (2013) Low-dose aspirin and ibuprofen's sterilizing effects on *Mycobacterium tuberculosis* suggest safe new adjuvant therapies for tuberculosis. *J Infect Dis* 208:1925–1927
- Estrella JL, Kan-Sutton C, Gong X et al. (2011) A novel in vitro human macrophage model to study the persistence of *Mycobacterium tuberculosis* using vitamin D(3) and retinoic acid activated THP-1 macrophages. *Front Microbiol* 2:67 [PubMed: 21747789]
- Fernandez-Ruiz M, Aguado JM (2018) Risk of infection associated with anti-TNF-alpha therapy. *Expert Rev Anti Infect Ther* 16:939–956 [PubMed: 30388900]
- Fischer R, Debbabi H, Dubarry M et al. (2006) Regulation of physiological and pathological Th1 and Th2 responses by lactoferrin. *Biochem Cell Biol* 84:303–311 [PubMed: 16936801]
- Flynn JL, Goldstein MM, Chan J et al. (1995) Tumor necrosis factor-alpha is required in the protective immune response against *Mycobacterium tuberculosis* in mice. *Immunity* 2:561–572 [PubMed: 7540941]
- Frydecka I, Zimecki M, Bocko D et al. (2002) Lactoferrin-induced up-regulation of zeta (zeta) chain expression in peripheral blood T lymphocytes from cervical cancer patients. *Anticancer Res* 22:1897–1901 [PubMed: 12168890]
- Gupta A, Misra A, Deretic V (2016) Targeted pulmonary delivery of inducers of host macrophage autophagy as a potential host-directed chemotherapy of tuberculosis. *Adv Drug Deliv Rev* 102:10–20 [PubMed: 26829287]
- Hayford FEA, Dolman RC, Blaauw R et al. (2020) The effects of anti-inflammatory agents as host-directed adjunct treatment of tuberculosis in humans: a systematic review and meta-analysis. *Respir Res* 21:223 [PubMed: 32847532]
- Hossain MM, Norazmi MN (2013) Pattern recognition receptors and cytokines in *Mycobacterium tuberculosis* infection—the double-edged sword? *Biomed Res Int* 2013:179174 [PubMed: 24350246]
- Huang Z, Luo Q, Guo Y et al. (2015) *Mycobacterium tuberculosis*-induced polarization of human macrophage orchestrates the formation and development of tuberculous granulomas in vitro. *PLoS One* 10:e0129744 [PubMed: 26091535]
- Hunter RL, Olsen MR, Jagannath C et al. (2006a) Multiple roles of cord factor in the pathogenesis of primary, secondary, and cavitary tuberculosis, including a revised description of the pathology of secondary disease. *Ann Clin Lab Sci* 36:371–386 [PubMed: 17127724]
- Hunter RL, Venkataprasad N, Olsen MR (2006b) The role of trehalose dimycolate (cord factor) on morphology of virulent *M. tuberculosis* in vitro. *Tuberculosis* 86:349–356 [PubMed: 16343989]
- Hunter RL, Actor JK, Hwang SA et al. (2018) Pathogenesis and animal models of post-primary (bronchogenic) tuberculosis, a review. *Pathogens* 7:19
- Hwang SA, Actor JK (2009) Lactoferrin modulation of BCG-infected dendritic cell functions. *Int Immunol* 21:1185–1197 [PubMed: 19692539]
- Hwang SA, Wilk KM, Bangale YA et al. (2007) Lactoferrin modulation of IL-12 and IL-10 response from activated murine leukocytes. *Med Microbiol Immunol* 196:171–180 [PubMed: 17377816]
- Hwang SM, Kim DD, Chung SJ et al. (2008) Delivery of ofloxacin to the lung and alveolar macrophages via hyaluronan microspheres for the treatment of tuberculosis. *J Control Release* 129:100–106 [PubMed: 18538437]
- Hwang SA, Arora R, Kruzel ML et al. (2009a) Lactoferrin enhances efficacy of the BCG vaccine: comparison between two inbred mice strains (C57BL/6 and BALB/c). *Tuberculosis* 89 Suppl 1:S49–54 [PubMed: 20006305]
- Hwang SA, Kruzel ML, Actor JK (2009b) Influence of bovine lactoferrin on expression of presentation molecules on BCG-infected bone marrow derived macrophages. *Biochimie* 91:76–85 [PubMed: 18486627]
- Hwang SA, Wilk K, Kruzel ML et al. (2009c) A novel recombinant human lactoferrin augments the BCG vaccine and protects alveolar integrity upon infection with *Mycobacterium tuberculosis* in mice. *Vaccine* 27:3026–3034 [PubMed: 19428915]

- Hwang SA, Welsh KJ, Boyd S et al. (2011) Comparing efficacy of BCG/lactoferrin primary vaccination versus booster regimen. *Tuberculosis* 91 Suppl 1:S90–95 [PubMed: 22088320]
- Hwang SA, Kruzel ML, Actor JK (2015) Effects of CHO-expressed recombinant lactoferrins on mouse dendritic cell presentation and function. *Innate Immun* 21:553–561 [PubMed: 25537452]
- Hwang SA, Kruzel ML, Actor JK (2016) Recombinant human lactoferrin modulates human PBMC derived macrophage responses to BCG and LPS. *Tuberculosis* 101S:S53–S62 [PubMed: 27727130]
- Hwang SA, Kruzel ML, Actor JK (2017) Oral recombinant human or mouse lactoferrin reduces Mycobacterium tuberculosis TDM induced granulomatous lung pathology. *Biochem Cell Biol* 95:148–154 [PubMed: 28165282]
- Hwang SA, Byerly CD, Actor JK (2019) Mycobacterial trehalose 6,6'-dimycolate induced vascular occlusion is accompanied by subendothelial inflammation. *Tuberculosis* 116S:S118–S122 [PubMed: 31072690]
- Kaplan G (2020) Tuberculosis control in crisis-causes and solutions. *Prog Biophys Mol Biol* 152:6–9 [PubMed: 31626823]
- Kaplan G, Post FA, Moreira AL et al. (2003) Mycobacterium tuberculosis growth at the cavity surface: a microenvironment with failed immunity. *Infect Immun* 71:7099–7108 [PubMed: 14638800]
- Keane J, Gershon S, Wise RP et al. (2001) Tuberculosis associated with infliximab, a tumor necrosis factor alpha-neutralizing agent. *N Engl J Med* 345:1098–1104 [PubMed: 11596589]
- Khader SA, Rangel-Moreno J, Fountain JJ et al. (2009) In a murine tuberculosis model, the absence of homeostatic chemokines delays granuloma formation and protective immunity. *J Immunol* 183:8004–8014 [PubMed: 19933855]
- Kruzel ML, Harari Y, Mailman D et al. (2002) Differential effects of prophylactic, concurrent and therapeutic lactoferrin treatment on LPS-induced inflammatory responses in mice. *Clin Exp Immunol* 130:25–31 [PubMed: 12296849]
- Kruzel ML, Zimecki M, Actor JK (2017) Lactoferrin in a context of inflammation-induced pathology. *Front Immunol* 8:1438 [PubMed: 29163511]
- Kruzel ML, Olszewska P, Pazdrak B et al. (2021) New insights into the systemic effects of oral lactoferrin: transcriptome profiling. *Biochem Cell Biol* 99:47–53 [PubMed: 32476452]
- Kuhara T, Yamauchi K, Tamura Y et al. (2006) Oral administration of lactoferrin increases NK cell activity in mice via increased production of IL-18 and type I IFN in the small intestine. *J Interferon Cytokine Res* 26:489–499 [PubMed: 16800788]
- Legrand D (2012) Lactoferrin, a key molecule in immune and inflammatory processes." *Biochem Cell Biol* 90:252–268 [PubMed: 22136726]
- Lv S, Han M, Yi R et al. (2014) Anti-TNF-alpha therapy for patients with sepsis: a systematic meta-analysis. *Int J Clin Pract* 68:520–528 [PubMed: 24548627]
- Mantovani A, Sica A, Sozzani S et al. (2004) The chemokine system in diverse forms of macrophage activation and polarization. *Trends Immunol* 25:677–686 [PubMed: 15530839]
- Manzoni P, Rinaldi M, Cattani S et al. (2009) Bovine lactoferrin supplementation for prevention of late-onset sepsis in very low-birth-weight neonates: a randomized trial. *JAMA* 302:1421–1428 [PubMed: 19809023]
- Marino S, Cilfone NA, Mattila JT et al. (2015) Macrophage polarization drives granuloma outcome during Mycobacterium tuberculosis infection. *Infect Immun* 83:324–338 [PubMed: 25368116]
- Martin CJ, Carey AF, Fortune SM (2016) A bug's life in the granuloma. *Semin Immunopathol* 38:213–220 [PubMed: 26577238]
- McClellan CM, Tobin DM (2016) Macrophage form, function, and phenotype in mycobacterial infection: lessons from tuberculosis and other diseases. *Pathog Dis* 74:ftw068 [PubMed: 27402783]
- McQuin C, Goodman A, Chernyshev V et al. (2018) CellProfiler 3.0: Next-generation image processing for biology. *PLoS Biol* 16:e2005970 [PubMed: 29969450]
- Minchinton AI, Tannock IF (2006) Drug penetration in solid tumours. *Nat Rev Cancer* 6:583–592 [PubMed: 16862189]

- Monin L, Khader SA (2014) Chemokines in tuberculosis: the good, the bad and the ugly. *Semin Immunol* 26:552–558 [PubMed: 25444549]
- Ndlovu H, Marakalala MJ (2016) Granulomas and inflammation: host-directed therapies for tuberculosis. *Front Immunol* 7:434 [PubMed: 27822210]
- Nguyen TKT, d'Aigle J, China L et al. (2020) Mycobacterial trehalose 6,6'-dimycolate-induced M1-type inflammation. *Am J Pathol* 190:286–294 [PubMed: 31734231]
- Nguyen TKT, Niaz Z, d'Aigle J et al. (2021) Lactoferrin reduces mycobacterial M1-type inflammation induced with trehalose 6,6'-dimycolate and facilitates the entry of fluoroquinolone into granulomas. *Biochem Cell Biol* 99:73–80 [PubMed: 32402212]
- Ochoa TJ, Pezo A, Cruz K et al. (2012) Clinical studies of lactoferrin in children. *Biochem Cell Biol* 90:457–467 [PubMed: 22380791]
- Page MJ, Bester J, Pretorius E (2018) The inflammatory effects of TNF-alpha and complement component 3 on coagulation. *Sci Rep* 8:1812 [PubMed: 29379088]
- Paige C, Bishai WR (2010) Penitentiary or penthouse condo: the tuberculous granuloma from the microbe's point of view. *Cell Microbiol* 12:301–309 [PubMed: 20039878]
- Pisu D, Huang L, Grenier JK et al. (2020a) Dual RNA-Seq of Mtb-infected macrophages in vivo reveals ontologically distinct host-pathogen interactions. *Cell Rep* 30:335–350.e334 [PubMed: 31940480]
- Pisu D, Huang L, Rin BN et al. (2020b) Dual RNA-sequencing of Mycobacterium tuberculosis-infected cells from a murine infection model. *STAR Protoc* 1:100123 [PubMed: 33377017]
- Plessner HL, Lin PL, Kohno T et al. (2007) Neutralization of tumor necrosis factor (TNF) by antibody but not TNF receptor fusion molecule exacerbates chronic murine tuberculosis. *J Infect Dis* 195:1643–1650 [PubMed: 17471434]
- Rascon-Cruz Q, Espinoza-Sanchez EA, Siqueiros-Cendon TS et al. (2021) Lactoferrin: A glycoprotein involved in immunomodulation, anticancer, and antimicrobial processes. *Molecules* 26:205
- Rosa L, Cutone A, Lepanto MS et al. (2017) Lactoferrin: A natural glycoprotein involved in iron and inflammatory homeostasis. *Int J Mol Sci* 18:1985
- Russell DG (2007) Who puts the tubercle in tuberculosis? *Nat Rev Microbiol* 5:39–47 [PubMed: 17160001]
- Schito M, Migliori GB, Fletcher HA et al. (2015) Perspectives on advances in tuberculosis diagnostics, drugs, and vaccines. *Clin Infect Dis* 61 Suppl 3: S102–118
- Schmitz N, Kurrer M, Bachmann MF et al. (2005) Interleukin-1 is responsible for acute lung immunopathology but increases survival of respiratory influenza virus infection. *J Virol* 79:6441–6448 [PubMed: 15858027]
- Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9:671–675 [PubMed: 22930834]
- Sfeir RM, Dubarry M, Boyaka PN et al. (2004) The mode of oral bovine lactoferrin administration influences mucosal and systemic immune responses in mice. *J Nutr* 134:403–409 [PubMed: 14747680]
- Sienkiewicz M, Jaskiewicz A, Tarasiuk A et al. (2021) Lactoferrin: an overview of its main functions, immunomodulatory and antimicrobial role, and clinical significance. *Crit Rev Food Sci Nutr* 1–18. doi: 10.1080/10408398.2021.1895063
- Skerry C, Harper J, Klunk M et al. (2012) Adjunctive TNF inhibition with standard treatment enhances bacterial clearance in a murine model of necrotic TB granulomas. *PLoS One* 7:e39680 [PubMed: 22761866]
- Sotgiu G, Centis R, D'Ambrosio L et al. (2015) Tuberculosis treatment and drug regimens. *Cold Spring Harb Perspect Med* 5:a017822 [PubMed: 25573773]
- Spadaro M, Montone M, Arigoni M et al. (2014) Recombinant human lactoferrin induces human and mouse dendritic cell maturation via Toll-like receptors 2 and 4. *FASEB J* 28:416–429 [PubMed: 24088817]
- Takakura N, Wakabayashi H, Yamauchi K et al. (2006) Influences of orally administered lactoferrin on IFN-gamma and IL-10 production by intestinal intraepithelial lymphocytes and mesenteric lymph-node cells. *Biochem Cell Biol* 84:363–368 [PubMed: 16936808]

- Thiriot JD, Martinez-Martinez YB, Endsley JJ et al. (2020) Hacking the host: exploitation of macrophage polarization by intracellular bacterial pathogens. *Pathog Dis* 78:ftaa009 [PubMed: 32068828]
- Turin CG, Zea-Vera A, Pezo A et al. (2014) Lactoferrin for prevention of neonatal sepsis. *Biometals* 27:1007–1016 [PubMed: 24935001]
- Vilaplana C, Marzo E, Tapia G et al. (2013) Ibuprofen therapy resulted in significantly decreased tissue bacillary loads and increased survival in a new murine experimental model of active tuberculosis. *J Infect Dis* 208:199–202 [PubMed: 23564636]
- Wallis RS, Broder MS, Wong JY et al. (2004) Granulomatous infectious diseases associated with tumor necrosis factor antagonists. *Clin Infect Dis* 38:1261–1265 [PubMed: 15127338]
- Wang N, Liang H, Zen K (2014) Molecular mechanisms that influence the macrophage m1-m2 polarization balance. *Front Immunol* 5:614 [PubMed: 25506346]
- Welsh KJ, Abbott AN, Hwang SA et al. (2008) A role for tumour necrosis factor- α , complement C5 and interleukin-6 in the initiation and development of the mycobacterial cord factor trehalose 6,6'-dimycolate induced granulomatous response. *Microbiology* 154(Pt 6):1813–1824 [PubMed: 18524936]
- Welsh KJ, Hwang SA, Hunter RL et al. (2010) Lactoferrin modulation of mycobacterial cord factor trehalose 6-6'-dimycolate induced granulomatous response. *Transl Res* 156:207–215 [PubMed: 20875896]
- Welsh KJ, Hwang SA, Boyd S et al. (2011) Influence of oral lactoferrin on Mycobacterium tuberculosis induced immunopathology. *Tuberculosis* 91: S105–S113 [PubMed: 22138562]
- Wisgrill L, Wessely I, Spittler A et al. (2018) Human lactoferrin attenuates the proinflammatory response of neonatal monocyte-derived macrophages. *Clin Exp Immunol* 192:315–324 [PubMed: 29393509]
- World Health Organization (2020) Global tuberculosis report 2020. Geneva: World Health Organization
- Yeom M, Park J, Lee B et al. (2011) Lactoferrin inhibits the inflammatory and angiogenic activation of bovine aortic endothelial cells. *Inflamm Res* 60:475–482 [PubMed: 21161563]
- Zhang R, Xi X, Wang C et al. (2018) Therapeutic effects of recombinant human interleukin 2 as adjunctive immunotherapy against tuberculosis: A systematic review and meta-analysis. *PLoS One* 13:e0201025 [PubMed: 30024982]
- Zhang H, Shi N, Diao Z et al. (2021) Therapeutic potential of TNF α inhibitors in chronic inflammatory disorders: Past and future. *Genes Dis* 8:38–47 [PubMed: 33569512]
- Zimecki M, Mazurier J, Machnicki M et al. (1991) Immunostimulatory activity of lactotransferrin and maturation of CD4- CD8- murine thymocytes. *Immunol Lett* 30:119–123 [PubMed: 1683650]
- Zimecki M, Miedzybrodzki R, Mazurier J et al. (1999) Regulatory effects of lactoferrin and lipopolysaccharide on LFA-1 expression on human peripheral blood mononuclear cells. *Arch Immunol Ther Exp* 47:257–264




Fig. 1. *Mtb* infection and lactoferrin treatment scheme.

C57Bl/6 mice were aerosol challenged with *Mtb* and treated with rHLF administered every other day orally beginning on day 14 (prophylactic relative to granulomatous responses; six total doses), or beginning on day 21 post infection (therapeutic relative to established granulomas; three total doses). Mice were intravenous injected with ofloxacin 30 min prior to sacrifice, at 28 days post initial infectious challenge.




Fig. 2. Lactoferrin treatment reduces pulmonary inflammation post infectious challenge with *Mtb*.

Lungs from *Mtb* infected mice were assessed at day 28 post aerosol infection (A) and compared to animals given rHLF in the prophylactic group (B) or therapeutic group (C). Histologic assessment revealed primary granulomatous response with monocytic cell infiltration, dense cellular foci, and occluded vascular regions in control infected mice (4× magnification). Both prophylactic and therapeutic rHLF treatment reduced inflammatory response resulting in modest inflammatory foci and reduced pathological damage to lung tissue. Hematoxylin and eosin stained histographs represent formalin fixed lung sections at 10× magnification obtained from repeated studies with 8–10 mice in each group; study representative of repeat experiments.




Fig. 3. Decreased inflammatory response in lactoferrin treated mice.

Mtb infected mice were assessed by digital analysis for cellularity and inflammation. Area occupied of pulmonary scanned sections is shown for individual mice, with and without rHLF treatments. Results represent mean ± standard deviation of the mean. Similar data was obtained in repeated experiments; 6–10 mice were included per group, per experiment. P: prophylactic treatment; T: therapeutic treatment; ****p* 0.001, one way ANOVA, Tukey post hoc test for multiple comparisons.




Fig. 4. Increased fluoroquinolone penetration to primary granulomas after treatment with lactoferrin.

Lungs from *Mtb* infected mice alone (left), or rHLF treated prophylactically (P; middle) or therapeutically (T; right) were assessed for presence of ofloxacin within granulomatous inflammation. Ofloxacin penetration was primarily excluded from inflammatory foci in the *Mtb* infected alone group, while both rHLF treated mice permitted entry of ofloxacin into regions of pathology. Top panels represent hematoxylin & eosin (H&E) brightfield stained histographs (40× magnification) with matching fluorescence captured using multispectral imaging (bottom). Example populations of macrophage-like cells (white arrows) and lymphocytic cells (black arrows) are indicated.




Fig. 5. Relative fluorescence for individual inflammatory foci post lactoferrin treatment. Quantitative assessment of ofloxacin penetration into granulomas in lactoferrin treated mice are compared to non-treated infected animals. Individual scans are represented from multiple sections from 6–8 mice per group; average values and standard deviation included. P: prophylactic treatment; T: therapeutic treatment; *** p 0.001, one way ANOVA, Tukey post hoc test for multiple comparisons.




Fig. 6. Accumulation of ofloxacin signal in activated macrophages in lactoferrin treated *Mtb* infected mice.

H&E staining of inflammatory foci within rHLF therapeutically treated mouse lung revealed acute regions of highly activated “foamy” macrophage-phenotypic cells (white arrows). Multispectral imaging correlates the presence of fluorescence which overlaps presence of activated cells, located within the focal granulomatous regions. Representative section, 100× magnification.




Fig. 7. Retention of vascularized structures within regions of inflammation in lactoferrin treated *Mtb*-infected mice.

Serial sections compared presence of maintained vascular structure within inflammatory foci in *Mtb* infected mice (left side) or in *Mtb* infected mice treated therapeutically with rHLF (right side). Fluorescence patterns obtained using multispectral imaging (top panels) were compared with serial lung sections that were immunohistochemically stained for PECAM-1 to identify vascular endothelial populations within inflammatory foci (bottom panels). A: alveolus; VE: vascular endothelium; AE: alveolar capillary endothelium (arrows).




Fig. 8. Altered localization of M1-like and M2-like populations following lactoferrin treatment of *Mtb* infected mice.

Serial sections of formalin fixed lung tissue were reacted with antibody to CD38 (top) or CD206 (bottom), for either *Mtb* alone (left side) or *Mtb* infected mice treated therapeutically with rHLF (right side). *Mtb* alone infected mice demonstrate accumulation of both M1- and M2-like phenotypic cells in a cuffing pattern surrounding blood vessels adjacent to granulomatous inflammation. A different pattern appears in the rHLF treated group, with

minimal presence of M1-like phenotype and a primarily diffuse distribution of M2-like cells throughout the inflammatory foci.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript