

RESEARCH ARTICLE

Open Access



MiR-539-3p inhibited chondrogenic differentiation in human adipose stem cells by targeting Sox9

Feng Qin¹, Fang Wang², Xiao-Ping Wang³, Jie Chen⁴, Feng-hua Zeng⁴, Cui-Lan Sun⁵, Jia-Cuo Peng Mao⁶ and Chun-Liang Li^{6*} 

Abstract

Background: Mesenchymal stem cells (MSCs) have emerged as the attractive candidates for cell therapy for cartilage repair in clinical therapy of osteoarthritis (OA). MiR-539-3p was reported to differentially express during chondrogenic differentiation of adipose stem cells (ASCs) by miRNA microarrays. The aim of the study was to investigate the effects and underlying mechanisms of miR-539-3p on chondrogenic differentiation of ASCs.

Methods: Human ASCs (hASCs) were obtained from liposuction and transfected with miR-539-3p mimic or inhibitor. Then, the cells were cultured in chondrogenic differentiation medium including transforming growth factor- β 1 (TGF- β 1).

Results: Our results found that miR-539-3p was gradually down-regulated during chondrogenic differentiation of hASCs. MiR-539-3p overexpression inhibited TGF- β 1-induced chondrogenic differentiation of hASCs, as supported by reducing the gene and protein expression of chondrogenic differentiation markers type II collagen alpha 1 (COL2A1), aggrecan (ACAN), and type II collagen. In contrast, miR-539-3p inhibitor significantly promoted the chondrogenic differentiation of hASCs. Dual luciferase reporter assay demonstrated that Sox9 was a direct target gene of miR-539-3p. The expression of SRY-box transcription factor 9 (Sox9) was up-regulated progressively over time during chondrogenic differentiation of hASCs. Additionally, Sox9 overexpression notably reversed chondrogenic differentiation of hASCs inhibited by miR-539-3p mimic, as demonstrated by the decreased expression of COL2A1, ACAN, and type II collagen.

Conclusions: Altogether, miR-539-3p inhibited chondrogenic differentiation of hASCs by targeting Sox9. MiR-539-3p may have significant clinical applications for use as a targeted therapy of OA.

Keywords: Adipose stem cells, Chondrogenic differentiation, miR-539-3p, Sox9

Background

Osteoarthritis (OA) was the most common skeletal disorder in middle-aged and older people which resulted from articular cartilage degradation and imbalance of subchondral bone remodeling [1]. Repair of the cartilage

defect and inhibition of cartilage degradation was most important for OA patients [2]. However, the majority of the drugs for the treatment of OA have been inadequate, as they only treated the symptoms of pain and inflammation [3]. Consequently more effective function-modifying therapeutic strategies would need to be introduced for the clinical treatment of OA.

In recent years, range of methods have been developed including osteochondral transplantation and autologous

*Correspondence: lichunliang403@126.com

⁶ Department of Orthopedic, Qinghai Provincial People's Hospital, Chengdong District, No. 2 Gonghe Road, Xining 810007, Qinghai, China
Full list of author information is available at the end of the article



chondrocyte transplantation (ACT) [4]. However, ACT still faced challenges in treating cartilage defects, such as the paucity of the cell source and the damage to natural tissues caused by cell harvesting [5]. Recent researched on mesenchymal stem cells (MSCs) have provided a new and exciting opportunity for bone and cartilage tissue engineering [6, 7]. Thus far, MSCs have been isolated from bone marrow, periosteum, trabecular bone, adipose tissue, synovium, skeletal muscle, and deciduous teeth [8]. Among them, adipose stem cells (ASCs) have the advantage of being easy to obtain. ASCs possessed the capacity to differentiate into adipocytes, chondrocytes, osteoblasts, endothelial cells, cardiac myocyte and so on [8, 9]. New strategies have to center around promoting the chondrogenic potential of the chondrocytes using MSCs therapy.

MicroRNAs (miRNAs) were single-stranded RNAs of 19–23 nucleotides and were found in a wide variety of organisms [10, 11]. MiRNAs regulated gene expression in the post-transcriptional level through its complementary pair of target mRNAs, which leading to mRNA translation inhibition and degradation. MiRNAs were involved in the regulation of growth and development of various organisms, signal transduction, differentiation, and diseases [12]. Importantly, many miRNAs, such as miR-130a [13], miR-31a-5p [14], and miR-101-3p [15] were abnormally expressed in stem cells and played an important role in self-renewal and differentiation of stem cells by regulating the expression of certain key genes [16]. Previous miRNA microarrays analysis of hASCs during chondrogenic differentiation found that miR-539-3p was differentially expressed [17]. MiR-539-3p was recently reported to play a role in inhibiting proliferation and promoting apoptosis in a variety of cancers [18, 19], but there was no report on its role in regulating stem cell chondrogenic differentiation.

In this study, we explored the expression of miR-539-3p in the chondrogenic differentiation of ASCs. Furthermore, the study was performed to explore the biological functions and underlying mechanisms of miR-539-3p as a target gene expression inhibitor in the chondrogenic differentiation of ASCs.

Materials and methods

Isolation and culture of human ASCs (hASCs)

Human adipose was obtained from 5 healthy female donors, aged 33–42 years, during the abdominal liposuction procedure at the Qinghai Provincial People's Hospital (Qinghai, China). Written informed consents were obtained from all donors. The research protocol was reviewed and approved by the Ethical Committee of Qinghai University Affiliated Hospital (SL-2021052). Adipose were digested with collagenase I to isolate ASCs,

which were incubated in high-glucose DMEM with 1% fetal bovine serum (FBS) for primary culture. The third-passage ASCs were cultured in high-glucose DMEM supplemented with 1% FBS, 10 µg/L transforming growth factor-β1 (TGF-β1), 50 nmol/L ascorbate-2-phosphate, 6.25 mg/L insulin, and 1% penicillin. The appearance and generation of ASCs were observed under inverted microscope, and the differentiations to chondrocyte were verified by alcian blue staining and immunocytochemical staining.

Cell transfection

NC mimic, miR-539-3p mimic, NC inhibitor, miR-539-3p inhibitor, pcDNA-NC, and pcDNA-SRY-box transcription factor 9 (Sox9) were designed and synthesized by RiboBio (Wuhan, China). Lipofectamine™ 3000 (L3000-008; Invitrogen, Grand Island, NY) transfection reagent was used for the transient transfection of according to the manufacturer's instructions.

Cell proliferation assays

The 1–9 days growth curve of ASCs at the passage 3 was confirmed by cell counting kit-8 (CCK-8) assay. The cultures were supplemented with CCK-8 solution (Dojindo, Japan) and incubated for 1 h at 37 °C. Cell proliferation was measured at a wavelength of 450 nm by a microplate reader.

Flow cytometry assays

Expression of ASC-related cell surface markers was evaluated by flow cytometry. Briefly, 5.0×10^5 cells at the passage 3 were incubated with phycoerythrin (PE) conjugated monoclonal antibodies for human CD29, CD90, CD105, CD34, or CD45 (eBioscience, San Diego, CA, USA) for 1 h at 4 °C. The IgG isotype antibodies were used as negative controls for each experiment. After washing with PBS, cells were subjected to flow cytometric analysis using an FACS Canto II flow cytometer system (BD Biosciences).

Alcian blue staining

Chondrogenic differentiation of ASCs was identified using alcian blue staining. ASCs were fixed using 4% paraformaldehyde for 10 min at room temperature. Then the cells were washed with PBS and performed using alcian blue solution (0.1% alcian blue, 0.1N HCl; Sigma-Aldrich, St. Louis, Missouri, USA) for 4 h at room temperature and images were captured.

Immunofluorescence (IF) assays

ASCs (1.0×10^5 cells) were fixed with 4% paraformaldehyde for immunofluorescence assay. For detection of collagen II expression, ASCs were blocked with 5%

goat serum for 30 min and then incubated with primary anti-type II collagen (Abcam, USA, No. ab34712; 1/500) overnight at 4 °C. The slides were washed with PBS, secondary antibody was added and incubated for 1 h at room temperature before stained with 4',6-diamidino-2-phenylindole (DAPI; Invitrogen, Grand Island, NY; 1/1000). Finally, the stained samples were observed using an inverted microscope eclipse Ti (Nikon Instruments Inc., New York, USA).

Quantitative Real-time reverse transcriptase-polymerase chain reaction (RT-qPCR)

RT-qPCR was performed to calculate the RNA expression of type II collagen alpha 1 (COL2A1), aggrecan (ACAN), Sox9, and miR-539-3p in ASCs (1.0×10^6 cells) which were grown in 6-well cell culture plates. TRIzol protocol was used to extract total RNA from ASCs. 2000 ng RNA was then reverse transcribed into cDNA using GoScript™ reverse transcription system (Promega, USA). 1 µg cDNA was subjected to RT-qPCR using a SYBR Green Super mix (Bio-Rad, Hercules, CA, USA). The $2^{-\Delta\Delta CT}$ method was used to calculate the relative expression. β -actin was used as an internal control to normalize COL2A1, ACAN, and Sox9 expression. The levels of miR-539-3p expression were compared to that of the u6 control group. Primer sequences are shown in Table 1.

Dual-luciferase reporter assays

The wild type pSI-Check2-Sox9 (Sox9-3'UTR-Wt) and the mutant pSI-Check2-Sox9 (Sox9-3'UTR-Mut) recombinant dual luciferase reporter plasmid were designed and synthesized based on the binding region of miR-539-3p and Sox9 3'UTR sequence (Fubio Biological Technology Co., Shanghai, China). Sox9 Wt or Mut was co-transfected along with miR-539-3p mimic or NC into HEK293T cells by Lipofectamine™ 3000 (L3000-008; Invitrogen, Grand Island, NY). After 48 h, the luciferase activities were recorded with a plate style luminometer using Promega Dual-Luciferase system (Promega, Madison, Wisconsin, WI, USA).

Western blot analysis

The total proteins of ASC lysates were collected RIPA buffer (Signaling Technology, Inc.). The protein concentration was determined using a BCA kit (Beyotime, Shanghai, China). Total protein (30 µg/sample) was separated via 10% SDS-PAGE and to nitrocellulose membranes. Use 5% skimmed milk powder to block the membranes. The corresponding protein antibodies were as follows: COL2A1 (Abclonal, China, No. A1560; 1/500), ACAN (Abcam, USA, No. ab3778; 1/1000), Sox9 (Abcam, USA, No. ab285966; 1/1000), and β -actin (Boster, Chian, No. BM0627; 1/1000). Then, the membrane washing was performed with Tris-buffered saline/0.1% Tween (TBST) and incubated for 1.5 h with a HRP Goat anti-Rabbit IgG (Abcam, No. ab6721; 1/1000). The band visualization was carried out using the ECL system (Thermo Scientific, Rockford, IL, USA). β -actin used as internal control.

Statistical analysis

For isolation and identification of hASCs, the experiments were performed 5 times independently, using ASC from 5 different people. For additional cell study, each experiment was repeated at least 3 times independently. The data were represented as means \pm standard deviation (SD). Statistical analysis was performed with the SPSS software (version 19.0, SPSS Inc., Chicago, IL, USA). One-way analysis of variance (ANOVA) followed by Tukey's post hoc test and Student's unpaired t test were used for comparison between groups. *P* value < 0.05 was considered statistically significant.

Results

Isolation and identification of ASCs

ASCs were isolated from human abdominal adipose tissue as described in the Methods section. At passage 3 (P3), ASCs were identified and used in subsequent experiments. After reaching 80–90% confluence, cells adopted a spindle-like shape (Fig. 1A). And CCK-8 assay showed that the cell viability of ASCs was significantly increased during 10 days of continuous culture (Fig. 1A). Meanwhile, ASCs were characterized by flow cytometry after staining surface markers. These cells were found to be positive for CD29, CD90, and CD105, but did not express

Table 1 Primers for quantitative RT-PCR

Gene	Forward (5'-3')	Reverse (5'-3')
COL2A1	CATTGCTGGTCTCCCGCT	ACCAGCATCACCAGGGCGAC
ACAN	GGAGGAGCAGGAGTTGTCAA	TGTCCATCCGACCAGCGA AA
Sox9	AAGCTCTGGAGACTTCTGAACG	CGTTCTTACCAGACTTCTCC
β -actin	CTCCATCGTCCACCGCAAATGCTTCT	GCTCCAACCGACTGCTGCACCTTC
u6	CCAAGGTCATCCATGACAAC	TGTCATACCAGGAAATGAGC




Fig. 1 Classification of human adipose stem cells (hASCs). **A** hASCs exhibited a representative spindle-like morphology. Scale bar: 100 μm. The cell proliferation of hASCs was tested by CCK-8 assay. **B** Flow cytometric analysis of characteristic cell surface markers of hASCs (CD29, CD90, CD105, CD45, and CD34). **C** Alcian blue staining of hASCs. Scale bar: 100 μm. **D-E** The mRNA levels of COL2A1 and ACAN were quantified by RT-qPCR. **F-H** The protein levels of COL2A1 and ACAN were measured using western blot. β-actin expression levels were detected as an endogenous control. **I** Immunofluorescence assay of Type II collagen. Type II collagen fluorescence (green) and DAPI (blue). Scale bar: 50 μm and 20 μm. The experiment was repeated independently 5 times. The data are presented as mean ± standard deviation (SD)

CD45 and CD34 (Fig. 1B). hASCs were cultured in chondrogenic differentiation medium that were evaluated by alcian blue staining (Fig. 1C). The resulted showed that alcian blue staining was positive (Fig. 1C). Moreover, the

expression of chondrogenic differentiation markers, such as COL2A1 and ACAN were analyses to confirmed chondrogenesis. The mRNA and protein levels of COL2A1 and ACAN were increased in chondrocyte-ASCs




Fig. 2 MiR-539-3p regulated ASC chondrogenic differentiation. hASCs were transfected with miR-539-3p mimic or miR-539-3p inhibitor, which were cultured in chondrogenic differentiation medium. hASCs cultured in high-glucose DMEM with 1% fetal bovine serum (FBS) were used as control. **A–C** The expression levels of miR-539-3p in hASCs or chondrogenic differentiation hASCs (CD-ASCs) were detected by RT-qPCR. Gene expression levels of COL2A1 and ACAN in hASCs or CD-ASCs were determined by RT-qPCR (**D** and **E**) and western blot (**F–H**). beta-actin expression levels were detected as an endogenous control. ** $P < 0.01$ (VS. Day 0/NC mimic/ASCs), ## $P < 0.01$ (VS. CD-ASCs). (**I**) Immunofluorescence assay of Type II collagen. Type II collagen fluorescence (green) and DAPI (blue). Scale bar: 100 μ m and 50 μ m. The experiment was repeated at least 3 times independently. The data are presented as mean \pm standard deviation (SD)

(CD-ASCs, Fig. 1D–H, and Type II collagen in IF staining was appeared positive (Fig. 1I).

MiR-539-3p regulated ASC chondrogenic differentiation

As shown in Fig. 2A, the expression of miR-539-3p gradually decreased at day 7, 14 and 21 of ASC chondrogenic differentiation, implying that miR-539-3p expression may relate with ASC chondrogenic differentiation. To confirm whether miR-539-3p modulates chondrogenic differentiation in ASCs, miR-539-3p mimic, miR-539-3p inhibitor, and their negative controls were designed and synthesized by RiboBio to stably overexpress miR-539-3p and inhibit the function of miR-539-3p, respectively (Fig. 2B and C). We tested COL2A1, ACAN, and Type II collagen expression to explore the effect of miR-539-3p on hASC chondrogenic differentiation. As shown in Fig. 2D and E, the down-regulated COL2A1 and ACAN mRNA expression were observed in miR-539-3p mimic transfected CD-ASCs. And the treatment of miR-539-3p inhibitor showed the opposite results (Fig. 2D and H). Meanwhile, COL2A1 and ACAN were lower expressed in miR-539-3p mimic transfected CD-ASCs and higher expressed in miR-539-3p inhibitor treated CD-ASCs (Fig. 2F–H). IF assay showed that miR-539-3p overexpression decreased Type II collagen expression and miR-539-3p inhibitor increased Type II collagen expression in CD-ASCs. These results demonstrated that miR-539-3p diminished the chondrogenic differentiation of ASCs.

Sox9 is a target gene of miR-539-3p

Subsequently, the putative binding site of miR-539-3p to the 3'-UTR of Sox9 was predicted using the TargetScan and Starbase v2.0 (Fig. 3A). Luciferase assay demonstrated that the luciferase activity of wild type pSI-Check2-Sox9 was attenuated by miR-338-3p in HEK293T cells (Fig. 3B). Conversely, no significant differences were observed in the luciferase activity of the mutant pSI-Check2-Sox9 (Fig. 3B). Our results further showed that miR-539-3p mimic significantly inhibited Sox9 expression (Fig. 3C–E). Moreover, the mRNA and protein levels of Sox9 were increased with time following chondrogenic differentiation in ASCs (Fig. 3F–H). These findings indicated that miR-539-3p suppressed Sox9 expression by directly binding to the 3'-UTR of Sox9.

Sox9 overexpression promoted ASC chondrogenic differentiation by blocking the effect of miR-539-3p mimic

To verify the regulatory function of Sox9 on miR-539-3p-mediated chondrogenic differentiation, Sox9 overexpression vector (pcDNA-Sox9) was synthesized and co-transfected with miR-539-3p mimic into CD-ASCs. As shown in Fig. 3I–K, the mRNA and protein levels of Sox9 were both enhanced by pcDNA-Sox9. After

chondrogenic differentiation was induced in media for 21 days, Sox9 overexpression reversed miR-539-3p-reduced COL2A1 and ACAN expression compared with the alone transfection of miR-539-3p mimic group. IF detection also showed that the expression of Type II collagen was decreased by miR-539-3p mimic, which was significantly blocked by Sox9 overexpression. These data indicated that miR-490-5p inhibited the chondrogenic differentiation of ASCs by targeting Sox9 and decreasing Sox9 expression (Fig. 4).

Discussion

In the present study, the down-regulated of miR-539-3p was observed on day 7, 14 and 21 during TGF- β 1-induced chondrogenic differentiation of ASCs. Furthermore, bioinformatics prediction and dual-luciferase reporter assays verified Sox9 was a target gene of miR-539-3p. Overexpression of miR-539-3p inhibited the expression of chondrogenic differentiation markers, such as COL2A1 and ACAN, as well as Type II collagen expression. Sox9 overexpression significantly reversed the anti-chondrogenic effect of miR-539-3p in ASCs.

MiRNAs regulate gene expression at the posttranscriptional level in many developmental and metabolic processes. Emerging evidence demonstrates that miRNAs also play an essential role in stem cell self-renewal and differentiation [20, 21]. Some miRNAs are abnormally expressed in stem cells, control stem cell self-renewal, and differentiation through negatively regulating the expression of certain key genes in stem cells [22]. The expression profile of miRNAs during chondrogenic differentiation of hASCs showed that 8 miRNAs were down-regulated and 12 miRNAs were up-regulated [17]. Among them, miR-490-5p facilitated the chondrogenic differentiation of hASCs by targeting PITPNM1 3'UTR and regulating PI3K/AKT axis [23]. MiR-1307-3p overexpression suppressed the transition of hASCs from chondrogenesis to osteogenesis, and further mechanistic evaluation showed that it functioned by targeting BMPR2 and inhibiting Smad5/8 signaling pathway [24]. An additional study found that miR-194 was a novel modulator of hASCs-mediated chondrogenesis by targeting Sox5 [25]. MiRNA let-7i-3p inhibited the chondrogenic differentiation of hASCs, and inhibiting the function of let-7i-3p activated the Wnt/ β -catenin pathway. Our RT-qPCR data indicated that miR-539-3p was obviously decreased during process of chondrogenic differentiation of hASCs. Furthermore, miR-539-3p mimic inhibited hASC chondrogenic differentiation by reduced the expression of chondrogenic markers Type II collagen, COL2A1, and ACAN. The inhibited of miR-539-3p function showed the opposite trends.




Fig. 3 Sox9 is a target gene of miR-539-3p. **A** Alignment of miR-539-3p and the 3'-UTR of Sox9, a potential miR-539-3p target. **B** A luciferase reporter carrying the 3'-UTR of wild-type (Sox9 3'UTR-WT) or mutant (Sox9 3'UTR-MUT) Sox9 was introduced into HEK-293T cells along with NC mimic or miR-539-3p mimic. **C, F,** and **I** The expression level of Sox9 in hASCs was determined by RT-qPCR. **D, E, G, H, J,** and **K** The protein expression level of Sox9 in hASCs was detected by western blot. β-actin expression levels were detected as an endogenous control. ***P* < 0.01 (VS. 0 Day/NC mimic/pcDNA-NC). The experiment was repeated at least 3 times independently. The data are presented as mean ± standard deviation (SD)




Fig. 4 Sox9 overexpression promoted hASC chondrogenic differentiation by blocking the effect of miR-539-3p mimic. hASCs were transfected with miR-539-3p mimic or/and pcDNA-Sox9, which were cultured in chondrogenic differentiation medium. hASCs cultured in high-glucose DMEM with 1% fetal bovine serum (FBS) were used as control. Gene expression levels of COL2A1 and ANCA in hASCs or CD-ASCs were determined by RT-qPCR (A and B) and western blot (C–E). β -actin expression levels were detected as an endogenous control. $**P < 0.01$ (VS. ASCs), $##P < 0.01$ (VS. ASCs), $\&P < 0.05$ (VS. miR-539-3p mimic), $\&\&P < 0.05$ (VS. miR-539-3p mimic), $\&\&P < 0.01$ (VS. pcDNA-Sox9). (I) Immunofluorescence assay of Type II collagen. Type II collagen fluorescence (green) and DAPI (blue). Scale bar: 100 μ m and 50 μ m. The experiment was repeated at least 3 times independently. The data are presented as means \pm standard deviation (SD)

These data demonstrated that miR-539-3p functions as a suppressor during chondrogenic differentiation of hASCs. Several previous reports of miR-539-3p were mostly focused on regulating cancer progression. MiR-539-3p was reported to promote the proliferation and invasion of epithelial ovarian cancer and inhibit the progression of gastric cancer and colon cancer [18, 19, 26]. The mechanism of miR-539-3p-mediated tumorigenesis was targeted SPARCL1 [18], RNF2 [27], CTBP1 [19], or CDK14 [26] and inhibited the expression of these proteins in tumor cells. In addition, another study revealed that in a model of ischemia–reperfusion (I/R) cardiac injury, miR-539-3p enhanced infarction area and stimulated oxidative stress, myocardial cell apoptosis, and reduced cardiac function recovery via targeting ErbB4 [28]. Accordingly, to the best of our knowledge, this study was the first to focus on miR-539-3p function in chondrogenic differentiation of hASCs. Further in vivo functional analyses will be needed to clarify the potential of miR-539-3p to treat OA with stem cell transplantation.

Sox9 was an important transcription factor for chondrocyte differentiation and played an important role in the transformation of stem cells into chondrocytes and in the process of chondrogenesis. Previous report showed that a denovirus-mediated Sox9 overexpression vector delivery induced chondrogenic differentiation and inhibited osteogenic and adipogenic differentiation of hMSCs [29]. The study by Magali Cucchiari found that application of recombinant adeno-associated virus (rAAV) Sox9 vector significantly treated rabbit knee articular cartilage defects and delayed premature terminal differentiation in the newly formed cartilage [30]. More importantly, in a rabbit model with full-thickness cartilage defects, bone marrow mesenchymal stem cells (BMSCs) transfected with Sox9 gene adenovirus were loaded on polyglycolic acid (PGA) scaffolds to repair chondral defects. The results revealed that this Sox9-BMSCs-PGA scaffold composition promoted cartilage matrix formation and chimerism of new cartilage with surrounding tissue [31]. Further, it was known that a variety of miRNAs, such as miR-30b [32], miR-495 [33], and miR-574-3p [34] could negatively regulate Sox9 in chondrogenic differentiation of stem cells. In the present study, Sox9 was identified as a target gene of miR-539-3p by luciferase reporter assay, which plays a role in promoting the chondrogenic differentiation of hASCs. Previous studies also showed that Sox9 target genes included COL2A1, COL11A1, COL9A2, Aggrecan (Agc1), cartilage link protein (CRTL1), and so on [35]. We found that Sox9 overexpression violently reversed the inhibitory effect of miR-539-3p mimic on hASCs executing the chondrogenic differentiation program,

indicated by increasing expression of Type II collagen, COL2A1, and ACAN.

Conclusion

Our in vitro study using hASCs is the first demonstration that miR-539-3p was down-regulated during chondrogenic differentiation. Our findings are also found miR-539-3p inhibited chondrogenic differentiation of hASCs by targeting Sox9. Our results may provide a potential reference target for the treatment of OA. Further experimentation will be needed to investigate whether miR-539-3p inhibitor-modified ASCs could also promote chondrogenesis in vivo.

Abbreviation

MSCs: Mesenchymal stem cells; OA: Osteoarthritis; ASCs: Adipose stem cells; TGF- β 1: Transforming growth factor- β 1; COL2A1: Type II collagen alpha 1; ACAN: Aggrecan; Sox9: SRY-box transcription factor 9; ACT: Autologous chondrocyte transplantation; miRNAs: MicroRNAs; IF: Immunofluorescence; RT-qPCR: Quantitative Real-time reverse transcriptase-polymerase chain reaction; Agc1: Aggrecan; CRTL1: Cartilage link protein.

Author's contributions

All authors read and approved the final manuscript.

Funding

This study was supported by a grant from the Qinghai Science and Technology Department Project (2019-ZJ-7067).

Availability of data and materials

The datasets used or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Written informed consent was obtained from all patients after approval by the ethics committee of Qinghai Provincial People's Hospital (2020-77).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Endocrinology, Qinghai University Affiliated Hospital, Xining 810006, China. ²Department of Gastroenterology, Qinghai University Affiliated Hospital, Xining 810006, China. ³Department of the Image Center, Qinghai University Affiliated Hospital, Xining 810006, China. ⁴Department of Anesthesia, Qinghai Provincial People's Hospital, Xining 810007, China. ⁵Department of Clean Operating Room, Qinghai Provincial People's Hospital, Xining 810007, China. ⁶Department of Orthopedic, Qinghai Provincial People's Hospital, Chengdong District, No. 2 Gonghe Road, Xining 810007, Qinghai, China.

Received: 19 January 2022 Accepted: 2 March 2022

Published online: 18 March 2022

References

1. Sacitharan PK. Ageing and Osteoarthritis. *Subcell Biochem.* 2019;91:123–59.

2. Messina OD, Vidal Wilman M. Nutrition, osteoarthritis and cartilage metabolism. *Aging Clin Exp Res*. 2019;31(6):807–13.
3. Ragle RL, Sawitzke AD. Nutraceuticals in the management of osteoarthritis : a critical review. *Drugs Aging*. 2012;29(9):717–31.
4. Messina JC, Torretta F, Randelli PS. Autologous chondrocyte transplantation in the treatment of thumb CMC joint osteoarthritis. *Hand Surg Rehab*. 2021;40s:S21–8.
5. Brittberg M. Autologous chondrocyte transplantation. *Clin Orthopaedics Relat Res*. 1999;367(Suppl):S147–155.
6. Zhu Y, Wang Y, Zhao B, Niu X, Hu B, Li Q, Zhang J, Ding J, Chen Y, Wang Y. Comparison of exosomes secreted by induced pluripotent stem cell-derived mesenchymal stem cells and synovial membrane-derived mesenchymal stem cells for the treatment of osteoarthritis. *Stem Cell Res Ther*. 2017;8(1):64.
7. De Bari C, Roelofs AJ. Stem cell-based therapeutic strategies for cartilage defects and osteoarthritis. *Curr Opin Pharmacol*. 2018;40:74–80.
8. Ding DC, Shyu WC, Lin SZ. Mesenchymal stem cells. *Cell Transplant*. 2011;20(1):5–14.
9. Fu X, Liu G, Halim A, Ju Y, Luo Q, Song AG. Mesenchymal stem cell migration and tissue repair. *Cells*. 2019;8(8):784.
10. Saliminejad K, Khorram Khorshid HR, Soleymani Fard S, Ghaffari SH. An overview of microRNAs: biology, functions, therapeutics, and analysis methods. 2019;**234**(5):5451–65
11. Lu TX, Rothenberg ME. MicroRNA. *J Allergy Clin Immunol*. 2018;141(4):1202–7.
12. Mohr AM, Mott JL. Overview of microRNA biology. *Semin Liver Dis*. 2015;35(1):3–11.
13. Lin Z, He H, Wang M, Liang J. MicroRNA-130a controls bone marrow mesenchymal stem cell differentiation towards the osteoblastic and adipogenic fate. *Cell Prolif*. 2019;52(6):e12688.
14. Xu R, Shen X, Si Y, Fu Y, Zhu W, Xiao T, Fu Z, Zhang P, Cheng J, Jiang H. MicroRNA-31a-5p from aging BMSCs links bone formation and resorption in the aged bone marrow microenvironment. *Aging Cell*. 2018;17(4):e12794.
15. Kaneda-Ikeda E, Iwata T. Regulation of osteogenesis via miR-101-3p in mesenchymal stem cells by human gingival fibroblasts. *J Bone Mineral Matab*. 2020;38(4):442–55.
16. Huang C, Geng J, Jiang S. MicroRNAs in regulation of osteogenic differentiation of mesenchymal stem cells. *Cell Tissue Res*. 2017;368(2):229–38.
17. Yang Z, Hao J, Hu ZM. MicroRNA expression profiles in human adipose-derived stem cells during chondrogenic differentiation. *Int J Mol Med*. 2015;35(3):579–86.
18. Gong YB, Fan XH. MiR-539-3p promotes the progression of epithelial ovarian cancer by targeting SPARCL1. *Eur Rev Med Pharmacol Sci*. 2019;23(6):2366–73.
19. Zhou J, Su M, Zhang H, Wang J, Chen Y. miR-539-3P inhibits proliferation and invasion of gastric cancer cells by targeting CTBP1. *Int J Clin Exp Pathol*. 2019;12(5):1618–25.
20. Martin EC, Qureshi AT, Dasa V, Freitas MA, Gimble JM, Davis TA. MicroRNA regulation of stem cell differentiation and diseases of the bone and adipose tissue: perspectives on miRNA biogenesis and cellular transcriptome. *Biochimie*. 2016;124:98–111.
21. Kumar S, Curran JE, DeLeon E. Role of miRNA-mRNA interaction in neural stem cell differentiation of induced pluripotent stem cells. *Int J Mol Sci*. 2020;21(19):6980.
22. Rajabi H, Aslani S, Abhari A, Sanajou D. Expression profiles of MicroRNAs in stem cells differentiation. *Curr Pharm Biotechnol*. 2020;21(10):906–18.
23. Li H, Zhao X, Wen X, Zeng A, Mao G, Lin R, Hu S, Liao W, Zhang Z. Inhibition of miR-490-5p promotes human adipose-derived stem cells chondrogenesis and protects chondrocytes via the PTPN1/PI3K/AKT Axis. *Front Cell Develop Biol*. 2020;8:573221.
24. Yang Z, Li R, Ao J, Wa QD, Zhang Y, Chen L, Wen J, Chen B, Pan W, Li B, et al. miR-1307-3p suppresses the chondrogenic differentiation of human adipose-derived stem cells by targeting BMP2. *Int J Mol Med*. 2018;42(6):3115–24.
25. Xu J, Kang Y, Liao WM, Yu L. MiR-194 regulates chondrogenic differentiation of human adipose-derived stem cells by targeting Sox5. *PLoS ONE*. 2012;7(3):e31861.
26. Wang Z, Hu T, Jin C, Yu J, Zhu D, Liu J. The anti-tumor effect of miR-539-3p on colon cancer via regulating cell viability, motility, and nude mouse tumorigenicity with CDK14 inhibition. *J Gastrointest Oncol*. 2020;11(5):899–910.
27. Yue F, Peng K, Zhang L, Zhang J. Circ_0004104 accelerates the progression of gastric cancer by regulating the miR-539-3p/RNF2 axis. *Dig Dis Sci*. 2021;66(12):4290–301.
28. Tang Y, Tang Y, Xiang Y, Yan J, Guo K. AK003290 Protects myocardial cells against apoptosis and promotes cardiac function recovery via miR-539-3p/ErbB4 axis in ischemic-reperfusion injury. *DNA Cell Biol*. 2021;40(12):1528–38.
29. Venkatesan JK, Ekici M, Madry H, Schmitt G, Kohn D, Cucchiari M. SOX9 gene transfer via safe, stable, replication-defective recombinant adeno-associated virus vectors as a novel, powerful tool to enhance the chondrogenic potential of human mesenchymal stem cells. *Stem Cell Res Ther*. 2012;3(3):22.
30. Cucchiari M, Orth P, Madry H. Direct rAAV SOX9 administration for durable articular cartilage repair with delayed terminal differentiation and hypertrophy in vivo. *J Mol Med (Berl)*. 2013;91(5):625–36.
31. Cao L, Yang F, Liu G, Yu D, Li H, Fan Q, Gan Y, Tang T, Dai K. The promotion of cartilage defect repair using adenovirus mediated Sox9 gene transfer of rabbit bone marrow mesenchymal stem cells. *Biomaterials*. 2011;32(16):3910–20.
32. Wa Q, He P, Huang S, Zuo J, Li X, Zhu J, Hong S, Lv G, Cai D, Xu D, et al. miR-30b regulates chondrogenic differentiation of mouse embryo-derived stem cells by targeting SOX9. *Exp Ther Med*. 2017;14(6):6131–7.
33. Lee S, Yoon DS, Paik S, Lee KM, Jang Y, Lee JW. microRNA-495 inhibits chondrogenic differentiation in human mesenchymal stem cells by targeting Sox9. *Stem cells and development*. 2014;23(15):1798–808.
34. Guérit D, Philipot D, Chuchana P, Toupet K, Brondello JM, Mathieu M, Jorgensen C, Noël D. Sox9-regulated miRNA-574-3p inhibits chondrogenic differentiation of mesenchymal stem cells. *PLoS ONE*. 2013;8(4):e62582.
35. Akiyama H. Control of chondrogenesis by the transcription factor Sox9. *Mod Rheumatol*. 2008;18(3):213–9.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

