

# Determinants of Face Mask Utilization to Prevent Covid-19 Pandemic among Quarantined Adults in Tigray Region, Northern Ethiopia, 2020

Clinical Nursing Research  
2021, Vol. 30(7) 1107–1112  
© The Author(s) 2021



Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/10547738211013219  
journals.sagepub.com/home/cnr



Mekonnen Haftom, MSc<sup>1</sup>  and Pammla M. Petrucka, PhD<sup>2</sup>

## Abstract

A face mask is a vital component of personal protective equipment to prevent potentially contagious respiratory infections. There was a lack of evidence showing the proportion and determinants of face mask use in Ethiopia. Therefore, this study aimed to identify face mask utilization determinants to prevent spread of the Covid-19 pandemic among quarantined adults in Tigray region, northern Ethiopia. A total of 331 participants selected using a systematic random sampling method were included in the study. An interviewer-administered questionnaire was employed. After describing the variables using frequencies, means, and standard deviations, multivariable logistic regression determined factors associated with face mask utilization to prevent COVID-19 spread. The study participants were primarily males (70%) and mean age was 30.5 (SD = 11) years. Nearly half of the participants reported they did not wear a face mask when leaving home. Face mask utilization was significantly associated with knowledge score, employment status, gender, age, and educational status of the study participants.

## Keywords

determinants, face masks, Covid-19 pandemic, personal protection equipment, pandemics, adult, Ethiopia

## Introduction

Corona virus disease—2019 (Covid-19) was declared a public health emergency of international concern in January 2020 (Cucinotta & Vanelli, 2020; Gagliano et al., 2020). As of May 27, 2020, the disease has caused 5,488,825 cases and 349,095 deaths worldwide. In Africa, 85,815 cases and 2,308 deaths have been reported (World Health Organization [WHO], 2020a). This disease affects all ages, but its severity was higher among the aged and those with underlying chronic diseases (Davies et al., 2020). For instance, the death rate among older people was 48% compared to younger people (Li et al., 2020). The pandemic caused rapid loss of life, joblessness, and deterioration of the healthcare delivery systems, national, and global economies (Lzvorski et al., 2020). In September 2019, it was projected that there would be a catastrophic respiratory disease that is rapidly moving and could result in the deaths of between 50 and 80 million people while impacting the world economy by 5% (Broberg, 2020). In Ethiopia, the first case of Covid-19 was confirmed on March 13, 2020, in Addis Ababa.

The pandemic has since spread to all parts of the country. As of May 27, 2020, the Africa news reported 731 confirmed cases and six deaths (Addis News, 2020). The use of a face

mask is one of the critical measures to prevent the virus' spread. However, it should be integrated with physical distancing, hand hygiene, and related preventive measures (WHO, 2020b, 2020c, 2020d). Infected individuals can be asymptomatic or may present with mild to severe manifestation. The risk of transmission from asymptomatic individuals is high, making the disease difficult to control (Howard et al., 2020; Wang et al., 2020; WHO, 2020e). Anticipating the disease's pandemic spread continues in developing countries, its impact will be more severe than that seen in the world's developed regions due to weak or unresponsive health systems (Moorthy et al., 2020).

A face mask is a vital component of personal protective equipment to prevent potentially contagious respiratory infections (Wei et al., 2020). There were controversies in using a mask. Issues regarding its effectiveness and fear of shortage of

<sup>1</sup>Adigrat University, Ethiopia

<sup>2</sup>University of Saskatchewan, Regina, Canada

## Corresponding Author:

Mekonnen Haftom, MSc, Department of Nursing, College of Medicine and Health Sciences, Adigrat University, Adigrat 50, Ethiopia.

Email: haftommekonnen@gmail.com

resources were some of the claimed reasons some experts recommended not to use a face mask (Madad et al., 2020). However, pieces of evidence also showed that mask use could decrease disease transmission and mortality (Chu et al., 2020; Eikenberry et al., 2020; Feng et al., 2020; Klompas et al., 2020; WHO, 2020b). No research studies have been identified to determine the proportions and determinants of face mask utilization to prevent Covid-19 in Ethiopia. Therefore, this study aimed to identify face mask utilization determinants to prevent the Covid-19 pandemic among quarantined adults in the Tigray region, northern Ethiopia.

## Methods and Materials: Study Design, Setting, and Population

This manuscript is a part of a broader study considering knowledge, attitude, and practice towards the Covid-19 pandemic in Tigray region. An institutional-based cross-sectional study was conducted among adults quarantined in Tigray Region, Northern Ethiopia. The criteria for quarantine were international travel history, travel history from Addis Ababa (where community transmission started), and exposure to positive individuals.

A total of 331 randomly selected participants aged 18 to 69 were interviewed from May 15 to 27, 2020. Using a single population proportion formula to calculate sample size assuming a 95% confidence level, 5% margin of error, and 50% of the population wear a face mask when leaving home ( $p=.5$ ), it yielded a calculated sample size of  $n=384$ . Since the total source population was less than 10,000 ( $N=2102$ ), the estimated sample size ( $n=384$ ) was reconsidered using a correction formula. The final sample size after correction ( $nf=326$ ) with the addition of a 5% non-response rate yielding a sample size of 343.

## Recruitment Procedure

Multi-stage sampling was used. The region is clustered into zones ( $n=7$ ). Three quarantine centers in each zone were randomly selected. The total sample size is proportionally allocated to the number of confined and isolated individuals over the most recent 2 weeks. The sampling interval was determined by dividing the number of individuals admitted to the centers in the selected 2 weeks by the sample allocated to that center, which yield a  $k$  of 7. A systematic random sampling method was used to choose the study participants. The first participant (between 1 and 7) was selected using a lottery method. Therefore, every seventh individual was enrolled in the study until the calculated sample size was achieved.

## Study Instrument and Data Collection Procedure

The instrument is adapted from similar studies done in Tanzania and China. The questionnaire was found to be valid,

reliable, and used in other studies (Advani et al., 2020; Sommerstein et al., 2020). The tool had three sections: the socio-demographic profile (sex, age, educational status, and employment status); knowledge about clinical characteristics of COVID-19 (items 1–4); knowledge about modes of transmission of COVID-19 (items 5–8); and knowledge about prevention and control of COVID-19 (items 9–13). Each item has “true” (coded as 0), “false” (coded 1), and “I don’t know” (coded 2) options. For analysis, it was again re-coded to sum the total scores as 1 for correct answers and 0 for both incorrect and I don’t know responses. The possible score ranges from 0 to 13; with the higher scores indicating good knowledge. For the purposes of this study, a score  $\geq 10$  being acceptable levels of knowledge (Zhong et al., 2020). The third section was about face mask utilization when leaving home. We collected data using a self-administered questionnaire for the literate participants. If the respondent was illiterate, the data were collected using face-to-face interviews by trained data collectors.

As part of the quarantine protocol, only one adult was allowed to be in a single room. Therefore, by default privacy was kept by keeping distance between the interviewer and the interviewee. Interviews were based on convenience for the individual participant.

## Data Quality Assurance and Control

The questionnaire was prepared in English and back-translated to the local language Tigrigna. Also, the tool’s accuracy was checked by back translating to English by experts who were blind to the original instrument.

Before starting the data collection, a pre-test was done on 10% (34 individuals) of the total sample size, and an amendment was made accordingly. Also, data were collected by trained data collectors, and the completed questionnaires’ completeness was ensured thorough manual checking daily. Data collectors were trained for 3 days regarding the study’s objectives, data collection procedure, and ethical considerations a week ahead of the actual data collection.

## Data Processing and Analysis

Data were entered and cleaned using Epi-data manager™ version 3.1, and Statistical Package for the Social Sciences (SPSS)™ version was used for analysis. Frequency, means, and standard deviations (SDs) of participants’ demographic characteristics were calculated to identify possible baseline differences and presented using tables and text. The binary logistic regression model was used to determine the magnitude, direction, and strength of association between demographic profiles, knowledge scores, and face mask utilization. Variables significant at  $p < .25$  with a face mask utilization were selected for a multivariate logistical regression model analysis. An odds ratio with 95% confidence level was computed, and  $p$ -value  $< .05$  was considered a significant association.

## Ethical Consideration

The study considered the Helsinki Declaration. The Tigray Regional Health Bureau (TRHB) approved the study. A support letter was submitted to the region's selected quarantine centers, and letters of permission were secured from each center's administrative bodies and coordinators. Participants recruited to the study received a verbal explanation of the research and were provided a written information sheet. They were informed regarding compensation (no compensation was provided). All potential participants who agreed to participate provided written consent to continue with the interviews. The confidentiality of information obtained was kept, and respondents' names were not recorded.

## Results

### Demographic Characteristics

Of the 343 approached participants, 331 completed the questionnaire, making the response rate 96.5%. Table 1 shows the socio-demographic profiles and knowledge scores of the study participants. The mean age was 30.5 years ( $SD=11$ ). Seventy percent of the participants were males, and more than half of the study participants (59.8%) were in the age range 18 to 29. About one-third of the study participants were self-employed (31.1%) and have completed secondary school (36%). The mean (standard deviation) knowledge score was 8.73 ( $\pm 2.64$ ). The rate of the correct answer was 67.2% ( $8.73/13 \times 100$ ). Less than half of participants [142 (42.9%) (95% confidence interval (CI): 37.5%–48%)] were knowledgeable (scored  $\geq 10$ ). The remaining 189 (57.1%) (95% CI: 52%–62.5%) were not knowledgeable. Forty-six percent (95% CI: 40.2%–51%) of the participants did not use a face mask when leaving home.

### Determinants of Low Utilization of Face Mask

Multivariable logistic regression showed that males were 2 (95% CI: 1.1–3.2) times more likely to wear masks when leaving home. The odds of mask utilization amongst the age group of  $\geq 50$  years and 30 to 49 years were 12 (95% CI: 3.2–45.1) and 2.5 (95% CI: 1.4–4.6) times higher than those aged 18 to 29. Controlling the effect of being a student, the odds of mask use among public and private sector employees were 5.2 (95% CI: 1.2–23) and 3.6 (95% CI: 1.1–12.5). The odds of mask use among farmers were 68% [0.32(95% CI: 0.1–0.9)] lower than students. Those who completed college or university were 8.1 (95% CI: 2.1–31.7) times more likely to use a mask when leaving home than those who cannot read and write. The odds of mask use amongst knowledgeable participants were 2 (95% CI: 1.2–3.3) times higher than the non-knowledgeable counterparts (Table 2).

**Table 1.** Demographic Profiles of Study Participants ( $n=331$ ).

Variables	Category	<i>n</i>	%
Gender	Male	230	69.5
	Female	101	30.5
Age	18–29	198	59.8
	30–49	104	31.4
	Above 50	29	8.8
Occupation	Student	39	11.8
	Public sector	28	8.5
	Private sector	24	7.3
	Self-employed	103	31.1
	Not employed	80	24.2
	Farmer	38	11.5
	Retired	4	1.2
Educational status	Others	15	4.5
	Illiterate	37	11.2
	Read and write	21	6.3
	Primary school	90	27.2
	Secondary school	119	36
Knowledge score	Illiterate	37	11.2
	Acceptable	142	42.9
	Not acceptable	189	57.1

## Discussion

After the virus emerged in China, it is argued that information was suppressed and erroneous. China initially reported the novel virus showed no evidence of human-to-human transmission (Azlan et al., 2020; Byanaku & Ibrahim, 2020). As per the authors' knowledge, this is the first study of persons in quarantine centers to assess mask utilization determinants to prevent Covid-19 in Ethiopia. This study has provided evidence-based data regarding face mask utilization. Nearly half of the study participants have used face masks when leaving home in recent days.

The lack of use of face masks is not consistent with the recommendations (Davidson, 2020; Greenhalgh et al., 2020; Ogden, 2020) that everyone should wear a mask when going out. Some countries (Mahase, 2020) have declared state mandates on mask use when leaving home, while using face mask was underemphasized in other countries (Greenhalgh et al., 2020). However, Ethiopia did not declare mandates on face mask use until the time of data collection. This position could be due to controversies to use or not to use face masks (Lyu & Wehby, 2020; Tso & Cowling, 2020; Wang et al., 2020) and may have contributed to the low report of mask used in our study area.

Our findings indicated that educational status, knowledge score, age, gender, and occupation of study participants were significant predictors of mask use. The results of this study demonstrate that a higher level of education and acceptable level of knowledge towards COVID-19 (Advani et al., 2020; Chen et al., 2020; Fisman et al., 2020; Kuo et al., 2011;

**Table 2.** Multivariable Logistic Regression on Socio-demographic Characteristics, Knowledge Score, and Utilization Face Mask (n = 331).

Variables	Category	Face mask used (%)	Face mask not used (%)	COR (95% CI)	AOR (95% CI)
Gender	Male	132 (39.9)	98 (29.6)	2.7 (1.7–4.3)	2 (1.1–3.2)**
	Female	48 (14.5)	53 (16)	Ref	Ref
Age	18–29	110 (33.2)	88 (26.6)	Ref	Ref
	30–49	57 (17.2)	47 (14.2)	2 (1.2–3.2)	2.5 (1.4–4.6)***
	Above 50	13 (3.9)	16 (4.8)	2.9 (1.2–6.9)	12 (3.2–45.1)***
Occupation	Student	22 (6.6)	17 (5.1)	Ref	Ref
	Public sector	19 (5.7)	9 (2.7)	6.9 (1.8–26.9)	5.2 (1.2–23)**
	Private sector	11 (3.3)	13 (3.9)	3.6 (1.1–11.5)	3.6 (1.1–12.5)**
	Self-employed	68 (20.5)	35 (10.6)	1.7 (0.8–3.5)	1.7 (0.8–3.9)
	Not employed	39 (11.8)	41 (12.4)	0.7 (0.3–1.4)	0.9 (0.4–2)
	Farmer	15 (4.5)	23 (6.9)	0.3 (0.1–0.7)	0.32 (0.1–0.9)**
	Retired	3 (0.9)	1 (0.3)	2.7 (0.3–28)	1.7 (0.1–19.5)
Educational status	Others#	3 (0.9)	12 (3.6)	0.2 (0.1–0.9)	0.4 (0.1–2)
	Illiterate	11 (3.3)	26 (7.9)	Ref	Ref
	Read and write	8 (2.4)	13 (3.9)	3.3 (1–10.6)	4.9 (1.2–20.5)**
	Primary school	53 (16)	37 (11.2)	5.4 (2.1–13.5)	6.3 (1.7–22.6)***
	Secondary school	71 (21.5)	48 (14.5)	5.9 (2.4–14.5)	6.9 (2–24.4)***
Knowledge score	College or university	37 (11.2)	27 (8.2)	10.5 (4–28.2)	8.1 (2.1–31.7)***
	Acceptable	94 (24.4)	48 (14.5)	2.8 (1.7–4.3)	2 (1.2–3.3)*
	Not acceptable	86 (26)	103 (31)	Ref	Ref

Note. AOR = adjusted odds ratio; COR = crude odds ratio;

#Others = driver, housewife, daily laborers.

\* $p < .05$ . \*\* $p < .01$ . \*\*\* $p < .001$ .

Taylor et al., 2009; Wong et al., 2020) increases face mask use by participants when leaving home. These findings indicate younger participants were less likely to use a face mask, which aligns with other studies (Chen et al., 2020; Lau et al., 2007) that older people tended to use a mask. COVID-19 is less common amongst youngsters (Bai et al., 2020; Lee et al., 2020; Tang & Wong, 2004; Tomar et al., 2020) compared to their older adult counterparts, which may contribute to the low utilization of face masks among those aged 18 to 29 compared to their elder counterparts.

Some studies (Brodin, 2020; Lee et al., 2020) were inconsistent with our finding that females were more likely to use face masks. Participants employed in private and public sectors are more likely to use face masks than students. However, farmers were least likely to use a face mask. The reason for this variability could be explained as employed participants might be obliged to use masks when they are on duty, and limited access to face masks and costs might cause farmers not to use them.

## Limitation

This is a small-scale cross-sectional study, which may limit the generalizability of the results across different periods and different communities. Therefore, a large-scale country level study would have provided more generalizable data. When the rapid response team traced the study participants, they

provided education regarding the pandemic on the way to the quarantine centers before the entry interview (data collection), which might have made study participants more aware of mask use compared to the community as a whole. Social desirability bias was likely regarding face mask use.

## Conclusion

Nearly half of the total participants did not use a face mask when leaving home. Employment status level of education, gender, age, and knowledge score were significantly associated with face mask utilization by adults in Tigray Region, Northern Ethiopia.

## Author Contributions

HM Co-conceptualized the research; conducted on ground research; co-analysis of data; co-writing of all copies of the contribution. PMP Co-conceptualized the research; co-analysis and co-writing of the contribution.

## Availability of Data and Material

The datasets used and analyzed during the study are available from the corresponding author on reasonable request.

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.



## Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

## Ethics Approval and Consent to Participate

Tigray Regional Health Bureau approved the study. Written informed consent was obtained from each participant.

## ORCID iD

Mekonnen Haftom  <https://orcid.org/0000-0002-8582-0280>

## References

- Addis News. (2020). Coronavirus-Ethiopia: Notification Note on COVID-19 Situational Update 27 May 2020.
- Advani, S. D., Smith, B. A., Lewis, S. S., Anderson, D. J., & Sexton, D. J. (2020). Universal masking in hospitals in the COVID-19 era: Is it time to consider shielding? *Infection Control & Hospital Epidemiology*, *1*(9), 1066–1067.
- Azlan, A. A., Hamzah, M. R., Sern, T. J., Ayub, S. H., & Mohamad, E. (2020). Public knowledge, attitudes and practices towards COVID-19: A cross-sectional study in Malaysia. *PLoS One*, *15*(5), e0233668.
- Bai, Y., Yao, L., Wei, T., Tian, F., Jin, D.-Y., Chen, L., & Wang, M. (2020). Presumed asymptomatic carrier transmission of COVID-19. *JAMA*, *323*(14), 1406–1407.
- Broberg, M. (2020). A critical appraisal of the World Health Organisation's international health regulations (2005) in times of pandemic—it is time for revision. *European Journal of Risk Regulation*, *1*(2), 202–209.
- Brodin, P. (2020). Why is COVID-19 so mild in children? *Acta Paediatrica*, *109*(6), 1082–1083.
- Byanaku, A., & Ibrahim, M. (2020). Knowledge, attitudes, and practices (KAP) towards COVID-19: A quick online cross-sectional survey among Tanzanian residents. *medRxiv*.
- Chen, X., Ran, L., Liu, Q., Hu, Q., Du, X., & Tan, X. (2020). Hand hygiene, mask-wearing behaviors and its associated factors during the COVID-19 epidemic: A cross-sectional study among primary school students in Wuhan, China. *International Journal of Environmental Research and Public Health*, *17*(8), 2893.
- Chu, D. K., Akl, E. A., Duda, S., Solo, K., Yaacoub, S., Schunemann, H. J., El-harakeh, A., Bognanni, A., Lotfi, T., Loeb, M., & Hajjzadeh, A. (2020). Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and Covid-19: A systematic review and meta-analysis. *The Lancet*, *395*(10242), 1973–1987.
- Cucinotta, D., & Vanelli, M. (2020). WHO declares COVID-19 a pandemic. *Acta Bio Medica: Atenei Parmensis*, *91*(1), 157.
- Davidson, H. (2020). Chinese inquiry exonerates coronavirus whistleblower doctor. *The Guardian*, 20.
- Davies, N. G., Klepac, P., Liu, Y., Prem, K., Jit, M., Eggo, R. M., & Eggo, R. M. (2020). Age-dependent effects in the transmission and control of COVID-19 epidemics. *Nature Medicine*, *26*(8), 1205–1211.
- Eikenberry, S. E., Mancuso, M., Iboi, E., Phn, T., Eikenberry, K., Kuang, Y., Kostelich, E., & Gumel, A. B. (2020). To mask or not to mask: Modeling the potential for face mask use by the general public to curtail the Covid-19 pandemic. *Infectious Disease Modelling*, *5*, 293–308.
- Feng, S., Shen, C., Xia, N., Song, W., Fan, M., & Cowling, B. J. (2020). Rational use of face masks in the Covid-19 pandemic. *The Lancet Respiratory Medicine*, *8*, 434–436.
- Fisman, D. N., Greer, A. L., & Tuite, A. R. (2020). Bidirectional impact of imperfect mask use of reproduction number of Covid-19: A next generation matrix approach. *Infectious Disease Modelling*, *5*, 405–408.
- Gagliano, A., Villani, P. G., Manelli, A., Paglia, S., Bisagni, P. A., Perotti, G. M., Storti, E., & Lombardo, M. (2020). COVID-19 epidemic in the middle province of Northern Italy: Impact, logistics, and strategy in the first line hospital. *Disaster Medicine and Public Health Preparedness*, *14*(3), 372–376.
- Greenhalgh, T., Chan, X. H., Khunti, K., Durand-Moreau, Q., Straube, S., Devane, D., Toomey, E., Ireland, E. S., & Ireland, C. (2020). *What is the efficacy of standard face masks compared to respirator masks in preventing Covid-type respiratory illnesses in primary care staff*. Centre for Evidence-Based Medicine, Nuffield Department of Primary Care Health Sciences, University of Oxford.
- Greenhalgh, T., Schmid, M. B., Cponka, T., Bassler, D., & Gruer, L. (2020). Face masks for the public during the covid-19 crisis. *BMJ (Clinical research ed)*, *369*, m1435.
- Howard, J., Huang, A., Li, Z., Tufekci, Z., Zdimal, V., van der Westhuizen, H.-M., von Delft, A., Price, A., Fridman, L., Tang, L. H., & Tang, V. (2020). Face masks against Covid-19: An evidence review. *Proceedings of the National Academy of Sciences*, *118*(4), e20146118.
- Klompas, M., Morris, C. A., Sinclair, J., Pearson, M., & Shenoy, E. S. (2020). Universal masking in hospitals in the Covid-19 era. *New England Journal of Medicine*, *382*(21), e63.
- Kuo, P.-C., Huang, J.-H., & Liu, M.-D. (2011). Avian influenza risk perception and preventive behavior among traditional market workers and shoppers in Taiwan: Practical implications for prevention. *PLoS One*, *6*(9), e24157.
- Lau, J. T., Kim, J. H., Tsui, H. Y., & Griffiths, S. (2007). Anticipated and current preventive behaviors in response to an anticipated human-to-human H5N1 epidemic in the Hong Kong Chinese general population. *BMC Infectious Diseases*, *7*(1), 18.
- Lee, P.-I., Hu, Y.-L., Chen, P.-Y., Huang, Y.-C., & Hsueh, P.-R. (2020). Are children less susceptible to COVID-19? *Journal of Microbiology, Immunology, and Infection*, *53*(3), 371.
- Li, H., Wang, S., Zhong, F., Bao, W., Li, Y., Liu, L., Wang, H., & He, Y. (2020). Age-dependent risks of Incidence and mortality of COVID-19 in Hubei Province and other parts of China. *Frontiers in Medicine*, *7*, 190.
- Lyu, W., & Wehby, G. L. (2020). Community use of face masks and COVID-19: Evidence from a natural experiment of state mandates In the US: Study examines impact on COVID-19 growth rates associated with state government mandates requiring face mask use in public. *Health Affairs*, *39*(8), 1419–1425. <https://doi.org/10.1377/hlthaff.2020.00818>
- Lzvorski, I., Mahajan, S., Moorthy, L., & Vincelette, G. A. (2020). *A policy framework for mitigating the economic impact of COVID-19*. <https://www.brookings.edu/blog/future-development/2020/04/20/a-policy-framework-for-mitigating-the-economic-impact-of-covid-19/>

- Madad, S., Moskovitz, J., Boyce, M. R., Cagliuso, N. V., & Katz, R. (2020). Ready or not, patients will present: Improving urban pandemic preparedness. *Disaster Medicine and Public Health Preparedness*, 16, 1–4.
- Mahase, E. (2020). Covid-19: What is the evidence for cloth masks? *BMJ*, 369, m1435.
- Moorthy, V., Restrepo, A. M. H., Preziosi, M.-P., & Swaminathan, S. (2020). Data sharing for novel coronavirus (COVID-19). *Bulletin of the World Health Organization*, 98(3), 150.
- Ogden, C. (2020). The role of competing narratives in China and the West's response to Covid-19. *British Journal of Chinese Studies*, 10.
- Sommerstein, R., Fux, C. A., Vuichard-Gysin, D., Abbas, M., Marschall, J., Balmelli, C., Troillet, N., Harbarth, S., Schlegel, M., & Widmer, A. (2020). Risk of SARS-CoV-2 transmission by aerosols, the rational use of masks, and protection of healthcare workers from COVID-19. *Antimicrobial Resistance & Infection Control*, 9(1), 1–8.
- Tang, C. S.-K., & Wong, C.-Y. (2004). Factors influencing the wearing of facemasks to prevent the severe acute respiratory syndrome among adult Chinese in Hong Kong. *Preventive Medicine*, 39(6), 1187–1189.
- Taylor, M., Raphael, B., Barr, M., Agho, K., Stevens, G., & Jorm, L. (2009). Public health measures during an anticipated influenza pandemic: Factors influencing willingness to comply. *Risk Management a Healthcare Policy*, 2, 9.
- Tomar, B. S., Singh, P., Nathiya, D., Suman, S., Raj, P., Tripathi, S., & Chauhan, D. S. (2020). Indian community's knowledge, attitude & practice towards COVID-19. *medRxiv*.
- Tso, R. V., & Cowling, B. J. (2020). Importance of face masks for COVID-19—a call for effective public education. *Clinical Infectious Diseases*, 71(16), 2195–2198.
- Wang, J., Pan, L., Tang, S., Ji, J. S., & Shi, X. (2020). Mask use during COVID-19: A risk adjusted strategy. *Environmental Pollution*, 266, 115099.
- Wang, L., Shi, Y., Xiao, T., Fu, J., Feng, X., Mu, D., Feng, Q., Hei, M., Hu, X., Li, Z., & Lu, G. (2020). Chinese expert consensus on the perinatal and neonatal management for the prevention and control of the 2019 novel coronavirus infection. *Annals of Translational Medicine*, 8(3), 47.
- Wei, W. E., Li, Z., Chiew, C. J., Yong, S. E., Toh, M. P., & Lee, V. J. (2020). Presymptomatic transmission of SARS-CoV-2—Singapore, January 23–March 16, 2020. *Morbidity and Mortality Weekly Report*, 69(14), 411.
- Wong, S. H., Teoh, J. Y., Leung, C.-H., Wu, W. K., Yip, B. H., Wong, M. C., & Hui, D. S. (2020). Covid-19 and public interest in face mask use. *American Journal of Respiratory and Critical Care Medicine*, 202(3), 453–455.
- World Health Organization. (2020a). *Coronavirus disease (COVID-19): Situation report, 182*. Author.
- World Health Organization. (2020b, June 5). *Advice on the use of masks in the context of COVID-19: Interim guidance*. Author.
- World Health Organization. (2020c, March 19). *Advice on the use of masks in the community, during home care, and in health care settings in the context of COVID-19: Interim guidance*. Author.
- World Health Organization. (2020d). *Rational use of personal protective equipment (PPE) for coronavirus disease and considerations during severe shortages: Interim guidance*. Author.
- World Health Organization. (2020e, March 19). *Rational use of personal protective equipment (PPE) for coronavirus disease (COVID-19): Interim guidance*. Author.
- Zhong, B.-L., Luo, W., Li, H.-M., Zhang, Q.-Q., Liu, X.-G., Li, W.-T., & Li, Y. (2020). Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: A quick online cross-sectional survey. *International Journal of Biological Sciences*, 16(10), 1745.

### Author Biographies

**Mekonnen Haftom** holds a MSc from Addis Ababa University School of Medicine. He currently holds a position as lecturer at Adigrat University, Adigrat, Ethiopia.

**Pammla M. Petrucka** holds a PhD from the University of Alberta, Canada. She currently is a Professor at the College of Nursing, University of Saskatchewan, holds an adjunct appointment at the Nelson Mandela African Institute of Science and Technology.