



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

- 1 Blood Pressure Lowering Treatment Trialists' Collaboration. Age-stratified and blood-pressure-stratified effects of blood-pressure-lowering pharmacotherapy for the prevention of cardiovascular disease and death: an individual participant-level data meta-analysis. *Lancet* 2021; **398**: 1053–64.
- 2 Law MR, Wald NJ, Morris JK, Jordan RE. Value of low dose combination treatment with blood pressure lowering drugs: analysis of 354 randomised trials. *BMJ* 2003; **326**: 1427–31.
- 3 Law MR, Morris JK, Wald NJ. Use of blood pressure lowering drugs in the prevention of cardiovascular disease: meta-analysis of 147 randomised trials in the context of expectations from prospective epidemiological studies. *BMJ* 2009; **338**: b1665.
- 4 Law MR, Wald NJ, Rudnicka AR. Quantifying effect of statins on low density lipoprotein cholesterol, ischaemic heart disease, and stroke: systematic review and meta-analysis. *BMJ* 2003; **326**: 1423–27.
- 5 Wald NJ, Law MR. A strategy to reduce cardiovascular disease by more than 80%. *BMJ* 2003; **326**: 1419–23.

Authors' reply

We thank Nicholas Wald and colleagues for their interest in our work.¹ They propose that our findings merely confirmed what has been previously known, as reported by them.

We summarised¹ relevant previous research and the evidence gap around the effects of blood pressure reduction in the very young and the very old, and whether the recommended use of age-dependent blood pressure thresholds for initiation or intensification of treatment by some clinical practice guidelines^{2,3} are justified. Wald and colleagues' work was very valuable but did not investigate these particular questions, and we could therefore neither confirm nor refute their findings. We reviewed and cited other meta-analyses that attempted to investigate these questions, and we discuss the novelty and added value of our work to clinical decision making in their context.

In comparison with earlier reports, our meta-analysis benefited from access to the largest source of individual participant-level data.^{4,5} This meant that we were able to simultaneously stratify effects by age (in 10-year increments from age <55 years to ≥85 years) and blood pressure (in 10 mm Hg increments from <120 mm Hg to ≥170 mm Hg

systolic, and from <70 mm Hg to ≥110 mm Hg diastolic) at baseline, while standardising for differing intensities of blood pressure reduction in individual trials. No meta-analysis has previously provided this level of detail. This advantage is analogous to looking at a dataset through a microscope; the higher resolution analysis helps reveal previously hidden relationships (even if theoretically assumed by some). For instance, although beneficial effects in lowering cardiovascular events were seen in most age groups (and blood pressure levels), the relative risk reductions per 5 mm Hg reduction in systolic blood pressure appeared to diminish with age and in people aged 85 years or older, the effect was highly uncertain (hazard ratio 0.99, 95% CI 0.87–1.12). These results do not seem to support Wald and colleagues' generalisation that any aged 50 years or older stands to benefit similarly from blood pressure-lowering treatment.

We do not dispute the need for better solutions that help overcome the implementation gap, as Wald and colleagues have suggested. However, providing reliable and precise quantification of expected treatment effects in a wide range of at-risk groups,¹ as we have done, is an important and necessary step towards informing better decisions and addressing those implementation gaps.

KR reports grants from the British Heart Foundation (BHF), the UK Research Innovation Global Challenges Research Fund, the Oxford Martin School, and the National Institute for Health Research Oxford Biomedical Research Centre; and personal fees from *British Medical Journal Heart* and for delivering Medscape lectures. RM declares working with Omron on the development and evaluation of a blood pressure telemonitoring system for which consultancy and licensing fees are paid to his institution. JC reports grants from the National Health and Medical Research Council of Australia. DC reports grants from the BHF. MW declare no competing interests.

*Kazem Rahimi, Richard McManus, Malgorzata Wamil, John Chalmers, Dexter Canoy

kazem.rahimi@wrh.ox.ac.uk

Deep Medicine, Nuffield Department of Women's and Reproductive Health (KR, MW, DC) and Nuffield Department of Primary Care Health Sciences (RM),

University of Oxford, Oxford OX1 2BQ, UK; NIHR Oxford Biomedical Research Centre, Oxford University Hospitals NHS Foundation Trust, Oxford, UK (KR, DC); The George Institute for Global Health, University of New South Wales, Sydney, NSW, Australia (JC)

- 1 Blood Pressure Lowering Treatment Trialists' Collaboration. Age-stratified and blood-pressure-stratified effects of blood-pressure-lowering pharmacotherapy for the prevention of cardiovascular disease and death: an individual participant-level data meta-analysis. *Lancet* 2021; **398**: 1053–64.
- 2 National Institute for Health and Care Excellence. Hypertension in adults: diagnosis and management. Aug 28, 2019. <https://www.nice.org.uk/guidance/ng136> (accessed Oct 26, 2021).
- 3 Williams B, Mancia G, Spiering W, et al. 2018 ESC/ESH guidelines for the management of arterial hypertension. *Eur Heart J* 2018; **39**: 3021–104.
- 4 Rahimi K, Canoy D, Nazarzadeh M, et al. Investigating the stratified efficacy and safety of pharmacological blood pressure-lowering: an overall protocol for individual patient-level data meta-analyses of over 300 000 randomised participants in the new phase of the Blood Pressure Lowering Treatment Trialists' Collaboration (BPLTTC). *BMJ Open* 2019; **9**: e028698.
- 5 Canoy D, Copland E, Nazarzadeh M, et al. Antihypertensive drug effects on long-term blood pressure: an individual-level data meta-analysis of randomised clinical trials. *Heart* 2022; published online Jan 20. <https://doi.org/10.1136/heartjnl-2021-320171>.

An open letter to the Executive Board of WHO from the surgical and anaesthesia community

An estimated 28 million surgeries were cancelled worldwide during the first 3 months of the COVID-19 pandemic, a number which might now be as high as 115 million—more than a third of all surgeries annually.^{1,2} We applaud the work of WHO over the past 2 years in managing and reducing the impacts of the pandemic, but are critical about the efforts taken to mitigate the impact of cancelled surgeries despite the recognition of surgery as an essential part of universal health coverage. Given the unequal impacts the pandemic has had across clinical disciplines, countries, and populations, we strongly support the development of a binding worldwide pandemic treaty.



Published Online
March 28, 2022
[https://doi.org/10.1016/S0140-6736\(22\)00521-9](https://doi.org/10.1016/S0140-6736(22)00521-9)

With the sharp rise in non-communicable diseases worldwide, the demand for surgical care is rapidly increasing. Surgical interventions are needed for 80% of injured patients and more than half of patients with cancer.³ Given the high versatility and the clear value surgical teams and infrastructure have added to the pandemic response, it is surprising that surgical system strengthening has not received more attention as part of pandemic preparedness initiatives thus far. We suggest a paradigm shift where access to surgical care is mainstreamed into pandemic preparedness policies. A mainstreaming approach would entail that the pandemic treaty assures that every policy adopted or implemented from this treaty has been evaluated for its impact on national-level surgical care provision. Only policies that do not harm surgical care provision should be included in the final version of the treaty so as to avoid detrimental impact. While novel in global surgery discourse, this would entail a policy approach similar to WHO's Health in All Policies.

As defined by the UN General Assembly at the beginning of the pandemic, this approach involves considering the systemic impact of policy decisions on health and making those decisions across different sectors to achieve synergy, equity, and improved health outcomes.⁴ By mainstreaming surgery into pandemic preparedness policy, we suggest that surgical care provision is taken into consideration for every policy recommendation or operative paragraph. This way, true health systems strengthening can take place to achieve a system where no-one is left behind before, during, or after a pandemic.

The 150th WHO Executive Board established a Standing Committee on Pandemic and Emergency Preparedness and Response to draft and negotiate the Pandemic Treaty under the sole auspices of the Member States.⁵ We urge the committee to recognise the importance of surgery as an essential

part of health systems by: (1) ensuring participation of civil society, including the surgical community, as key stakeholders in the treaty negotiation process; and (2) mainstreaming surgical care into the final draft of the Pandemic Preparedness Treaty.

The surgical and anaesthesia communities stand ready to be involved in this process.

We declare no competing interests. Signatories of this Correspondence are listed in the appendix.

Anam N Ehsan, Kee B Park,
*Manon Pigeolet
manon.pigeolet@outlook.com

Program in Global Surgery and Social Change, Department of Global Health and Social Medicine, Harvard Medical School, Boston, MA 02115, USA (ANE, KBP, MP); Medical College, The Aga Khan University, Karachi, Pakistan (ANE); World Federation of Neurosurgical Societies (WFNS), WFNS Central Office, Nyon, Vaud, Switzerland (KBP); Faculty of Medicine, Université Libre de Bruxelles, Brussels, Belgium (MP)

- 1 COVIDSurg Collaborative. Elective surgery cancellations due to the COVID-19 pandemic: global predictive modelling to inform surgical recovery plans. *Br J Surg* 2020; **107**: 1440–49.
- 2 Simoes J, Bhangu A, COVIDSurg Collaborative. Should we be re-starting elective surgery? *Anaesthesia* 2020; **75**: 1563–65.
- 3 Pigeolet M, Marks I, Bentounsi Z, Wanjau W. Letter to Editor: can mainstreaming surgery advocacy into NCD advocacy help us overcome the NCD epidemic? *World J Surg* 2019; **43**: 2114–15.
- 4 UN General Assembly. Resolution adopted by the General Assembly on 11 December 2019. Global health and foreign policy A/RES/74/20. New York, NY: United Nations, 2020.
- 5 WHO. Executive Board 150th session EB150/17: Standing Committee on Pandemic and Emergency Preparedness and Response. Geneva: World Health Organization, 2021.

Host-targeting oral antiviral drugs to prevent pandemics

Unlike bacteria, viruses must use host cells to replicate. This has enabled us to identify the Achilles heel of many viruses. We want to exploit this knowledge for the therapeutic targeting of current major human pathogens, such as coronaviruses and influenza for which there is a great unmet need. The orally available small molecule broad-spectrum antiviral


compounds we have developed over many years target the host¹ and are thus resistant to viral mutations. Host-targeting drugs can also be employed against newly emerging viruses even before detailed information about the virus becomes available.^{2,3} This will be crucial in preventing the inevitable new epidemics from turning into pandemics.

Most enveloped viruses need to use a sugar-mediated pathway in the infected human host cell to form their correct three-dimensional structures, which involves adding and processing glycans on viral envelope glycoproteins.^{2,3} The glycosylation process involves enzymes in the cell trimming the sugars of the viral glycoproteins for entry into this protein folding quality control pathway. Drugs that partially inhibit these enzymes prevent the virus from making proper use of this folding pathway and lead to inhibition of secretion of infectious virus.^{4–8}

Over the past 25 years, we have developed a class of drugs called iminosugars—orally available sugar mimetics that are recognised by and inhibit these sugar processing enzymes that most enveloped viruses rely on.¹ The family of iminosugars derive from the parent compound initially isolated from the leaves of the mulberry tree.

Safety and efficacy data in animals accumulated over the past 20 years show that iminosugar derivatives reduce viral levels and increase survival in animal models of chronic hepatitis B infection,⁶ hepatitis C, Japanese encephalitis, influenza,⁹ and dengue.^{7,8} When tested in vitro against over 31 clinical HIV isolates, including HIV-1, HIV-2, and multidrug resistant strains, iminosugars are active against a diverse panel of HIV-1 from different genetic subtypes and geographical regions, and against HIV-2 isolates and mutants resistant to antiretrovirals.¹⁰ All HIV isolates tested were rendered non-infectious by iminosugar treatment. Similarly,

See Online for appendix



Kateryna Konr / Science Photo Library

Published Online
March 25, 2022
[https://doi.org/10.1016/S0140-6736\(22\)00454-8](https://doi.org/10.1016/S0140-6736(22)00454-8)