




BMJ Open Technology-based group exercise interventions for people living with dementia or mild cognitive impairment: a scoping review protocol

Lillian Hung ^{1,2}, Hannah Levine ³, Paavan Randhawa,⁴ Juyoung Park ⁵

To cite: Hung L, Levine H, Randhawa P, *et al.* Technology-based group exercise interventions for people living with dementia or mild cognitive impairment: a scoping review protocol. *BMJ Open* 2022;**12**:e055990. doi:10.1136/bmjopen-2021-055990

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-055990>).

Received 02 August 2021
Accepted 18 February 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Hannah Levine;
hlevine2016@health.fau.edu

ABSTRACT

Introduction More than 50 million people worldwide are living with dementia in 2020, and this number is expected to double every 20 years. Physical exercise is a growing field in non-pharmacological interventions for dementia care. Due to public health measures during the COVID-19 pandemic, more people have considered adapting to technology-based exercise via digital devices. This scoping review will explore evidence relating to the use of technology-based group exercise by people with dementia or mild cognitive impairment.

Methods and analysis This review will follow the Joanna Briggs Institute scoping review methodology to review literature published between June and December 2021. This review is designed to identify existing types of technology-based group exercise interventions for people with dementia. The review will provide a synthesis of current evidence on the outcome and impacts of technology-based group exercise. The context of this review will include homes, assisted living facilities and memory care services but exclude hospitals. The review will include a three-step search strategy: (a) identify keywords from MEDLINE and Embase, (b) search using the identified keywords in databases (MEDLINE/PubMed, CINAHL, Web of Science, Embase, Cochrane Library, PsychInfo and Google) and (c) review references from included studies to identify additional studies. Only studies in English will be included. Four researchers will independently assess titles and abstracts and then review the full text of the selected articles, applying the inclusion criteria. The extracted data will be presented in tables and summarised narratively.

Ethics and dissemination Scoping review data will be collected from publicly available articles; research ethics approval is not required. The findings will be disseminated to healthcare practitioners and the public through a peer-reviewed publication and conference presentations.

INTRODUCTION

Dementia is a neurodegenerative condition characterised by progressive deterioration of cognitive function, often affecting memory, thinking, judgement, personality and visuospatial skills, among many other possible deficits.¹ In addition to cognitive impairment, due to its progressive degenerative nature,

Strengths and limitations of this study

- This scoping review will follow the Joanna Briggs Institute scoping review manual to ensure scientific rigour.
- Patients and family partners will be involved in the scoping review study.
- The review will be limited to literature published in English.
- Because the focus of this review is on technology-based group exercise for people with dementia or mild cognitive impairment living in the community, exercise programmes provided by acute care services will not be captured.

dementia leads to physical decline in balance, gait and movement coordination, behavioural and psychological symptoms of dementia (BPSD) and social disability.^{2 3} Dementia can be categorised based on aetiology. Alzheimer's disease (AD) is the most common type of dementia, followed by other types, such as vascular dementia, Lewy body dementia, frontotemporal dementia and Parkinson's disease. Dementia is a widespread health issue, affecting more than 50 million people globally, with more than 10 million new cases every year.⁴ Mild cognitive impairment (MCI) is noticeable decline in cognitive function that is not severe enough to require help with daily activities.^{5 6} Although considered to be distinct from dementia, MCI is often a precursor to dementia.⁷ It has been estimated that 10%–15% of people with MCI progress to AD.⁸ Research on MCI can provide knowledge for preventing progression of MCI into dementia.⁹

Although the number of people with dementia is increasing worldwide, no curative treatment is currently available and effective palliative treatments are lacking. Antipsychotic medications used to manage BPSD in people with dementia, in particular

older adults, can cause adverse events such as oversedation, psychomotor impairment, Parkinsonian symptoms, stroke or falls, all of which can contribute to admission to the emergency department or long-term care facilities and increased healthcare costs.^{10 11} Evidence-based, safe and effective non-pharmacologic treatments are needed to relieve dementia symptoms and/or delay progression of the severity of the disease. Group exercise programmes, such as chair yoga and tai chi, have shown positive effects in managing dementia symptoms.^{12–15}

Meta-analyses have shown overall benefits of various types of physical exercise on physical function and BPSD in people with dementia.^{16–18} However, people with dementia face challenges to participate in exercise groups due to restricted range of motion, declined cognition and balance, and risk of falls.^{2 13 14} Major barriers to physical exercise in this population could include lack of accessible opportunities, transportation to community sites where exercise classes are offered and limited support from caregivers.^{12 19 20} People with dementia are more likely to experience social isolation and loneliness, which have been linked to increased depression, cardiovascular risk and overall detrimental effects to quality of life.^{21 22} Group exercise programmes may not only help to improve mobility and function; they may support social engagement and reduce social isolation and loneliness. Given the rapid growth of technology use by older people, technology-based group exercise interventions offer opportunities for people with dementia to participate in exercise to manage dementia symptoms, improve social connections and enhance their quality of life.²³

In this scoping review study, technology-based exercise intervention will refer to any exercise intervention in which technology is essential to the activity. Exergames, also known as interactive exercise-based video games or virtual reality exercise games, have been used as an acceptable intervention to increase physical activity in older adults.^{24 25} In addition to exergames and virtual reality exercise games, recent technological advances have created the potential for an online exercise intervention with synchronous monitoring through a secure videoconferencing platform (eg, Zoom, WebEx) on a computer or tablet.^{23 26} Remotely delivered exercise interventions via video conferencing have been found to be feasible for improving physical activity in persons with disabilities.²³ Group-based exercise interventions could increase physical activity and improve emotional well-being. Synchronous online group exercise could develop social connections via the Internet among people with dementia, who are often isolated from society.²³

With technology-based interventions, people with dementia and their caregivers can experience socialisation without the need to go to a community centre.²³ Several systematic reviews or meta-analyses have analysed the effects of exergames²³ or virtual reality²⁵ in people with major cognitive neurocognitive disorder, MCI or dementia.^{27–29} However, a preliminary search of MEDLINE, the Cochrane Database of Systematic Reviews

and Joanna Briggs Institute (JBI) Evidence Synthesis was conducted as of July 2021, and no current systematic review, meta-analysis or scoping review has been conducted on the topic of technology-based group interventions that included virtual reality, exergame or other Internet-based intervention (synchronous or asynchronous) to support people with dementia. This gap will be addressed in this comprehensive review of the types of technology-based interventions across dementias and MCI and their effects. The format of the scoping review is specifically appropriate to mapping out the existing evidence and identifying gaps in knowledge on the topic of technology-based group interventions for people with dementia or MCI.³⁰

The main objective of the scoping review will be to understand and summarise the extent of evidence with respect to the existence of technology-based group exercise interventions for people with dementia or MCI and the outcome measures studied and reported in the literature.

Review questions

1. What technology-based group exercise interventions exist for people with dementia or MCI?
2. What outcomes and impacts of technology-based group exercise interventions for people with dementia or MCI have been reported in the literature?

METHODS

The proposed scoping review will be conducted in accordance with the JBI methodology for scoping reviews.³¹

Inclusion criteria

Participants

We will include studies that focus on people with all types of dementia (eg, AD, vascular dementia, Lewy body dementia, Parkinson's disease dementia, frontotemporal dementia) or MCI in people of all ages.

Concept

This review will focus on technology-based group exercise interventions for people with dementia. We will include studies that involved at least one group of people with dementia or MCI. A group is defined as any collection of two or more people, including participant-caregiver dyads. Technology-based exercise includes any form of exercise in which technology is integral to the execution of the activity, such as interventions through video-sharing or online platforms, exergaming and virtual reality exercises. We will consider exercise intervention to include any sort of physical activity. Any studies that focus solely on cognitive-based interventions will be excluded. Although MCI is considered to be distinct from dementia, due to the relative difficulty of studying people with higher levels of cognitive impairment, as well as the fact that MCI is often considered a precursor to dementia, studies that focus on technology-based group exercise interventions for people with MCI will be included in this review, as

implications learnt from MCI-focused studies could be relevant for populations with dementia.⁷

Context

We will include studies in which the intervention took place in homes, assisted living facilities or memory care facilities. We will not include studies in which interventions took place in a hospital or research facility.

Types of studies

This scoping review will consider both experimental and quasi-experimental study designs, including randomised controlled trials, non-randomised controlled trials, before-and-after studies and interrupted time-series studies. In addition, analytical observational studies, including prospective and retrospective cohort studies, case-control studies and analytical cross-sectional studies will be considered for inclusion. This review will also consider descriptive observational study designs, including case series and descriptive cross-sectional studies. Qualitative studies and systematic reviews that meet the inclusion criteria will be considered. Text and opinion papers will be considered. Protocols will not be included. Only studies published in English will be included. There will be no limitation based on publication date.

Search strategy

The scoping review will follow the three-step search strategy recommended by the JBI methodology.³¹ The search strategy is designed to locate both published and unpublished studies. An initial limited search of MEDLINE, CINAHL and Embase will be undertaken to identify articles on the topic. This preliminary search will use these selected keywords: (Dementia or (Alzheimer disease)) AND ((web-based intervention) OR (exergaming) OR (exergame) OR (virtual reality) OR (technology-based intervention) OR telehealth OR telemedicine) AND (Exercise OR (physical activity)). The text words contained in the titles and abstracts of relevant articles and the index terms used to describe the articles will be used to develop a full search strategy for selected databases such as MEDLINE/PubMed, CINAHL, Web of Science, Embase, Cochrane Library and PsychInfo (online supplemental appendix I). The search strategy, including all identified keywords and index terms, will be adapted for each included database. A full search strategy for MEDLINE/PubMed is included in online supplemental appendix I. The reference lists of all included sources of evidence will be screened for additional relevant studies. We have worked and will continue to work with a medical librarian to refine the search strategy in order to ensure that relevant and key articles are found.

Information sources

The databases to be searched will be MEDLINE/PubMed, CINAHL, Web of Science, Embase, Cochrane Library and PsychInfo. Sources of unpublished studies/grey literature to be searched will be Google, Web of Science and Embase.

Study/source of evidence selection

Following the search, all identified citations will be collated and uploaded into Mendeley and duplicates will be removed. Titles and abstracts will then be screened by four researchers for assessment against the inclusion criteria. Potentially relevant sources will be retrieved in full and their citation details imported into Mendeley. The full text of selected citations will be assessed in detail against the inclusion criteria by two independent reviewers. Reasons for exclusion of sources of evidence at the full text review that do not meet the inclusion criteria will be recorded and reported in the scoping review. Any disagreements between the reviewers at each stage of the selection process will be resolved through discussion with additional reviewers. The results of the search and the study inclusion process will be reported in full in the final scoping review and presented in a Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for scoping review flow diagram.³²

Data extraction

Data will be extracted from papers included in the scoping review by four independent reviewers using a data extraction tool developed by the reviewers. The extracted data will include specific details about the author(s), year and country of publication, setting, population and sample size, research design and methods, intervention type, outcome measures and impact and results.

A draft charting table is provided (online supplemental appendix II). The data extraction tool will be pilot tested by the two independent researchers by extracting data from three papers and comparing results. The draft data extraction tool will be modified and revised as necessary during the process of extracting data from each included evidence source. Modifications will be detailed in the scoping review. Any disagreements between the reviewers will be resolved through discussion. If appropriate, authors of papers will be contacted to request missing or additional data, where required.

Patient and public involvement

Several family and patient partners will join us for meetings in which we will discuss the data that have been extracted from the articles. The family partners are people who have a family member with dementia and may act as the primary caregiver to the person with dementia, whereas the patient partners are people with dementia. Prior to discussing extracted data in meetings, the patient and family partners may review extracted data and several full-text articles; the number of full-text articles will be determined by each individual partner. We anticipate that each person will review one to three articles. Each partner can choose to be a named co-author in the scoping review paper or remain anonymous (see acknowledgements of this paper for the full names of patient and family partners



who have chosen to be identified). The patient and family partners were recruited from various memory disorder clinics.

Ethics and dissemination

For this scoping review, data will be collected from publicly available articles. Thus, neither informed consent nor research ethics approval is required. The findings will be disseminated to healthcare practitioners and the public through a peer-reviewed publication and presentations at conferences.

Data analysis and presentation

The extracted data will be presented in a table to summarise the evidence. A narrative summary will accompany the charted results and will describe the qualities of the literature on technology-based group exercise interventions for people with dementia. The categories that will be used in the presentation of data are year of publication, author(s), country, setting, population and sample size, research design and methods, intervention type, outcome measures, and impact and results.

Author affiliations

¹School of Nursing, University of British Columbia, Vancouver, British Columbia, Canada

²School of Nursing, Vancouver General Hospital, Vancouver, British Columbia, Canada

³Charles E Schmidt College of Medicine, Marcus Institute of Integrative Health at FAU Medicine, Florida Atlantic University, Boca Raton, Florida, USA

⁴Faculty of General Sciences, Langara College, Vancouver, British Columbia, Canada

⁵College of Social Work and Criminal Justice, Phyllis and Harvey Sandler School of Social Work, Florida Atlantic University, Boca Raton, Florida, USA

Twitter Lillian Hung @nursellillian

Acknowledgements The authors acknowledge and thank Senior Medical Librarian Michelle Keba Knecht at Florida Atlantic University Charles E. Schmidt College of Medicine for her assistance. We thank the family and patient partners for their work and contribution to this scoping review. Patient partners are Betty Riley, Vincent Celeste and David Call; family partners are Nate Riley, Diane Celeste and Dierdre Lacativa.

Contributors LH and JP conceived the idea. All authors helped to refine the research question and develop the protocol and methods. HL drafted the manuscript with support from JP, PR and LH, who made meaningful contributions. LH edited the final manuscript. All authors approved the final manuscript.

Funding This work was supported by the Marcus Institute of Integrative Health at FAU Medicine, grant number 2485.

Competing interests None declared.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially,

and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Lillian Hung <http://orcid.org/0000-0002-7916-2939>

Hannah Levine <http://orcid.org/0000-0003-3441-8395>

Juyoung Park <http://orcid.org/0000-0001-6737-3109>

REFERENCES

- Emmady PD, Tadi P. Dementia. StatPearls Publishing, 2021. Available: <https://www.who.int/news-room/fact-sheets/detail/dementia> [Accessed 17 Jul 2021].
- Tible OP, Riese F, Savaskan E, et al. Best practice in the management of behavioural and psychological symptoms of dementia. *Ther Adv Neurol Disord* 2017;10:297–309.
- Lam FM, Huang M-Z, Liao L-R, et al. Physical exercise improves strength, balance, mobility, and endurance in people with cognitive impairment and dementia: a systematic review. *J Physiother* 2018;64:4–15.
- Cao Q, Tan C-C, Xu W, et al. The prevalence of dementia: a systematic review and meta-analysis. *Journal of Alzheimer's Disease* 2020;73:1157–66.
- Gillis C, Mirzaei F, Potashman M, et al. The incidence of mild cognitive impairment: a systematic review and data synthesis. *Alzheimers Dement* 2019;11:248–56.
- Langa KM, Levine DA. The diagnosis and management of mild cognitive impairment: a clinical review. *JAMA* 2014;312:2551–61.
- Plassman BL, Langa KM, Fisher GG, et al. Prevalence of cognitive impairment without dementia in the United States. *Ann Intern Med* 2008;148:427.
- Petersen RC, Smith GE, Waring SC, et al. Mild cognitive impairment: clinical characterization and outcome. *Arch Neurol* 1999;56:303–8.
- Campbell NL, Unverzagt F, LaMantia MA, et al. Risk factors for the progression of mild cognitive impairment to dementia. *Clin Geriatr Med* 2013;29:873–93.
- Ohno Y, Kunisawa N, Shimizu S. Antipsychotic treatment of behavioral and psychological symptoms of dementia (BPSD): management of extrapyramidal side effects. *Front Pharmacol* 2019;10:1045.
- Macfarlane S, O'Connor D. Managing behavioural and psychological symptoms in dementia. *Aust Prescr* 2016;39:123–5.
- Hobson N, Dupuis SL, Giangregorio LM, et al. Perceived facilitators and barriers to exercise among older adults with mild cognitive impairment and early dementia. *J Aging Phys Act* 2020;28:208–18.
- Morris JK, Vidoni ED, Johnson DK, et al. Aerobic exercise for Alzheimer's disease: A randomized controlled pilot trial. *PLoS One* 2017;12:e0170547.
- McCaffrey R, Park J, Newman D, et al. The effect of chair yoga in older adults with moderate and severe Alzheimer's disease. *Res Gerontol Nurs* 2014;7:171–7.
- Hsieh C-C, Lin P-S, Hsu W-C, et al. The effectiveness of a virtual Reality-Based tai chi exercise on cognitive and physical function in older adults with cognitive impairment. *Dement Geriatr Cogn Disord* 2018;46:358–70.
- Lee HS, Park SW, Park YJ. Effects of physical activity programs on the improvement of dementia symptom: a meta-analysis. *Biomed Res Int* 2016;2016:1–7.
- Li X, Guo R, Wei Z, et al. Effectiveness of exercise programs on patients with dementia: a systematic review and meta-analysis of randomized controlled trials. *Biomed Res Int* 2019;2019:1–16.
- Jia R-X, Liang J-H, Xu Y, et al. Effects of physical activity and exercise on the cognitive function of patients with Alzheimer disease: a meta-analysis. *BMC Geriatr* 2019;19:181.
- Boehm J, Franklin RC, Newitt R, et al. Barriers and motivators to exercise for older adults: a focus on those living in rural and remote areas of Australia. *Aust J Rural Health* 2013;21:141–9.
- Logsdon RG, McCurry SM, Pike KC, et al. Making physical activity accessible to older adults with memory loss: a feasibility study. *Gerontologist* 2009;49 Suppl 1:S94–9.
- Balouch S, Rifaat E, Chen HL, et al. Social networks and loneliness in people with Alzheimer's dementia. *Int J Geriatr Psychiatry* 2019;34:666–73.
- Courtin E, Knapp M, isolation S. Social isolation, loneliness and health in old age: a scoping review. *Health Soc Care Community* 2017;25:799–812.
- Ptomey LT, Willis EA, Greene JL, et al. The feasibility of group video Conferencing for promotion of physical activity in adolescents with

- intellectual and developmental disabilities. *Am J Intellect Dev Disabil* 2017;122:525–38.
- 24 Valenzuela T, Okubo Y, Woodbury A, *et al.* Adherence to technology-based exercise programs in older adults: a systematic review. *J Geriatr Phys Ther* 2018;41:49–61.
 - 25 Nawaz A, Skjæret N, Helbostad JL, *et al.* Usability and acceptability of balance exergames in older adults: a scoping review. *Health Informatics J* 2016;22:911–31.
 - 26 Galiano-Castillo N, Cantarero-Villanueva I, Fernández-Lao C, *et al.* Telehealth system: a randomized controlled trial evaluating the impact of an Internet-based exercise intervention on quality of life, pain, muscle strength, and fatigue in breast cancer survivors. *Cancer* 2016;122:3166–74.
 - 27 Swinnen N, Vandenbulcke M, Vancampfort D. *Exergames in people with major neurocognitive disorder: a systematic review. disability and rehabilitation: assistive technology*, 2020: 1–14.
 - 28 Zhao Y, Feng H, Wu X, *et al.* Effectiveness of Exergaming in improving cognitive and physical function in people with mild cognitive impairment or dementia: systematic review. *JMIR Serious Games* 2020;8:e16841.
 - 29 Kim O, Pang Y, Kim J-H. The effectiveness of virtual reality for people with mild cognitive impairment or dementia: a meta-analysis. *BMC Psychiatry* 2019;19.
 - 30 Peters M, Godfrey C, McInerney P. Chapter 11: Scoping reviews. In: *JBI manual for evidence synthesis*, 2020.
 - 31 Peters MDJ, Godfrey CM, Khalil H, *et al.* Guidance for conducting systematic scoping reviews. *Int J Evid Based Healthc* 2015;13:141–6.
 - 32 Page MJ, McKenzie JE, Bossuyt PM. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *The BMJ* 2021;372.