




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Preadmission use of antidiabetic medications and mortality among patients with COVID-19 having type 2 diabetes: A meta-analysis

Nam Nhat Nguyen^a, Dung Si Ho^{b,c}, Hung Song Nguyen^{a,d}, Dang Khanh Ngan Ho^e, Hung-Yuan Li^f, Chia-Yuan Lin^g, Hsiao-Yean Chiu^{h,i}, Yang-Ching Chen^{g,j,*}

^a College of Medicine, Taipei Medical University, Taipei, Taiwan

^b Department of Geriatric Medicine, Faculty of Medicine, Pham Ngoc Thach University of Medicine, Ho Chi Minh City, Viet Nam

^c Department of Pulmonology, Thong Nhat Hospital, Ho Chi Minh City, Viet Nam

^d Wellcome Trust Major Overseas Program, Oxford University Clinical Research Unit, Ho Chi Minh City, Viet Nam

^e School of Nutrition and Health Sciences, College of Nutrition, Taipei Medical University, Taipei, Taiwan

^f Department of Internal Medicine, National Taiwan University Hospital, Taipei, Taiwan

^g Department of Family Medicine, Taipei Medical University Hospital, Taipei, Taiwan

^h School of Nursing, College of Nursing, Taipei Medical University, Taipei, Taiwan

ⁱ Research Center of Sleep Medicine, College of Medicine, Taipei Medical University, Taipei, Taiwan

^j Department of Family Medicine, School of Medicine, College of Medicine, Taipei Medical University, Taipei, Taiwan

ARTICLE INFO

Keywords:

COVID-19
Type 2 diabetes mellitus
Antidiabetic medication

ABSTRACT

Background: Diabetes is an independent predictor of poor outcomes in patients with COVID-19. We compared the effects of the preadmission use of antidiabetic medications on the in-hospital mortality of patients with COVID-19 having type 2 diabetes.

Methods: A systematic search of PubMed, EMBASE, Scopus and Web of Science databases was performed to include studies (except case reports and review articles) published until November 30, 2021. We excluded papers regarding in-hospital use of antidiabetic medications. We used a random-effects meta-analysis to calculate the pooled OR (95% CI) and performed a sensitivity analysis to confirm the robustness of the meta-analyses.

Main findings: We included 61 studies (3,061,584 individuals), which were rated as having low risk of bias. The OR (95% CI) indicated some medications protective against COVID-related death, including metformin [0.54 (0.47–0.62), I² 86%], glucagon-like peptide-1 receptor agonist (GLP-1RA) [0.51 (0.37–0.69), I² 85%], and sodium–glucose transporter-2 inhibitor (SGLT-2i) [0.60 (0.40–0.88), I² 91%]. Dipeptidyl peptidase-4 inhibitor (DPP-4i) [1.23 (1.07–1.42), I² 82%] and insulin [1.70 (1.33–2.19), I² 97%] users were more likely to die during hospitalization. Sulfonylurea, thiazolidinedione, and alpha-glucosidase inhibitor were mortality neutral [0.92 (95% CI 0.83–1.01, I² 44%), 0.90 (95% CI 0.71–1.14, I² 46%), and 0.61 (95% CI 0.26–1.45, I² 77%), respectively]. The sensitivity analysis indicated that our findings were robust.

Conclusions: Metformin, GLP-1RA, and SGLT-2i were associated with lower mortality rate in patients with COVID-19 having type 2 diabetes. DPP-4i and insulin were linked to increased mortality. Sulfonylurea, thiazolidinedione, and alpha-glucosidase inhibitors were mortality neutral. These findings can have a large impact on the clinicians' decisions amid the COVID-19 pandemic.

1. Introduction

Since late 2019, SARS-CoV-2 has emerged as a novel pathogenic microbe, resulting in the COVID-19 pandemic. By the end of November 2021, more than 257 million people had been infected with SARS-CoV-2 globally, approximately 5.1 million of whom died [1]. Several risk

factors have been linked with the progression and deterioration of COVID-19, such as advanced age, diabetes, hypertension, cardiovascular diseases, and obesity [2]. Diabetes, with its increasing worldwide prevalence, has become major comorbidity in patients with COVID-19 and predisposes them to poor outcomes. Many potential pathways for this have been proposed, including increased inflammatory cascade,

* Corresponding author at: Department of Family Medicine, Taipei Medical University Hospital, Taipei, Taiwan.

E-mail address: melisa26@tmu.edu.tw (Y.-C. Chen).

<https://doi.org/10.1016/j.metabol.2022.155196>

Received 30 December 2021; Accepted 28 March 2022

Available online 31 March 2022

0026-0495/© 2022 Elsevier Inc. All rights reserved.

immunocompromised status, glucose homeostasis dysfunction, hypercoagulability, alveolar hyperpermeability, and vascular endothelial damage. These pathophysiological changes might lead to acute respiratory distress syndrome, thromboembolism events, and cytokine storms, thereby contributing to increased COVID-19-related deaths [3].

In the past two decades, many drugs have been approved for diabetic patients, leading to a noticeable change in the trend of medication use. Glucose-lowering therapies have also received much critical attention recently as potential host-directed therapies due to their mechanisms of action that may influence the natural course of SARS-CoV-2 infection. Many studies have evaluated whether the preadmission use of certain antidiabetic medications might improve outcomes in those participants. The results have remained controversial, partly because different classes of drugs may differ in their effectiveness and safety against SARS-CoV-2. The gap between preclinical research and real-world data must be bridged. For example, dipeptidyl peptidase-4 inhibitor (DPP-4i) has recently gained much attention due to its safety, cardiovascular neutrality, and potential mechanistic pathways that could alleviate the course of SARS-CoV-2 infection. Although the exact mechanisms underlying the effect of this class on the prognosis of COVID-19 remain unclear, several hypotheses may provide some insights. In addition to glucose homeostasis, DPP-4i inhibits the enzyme DPP-4, which is involved in many events of COVID-19 pathophysiology, including T-cell proliferation, nuclear factor kappa-light-chain-enhancer of activated B (NF- κ B) activation, CD86 expression, and inflammatory cytokines production [4]. However, many studies and meta-analyses have indicated no significant benefit of DPP-4i against COVID-19 [5,6]. Moreover, even for the same drug class, previous small meta-analyses have indicated inconsistent effects regarding the severity or mortality of patients with COVID-19, as in the case of the glucagon-like peptide-1 receptor agonist (GLP-1RA) [5,7]. Therefore, little is known about their true efficacy in the prognosis of that disease.

In this systematic review and meta-analysis, we (1) summarized the effects of every single antidiabetic medication on the mortality of patients with COVID-19 having diabetes and (2) evaluated the dose-responsiveness of the impacts of medications on mortality. By incorporating much more original papers, our findings would strengthen or reject the evidence for effects of each antidiabetic medication on COVID-19 mortality from inconsistent meta-analyses, and provided novel results regarding the effect of TZD and AGI, and the relationship between dosages and effects, which have not been previously reported.

2. Material and methods

2.1. Population, intervention, comparison, outcomes, and study design (PICOS)

Participants included patients with confirmed COVID-19 who had diabetes and were on prehospital medications extending to the pandemic. A confirmed case of COVID-19 was defined using a positive result on reverse transcription-polymerase chain reaction (RT-PCR) according to the diagnostic procedures of each center. Preexisting diabetes was ascertained through a diabetes diagnosis in medical records. The current use of antidiabetic medications was recorded at the time of recruitment. The interventional therapies considered were one of the following medications: metformin, sulfonylurea (SU), meglitinide (gli-nide), thiazolidinedione (TZD), alpha-glucosidase inhibitor (AGI), GLP-1RA, DPP-4i, sodium-glucose transporter-2 inhibitor (SGLT-2i), and insulin. Specific-agent users were defined as those who have been on a current prescription. The comparator included nonusers of specific antidiabetic medications. Our primary outcome was in-hospital mortality or mortality within 90 days, confirmed with the medical record.

We planned to include randomized and nonrandomized controlled trials and observational studies, including prospective and retrospective cohort studies and case-control studies, which were either peer-reviewed or published as abstracts or preprints. If an official

publication has already replaced a preprint, the publication was chosen instead of a preprint. We excluded case reports and review articles.

Based on the predetermined inclusion criteria, three independent reviewers (DSH, HSN, and DKNH) searched, screened, reviewed, extracted, and recorded data. In case of discrepancy, a fourth reviewer (NNN) was consulted to reach a final consensus. We verified transparent reporting following the Meta-analysis of Observational Studies in Epidemiology (MOOSE) checklist because we only found observational studies relevant to this topic.

2.2. Systematic review protocol

This systematic review and meta-analysis were registered in the PROSPERO International Prospective Register of Systematic Reviews (ID: CRD42021293064).

2.3. Search strategy and data sources

We systematically searched PubMed/MEDLINE, EMBASE, Scopus, and Web of Science databases for relevant articles up to November 30, 2021, without limiting the language or publication year. The following main keywords and related terms were used: "COVID-19," "diabetes," "antidiabetic medication," or the names of specific classes. The detailed search strategy is presented in Table A.1 (Supplementary appendix). We further identified additional articles through a manual search. We used Endnote (version 20; Clarivate, Philadelphia, PA, USA) to manage studies found.

2.4. Data extraction

The number of events, the number of observations, and other demographic variables, including race/ethnicity, sex, age, HbA1c, diabetes duration, BMI, and percentage of important comorbidities such as hypertension and chronic kidney disease, were documented for each group. OR was also extracted from the papers. The article's corresponding author was contacted through e-mail if raw data were required.

2.5. Data analysis

The risk of bias was assessed by two independent reviewers by using the Newcastle-Ottawa Scale [8].

Effect sizes were calculated as the natural logarithm of ORs. The logOR and standard error of the logOR were used as input for meta-analysis in statistical software. Forest plots were used to display the OR from each original study and the pooled findings. We used Cochran's Q test and I^2 statistics to assess heterogeneity between studies [9,10]. A random-effects model was chosen when the Cochran's Q test p-value of <0.1 or an I^2 of $>50\%$ was obtained. A fixed-effects model was preferred if there was no evidence of heterogeneity. Publication bias was statistically assessed using Egger's asymmetry test [11]. A publication bias was suspected if the p-value for Egger's test was <0.05 . Meta-regression and subgroup analysis were predefined to explore the source of heterogeneity further. We performed meta-regression on a set of pre-specified important characteristics, comorbidities, and chronic complications that are commonly found in diabetes patients, including age, gender, race/ethnicity, BMI, hypertension, and chronic kidney disease. We performed sensitivity analysis by outlier removal and trim-and-fill methods and then compared the original results with re-analyzed results to confirm the stability and robustness of our main meta-analyses.

A two-sided p-value of <0.05 was considered statistically significant. We analyzed data by using R software (version 4.0.2; R Foundation for Statistical Computing; Vienna, Austria).

2.6. Ethics

Formal ethics approval is not required because we only collect nonconfidential information from which the patients' identities could not be ascertained.

3. Results

3.1. Literature search and study selection

A total of 6920 articles were identified from the databases through a systematic search (Fig. 1). Next, 5790 articles remained after deduplication to be screened for their titles and abstracts. Of these articles, 5644 were excluded due to full-text inaccessibility (n = 173), duplication (n = 566), and irrelevancy (n = 4905); thus, 146 papers remained for eligibility assessment. The other 85 publications were further excluded because they did not include the outcome of interest; reported composite endpoint of intensive care unit admission, mechanical ventilation, and death; involved the same cohort; investigated inpatient use of antidiabetic drugs; or were irrelevant to our topic. Finally, 61 studies met our inclusion criteria for a systematic review. However, only 59 articles were pooled in the meta-analyses because one publication reported the hazard ratio instead of odds ratio, and one reported longer-term mortality (7 months) [12,13].

3.2. Study and participant characteristics

A total of 3,061,584 participants were recruited from studies [14–72]. Most of them were retrospective, except for two cross-sectional studies [26,52]. The antidiabetic drugs that were investigated included

metformin (42 articles), SU (21), TZD (8), AGI (8), GLP-1RA (12), DPP-4i (28), SGLT-2i (13), and insulin (33) (Table 1). Only two papers reported glinide-associated mortality in patients with COVID-19 with few users [34,50]. Therefore, we did not present this drug in our research. The Newcastle–Ottawa assessment results revealed that all studies were rated as having adequate quality (Table A.2). No publication bias was found using Egger's test (Table A.3).

3.3. Main findings

3.3.1. Mortality between medication users and nonusers

Compared with nonusers, metformin (OR 0.54, 95% CI 0.47–0.62, I² 86%), GLP-1RA (OR 0.51, 95% CI 0.37–0.69, I² 85%), and SGLT-2i (OR 0.60, 95% CI 0.40–0.88, I² 91%) use significantly reduced mortality among patients with COVID-19 with diabetes (Figs. 2–4). By contrast, DPP-4i (OR 1.23, 95% CI 1.07–1.42, I² 82%) and insulin (OR 1.70, 95% CI 1.33–2.19, I² 97%) were associated with an increased risk of in-hospital death (Figs. A.1, A.2). SU (OR 0.92, 95% CI 0.83–1.01, I² 44%), TZD (OR 0.90, 95% CI 0.71–1.14, I² 46%), and AGI (OR 0.61, 95% CI 0.26–1.45, I² 77%) were mortality neutral (Figs. A.3–A.5).

3.3.2. Meta-regression of confounding factors

Using meta-regression, we observed some significant variables that were significantly associated with mortality due to COVID-19, including continent, white race, male sex, age, BMI, HbA1C, hypertension, and CKD (Table 2).

3.3.3. Subgroup analysis

We performed subgroup analyses based on confounding factors identified through meta-regression to compare the effects of antidiabetic




Fig. 1. PRISMA flowchart summarizing the study selection process.

Table 1
Characteristics of studies (systematic review).

Study	Country	Number of patients	Race/ethnicity (%)	Male sex (%)	Age (years)	HbA1C (%)	Body mass index (kg/m ²) or obesity (%)	Hypertension (%)	CKD (%)	Mortality
<i>Metformin users/nonusers</i>										
An et al. [15]	Korea	423/598		39.9	45.0 ± 19.9			18.2	0.8	NS
Bliden et al. [16]	USA	34/41				8.9 ± 2.3 vs. 8.4 ± 1.8				NS
Bramante et al. [18]	USA	2333/3923		51.6 vs. 44.6	73.0 (66.0–80.0) vs. 76.0 (67.0–84.0)		Obesity: 4.8% vs. 9.0%	56.3 vs. 60.4	6.3 vs. 18.6	NS
Cernigliaro et al. [19]	Italy	82/90								Decreased
Chen et al. [20]	China	43/77		42.9	62.0 (56.0–69.0) vs. 67.0 (57.5–73.0)	7.7 (6.9–9.1) vs. 8.4 (7.4–10.7)		61.2	10.2	NS
Cheng et al. [21]	China	18/32		66.7 vs. 46.9	48.0 (40.5–56.5) vs. 58.0 (49.5–66.5)		25.9 (24.7–26.9) vs. 24.5 (21.3–28.0)	22.2 vs. 43.8	27.8 vs. 37.5	NS
Crouse et al. [22]	USA	76/144	White: 27.6, African American: 64.9	50.6		8.0 ± 2.6 vs. 7.0 ± 1.8	35.2 ± 9.4 vs. 33.6 ± 8.7	91.6		Decreased
Dave et al. [23]	Africa	4084/1624		39.3	55.0 (46.0–63.0)			55.5	9.2	Decreased
Do et al. [25]	Korea	469/1396		51.8 vs. 42.1	64.8 ± 11.4 vs. 67.4 ± 12.1			73.8 vs. 82.1	29.4 vs. 47.4	NS
Eliboi et al. [26]	Turkey	379/53		45.6	63.3 ± 10.3			74.1	4.6	NS
Ghany et al. [29]	USA	243/350	Black: 71.0 vs. 70.0	39.0 vs. 41.0	70.9 ± 8.9 vs. 71.2 ± 8.9	7.7 ± 1.5 vs. 6.4 ± 1.5	33.2 ± 7.7 vs. 31.7 ± 9.6	60.0 vs. 50.0		Decreased
Goodall et al. [31]	England	210/166	White: 25.5, Black: 13.9, Asian: 37.8	64.3	69.0 (56.0–80.0)			49.6		NS
Khunti et al. [34]	England	1,800,005/1,051,460	White: 64.5, Black: 4.8, Asian: 16.0	58.1	67.0 (57.0–77.0)				78.0	Decreased
Kim et al. [35]	Korea	113/122		45.1	68.3 ± 11.9		24.2 ± 3.2	62.6	7.7	NS
Lally et al. [37]	USA	127/172	White: 61.4, Black: 30.7	98.4 vs. 98.6	72.3 ± 8.3 vs. 75.6 ± 9.2	7.5 ± 1.4 vs. 6.5 ± 1.3	29.7 ± 6.6 vs. 28.0 ± 7.0		13.4	Decreased
Li et al. (1) [39]	China	37/94		59.5 vs. 55.3	64.6 ± 11.2 vs. 67.7 ± 11.7	9.2 ± 4.6 vs. 9.0 ± 4.6	24.2 ± 3.3 vs. 24.2 ± 3.7	62.2 vs. 58.5		Decreased
Li et al. (2) [38]	China	142/245		51.1	60.0 (49.0–68.0)			48.6	1.0	NS
Luk et al. [40]	China	737/254	Asian	55.0 vs. 51.6	65.6 (57.7–72.6) vs. 68.9 (61.3–79.7)	7.3 (6.6–8.5) vs. 6.6 (6.1–7.8)	24.1 (21.5–27.7) vs. 23.7 (22.2–27.0)	63.1 vs. 56.7	19.5 vs. 37.8	Decreased
Luo et al. (1) [41]	China	104/179		51.0 vs. 57.5	63.0 (55.8–68.3) vs. 65.0 (57.5–71.0)			59.6 vs. 57.0		Decreased
Luo et al. (2) [42]	China	54/137		54.0 vs. 54.0	61.0 (56.0–69.0) vs. 61.0 (57.8–68.3)	8.0 ± 1.9 vs. 6.7 ± 1.9		55.5	2.0 vs. 2.0	Decreased
Ma et al. [43]	USA	361/995	White: 72.6, Black: 12.2, Asian: 1.9	60.4 vs. 54.1				79.5 vs. 85.0		Decreased
Mirani et al. [46]	Italy	69/21		72.5 vs. 71.4	69.0 ± 13.0 vs. 75.0 ± 8.0		Obesity: 47.8% vs. 47.6%	72.5 vs. 90.5	11.6 vs. 42.9	NS
Mirsoleymani et al. [47]	Iran	36/69		72.4	59.8 ± 17.2			37.1		NS
Nafakhi et al. [48]	Iraq	35/32		43.0	60.0 ± 10.0		29.8 ± 5.0	66.0		Decreased
Nyland et al. [50]	USA	5077/24,439	White: 47.9, African American: 25.5, Asian: 3.1	48.2	60.9 ± 15.0	7.7 ± 2.1	32.8 ± 8.9	47.7	15.5	Decreased

(continued on next page)

Table 1 (continued)

Study	Country	Number of patients	Race/ethnicity (%)	Male sex (%)	Age (years)	HbA1C (%)	Body mass index (kg/m ²) or obesity (%)	Hypertension (%)	CKD (%)	Mortality
Oh et al. [51]	Korea	7204/20,289		44.7 vs. 38.2					3.4 vs. 4.1	NS
Ong et al. [52]	Philippines	186/169		58.6 vs. 52.7	61.6 ± 11.6 vs. 63.9 ± 12.8	7.0 ± 2.4 vs. 7.6 ± 1.9	Obesity: 62.0% vs. 65.1%	73.1 vs. 76.3	4.8 vs. 21.9	Decreased
Orioli et al. [53]	Belgium	45/23		48.0	69.0 ± 14.0	7.1 (6.6–8.3)	30.5 ± 5.3	80.8	34.2	Decreased
Perez-Belmonte et al. [54]	Spain	825/663		65.7 vs. 57.2	74.8 ± 7.9 vs. 77.1 ± 7.1		Obesity: 26.4% vs. 26.1%	74.2 vs. 79.5	4.7 vs. 29.0	NS
Philipose et al. [55]	England	100/59	White: 45.5, Afro-Caribbean: 20.2, Asian: 19.1	59.0				50.2		NS
Ramos-Rincon et al. [56]	Spain	421/369		47.1			Obesity: 17.7%	84.3	17.2	NS
Ravindra et al. [57]	India	53/313		63.2	46.7 ± 17.1			28.7	0.9	NS
Saygili et al. [60]	Turkey	432/154		49.8 vs. 50.2	65.0 ± 11.2 vs. 68.9 ± 13.5	8.0 (6.8–9.9) vs. 7.7 (6.7–10.1)		67.1 vs. 70.1	0.0	Decreased
Shetaskova et al. [61]	Russia	196/113								Decreased
Silverii et al. [62]	Italy	76/83		54.1	73.3 ± 12.7					Decreased
Sourij et al. [63]	Austria	77/103		63.9	67.6 ± 14.0	6.7 (1.9)	29.4 ± 5.7	77.0	23.1	NS
Tamura et al. [65]	Brazil	116/72		63.5 vs. 61.6	62.1 ± 15.1 vs. 68.6 ± 17.3		29.2 ± 5.3 vs. 29.4 ± 5.1	60.3 vs. 76.4	1.7 vs. 12.3	Decreased
Wander et al. [66]	USA	29,685/64,892	White: 66.0, Black: 27.0, Hispanic: 9.0	64.0	67.7			89.0	36.0	Decreased
Wang et al. (1) [67]	USA	9/7	African American: 23.0, Hispanic: 16.0	52.0	67 (12.5)		27.6	64.0	24.0	NS
Wang et al. (2) [68]	England	110/54		61.9 vs. 52.0	64.8 ± 11.7 vs. 67.8 ± 13.1		32.1 ± 6.7 vs. 31.8 ± 6.8	59.1 vs. 60.7	13.0 vs. 24.2	NS
Wargny et al. [69]	France	1553/1241	White: 58.1, African: 17.4, Asian: 3.6	36.3	69.7 ± 13.2	7.7 (6.8–9.0)	28.4 (25.0–32.4)	76.8		Decreased
Cheng et al. [12]	China	678/553		53.8 vs. 49.9	62.0 (55.0–68.0) vs. 64.0 (58.0–70.0)	8.1 (7.0–9.9) vs. 7.6 (6.7–8.9)	24.3 (22.0–25.9) vs. 24.5 (22.6–26.2)		2.4 vs. 2.6	NS
Yuan et al. [72]	China	73/109		52.1	62.0 (55.0–70.0)	8.3 (7.2–9.9)	23.7 (22.0–25.4)	52.1	0.0	Decreased
Pazoki et al. [13]	Iran	177/216		56.2	65.4 ± 11.6		28.0 ± 5.1	65.4	7.9	NS
<i>SU users/nonusers</i>										
An et al. [15]	Korea	212/809		39.9	45.0 ± 19.9			18.2	0.8	NS
Cernigliaro et al. [19]	Italy	35/137								NS
Chen et al. [20]	China	53/67		42.9	66.0 (60.0–72.5) vs. 65.0 (55.0–73.0)	8.3 (7.4–9.5) vs. 7.7 (7.1–10.4)		61.2	10.2	NS
Dave et al. [23]	Africa	2110/3598		39.3	55.0 (46.0–63.0)			55.5	9.2	NS
Eliboi et al. [26]	Turkey	66/366		45.6	63.3 ± 10.3			74.1	4.6	NS
Khunti et al. [34]	England	561,290/2,290,175	White: 63.7, Black: 5.0, Asian: 17.2	60.5	67.0 (57.0–77.0)			80.7		Decreased
Kim et al. [35]	Korea	60/175		45.1	68.3 ± 11.9		24.2 ± 3.2	62.6	7.7	NS
Li et al. (1) [39]	China	22/109		56.5	66.8 ± 11.6	7.9 ± 1.9	24.2 ± 3.4	59.5		NS
Li et al. (2) [38]	China	91/296		51.1	60.0 (49.0–68.0)			48.6	1.0	NS
Luk et al. [40]	China	385/679	Asian	57.7 vs. 51.5	66.0 (58.5–73.1) vs.	7.7 (6.9–9.1) vs. 6.9 (6.4–8.2)	24.4 (21.8–27.8) vs.	69.4 vs. 48.5	25.5 vs. 19.9	NS

(continued on next page)

Table 1 (continued)

Study	Country	Number of patients	Race/ethnicity (%)	Male sex (%)	Age (years)	HbA1C (%)	Body mass index (kg/m ²) or obesity (%)	Hypertension (%)	CKD (%)	Mortality
Luo et al. [42]	China	37/154			65.3 (57.3–73.6)		23.5 (21.5–27.0)			
Mirani et al. [46]	Italy	10/80		56.5	62.7 ± 11.0	7.9 (6.3–9.0)		55.5	3.0	Decreased
Nyland et al. [50]	USA	1889/27,627	White: 47.9, African American: 25.5, Asian: 3.1	60.0 vs. 73.8	75.0 ± 8.0 vs. 70.0 ± 12.0		Obesity: 50.0% vs. 47.5%	80.0 vs. 76.3	0.0 vs. 21.4	NS
Oh et al. [51]	Korea	3680/23,813		48.2	60.9 ± 15.0	7.7 ± 2.1	32.8 ± 8.9	47.7	15.5	Decreased
Orioli et al. [53]	Belgium	19/49		48.0	69.0 ± 14.0	7.1 (6.6–8.3)	30.5 ± 5.3	80.8	34.2	NS
Shetaskova et al. [61]	Russia	129/180								NS
Silverii et al. [62]	Italy	33/126		54.1	73.3 ± 12.7					NS
Sourij et al. [63]	Austria	14/166		63.9	67.6 ± 14.0	6.7 (1.9)	29.4 ± 5.7	77.0	23.1	NS
Wander et al. [66]	USA	12,298/52,594	White: 66.0, Black: 27.0, Hispanic: 9.0	64.0	67.7			89.0	36.0	NS
Wargny et al. [69]	France	782/2012	White: 58.1, African: 17.4, Asian: 3.6	36.3	69.7 ± 13.2	7.7 (6.8–9.0)	28.4 (25.0–32.4)	76.8		NS
Yuan et al. [72]	China	43/139		55.8	67.0 (60.0–73.0)	8.5 (7.0–9.5)	23.7 (22.0–25.4)	48.8	0.0	Decreased
Pazoki et al. [13]	Iran	72/321		56.2	65.4 ± 11.6		28.0 ± 5.1	65.4	7.9	NS
<i>TZD users/nonusers</i>										
Cernigliaro et al. [19]	Italy	10/162								NS
Eliboi et al. [26]	Turkey	27/405		45.6	63.3 ± 10.3			74.1	4.6	NS
Khunti et al. [34]	England	60,085/2,791,380	White: 63.5, Black: 3.7, Asian: 18.4	63.4	67.0 (57.0–77.0)			80.5		NS
Luo et al. [42]	China	7/184		56.5	62.7 ± 11.0	7.9 (6.3–9.0)		55.5	3.0	NS
Nyland et al. [50]	USA	469/23,714	White: 52.4, African American: 23.2, Asian: 3.5	53.3 vs. 48.8	63.1 ± 12.5 vs. 60.9 ± 15.3	8.2 ± 2.0 vs. 7.5 ± 2.1	34.3 ± 9.0 vs. 32.3 ± 8.7	52.1 vs. 44.9	17.4 vs. 14.9	NS
Oh et al. [51]	Korea	1264/26,229								NS
Silverii et al. [62]	Italy	8/151		54.1	73.3 ± 12.7					NS
Wander et al. [66]	USA	2075/62,817	White: 66.0, Black: 27.0, Hispanic: 9.0	64.0	67.7			89.0	36.0	NS
<i>AGI users/nonusers</i>										
An et al. [15]	Korea	7/1014		39.9	45.0 ± 19.9			18.2	0.8	Increased
Chen et al. [20]	China	69/51		42.9	66.0 (57.5–73.0) vs. 65.0 (56.0–72.0)	8.4 (7.4–10.3) vs. 7.9 (6.9–9.1)		61.2	10.2	NS
Khunti et al. [34]	England	1665/2,849,800	White: 56.5, Black: 7.5, Asian: 23.4	56.8	67.0 (57.0–77.0)			87.4		NS
Li et al. (1) [39]	China	38/93		56.5	66.8 ± 11.6	7.9 ± 1.9	24.2 ± 3.4	59.5		NS
Li et al. (2) [38]	China	140/247		51.1	60.0 (49.0–68.0)			48.6	1.0	NS
Luo et al. [42]	China	77/114		65.0 vs. 63.0	62.3 ± 9.6 vs. 61.9 ± 9.4	7.9 ± 1.8 vs. 8.3 ± 2.0		55.5	2.2 vs. 0.0	Decreased
Nyland et al. [50]	USA	16/29,500	White: 47.9, African American: 25.5, Asian: 3.1	48.2	60.9 ± 15.0	7.7 ± 2.1	32.8 ± 8.9	47.7	15.5	NS
Yuan et al. [72]	China	88/94		51.1		8.2 (7.0–9.2)		58.0	1.1	Decreased

(continued on next page)

Table 1 (continued)

Study	Country	Number of patients	Race/ethnicity (%)	Male sex (%)	Age (years)	HbA1C (%)	Body mass index (kg/m ²) or obesity (%)	Hypertension (%)	CKD (%)	Mortality
					66.0 (57.0–72.0)		23.7 (22.0–25.4)			
<i>GLP-1RA users/nonusers</i>										
Cernigliaro et al. [19]	Italy	8/164								NS
Israelsen et al. [32]	Denmark	370/558		53.0	59.0 (51.0–70.0)		Obesity: 29.2%	56.2		NS
Kahkoska et al. [33]	US	6692/5854	White: 64.1	40.9	55.7 ± 12.6	8.0 ± 2.0	37.2 ± 8.1	74.9 vs. 76.0	18.5	Decreased
Khunti et al. [34]	England	100,820/ 2,750,645	White: 76.3, Black: 3.3, Asian: 7.9	51.7	67.0 (57.0–77.0)				83.1	NS
Nyland et al. [50]	USA	1774/ 23,714	White: 52.3, Black: 28.7, Asian: 0.9	39.2 vs. 48.8	55.0 ± 12.7 vs. 60.9 ± 15.3	8.4 ± 2.2 vs. 7.5 ± 2.1	37.5 ± 9.3 vs. 32.3 ± 8.7	55.9 vs. 44.9	12.9 vs. 14.9	Decreased
Orioli et al. [53]	Belgium	5/63		48.0	69.0 ± 14.0	7.1 (6.6–8.3)	30.5 ± 5.3	80.8	34.2	NS
Ramos-Rincon et al. [56]	Spain	37/753		47.1			Obesity: 17.7%	84.3	17.2	NS
Shetaskova et al. [61]	Russia	1/308								NS
Silverii et al. [62]	Italy	7/152		54.1	73.3 ± 12.7					NS
Sourij et al. [63]	Austria	3/177		63.9	67.6 ± 14.0	6.7 (1.9)	29.4 ± 5.7	77.0	23.1	NS
Wander et al. [66]	USA	4737/ 60,155	White: 66.0, Black: 27.0, Hispanic: 9.0	64.0	67.7			89.0	36.0	NS
Wargny et al. [69]	France	254/2540	White: 58.1, African: 17.4, Asian: 3.6	36.3	69.7 ± 13.2	7.7 (6.8–9.0)	28.4 (25.0–32.4)	76.8		NS
<i>DPP-4i users/nonusers</i>										
An et al. [15]	Korea	229/792		39.9	45.0 ± 19.9			18.2	0.8	NS
Cernigliaro et al. [19]	Italy	13/159								NS
Chen et al. [20]	China	20/100		42.9	66.0 (56.0–73.0) vs. 65.0 (57.0–72.0)	7.8 (6.8–10.6) vs. 8.3 (7.3–9.5)		61.2	10.2	NS
Eliboi et al. [26]	Turkey	246/186		45.6	63.3 ± 10.3			74.1	4.6	NS
Emral et al. [27]	Turkey	6846/ 26,632		42.0 vs. 41.3	60.0 (16.0) vs. 52.0 (24.0)	8.1 (2.7) vs. 6.4 (1.6)	30.8 (6.7) vs. 29.4 (7.3)	85.6 vs. 64.0		Decreased
Fanidi et al. [28]	Italy	9/72			72.2 ± 12.8 vs. 70.1 ± 13.3	7.5 ± 3.3 vs. 7.6 ± 4.3		88.9 vs. 67.1	11.2 vs. 15.8	NS
Israelsen et al. [32]	Denmark	284/644		60.9	67.0 (57.0–76.0)		Obesity: 12.3%	61.6		NS
Kahkoska et al. [33]	USA	3511/8935	White: 57.4	49.9	64.1 ± 12.9 vs. 58.6 ± 13.1	7.8 ± 1.9 vs. 8.0 ± 1.9	36.0 ± 6.2 vs. 35.4 ± 8.2	78.7 vs. 76.0	31.6	Increased
Khunti et al. [34]	England	479,555/ 2,371,910	White: 65.5, Black: 4.7, Asian: 15.7	58.3	67.0 (57.0–77.0)			81.6		Increased
Kim et al. [35]	Korea	85/150		45.1	68.3 ± 11.9		24.2 ± 3.2	62.6	7.7	NS
Kristan et al. [36]	USA	76/756	White: 32.7, African American: 52.0, Asian: 1.4	51.0	62.0 ± 15.0	7.9 ± 2.3	32.9 ± 8.6	78.4	21.3	NS
Luk et al. [40]	China	199/952		59.3 vs. 53.2	67.0 (58.4–75.5) vs. 65.1 (56.8–72.2)	7.6 (6.8–8.9) vs. 7.2 (6.5–8.9)	25.0 (18.7–27.0) vs. 23.3 (21.6–27.4)	61.8 vs. 52.3	36.2 vs. 17.2	NS
Luo et al. [42]	China	11/180		56.5	62.7 ± 11.0	7.9 (6.3–9.0)		55.5	3.0	NS
Meijer et al. [45]	Netherlands	28/537		60.7 vs. 64.2	66.9 ± 12.4 vs. 69.5 ± 12.5		29.1 ± 6.0 vs. 29.8 ± 6.3	66.7 vs. 70.0	25.9 vs. 14.4	NS
Mirani et al. [46]	Italy	11/79		90.9 vs. 69.6	70.0 ± 13.0 vs. 71.0 ± 12.0		Obesity: 27.3% vs. 50.6%	54.6 vs. 79.8	18.2 vs. 19.0	Decreased
Noh et al. [49]	Korea	453/133								NS

(continued on next page)

Table 1 (continued)

Study	Country	Number of patients	Race/ethnicity (%)	Male sex (%)	Age (years)	HbA1C (%)	Body mass index (kg/m ²) or obesity (%)	Hypertension (%)	CKD (%)	Mortality
Nyland et al. [50]	USA	2264/23,714	White: 49.2, African: 36.6, Asian: 5.1	49.2 vs. 55.6 49.1 vs. 48.8	64.6 ± 13.5 vs. 60.9 ± 15.3	8.0 ± 2.0 vs. 7.5 ± 2.1	31.4 ± 8.1 vs. 32.3 ± 8.7	55.9 vs. 44.9	21.2 vs. 18.0 22.4 vs. 14.9	Increased
Oh et al. [51]	Korea	4132/23,361								NS
Orioli et al. [53]	Belgium	4/64		48.0	69.0 ± 14.0	7.1 (6.6–8.3)	30.5 ± 5.3	80.8	34.2	NS
Perez-Belmonte et al. [54]	Spain	180/1409		59.4 vs. 62.9	78.8 ± 7.1 vs. 74.7 ± 8.2		Obesity: 30.6% vs. 28.1%	55.6 vs. 56.5	32.2 vs. 11.9	Increased
Ramos-Rincon et al. [56]	Spain	266/524		47.1			Obesity: 17.7%	84.3	17.2	Decreased
Shetaskova et al. [61]	Russia	26/283								NS
Silverii et al. [62]	Italy	13/146		54.1	73.3 ± 12.7					NS
Sourij et al. [63]	Austria	42/138		63.9	67.6 ± 14.0	6.7 (1.9)	29.4 ± 5.7	77.0	23.1	NS
Strollo et al. [64]	Italy	30/163		54.9	76.7 ± 11.8					NS
Wander et al. [66]	USA	5810/59,082	White: 66.0, Black: 27.0, Hispanic: 9.0	64.0	67.7			89.0	36.0	NS
Wargny et al. [69]	France	615/2179	White: 58.1, African: 17.4, Asian: 3.6	36.3	69.7 ± 13.2	7.7 (6.8–9.0)	28.4 (25.0–32.4)	76.8		NS
Wong et al. [70]	China	107/1107		60.7 vs. 53.7	66.3 ± 11.7 vs. 65.1 ± 13.0	7.8 ± 2.3 vs. 7.4 ± 2.5	Obesity: 15% vs. 11.3%	88.8 vs. 75.2	30.8 vs. 11.3	NS
Pazoki et al. [13]	Iran	20/373		56.2	65.4 ± 11.6		28.0 ± 5.1	65.4	7.9	NS
<i>SGLT-2i users/nonusers</i>										
Cernigliaro et al. [19]	Italy	4/168								Decreased
Eliboi et al. [26]	Turkey	56/376		45.6	63.3 ± 10.3			74.1	4.6	NS
Israelsen et al. [32]	Denmark	274/654		61.8	59.0 (52.0–68.0)		Obesity: 15.4%	49.6		NS
Kahkoska et al. [33]	USA	3665/8781	White: 33.9	55.2	57.9 ± 11.7	8.2 ± 1.8	35.2 ± 7.8	77.3	16.3	Decreased
Khunti et al. [34]	England	266,505/2,584,960	White: 66.8, Black: 3.6, Asian: 15.2	60.8	67.0 (57.0–77.0)			75.4		Decreased
Kim et al. [35]	Korea	8/227		45.1	68.3 ± 11.9		24.2 ± 3.2	62.6	7.7	NS
Nyland et al. [50]	USA	792/28,724	White: 47.9, African: 25.5, Asian: 3.1	48.2	60.9 ± 15.0	7.7 ± 2.1	32.8 ± 8.9	47.7	15.5	Decreased
Orioli et al. [53]	Belgium	4/64		48.0	69.0 ± 14.0	7.1 (6.6–8.3)	30.5 ± 5.3	80.8	34.2	NS
Ramos-Rincon et al. [56]	Spain	45/745		47.1			Obesity: 17.7%	84.3	17.2	NS
Shetaskova et al. [61]	Russia	13/296								NS
Silverii et al. [62]	Italy	4/155		54.1	73.3 ± 12.7					NS
Sourij et al. [63]	Austria	24/156		63.9	67.6 ± 14.0	6.7 (1.9)	29.4 ± 5.7	77.0	23.1	NS
Wander et al. [66]	USA	5542/59,350	White: 66.0, Black: 27.0, Hispanic: 9.0	64.0	67.7			89.0	36.0	Decreased
<i>Insulin users/nonusers</i>										
Agarwal et al. [14]	USA	531/661	White: 15.5, African: 74.5	49.3	67.9 ± 13.7	7.5 ± 2.0	30.1 ± 7.5	90.9	42.5	Increased
Boye et al. [17]	USA	3461/6070		46.0	71.6 ± 12.5	7.2			37	Increased

(continued on next page)

Table 1 (continued)

Study	Country	Number of patients	Race/ethnicity (%)	Male sex (%)	Age (years)	HbA1C (%)	Body mass index (kg/m ²) or obesity (%)	Hypertension (%)	CKD (%)	Mortality
Cernigliaro et al. [19]	Italy	42/130	Caucasian: 47, African American: 31, Asian: 2, Hispanic: 5							NS
Chen et al. [20]	China	7/49		42.9	65.0	8.8		61.2	10.2	Increased
Cheng et al. [12]	China	11/39		54.5 vs. 53.8	58.0 vs. 52.0 (54.0–60.0) vs. (44.0–65.0)	65.0 vs. 7.5 (6.8–8.3)	24.8 (19.9–25.6) vs. 26.0 (23.8–27.2)	27.3 vs. 38.5	27.3 vs. 35.9	NS
Crouse et al. [22]	USA	87/133	White: 27.6, African American: 64.9	50.6			Obesity: 74.5%	91.6		NS
Dave et al. [23]	Africa	2073/3635		39.3	55.0 (46.0–63.0)			55.5	9.2	Increased
Deng et al. [24]	China	29/56		57.6	65.0 (34.0–91.0)			68.2	7.1	NS
Giorda et al. [30]	Italy	656/1226		50.9				84.4	60.1	NS
Khunti et al. [34]	England	350,960/2,500,505	White: 71.1, Black: 4.7, Asian: 12.3	54.5	67.0 (57.0–77.0)			85.2		Increased
Kim et al. [35]	Korea	19/216		45.1	68.3 ± 11.9		24.2 ± 3.2	62.6	7.7	NS
Kristan et al. [36]	USA	281/551	White: 32.7, African American: 52.0, Asian: 1.4	51.0	62.0 ± 15.0	7.9 ± 2.3	32.9 ± 8.6	78.4	21.3	NS
Lally et al. [37]	USA	103/190	White: 54.4, Black: 40.8	97.1	73.3 ± 9.4 vs. 75.6 ± 9.2	7.7 ± 1.5 vs. 6.5 ± 1.3	29.3 ± 3.0 vs. 28.0 ± 7.0		52.4	NS
Li et al. (1) [39]	China	26/105		56.5	66.8 ± 11.6	7.9 ± 1.9	24.2 ± 3.4	59.5		NS
Li et al. (2) [38]	China	102/285		51.1	60.0 (49.0–68.0)			48.6	1.0	NS
Luk et al. [40]	China	385/679	Asian	57.7 vs. 51.5	66.0 (58.5–73.1) vs. 65.3 (57.3–73.6)	7.7 (6.9–9.1) vs. 6.9 (6.4–8.2)	22.9 (19.8–25.9) vs. 24.4 (22.2–27.4)	69.4 vs. 48.5	25.5 vs. 19.9	Increased
Luo et al. [42]	China	88/103		56.5	62.7 ± 11.0	7.9 (6.3–9.0)		55.5	3.0	NS
Mansour et al. [44]	Iran	25/86		55.9	63.6 ± 13.3		28.2 ± 5.6	57.7	9.0	NS
Mirani et al. [46]	Italy	29/61		72.4 vs. 72.1	72.0 ± 10.0 vs. 70.0 ± 13.0		Obesity: 51.7% vs. 45.9%	79.3 vs. 75.4	31.0 vs. 13.1	Increased
Nyland et al. [50]	USA	9149/20,367	White: 47.9, African American: 25.5, Asian: 3.1	48.2	60.9 ± 15.0	7.7 ± 2.1	32.8 ± 8.9	47.7	15.5	Increased
Oh et al. [51]	Korea	914/26,579								NS
Orioli et al. [53]	Belgium	31/37		48.0	69 ± 14	7.1 (6.6–8.3)	30.5 ± 5.3	80.8	34.2	NS
Perez-Belmonte et al. [54]	Spain	292/1458			77.9 ± 9.0		Obesity: 20.9% vs. 28.8%	50.0 vs. 57.8		Increased
Ramos-Rincon et al. [56]	Spain	225/565		47.1			Obesity: 17.7%	84.3	17.2	NS
Riahl et al. [58]	USA	88/78	White: 6.0, African American: 71.0	52.0	66.4 ± 12.7	8.6 ± 2.5 vs. 7.0 ± 1.7	31.1 ± 8.5	91.0	25.0	Increased
Satman et al. [59]	Turkey	3340/15,318		42.3	53.0 (22.0)	6.9 (2.3)	30.0 (7.1)	66.4	18.9	Increased
Shetaskova et al. [61]	Russia	115/194								Increased
Silverii et al. [62]	Italy	43/116		54.1	73.3 ± 12.7					NS
Sourij et al. [63]	Austria	41/139		63.9	67.6 ± 14.0	6.7 (1.9)	29.4 ± 5.7	77.0	23.1	NS
Wander et al. [66]	USA	18,521/46,371	White: 66.0, Black: 27.0, Hispanic: 9.0	64.0	67.7			89.0	36.0	Increased
Wargny et al. [69]	France	1039/1757		36.3	69.7 ± 13.2	7.7 (6.8–9.0)	28.4 (25.0–32.4)	76.8		Increased

(continued on next page)

Table 1 (continued)

Study	Country	Number of patients	Race/ethnicity (%)	Male sex (%)	Age (years)	HbA1C (%)	Body mass index (kg/m ²) or obesity (%)	Hypertension (%)	CKD (%)	Mortality
			White: 58.1, African: 17.4, Asian: 3.6							
Yan et al. [71]	China	4/30		68.8	69.4 ± 9.9	7.2 (6.7–8.3)		50.0	0.0	Increased
Yuan et al. [72]	China	76/106		47.4	66.0 (61.0–72.0)	8.6 (7.9–10.0)	23.7 (22.0–25.4)	57.9	2.6	Increased
Pazoki et al. [13]	Iran	53/340		56.2	65.4 ± 11.6		28.0 ± 5.1	65.4	7.9	NS

Data are presented as mean ± SD or median (IQR).

Abbreviation: AGI, alpha-glucosidase inhibitor; CKD, chronic kidney disease; DPP-4i, dipeptidyl peptidase inhibitor; GLP-1RA, glucagon-like peptide-1 receptor agonist; NS, not significant; SGLT-2i, sodium–glucose transporter-2 inhibitor; SU, sulfonylurea; TZD, thiazolidinedione.




Fig. 2. Forest plot of the relationship between metformin and mortality in patients with COVID-19 having type 2 diabetes.

medications in more homogenous populations. The results of metformin and insulin were consistently confirmed among various groups in terms of vulnerability, including advanced age, high BMI, and high rate of CKD

(Figs. A.6–A.8 and A.25–A.27, respectively). Meanwhile, GLP-1RA and SGLT-2i were still beneficial compared to nonusers, albeit less pronounced in populations with a higher rate of comorbidities and older




Fig. 3. Forest plot of the relationship between GLP-1RA and mortality in patients with COVID-19 having type 2 diabetes.




Fig. 4. Forest plot of the relationship between SGLT-2i and mortality in patients with COVID-19 having type 2 diabetes.

patients, respectively (Figs. A.15, A.17, and A.21). Despite overall mortal neutrality, SU might have mild benefits in younger and less vulnerable populations (Figs. A.9–A.12). In contrast, DPP-4i showed harm or at least no benefit (A.18–A.20).

3.3.4. Sensitivity analysis

We further performed a sensitivity analysis by using two methods. First, we identified outliers by implementing the outlier removal algorithm in the *dmetar* package to explore the influence of individual studies on pooled effects. After outliers were removed, the pooled OR did not significantly change (all $p > 0.05$). Next, we conducted the trim-and-fill method to impute missing effects and concluded that our main results were stable after extending additional effects (all $p > 0.05$; Table 3).

3.3.5. Dose-response meta-analysis

Metformin was the only medication that was reported the daily dosage in these original papers. Therefore, we performed a dose-response meta-analysis for metformin. We observed a significant linear dose-response association between metformin dose and odds ratio of mortality (estimate: -0.88 , standard error: 0.22 , $p < 0.001$) and no evidence of heterogeneity among studies ($I^2 = 0\%$, $p = 0.46$; Fig. 5).

3.3.6. Comparison with previous meta-analyses

We next compared our results with those from other publications [4–7,73–88]. No published meta-analysis has analyzed the association between TZD or AGI and COVID-19-related mortality (Table 4).

4. Discussion

4.1. Summary of main findings

To the best of our knowledge, this timely study has been the most extensive systematic review and meta-analysis confirming that different antidiabetic medications could predispose individuals with COVID-19 to different prognoses. Compared with a previous publication [5], we observed significant roles of GLP-1RA and SGLT-2i, besides metformin, in protecting individuals from COVID-19-related death. Similar to most studies, we also identified a positive association between DPP-4i usage and mortality. Moreover, we are the first to report the pooled effect of TZD and the pooled effect of AGI. Similar to smaller meta-analyses [5,75,85], our data also indicated the inconsistent impact of SU. Finally, we are the first to perform a dose-response meta-analysis regarding the daily dose of metformin to predict the magnitude of the effect on mortality in patients with COVID-19 having diabetes. These findings can have a large impact on the outpatient management strategy

Table 2
Meta-regression analysis on potentially confounding factors.

Medication	Confounding factor	Estimate	SE	p-Value	
Metformin	Continent (vs. America)				
	Africa	0.274	0.482	0.57	
	Asia	-0.076	0.227	0.74	
	Europe	0.096	0.235	0.68	
	White race (%)	0.004	0.006	0.53	
	Age (years)	-0.003	0.013	0.81	
	Male sex (%)	-0.001	0.007	0.87	
	HbA1C (%)	-0.100	0.181	0.59	
	Body mass index (kg/m ²)	0.043	0.037	0.26	
	Hypertension (%)	-0.001	0.006	0.87	
	Chronic kidney disease (%)	0.001	0.005	0.89	
	Sulfonylurea	Continent (vs. America)			
		Africa	-0.123	0.204	0.56
		Asia	0.075	0.185	0.69
		Europe	0.076	0.158	0.64
		White race (%)	0.017	0.003	0.02
Age (years)		0.015	0.007	0.03	
Male sex (%)		0.009	0.003	0.01	
HbA1C (%)		-0.753	0.551	0.55	
Body mass index (kg/m ²)		-0.045	0.030	0.19	
Hypertension (%)		0.006	0.002	0.01	
Chronic kidney disease (%)		0.009	0.003	0.02	
Thiazolidinedione		Continent (vs. America)			
		Asia	0.389	0.398	0.37
		Europe	0.182	0.350	0.62
		White race (%)	0.071	0.026	0.22
		Age (years)	0.099	0.063	0.19
	Male sex (%)	-0.001	0.030	0.97	
	HbA1C (%)	Insufficient data for analysis			
	Body mass index (kg/m ²)	Insufficient data for analysis			
	Hypertension (%)	0.025	0.008	0.05	
	Chronic kidney disease (%)	0.005	0.025	0.87	
	Alpha-glucosidase inhibitor	Continent (vs. America)			
		Asia	0.073	1.966	0.97
		Europe	1.452	2.234	0.54
		White race (%)	Insufficient data for analysis		
		Age (years)	-0.078	0.067	0.28
		Male sex (%)	-0.090	0.054	0.15
HbA1C (%)		1.845	1.991	0.42	
Body mass index (kg/m ²)		0.108	0.174	0.65	
Hypertension (%)		-0.007	0.027	0.81	
Chronic kidney disease (%)		0.023	0.124	0.86	
Glucagon-peptide like-1 receptor agonist		Continent (vs. America)			
		Asia	1.707	1.459	0.27
		Europe	-0.004	0.283	0.99
		White race (%)	0.033	0.027	0.30
		Age (years)	0.043	0.021	0.08
		Male sex (%)	0.032	0.010	0.01
	HbA1C (%)	-1.000	0.361	0.07	
	Body mass index (kg/m ²)	-0.053	0.038	0.25	
	Hypertension (%)	0.029	0.010	0.02	
	Chronic kidney disease (%)	0.008	0.007	0.32	
Dipeptidyl peptidase-4 inhibitor	Continent (vs. America)				
	Asia	-0.183	0.247	0.47	
	Europe	-0.260	0.243	0.30	
	White race (%)	-0.003	0.018	0.90	
	Age (years)	-0.005	0.014	0.74	

Table 2 (continued)

Medication	Confounding factor	Estimate	SE	p-Value
Sodium-glucose transporter-2 inhibitor	Male sex (%)	0.000	0.012	1.00
	HbA1C (%)	0.005	0.347	0.99
	Body mass index (kg/m ²)	0.087	0.030	0.02
	Hypertension (%)	-0.001	0.006	0.87
	Chronic kidney disease (%)	0.009	0.010	0.39
	Continent (vs. America)			
	Asia	0.675	0.381	0.11
	Europe	-0.500	0.218	0.04
	White race (%)	-0.006	0.017	0.77
	Age (years)	0.029	0.048	0.56
Insulin	Male sex (%)	-0.031	0.023	0.21
	HbA1C (%)	0.565	0.128	0.05
	Body mass index (kg/m ²)	-0.069	0.107	0.57
	Hypertension (%)	0.011	0.012	0.38
	Chronic kidney disease (%)	-0.008	0.018	0.66
	Continent (vs. America)			
	Africa	-0.217	0.576	0.71
	Asia	0.009	0.285	0.98
	Europe	-0.221	0.280	0.44
	White race (%)	-0.000	0.011	0.98
Age (years)	-0.032	0.020	0.12	
Male sex (%)	-0.001	0.011	0.97	
HbA1C (%)	0.029	0.347	0.93	
Body mass index (kg/m ²)	0.115	0.061	0.08	
Hypertension (%)	-0.011	0.008	0.19	
Chronic kidney disease (%)	-0.002	0.009	0.87	


Abbreviations: SE, standard error.

of diabetes patients amid the COVID-19 pandemic. These results can be helpful for clinicians in terms of choosing proper glucose-lowering regimens and dosage for those patients to reduce the risk of in-hospital death, i.e. by promoting the prescription of metformin, GLP-1RA, and SGLT-2i in the absence of any contraindications. In contrast, caution should be exercised in long-term insulin use.

Metformin might decrease or did not significantly affect COVID-19 death in the original studies. However, when performing meta-analyses, it yielded the most consistent result, even in vulnerable patients. Our study corroborated previous publications highlighting the potential benefits of metformin in patients with COVID-19 and diabetes. Several mechanisms might explain the lower mortality from SARS-CoV-2 infections in individuals taking metformin. First, beyond the hypoglycemic effect, metformin could reduce the release of inflammatory cytokines such as interleukin-6 and tumor necrosis factor-alpha, which play a vital role in COVID-19 pathophysiology [89]. Second, metformin is also involved in other pathways: angiotensin-converting enzyme-2 (ACE-2) modulation through adenosine monophosphate-activated protein kinase, decreased coagulation and thrombosis formation, mast cell stabilization, and improved endothelial function [18,90]. Therefore, several researchers are currently investigating metformin as a host-directed medication in patients with COVID-19 [91]. Our current study indicated that metformin is effective among different races, sexes, weight status, and levels of glucose control. The dosage of metformin also affected the risk of mortality. First, Cheng et al. indicated that preadmission metformin usage was associated with better outcomes in a dose-response manner. In that study, metformin median dose was 1000 (890–1220) mg/day [21]. Ghany et al. reported that individuals using metformin at a dose of ≥1000 mg/day had lower mortality than those on 500–850 mg/day [29]. Referenced to nonusers, Ong et al. reported the greatest benefit on mortality with the dose from 1000 to <2000 mg/day [52]. Our findings were consistent with these studies. Specifically, every

Table 3
Sensitivity analysis.

Medication	Main meta-analysis		Sensitivity analysis				
	OR (95% CI)	I ²	Outlier removal method			Trim-and-fill method	
			OR (95% CI)	I ²	p value ^a	OR (95% CI)	p value ^b
Metformin	0.54 (0.47–0.62)	86%	0.50 (0.45–0.55)	41%	0.37	0.61 (0.54–0.70)	0.17
Sulfonylurea	0.92 (0.83–1.01)	44%	0.98 (0.90–1.06)	18%	0.31	0.93 (0.84–1.04)	0.80
Thiazolidinedione	0.90 (0.71–1.14)	46%	No outlier			0.88 (0.70–1.11)	0.91
Alpha-glucosidase inhibitor	0.61 (0.26–1.45)	77%	1.13 (0.60–2.11)	47%	0.26	1.45 (0.57–3.74)	0.18
Glucagon-like peptide-1 receptor agonist	0.51 (0.37–0.69)	85%	0.54 (0.49–0.60)	0%	0.70	0.62 (0.45–0.84)	0.40
Dipeptidyl peptidase-4 inhibitor	1.23 (1.07–1.42)	82%	1.25 (1.14–1.37)	37%	0.86	1.29 (1.12–1.48)	0.67
Sodium-glucose transporter-2 inhibitor	0.60 (0.40–0.88)	91%	0.67 (0.52–0.85)	47%	0.63	0.54 (0.37–0.79)	0.72
Insulin	1.70 (1.33–2.19)	97%	1.60 (1.41–1.81)	60%	0.65	2.00 (1.58–2.52)	0.37

^a Comparison of OR before vs. after removing outliers.^b Comparison of OR before vs. after trimming and filling.**Fig. 5.** Dose–response meta-analysis between daily metformin dosage and mortality in patients with COVID-19 with diabetes.

250 mg/day increase in metformin use was associated with a 19.7% lower odds of mortality. In summary, the minimum metformin dosage that was found beneficial was 500 mg/day, and the higher the dose, the higher the effect. However, notably, the maximum approved dose for metformin is only 2550 mg/day (immediate-release form) and 2000 mg/day (extended-release form).

GLP-1RA and SGLT-2i are two novel classes of antidiabetic medications that have been approved for cardiorenal protection in type 2 diabetes patients. In the COVID-19 scenario, GLP-1RA can help reduce cytokine-induced lung injury by interfering with the NF-κB pathway or exerting anti-inflammatory effects [92,93]. Meanwhile, when hypoxemia and hypoxia occur, SGLT-2i reverses the acid-base cytokine balance by decreasing lactic acid accumulation, thereby inhibiting the lowering of cytosolic pH and preventing cell damage during COVID-19-induced cytokine storm [94]. These cardiorenal benefits can synergistically offer protection to vital organs to reduce the risk of severity progression and death in the context of SARS-CoV-2 infection. It was not surprised from our findings that SGLT-2i might have more obvious impacts on those with high baseline BMI or history of CKD due to the renal-metabolic benefits of this class. Consistently, SGLT-2i was more beneficial in a subgroup with a history of cardiovascular disease [34]. It should be noted that, however, this benefit might be less pronounced in vulnerable patients.

In contrast to previous smaller meta-analyses reporting that DPP-4i had no significant effect on COVID-19-related death [6,75,80,85], after incorporating a larger number of studies, we observed that pre-admission DPP-4i users were associated with higher odds of in-hospital

mortality. DPP-4i has yielded both putative protective and harmful effects on the underlying mechanisms of SARS-CoV-2 infection and progression from preclinical studies [4,95]. Moreover, the controversial results of DPP-4i from various original studies and meta-analyses up to the present might be explained by the fact that the authors could not entirely exclude potential confounders, even with multivariate adjustment or propensity score matching. For example, we observed a trend toward higher use of DPP-4i in older fragile people and in patients with several comorbidities who had a compelling need to minimize hypoglycemia. These characteristics promoted the prescription of DPP-4i and limited the indication of other antidiabetic medications [33,50,54]. On the other hand, our subgroup analyses showed that DPP-4i might have little or no benefit among patient groups differed by vulnerability, suggesting that DPP-4i might not be associated with favorable COVID-19-related outcomes. To summarize, higher mortality rates in DPP-4i users should be cautiously interpreted.

The association between insulin treatment and severity or mortality is more complex. This result may still be affected by a confounding factor regarding the late commencement of insulin at an advanced stage of diabetes and the heterogeneous effectiveness of different insulin regimens, such as basal, basal-bolus, or premixed therapies. We speculate that insulin therapy is likely a surrogate indicator of diabetes progression accompanied by beta-cell dysfunction. Therefore, it was not insulin therapy, per se, that was associated with poor prognosis of patients with COVID-19 having type 2 diabetes, but rather that it represented a proxy of severity and duration of diabetes. However, notably, iatrogenic hyperinsulinemia caused by exogenous insulin use might lead to adverse effects, including insulin resistance due to downregulation of insulin receptors, vascular changes, and subsequent adverse cardiovascular outcomes [96]. Moreover, our subgroup analyses as well as those from previous publications controlling for severity markers did not eliminate the association, raising concerns about the actual harmful effects of insulin [17]. Like DPP-4i, the increased risk of death among insulin users should be cautiously interpreted.

Unlike two smaller meta-analyses demonstrating that SU could reduce mortality risk [5,75], our results indicated that SU was not significantly associated with COVID-19-related mortality. Moreover, our study conducted a meta-analysis of AGI, which has not been reported previously. Traditionally, these drugs were often considered cardiovascular neutral. This characteristic makes them not a first-line treatment in patients with type 2 diabetes in general. Therefore, it is reasonable that they did not affect mortality in the COVID-19 setting, where cardiovascular events caused by hyperinflammation and hypercoagulation were the leading causes of intensive care unit admission, mechanical ventilation, and death. Although TZD could alleviate the long-term progressive atherosclerosis and inhibit the macrophage training, both of which were associated with the development of severe COVID-19, its benefit might be counteracted by the putative harmful effect regarding the fluid retention that could exacerbate pulmonary congestion in acute

Table 4
Comparison with previous meta-analyses.

Medication	Study	Medication use setting	Number of studies	OR/RR	Conclusion	
Metformin	Our current study	Preadmission	42	0.54 (0.47–0.62)	Decreased	
	Han et al. [5]	Preadmission + in-hospital	20	0.62 (0.50–0.76)	Decreased	
	Hariyanto et al. [74]	Preadmission	5	0.54 (0.32–0.90)	Decreased	
	Kan et al. [75]	Preadmission + in-hospital	15	0.69 (0.55–0.86)	Decreased	
	Kow et al. [76]	Preadmission	5	0.62 (0.43–0.89)	Decreased	
	Li et al. [77]	Preadmission + in-hospital	19	0.66 (0.56–0.78)	Decreased	
	Lukito et al. [78]	Preadmission	6	0.64 (0.43–0.97)	Decreased	
	Oscanoa et al. [79]	Preadmission + in-hospital	22	0.56 (0.45–0.68)	Decreased	
	Poly et al. [82]	Preadmission + in-hospital	16	0.66 (0.54–0.80)	Decreased	
	Scheen et al. [84]	Preadmission	4	0.75 (0.67–0.85)	Decreased	
	Schlesinger et al. [85]	ND	4	0.50 (0.28–0.90)	Decreased	
	Sun et al. [86]	Preadmission	7	0.54 (0.35–0.84)	Decreased	
	Yang et al. [87]	Preadmission + in-hospital	17	0.63 (0.51–0.79)	Decreased	
	Sulfonylurea	Our current study	Preadmission	21	0.92 (0.83–1.01)	NS
		Han et al. [5]	Preadmission + in-hospital	4	0.93 (0.89–0.98)	Decreased
Kan et al. [75]		Preadmission + in-hospital	5	0.80 (0.66–0.96)	Decreased	
Schlesinger et al. [85]		ND	2	0.73 (0.49–1.09)	NS	
Thiazolidinedione	Our current study	Preadmission	8	0.90 (0.71–1.14)	NS	
	No published meta-analysis					
Alpha-glucosidase inhibitor	Our current study	Preadmission	8	0.61 (0.26–1.45)	NS	
	No published meta-analysis					
Glucagon-like peptide-1 receptor agonist	Our current study	Preadmission	12	0.51 (0.37–0.69)	Decreased	
	Han et al. [5]	Preadmission + in-hospital	3	0.92 (0.80–1.04)	NS	
	Hariyanto et al. [7]	Preadmission	9	0.53 (0.43–0.66)	Decreased	
Dipeptidyl peptidase-4 inhibitor	Our current study	Preadmission	28	1.23 (1.07–1.42)	Increased	
	Bonora et al. [73]	Preadmission	7	0.81 (0.57–1.15)	NS	
	Han et al. [5]	Preadmission + in-hospital	11	0.95 (0.72–1.26)	NS	
	Hariyanto et al. [6]	Preadmission	7	1.14 (0.87–1.51)	NS	
	Kan et al. [75]	Preadmission + in-hospital	8	0.72 (0.51–1.51)	NS	
	Pal et al. [80]	Preadmission	4	1.21 (0.72–2.03)	NS	
	Patoulas et al. [81]	Preadmission	8	1.14 (0.78–1.66)	NS	
	Rakhmat et al. [83]	Preadmission + in-hospital	9	0.76 (0.60–0.97)	Decreased	
	Schlesinger et al. [85]	ND	2	0.90 (0.59–1.36)	NS	
	Yang et al. [4]	Preadmission + in-hospital	4	0.58 (0.34–0.99)	Decreased	
Sodium–glucose transporter-2 inhibitor	Our current study	Preadmission	13	0.60 (0.40–0.88)	Decreased	
	Han et al. [5]	Preadmission + in-hospital	3	1.04 (0.56–1.92)	NS	
	Insulin	Our current study	Preadmission	33	1.70 (1.33–2.19)	Increased
Kan et al. [75]		Preadmission + in-hospital	7	2.20 (1.34–3.60)	Increased	
Schlesinger et al. [85]		ND	5	1.75 (1.01–3.03)	Increased	
Yang et al. [88]		Preadmission + in-hospital	12	2.10 (1.51–2.93)	Increased	

Abbreviations: ND, not defined; NS, not significant.

lung disease [97]. Moreover, evidence has shown that a TZD could downregulate A Disintegrin and Metalloproteinase-17 (ADAM-17), an ACE2 cleaving enzyme in human skeletal muscles [98]. This event, in turn, increased membrane ACE2 and facilitated cellular viral entry, raising concerns about increased susceptibility to SARS-CoV-2 infection. These hypotheses partially explained why TZD did not improve the mortality outcomes of patients with COVID-19 with diabetes in our study.

4.2. Strengths and limitations

Our study has several strengths. Despite the high heterogeneity related to some analyses, the robustness of our findings was confirmed through meta-regression, subgroup analysis, and sensitivity analysis. First, after outliers were identified and removed, the heterogeneity of all remaining studies drastically decreased without a significant change in OR (all $p > 0.05$). Second, after the trim-and-fill method was performed, the OR did not significantly change (all $p > 0.05$), indicating that our pooled odds ratio still reflected the actual effect size. In other words, our results were reliable and stable, even in the presence of high heterogeneity. Third, we only included preadmission-usage studies instead of combining both preadmission and in-hospital use like some meta-analyses, leading to a more consistent data interpretation. Moreover, unlike some publications, we updated the most recent and completed data instead of using ongoing data or pooling two studies from the same cohort. Next, we recruited relatively diverse samples from multicenter and multinational cohorts, thus increasing the ability to generalize to a

larger population. Finally, we could present a dose-response meta-analysis to predict the effect of daily metformin doses on COVID-19 mortality.

Our study nevertheless has some limitations. First, we could include only observational studies because no randomized controlled trial was conducted on the topic of interest at the time of analysis. Any conclusions, therefore, should be cautiously drawn (considering indication bias). However, we recruited the largest number of participants from various papers of acceptable quality, making our systematic review and meta-analysis have high internal validity. Second, due to the observational nature of the studies, the multidrug issue could not be excluded. An investigation of specific combination therapies was necessary because a large proportion of diabetic patients need two or more glucose-lowering agents (either oral or injectable medications) to achieve glycemic targets. However, it was not feasible to perform such analysis due to limited raw data from original studies, even after we contacted the authors, because of the complexity of current diabetes treatment algorithms that would require additional mining of the original sources. Third, we were also unable to exclude the possibility of using medications beyond the hospital admission. However, our findings still reflected effects received before admission rather than short-term in-hospital effects because several included studies predefined medication users as those who had received a prescription that lasted at least 90–180 days, which is considered enough to exert their long-term effects. Fourth, because the COVID-19 treatment protocol has not been published as an international consensus among medical centers and countries, we lacked standardized severity assessment and concomitant

drugs used during hospitalization, both of which are especially critical for mortality modeling. Fifth, it is impossible to completely rule out unmeasured confounders, such as smoking or socioeconomic status, although the original studies tried to adjust for these factors to a certain extent. Therefore, further studies with a strictly controlled design are warranted to confirm the relationships between therapies and mortality among patients with COVID-19 having type 2 diabetes. Last, because of the high publication rate regarding the COVID-19 topic within the past three years, there is a possibility that some studies might have been missed and therefore were not included in our current review. Although it is unavoidable, we minimized that issue by assigning three researchers to systematically search and select studies and another reviewer to be consulted to reach a final decision if needed.

5. Conclusions

The preadmission prescription of glucose-lowering therapies was associated with different outcomes in patients with COVID-19 having type 2 diabetes. Specifically, metformin, GLP-1RA, and SGLT-2i were more likely to be beneficial regarding in-hospital death. By contrast, DPP-4i and insulin were associated with increased mortality. However, SU, TZD, and AGI were mortality neutral.

Abbreviations

ACE-2	Angiotensin-converting enzyme-2
ADAM-17	A Disintegrin And Metalloproteinase-17
AGI	Alpha-glucosidase inhibitor
CKD	Chronic kidney disease
COVID-19	Coronavirus disease of 2019
DPP-4i	Dipeptidyl peptidase-4 inhibitor
GLP-1RA	Glucagon-like peptide-1 receptor agonist
NF- κ B	Nuclear factor kappa-light-chain-enhancer of activated B
SARS-CoV-2	Severe Acute Respiratory Syndrome Coronavirus 2
SGLT-2i	Sodium-glucose transporter-2 inhibitor
SU	Sulfonylurea
TZD	Thiazolidinedione

CRedit authorship contribution statement

NN conceived of the original idea, performed meta-analyses, meta-regression, sensitivity analyses, interpreted data, and wrote the first manuscript. DSH, HSN, and DKNH performed the systematic search, study selection, risk of bias assessment, and data extraction. HYC and YCC verified the analytical methods, supervised the findings of this work, and contributed to the revisions of the final manuscript. HYL and CYL provided clinical advice on the interpretation of the data and contributed to the revisions of the final manuscript. All authors approved the final manuscript as submitted and have agreed to be accountable for all aspects of the work. YCC is the guarantor of this work.

Declaration of competing interest

The authors have no conflicts of interest relevant to this article to disclose. All authors declare that there are no relationships or activities that might bias, or be perceived to bias, their work.

Acknowledgement

This manuscript was edited by Wallace Academic Editing. We gratefully acknowledge Ngan Khanh Nguyen for her expertise and assistance in designing the graphical abstract.

Data availability

Data were extracted from published research papers, all of which are available and accessible. All datasets generated during the current study are available upon reasonable request from the corresponding authors. The study protocol has been published (PROSPERO ID: CRD42021293064; www.crd.york.ac.uk/PROSPERO/) and is unrestrictedly available.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.metabol.2022.155196>.

References

- [1] WHO n.d. Coronavirus disease (COVID-19) pandemic. Updated November 23. Accessed November 30, 2021. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>.
- [2] Hu J, Wang Y. The clinical characteristics and risk factors of severe COVID-19. *Gerontology* 2021;67(3):255–66. <https://doi.org/10.1159/000513400>.
- [3] Lim S, Bae JH, Kwon HS, Nauck MA. COVID-19 and diabetes mellitus: from pathophysiology to clinical management. *Nat Rev Endocrinol* Jan 2021;17(1):11–30. <https://doi.org/10.1038/s41574-020-00435-4>.
- [4] Yang Y, Cai Z, Zhang J. DPP-4 inhibitors may improve the mortality of coronavirus disease 2019: a meta-analysis. *PLoS One* 2021;16(5):e0251916. <https://doi.org/10.1371/journal.pone.0251916>.
- [5] Han T, Ma S, Sun C, et al. The association between anti-diabetic agents and clinical outcomes of COVID-19 in patients with diabetes: a systematic review and meta-analysis. *Arch Med Res* 2021. <https://doi.org/10.1016/j.arcmed.2021.08.002>. 2021/12/01/.
- [6] Hariyanto TI, Kurniawan A. Dipeptidyl peptidase 4 (DPP4) inhibitor and outcome from coronavirus disease 2019 (COVID-19) in diabetic patients: a systematic review, meta-analysis, and meta-regression. *J Diabetes Metab Disord* 2021;20(1):1–8. <https://doi.org/10.1007/s40200-021-00777-4>.
- [7] Hariyanto TI, Intan D, Hananto JE, Putri C, Kurniawan A. Pre-admission glucagon-like peptide-1 receptor agonist (GLP-1RA) and mortality from coronavirus disease 2019 (Covid-19): a systematic review, meta-analysis, and meta-regression. *Diabetes Res Clin Pract* Sep 2021;179:109031. <https://doi.org/10.1016/j.diabres.2021.109031>.
- [8] Wells G, Shea B, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp. [Accessed 30 November 2021].
- [9] Wg C. The combination of estimates from different experiments. *Biometrics* 1954;10:101–29.
- [10] Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21(11):1539–58. <https://doi.org/10.1002/sim.1186>.
- [11] Sterne JA, Gavaghan D, Egger M. Publication and related bias in meta-analysis: power of statistical tests and prevalence in the literature. *J Clin Epidemiol* Nov 2000;53(11):1119–29. [https://doi.org/10.1016/s0895-4356\(00\)00242-0](https://doi.org/10.1016/s0895-4356(00)00242-0).
- [12] Cheng X, Liu YM, Li H, et al. Metformin is associated with higher incidence of acidosis, but not mortality, in individuals with COVID-19 and pre-existing type 2 diabetes. *Cell Metab* 2020;32(4):537–547.e3. <https://doi.org/10.1016/j.cmet.2020.08.013>.
- [13] Pazoki M, Chichagi F, Hadadi A, et al. Association of clinical characteristics, antidiabetic and cardiovascular agents with diabetes mellitus and COVID-19: a 7-month follow-up cohort study. *J Diabetes Metab Disord* Nov 8 2021:1–11. <https://doi.org/10.1007/s40200-021-00901-4>.
- [14] Agarwal S, Schechter C, Southern W, Crandall JP, Tomer Y. Preadmission diabetes-specific risk factors for mortality in hospitalized patients with diabetes and coronavirus disease 2019. *Diabetes Care* Oct 2020;43(10):2339–44. <https://doi.org/10.2337/dc20-1543>.
- [15] An C, Lim H, Kim DW, Chang JH, Choi YJ, Kim SW. Machine learning prediction for mortality of patients diagnosed with COVID-19: a nationwide Korean cohort study. *Sci Rep* Oct 30 2020;10(1):18716. <https://doi.org/10.1038/s41598-020-75767-2>.
- [16] Blden K, Tantry U, Usman A, et al. Abstract 12228: metformin use in patients hospitalized with COVID-19: lower inflammation, oxidative stress, and thrombotic risk markers and better clinical outcomes. *Circulation* 2021;144(Suppl.1). https://doi.org/10.1161/circ.144.suppl_1.12228. A12228-A12228.
- [17] Boye KS, Tokar Erdemir E, Zimmerman N, et al. Risk factors associated with COVID-19 hospitalization and mortality: a large claims-based analysis among people with type 2 diabetes mellitus in the United States. *Diabetes Ther* Aug 2021;12(8):2223–39. <https://doi.org/10.1007/s13300-021-01110-1>.

- [18] Bramante CT, Ingraham NE, Murray TA, et al. Metformin and risk of mortality in patients hospitalised with COVID-19: a retrospective cohort analysis. *Lancet Health Longev*. Jan 2021;2(1):e34–41. [https://doi.org/10.1016/s2666-7568\(20\)30033-7](https://doi.org/10.1016/s2666-7568(20)30033-7).
- [19] Cernigliaro A, Allotta AV, Scondotto S. Diabete e trattamento farmacologico con ipoglicemizzanti possono essere considerati fattori di rischio per gli esiti di salute in soggetti COVID-19? I risultati di uno studio nella popolazione residente in Sicilia [Can diabetes and its related hypoglycemic drug treatment be considered risk factors for health outcomes in COVID-19 patients? The results of a study in the population residing in Sicily Region (Southern Italy)]. *Epidemiol Prev Sep-Dec 2020*;44(5-6 Suppl 2):315–22. <https://doi.org/10.19191/ep20.5-6.S2.132>.
- [20] Chen Y, Yang D, Cheng B, et al. Clinical characteristics and outcomes of patients with diabetes and COVID-19 in association with glucose-lowering medication. *Diabetes Care Jul 2020*;43(7):1399–407. <https://doi.org/10.2337/dc20-0660>.
- [21] Cheng X, Xin S, Chen Y, et al. Effects of metformin, insulin on COVID-19 patients with pre-existed type 2 diabetes: A multicenter retrospective study. *Life Sci Jun 15 2021*;275:119371. <https://doi.org/10.1016/j.lfs.2021.119371>.
- [22] Crouse AB, Grimes T, Li P, Might M, Ovalle F, Shalev A. Metformin use is associated with reduced mortality in a diverse population with COVID-19 and diabetes. *Front Endocrinol (Lausanne) 2020*;11:600439. <https://doi.org/10.3389/fendo.2020.600439>.
- [23] Dave JA, Tamuhla T, Tiffin N, et al. Risk factors for COVID-19 hospitalisation and death in people living with diabetes: a virtual cohort study from the Western Cape Province South Africa. *Diabetes Res Clin Pract Jul 2021*;177:108925. <https://doi.org/10.1016/j.diabres.2021.108925>.
- [24] Deng YP, Xie W, Liu T, et al. Association of diabetes with severity and mortality in hospitalized patients with COVID-19 in Wuhan, China: a single-centered, retrospective study. *Arch Endocrinol Metab 2021*;65(5):596–608. <https://doi.org/10.20945/2359-3997000000384>.
- [25] Do JY, Kim SW, Park JW, Cho KH, Kang SH. Is there an association between metformin use and clinical outcomes in diabetes patients with COVID-19? *Diabetes Metab Jul 2021*;47(4):101208. <https://doi.org/10.1016/j.diabet.2020.10.006>.
- [26] Elibol A, Eren D, Erdoğan MD, et al. Factors influencing on development of COVID-19 pneumonia and association with oral anti-diabetic drugs in hospitalized patients with diabetes mellitus. *Prim Care Diabetes Oct 2021*;15(5):806–12. <https://doi.org/10.1016/j.pcd.2021.08.001>.
- [27] Emral R, Haymana C, Demirci I, et al. Lower COVID-19 mortality in patients with type 2 diabetes mellitus taking dipeptidyl peptidase-4 inhibitors: results from a Turkish nationwide study. *Diabetes Ther Nov 2021*;12(11):2857–70. <https://doi.org/10.1007/s13300-021-01133-8>.
- [28] Fadini GP, Morieri ML, Longato E, et al. Exposure to dipeptidyl-peptidase-4 inhibitors and COVID-19 among people with type 2 diabetes: a case-control study. *Diabetes Obes Metab*. Oct 2020;22(10):1946–50. <https://doi.org/10.1111/dom.14097>.
- [29] Ghany R, Palacio A, Dawkins E, et al. Metformin is associated with lower hospitalizations, mortality and severe coronavirus infection among elderly medicare minority patients in 8 states in USA. *Diabetes Metab Syndr 2021*;15(2):513–8. <https://doi.org/10.1016/j.dsx.2021.02.022>. Mar-Apr.
- [30] Giorda CB, Picariello R, Tartaglino B, et al. From swab testing to health outcomes within the T2DM population: impact of diabetes background on COVID19 progression. *Diabetes Res Clin Pract Oct 2021*;180:109021. <https://doi.org/10.1016/j.diabres.2021.109021>.
- [31] Goodall JW, TAN Reed, Ardissino M, et al. Risk factors for severe disease in patients admitted with COVID-19 to a hospital in London, England: a retrospective cohort study. *Epidemiol Infect Oct 13 2020*;148:e251. <https://doi.org/10.1017/s0950268820002472>.
- [32] Israelsen SB, Pottegård A, Sandholdt H, Madsbad S, Thomsen RW, Benfield T. Comparable COVID-19 outcomes with current use of GLP-1 receptor agonists, DPP-4 inhibitors or SGLT-2 inhibitors among patients with diabetes who tested positive for SARS-CoV-2. *Diabetes Obes Metab Jun 2021*;23(6):1397–401. <https://doi.org/10.1111/dom.14329>.
- [33] Kahkoska AR, Abrahamson TJ, Alexander GC, et al. Association between glucagon-like peptide 1 receptor agonist and sodium-glucose cotransporter 2 inhibitor use and COVID-19 outcomes. *Diabetes Care Jul 2021*;44(7):1564–72. <https://doi.org/10.2337/dc21-0065>.
- [34] Khunti K, Knighton P, Zaccardi F, et al. Prescription of glucose-lowering therapies and risk of COVID-19 mortality in people with type 2 diabetes: a nationwide observational study in England. *Lancet Diabetes Endocrinol May 2021*;9(5):293–303. [https://doi.org/10.1016/s2213-8587\(21\)00050-4](https://doi.org/10.1016/s2213-8587(21)00050-4).
- [35] Kim MK, Jeon JH, Kim SW, et al. The clinical characteristics and outcomes of patients with moderate-to-severe coronavirus disease 2019 infection and diabetes in Daegu, South Korea. *Diabetes Metab J Aug 2020*;44(4):602–13. <https://doi.org/10.4093/dmj.2020.0146>.
- [36] Kristan MM, Kim YK, Nelson T, et al. Predictors of severe COVID-19 in patients with diabetes: a multicenter review. *Endocr Pract Aug 2021*;27(8):842–9. <https://doi.org/10.1016/j.eprac.2021.05.011>.
- [37] Lally MA, Tsoukas P, Halladay CW, O'Neill E, Gravenstein S, Rudolph JL. Metformin is associated with decreased 30-day mortality among nursing home residents infected with SARS-CoV2. *J Am Med Dir Assoc Jan 2021*;22(1):193–8. <https://doi.org/10.1016/j.jamda.2020.10.031>.
- [38] Li F, Cai Y, Gao C, et al. Effects of diabetes and blood glucose on COVID-19 mortality: a retrospective observational study. *medRxiv 2021*:2021.01.21.20202119. <https://doi.org/10.1101/2021.01.21.20202119>.
- [39] Li J, Wei Q, Li WX, et al. Metformin use in diabetes prior to hospitalization: effects on mortality in Covid-19. *Endocr Pract Oct 2020*;26(10):1166–72. <https://doi.org/10.4158/ep-2020-0466>.
- [40] AYO Luk, TCF Yip, Zhang X, et al. Glucose-lowering drugs and outcome from COVID-19 among patients with type 2 diabetes mellitus: a population-wide analysis in Hong Kong. *BMJ Open Oct 2021*;11(10):e052310. <https://doi.org/10.1136/bmjopen-2021-052310>.
- [41] Luo P, Qiu L, Liu Y, et al. Metformin treatment was associated with decreased mortality in COVID-19 patients with diabetes in a retrospective analysis. *Am J Trop Med Hyg Jul 2020*;103(1):69–72. <https://doi.org/10.4269/ajtmh.20-0375>.
- [42] Luo SK, Hu WH, Lu ZJ, et al. Diabetes patients with comorbidities had unfavorable outcomes following COVID-19: a retrospective study. *World J Diabetes 2021*;12(10):1789–808. <https://doi.org/10.4239/wjdv12.i10.1789>.
- [43] Ma Z, Patel N, Vemparala P, Krishnamurthy M. Metformin is associated with favorable outcomes in patients with COVID-19 and Type 2 diabetes mellitus. *medRxiv 2021*:2021.05.20.21257490. <https://doi.org/10.1101/2021.05.20.21257490>.
- [44] Mansour A, Sajjadi-Jazi SM, Kasaeian A, et al. Clinical characteristics and outcomes of diabetics hospitalized for COVID-19 infection: a single-centered, retrospective, observational study. *EXCLI J 2020*;19:1533–43. <https://doi.org/10.17179/excli2020-2988>.
- [45] Meijer RI, Hoekstra T, van den Oever NCG. Treatment with a DPP-4 inhibitor at time of hospital admission for COVID-19 is not associated with improved clinical outcomes: data from the COVID-PREDICT cohort study in The Netherlands. *J Diabetes Metab Disord Jun 26 2021*:1–6. <https://doi.org/10.1007/s40200-021-00833-z>.
- [46] Mirani M, Favacchio G, Carrone F, et al. Impact of comorbidities and glycemia at admission and dipeptidyl peptidase 4 inhibitors in patients with Type 2 diabetes with COVID-19: a case series from an academic hospital in Lombardy, Italy. *Diabetes Care*. Dec 2020;43(12):3042–9. <https://doi.org/10.2337/dc20-1340>.
- [47] Mirsoleymani S, Nekooghadam SM, Ahmadi Marzaleh M, et al. Assessment of risk factors for severe coronavirus disease 2019 among Iranian patients. *Iranian Red Crescent Medical Journal 2020*;22(9). <https://doi.org/10.32592/ircmj.2020.22.9.72>. 09/30.
- [48] Nafakhi H, Alareedh M, Al-Buthabak K, Shaghee F, Nafakhi A, Kasim S. Predictors of adverse in-hospital outcome and recovery in patients with diabetes mellitus and COVID-19 pneumonia in Iraq. *Diabetes Metab Syndr 2021*;15(1):33–8. <https://doi.org/10.1016/j.dsx.2020.12.014>. Jan-Feb.
- [49] Noh Y, Oh IS, Jeong HE, Filion KB, Yu OHY, Shin JY. Association between DPP-4 inhibitors and COVID-19-related outcomes among patients with type 2 diabetes. *Diabetes Care Apr 2021*;44(4):e64–6. <https://doi.org/10.2337/dc20-1824>.
- [50] Nyland JE, Raja-Khan NT, Bettermann K, et al. Diabetes, drug treatment and mortality in COVID-19: a multinational retrospective cohort study. *Diabetes Sep 27 2021*. <https://doi.org/10.2337/db21-0385>.
- [51] Oh TK, Song IA. Metformin use and risk of COVID-19 among patients with type II diabetes mellitus: an NHIS-COVID-19 database cohort study. *Acta Diabetol Jun 2021*;58(6):771–8. <https://doi.org/10.1007/s00592-020-01666-7>.
- [52] Ong AN, Tan CC, Cañete MT, Lim BA, Robles J. Association between metformin use and mortality among patients with type 2 diabetes mellitus hospitalized for COVID-19 infection. *Journal of the ASEAN Federation of Endocrine Societies 2021*;36(2):133–41. <https://doi.org/10.15605/jafes.036.02.20>.
- [53] Orioli L, Servais T, Belkhir L, et al. Clinical characteristics and short-term prognosis of in-patients with diabetes and COVID-19: a retrospective study from an academic center in Belgium. *Diabetes Metab Syndr 2021*;15(1):149–57. <https://doi.org/10.1016/j.dsx.2020.12.020>. Jan-Feb.
- [54] Pérez-Belmonte LM, Torres-Peña JD, López-Carmona MD, et al. Mortality and other adverse outcomes in patients with type 2 diabetes mellitus admitted for COVID-19 in association with glucose-lowering drugs: a nationwide cohort study. *BMC Med Nov 16 2020*;18(1):359. <https://doi.org/10.1186/s12916-020-01832-2>.
- [55] Philipose S, Smati N, Jefferson Wong CS, Aspey K, Mendall M. Obesity, old age, and frailty are the true risk factors for COVID-19 mortality and not chronic disease or ethnicity. *medRxiv 2020*:2020.08.12.20156257. <https://doi.org/10.1101/2020.08.12.20156257>.
- [56] Ramos-Rincón JM, Pérez-Belmonte LM, Carrasco-Sánchez FJ, et al. Cardiometabolic therapy and mortality in very old patients with diabetes hospitalized due to COVID-19. *J Gerontol A Biol Sci Med Sci 2021*;76(8):e102–9. <https://doi.org/10.1093/geron/glab124>.
- [57] Ravindra G, Chitra L, Madhur M, et al. Retrospective assessment of treatments of hospitalized Covid-19 patients. *medRxiv 2021*:2021.04.20.21255792. <https://doi.org/10.1101/2021.04.20.21255792>.
- [58] Riahi S, Sombra LRS, Lo KB, et al. Insulin use, diabetes control, and outcomes in patients with COVID-19. *Endocr Res 2021*;46(2):45–50. <https://doi.org/10.1080/07435800.2020.1856865>. Feb-May.
- [59] Satman I, Demirci I, Haymana C, et al. Unexpectedly lower mortality rates in COVID-19 patients with and without type 2 diabetes in Istanbul. *Diabetes Res Clin Pract Apr 2021*;174:108753. <https://doi.org/10.1016/j.diabres.2021.108753>.
- [60] Saygili ES, Karakiliç E, Mert E, Şener A, Mirici A. Preadmission usage of metformin and mortality in COVID-19 patients including the post-discharge period. *Ir J Med Sci Oct 29 2021*:1–7. <https://doi.org/10.1007/s11845-021-02823-9>.
- [61] Shestakova MV, Vikulova OK, Isakov M, Dedov II. Diabetes and COVID-19: analysis of the clinical outcomes according to the data of the Russian diabetes registry. *Probl Endokrinol (Mosk) 2020*;66(1):35–46. <https://doi.org/10.14341/probl12458>.
- [62] Silverii GA, Monami M, Cernigliaro A, et al. Are diabetes and its medications risk factors for the development of COVID-19? Data from a population-based study in Sicily. *Nutr Metab Cardiovasc Dis 2021*;31(2):396–8. <https://doi.org/10.1016/j.numecd.2020.09.028>.

- [63] Sourij H, Aziz F, Bräuer A, et al. COVID-19 fatality prediction in people with diabetes and prediabetes using a simple score upon hospital admission. *Diabetes Obes Metab*. Feb 2021;23(2):589–98. <https://doi.org/10.1111/dom.14256>.
- [64] Strollo R, Maddaloni E, Dauriz M, Pedone C, Buzzetti R, Pozzilli P. Use of DPP4 inhibitors in Italy does not correlate with diabetes prevalence among COVID-19 deaths. *Diabetes Res Clin Pract* Jan 2021;171:108444. <https://doi.org/10.1016/j.diabres.2020.108444>.
- [65] Tamura RE, Said SM, de Freitas LM, IGS Rubio. Outcome and death risk of diabetes patients with Covid-19 receiving pre-hospital and in-hospital metformin therapies. *Diabetol Metab Syndr* Jul 13 2021;13(1):76. <https://doi.org/10.1186/s13098-021-00695-8>.
- [66] Wander PL, Lowy E, Beste LA, et al. Prior glucose-lowering medication use and 30-day outcomes among 64,892 veterans with diabetes and COVID-19. *Diabetes Care* Oct 6 2021. <https://doi.org/10.2337/dc21-1351>.
- [67] Wang B, Van Oekelen O, Mouhieddine TH, et al. A tertiary center experience of multiple myeloma patients with COVID-19: lessons learned and the path forward. *J Hematol Oncol* Jul 14 2020;13(1):94. <https://doi.org/10.1186/s13045-020-00934-x>.
- [68] Wang J, Cooper JM, Gokhale K, et al. Association of Metformin with susceptibility to COVID-19 in people with type 2 diabetes. *J Clin Endocrinol Metab* 2021;106(5):1255–68. <https://doi.org/10.1210/clinem/dgab067>.
- [69] Wargny M, Potier L, Gourdy P, et al. Predictors of hospital discharge and mortality in patients with diabetes and COVID-19: updated results from the nationwide CORONADO study. *Diabetologia* Apr 2021;64(4):778–94. <https://doi.org/10.1007/s00125-020-05351-w>.
- [70] CKH Wong, DTW Lui, AYC Lui, et al. Use of DPP4i reduced odds of clinical deterioration and hyperinflammatory syndrome in COVID-19 patients with type 2 diabetes: propensity score analysis of a territory-wide cohort in Hong Kong. *Diabetes & Metabolism* 2021;101307. <https://doi.org/10.1016/j.diabet.2021.101307>. 2021/12/01/.
- [71] Yan Y, Yang Y, Wang F, et al. Clinical characteristics and outcomes of patients with severe covid-19 with diabetes. *BMJ Open Diabetes Res Care* Apr 2020;8(1). <https://doi.org/10.1136/bmjdr-2020-001343>.
- [72] Yuan S, Li H, Chen C, Wang F, Wang DW. Association of glycosylated haemoglobin HbA1c levels with outcome in patients with COVID-19: a retrospective study. *J Cell Mol Med* Apr 2021;25(7):3484–97. <https://doi.org/10.1111/jcmm.16431>.
- [73] Bonora BM, Avogaro A, Fadini GP. Disentangling conflicting evidence on DPP-4 inhibitors and outcomes of COVID-19: narrative review and meta-analysis. *J Endocrinol Invest* Jul 2021;44(7):1379–86. <https://doi.org/10.1007/s40618-021-01515-6>.
- [74] Hariyanto TI, Kurniawan A. Metformin use is associated with reduced mortality rate from coronavirus disease 2019 (COVID-19) infection. *Obes Med* Sep 2020;19:100290. <https://doi.org/10.1016/j.obmed.2020.100290>.
- [75] Kan C, Zhang Y, Han F, et al. Mortality risk of antidiabetic agents for type 2 diabetes with COVID-19: a systematic review and meta-analysis. *Front Endocrinol (Lausanne)* 2021;12:708494. <https://doi.org/10.3389/fendo.2021.708494>.
- [76] Kow CS, Hasan SS. Mortality risk with preadmission metformin use in patients with COVID-19 and diabetes: a meta-analysis. *J Med Virol* Feb 2021;93(2):695–7. <https://doi.org/10.1002/jmv.26498>.
- [77] Li Y, Yang X, Yan P, Sun T, Zeng Z, Li S. Metformin in patients with COVID-19: a systematic review and meta-analysis. *Front Med (Lausanne)* 2021;8:704666. <https://doi.org/10.3389/fmed.2021.704666>.
- [78] Lukito AA, Pranata R, Henrina J, Lim MA, Lawrensia S, Suastika K. The effect of metformin consumption on mortality in hospitalized COVID-19 patients: a systematic review and meta-analysis. *Diabetes Metab Syndr* 2020;14(6):2177–83. <https://doi.org/10.1016/j.dsx.2020.11.006>. Nov-Dec.
- [79] Oscanoa TJ, Amado J, Vidal X, Savarino A, Romero-Ortuno R. Metformin therapy and severity and mortality of SARS-CoV-2 infection: a meta-analysis. *Clin Diabetol* 2021;10(4):317–29. <https://doi.org/10.5603/DK.a2021.0035>.
- [80] Pal R, Banerjee M, Mukherjee S, Bhogal RS, Kaur A, Bhadada SK. Dipeptidyl peptidase-4 inhibitor use and mortality in COVID-19 patients with diabetes mellitus: an updated systematic review and meta-analysis. *Ther Adv Endocrinol Metab* 2021;12:2042018821996482. <https://doi.org/10.1177/2042018821996482>.
- [81] Patoulis D, Doumas M. Dipeptidyl Peptidase-4 inhibitors and COVID-19-related deaths among patients with type 2 diabetes mellitus: a meta-analysis of observational studies. *Endocrinol Metab (Seoul)* Aug 2021;36(4):904–8. <https://doi.org/10.3803/EnM.2021.1048>.
- [82] Poly TN, Islam MM, Li YJ, Lin MC, Hsu MH, Wang YC. Metformin use is associated with decreased mortality in COVID-19 patients with diabetes: evidence from retrospective studies and biological mechanism. *J Clin Med* Aug 9 2021;10(16). <https://doi.org/10.3390/jcm10163507>.
- [83] Rakhmat II, Kusmala YY, Handayani DR, et al. Dipeptidyl peptidase-4 (DPP-4) inhibitor and mortality in coronavirus disease 2019 (COVID-19) - a systematic review, meta-analysis, and meta-regression. *Diabetes Metab Syndr* 2021;15(3):777–82. <https://doi.org/10.1016/j.dsx.2021.03.027>. May-Jun.
- [84] Scheen AJ. Metformin and COVID-19: from cellular mechanisms to reduced mortality. *Diabetes Metab Nov* 2020;46(6):423–6. <https://doi.org/10.1016/j.diabet.2020.07.006>.
- [85] Schlesinger S, Neuwenschwander M, Lang A, et al. Risk phenotypes of diabetes and association with COVID-19 severity and death: a living systematic review and meta-analysis. *Diabetologia* Jul 2021;64(7):1480–91. <https://doi.org/10.1007/s00125-021-05458-8>.
- [86] Sun C, Cheng C, Kim KY, Ahmed MA, Manem R. Is metformin use associated with a decreased mortality for COVID-19 diabetic patients? A meta-analysis. *Journal of the Endocrine Society* 2021;5(Supplement 1). <https://doi.org/10.1210/jendso/bvab048.709>. A348-A348.
- [87] Yang W, Sun X, Zhang J, Zhang K. The effect of metformin on mortality and severity in COVID-19 patients with diabetes mellitus. *Diabetes Res Clin Pract* Aug 2021;178:108977. <https://doi.org/10.1016/j.diabres.2021.108977>.
- [88] Yang Y, Cai Z, Zhang J. Insulin treatment may increase adverse outcomes in patients with COVID-19 and diabetes: a systematic review and meta-analysis. *Front Endocrinol (Lausanne)* 2021;12:696087. <https://doi.org/10.3389/fendo.2021.696087>.
- [89] Sharma S, Ray A, Sadasivam B. Metformin in COVID-19: a possible role beyond diabetes. *Diabetes Res Clin Pract* Jun 2020;164:108183. <https://doi.org/10.1016/j.diabres.2020.108183>.
- [90] Zhou G, Myers R, Li Y, et al. Role of AMP-activated protein kinase in mechanism of metformin action. *J Clin Invest* Oct 2001;108(8):1167–74. <https://doi.org/10.1172/jci13505>.
- [91] Varghese E, Samuel SM, Liskova A, Kubatka P, Büsselberg D. Diabetes and coronavirus (SARS-CoV-2): molecular mechanism of metformin intervention and the scientific basis of drug repurposing. *PLoS Pathog* Jun 2021;17(6):e1009634. <https://doi.org/10.1371/journal.ppat.1009634>.
- [92] Drucker DJ. Coronavirus infections and Type 2 diabetes-shared pathways with therapeutic implications. *Endocr Rev* Jun 1 2020;41(3). <https://doi.org/10.1210/andrev/bnaa011>.
- [93] Lee YS, Jun HS. Anti-inflammatory effects of GLP-1-based therapies beyond glucose control. *Mediat Inflamm* 2016;2016:3094642. <https://doi.org/10.1155/2016/3094642>.
- [94] Cure E, Cumhur Cure M. Can dapagliflozin have a protective effect against COVID-19 infection? A hypothesis. *Diabetes Metab Syndr* 2020;14(4):405–6. <https://doi.org/10.1016/j.dsx.2020.04.024>. Jul-Aug.
- [95] Vankadari N, Wilce JA. Emerging Wuhan (COVID-19) coronavirus: glycan shield and structure prediction of spike glycoprotein and its interaction with human CD26. *Emerg Microbes Infect* 2020;9(1):601–4. <https://doi.org/10.1080/22221751.2020.1739565>.
- [96] Juan CC, Fang VS, Kwok CF, Perng JC, Chou YC, Ho LT. Exogenous hyperinsulinemia causes insulin resistance, hyperendothelinemia, and subsequent hypertension in rats. *Metabolism* Apr 1999;48(4):465–71. [https://doi.org/10.1016/s0026-0495\(99\)90105-1](https://doi.org/10.1016/s0026-0495(99)90105-1).
- [97] Erol A. Role of oxidized LDL-induced "trained macrophages" in the pathogenesis of COVID-19 and benefits of pioglitazone: a hypothesis. *Diabetes Metab Syndr* 2020;14(4):713–4. <https://doi.org/10.1016/j.dsx.2020.05.007>. Jul-Aug.
- [98] Tripathy D, Daniele G, Fiorentino TV, et al. Pioglitazone improves glucose metabolism and modulates skeletal muscle TIMP-3-TACE dyad in type 2 diabetes mellitus: a randomised, double-blind, placebo-controlled, mechanistic study. *Diabetologia* Oct 2013;56(10):2153–63. <https://doi.org/10.1007/s00125-013-2976-z>.