


The impact of the *Hsp67Bc* gene product on *Drosophila melanogaster* longevity, fecundity, and acute heat stress tolerance

D. Malkeyeva , S.A. Fedorova, E. Kiseleva


Institute of Cytology and Genetics of the Siberian Branch of the Russian Academy of Sciences, Novosibirsk, Russia

 malkeyeva@bionet.nsc.ru


Abstract. *Drosophila melanogaster* Hsp67Bc is a heat- and cold-inducible small heat shock protein that participates in the prevention of aggregation of misfolded proteins and in macroautophagy regulation. Overexpression of the *Hsp67Bc* gene has been shown to enhance macroautophagy in *Drosophila* S2 cells, and the deletion of this gene leads to the formation of a slightly increased number of autophagic vacuoles in the fruit fly brain neurons. Recently, we found that *Hsp67Bc*-null *D. melanogaster* flies have poor tolerance to cold stress (0 °C) of various durations. In the present work, we investigated how the *Hsp67Bc* gene deletion affects the fitness of fruit flies under normal conditions and their tolerance to elevated temperatures at different developmental stages. Larvae and pupae were not adversely affected by the *Hsp67Bc* gene deletion, and adult *Hsp67Bc*-null flies showed an extended lifespan in comparison with the control at normal (24–25 °C) and elevated temperature (29 °C), and after acute heat stress (37 °C, 2 h). At the same time, the fecundity of the mutant females was lower by 6–13 % in all tested environments, except for permanent maintenance at 29 °C, where the mean numbers of eggs laid by the mutant and control flies were equal. We explain this phenomenon by a reduced number of ovarioles in *Hsp67Bc*-null females and enhanced macroautophagy in their germaria, which promotes the death of forming egg chambers. In addition, short heat stress (37 °C, 2 h), which increased the control line's longevity (an effect common for a wide range of organisms), had a negative impact on the lifespan of *Hsp67Bc*-null flies. Therefore, *Hsp67Bc*-null *D. melanogaster* have an extended lifespan under normal and elevated temperature conditions, and reduced fecundity and thermal stress tolerance. Key words: *Drosophila* longevity; thermal stress tolerance; elevated temperature; heat stress; small heat shock proteins; autophagy.

For citation: Malkeyeva D., Fedorova S.A., Kiseleva E. The impact of the *Hsp67Bc* gene product on *Drosophila melanogaster* longevity, fecundity, and acute heat stress tolerance. *Vavilovskii Zhurnal Genetiki i Selekcii* = *Vavilov Journal of Genetics and Breeding*. 2022;26(2):169-178. DOI 10.18699/VJGB-22-21

Влияние продукта гена *Hsp67Bc* на продолжительность жизни, плодовитость и устойчивость *Drosophila melanogaster* к кратковременному тепловому стрессу

Д.А. Малькеева , С.А. Фёдорова, Е.В. Киселева

Федеральный исследовательский центр Институт цитологии и генетики Сибирского отделения Российской академии наук, Новосибирск, Россия

 malkeyeva@bionet.nsc.ru

Аннотация. Hsp67Bc *Drosophila melanogaster* – индуцируемый в ответ на тепловой и холодовой стресс малый белок теплового шока, участвующий в предотвращении агрегации поврежденных белков и в регуляции макроаутофагии. Было показано, что повышенная экспрессия гена *Hsp67Bc* стимулирует макроаутофагию в клетках S2 дрозофилы, а его делеция приводит к небольшому увеличению количества аутофагических вакуолей в нейронах мозга мух. Недавно нами обнаружено, что нуль-аллельные по гену *Hsp67Bc* особи *D. melanogaster* имеют сниженную устойчивость к холодовому стрессу (0 °C) различной длительности. В настоящей работе мы исследовали, как наличие делеции в гене *Hsp67Bc* повлияет на жизнеспособность *D. melanogaster* в нормальных условиях и на их устойчивость к повышенной температуре на разных стадиях развития. Делеция *Hsp67Bc* не оказала на личинок и куколок дрозофил неблагоприятного воздействия; нуль-аллельные по гену *Hsp67Bc* имаго имели увеличенную по сравнению с контролем продолжительность жизни при нормальной (24–25 °C) и повышенной (29 °C) температуре, а также после кратковременного теплового стресса (37 °C, 2 ч). В то же время плодовитость мутантных самок была снижена на 6–13 % по сравнению с контролем при всех исследованных температурных режимах, за исключением постоянного содержания при 29 °C, при котором среднее число откладываемых яиц не различалось между контрольной и мутантной линиями. Мы связываем этот феномен со сниженным количеством овариол у нуль-аллельных по гену *Hsp67Bc* самок, а также с усиленной макроаутофагией в их гермариях, приводящей к росту числа гибнущих формирующихся яйцевых камер.

Кроме того, кратковременный тепловой стресс (37 °C, 2 ч), приводивший к увеличению продолжительности жизни *D. melanogaster* контрольной линии (что является распространенной реакцией у живых организмов), отрицательно влиял на продолжительность жизни мух с делецией *Hsp67Bc*. Таким образом, *D. melanogaster* с делецией в гене *Hsp67Bc* имеют увеличенную продолжительность жизни в нормальных условиях и при повышенной температуре и сниженные плодовитость и устойчивость к температурному стрессу.

Ключевые слова: продолжительность жизни *Drosophila*; устойчивость к температурному стрессу; повышенная температура; тепловой стресс; малые белки теплового шока; аутофагия.

Introduction

During ontogenesis, all living organisms experience stress. The effects of stress-inducing agents on cells include oxidative modification of proteins, which leads to their misfolding (Jolly, Morimoto, 2000). Misfolded proteins are detrimental to the cell because they may gain deleterious biological functions and are prone to forming insoluble aggregates (Jolly, Morimoto, 2000). To maintain homeostasis, cells synthesize heat shock proteins (HSPs): a group of conservative proteins that ensure correct folding of peptides, prevent aggregation of denatured proteins, and resolubilize protein aggregates (Jolly, Morimoto, 2000). This response is universal among all the known pro- and eukaryotes (Lindquist, 1986). Expression of the majority of *HSP* genes is up-regulated in stressful conditions such as heat and cold stress, hypoxia, bacterial and viral infections, and oxidative stress (Lindquist, 1986; Sørensen et al., 2003). Some HSPs are constitutively expressed and are necessary for growth and development of organisms under normal conditions (Kampinga et al., 2009; Sarkar et al., 2011). In a study by Raut et al. (2017) on *Drosophila*, knockdown of 42 out of 95 tested *HSP* genes led to F1 lethality indicating their crucial role in the fly development.

Drosophila melanogaster Hsp67Bc belongs to the small heat shock protein family of HSPs (Vos et al., 2016). It shares a function of preventing damaged protein aggregation with other members of the family (Vos et al., 2016). In addition, *Hsp67Bc* was shown to be involved in the regulation of macroautophagy – a conservative catabolic process allowing the recycling of cytoplasm components – alongside Starvin protein (Carra et al., 2010; Parzych, Klionsky, 2014). Overexpression of the *Hsp67Bc* gene separately or together with *stv* resulted in protein synthesis inhibition and macroautophagy stimulation (Carra et al., 2010).

Our studies on the *Hsp67Bc* gene deletion in *D. melanogaster* revealed that in brain neurons of *Hsp67Bc*-null flies infected by a pathogenic *Wolbachia* bacteria strain *wMelPop*, the number of autophagosomes and autolysosomes (organelles formed in the process of macroautophagy that sequester cytoplasm components and digest them) was increased, and the cross-sectional area of autolysosomes was more than 1.5-fold larger than in the control line with the wild-type *Hsp67Bc* gene (Malkeyeva et al., 2021). These observations may indicate that in the absence of the *Hsp67Bc* gene product, macroautophagy is slightly enhanced and autophagosome maturation process is affected. Furthermore, we showed that the *Hsp67Bc* gene product plays an important role in tolerance to cold stress in the fruit fly (Malkeyeva et al., 2020). *Hsp67Bc*-null adult flies needed more time to recover from chill coma than the control flies, and adult females with the *Hsp67Bc* gene deletion had a 1.6–3-fold lower survival after cold stress of various dura-

tions (2, 4, and 12 h at 0 °C) as compared to the control line (Malkeyeva et al., 2020).

In this study, we investigated fitness of *Hsp67Bc*-null *D. melanogaster* under normal conditions (24–25 °C) and their tolerance to elevated temperatures (29 °C or 2 h at 37 °C) at different stages of ontogenesis (larva, pupa, and imago). We found that the adult mutant flies had an increased lifespan at all the tested temperatures, in comparison with the control line with an intact *Hsp67Bc* gene. *Hsp67Bc*-null adult flies, however, had slightly reduced fecundity under normal conditions and after heat stress (37 °C, 2 h) and were negatively affected by acute heat stress (37 °C, 2 h) that prolonged longevity of the control line. Thus, despite having extended lifespan in comparison with the control line under all tested conditions, *Hsp67Bc*-null flies had lower fecundity and were less tolerant to acute heat stress.

Materials and methods

***Drosophila melanogaster* lines.** In this study, we used *Hsp67Bc*-null *D. melanogaster* line *Hsp67Bc*-0 we created by an imprecise excision of a *P*-element located in proximity to the *Hsp67Bc* gene transcription start. Fly line *Hsp67Bc*-2 containing a wild-type variant of *Hsp67Bc* obtained by a precise cutting out of the mentioned *P*-element was used as a control. The procedure for obtaining the fly lines is described in our recent article (Malkeyeva et al., 2020).

Heat stress applied to larvae and pupae. For these experiments, wandering late 3rd instar (L3) larvae were transferred from their rearing vials to the walls of vials with fresh cornmeal-agar medium, at 20 per vial. The larvae were then either directly transferred to a 37 °C environment (water bath in an incubator) for 2 h incubation or allowed to first reach the developmental stage that was to be treated. In particular, these were pupal stages P1–P2 (white prepupae, 1–2 h after pupation), P5 (18–20 h after pupation), or P7–P8 (46–48 h after pupation) (Bainbridge, Bownes, 1981). The cottons sealing the vials were slightly moisturized with water before the start of heat treatment to prevent drying of the larvae and pupae. After the heat stress treatment, the flies were kept at 24–25 °C until eclosion. Survivors to the pupa stage (in case of late L3 treatment) and to the adult stage (for all treatment groups) were then counted. In each experiment, 39–107 flies of each genotype were used.

Analysis of the lifespan and fecundity of adult *D. melanogaster* kept at either normal or elevated temperature. The flies were collected from rearing vials on the 1st day after eclosion and placed into vials with a fresh cornmeal-agar medium, at eight males and eight females per vial. The flies then underwent one of four treatments. The 1st group was kept at 24–25 °C (normal conditions); the 2nd group was subjected

to heat treatment at 37 °C for 2 h at 1 day of age and then was returned to the 24–25 °C environment; the 3rd group was heat treated (37 °C, 2 h) at 7 days of age, then returned to the 24–25 °C environment; the 4th group was transferred to a 29 °C environment at 1 day of age and kept at the elevated temperature. Each experimental group contained 45–62 males and 51–62 females of relevant genotypes.

All the flies were kept under the specified conditions until the death of all individuals, with survivors transferred to fresh food daily or every other day. In parallel, fecundity was measured in these *Drosophila* starting from day 2 in the 1st, 2nd, and 4th groups and starting from day 8 in the 3rd group (one day after the heat treatment). *D. melanogaster* females were allowed to lay eggs for 24 h in vials with fresh medium, then the parents were transferred to new food, and the eggs were counted. The number of eggs in a vial was then divided by the number of females that oviposited in that very vial. The egg per female ratio was evaluated on days 2–11, 15–17, and 22–24 in the 1st, 2nd, and 4th experimental groups; in the 3rd group, the ratio was measured on days 8–10, 13–15, and 20–22.

Protein starvation assay. For this assay, newly eclosed adult *D. melanogaster* individuals were collected every 2 h from their rearing vials and transferred to vials with protein-free medium containing 100 g/L sucrose, 5 g/L agar and 0.78 g/L methyl 4-hydroxybenzoate. The flies were transferred to a fresh medium every other day.

LysoTracker Red (LTR) staining. On the 5th and 15th days of the protein starvation experiment, ovaries of the starved adult *D. melanogaster* females and females kept on standard food were dissected in 0.01 M PBS (Medigen) (pH 7.4) and stained with 100 nM LysoTracker Red DND-99 (Life Technologies) and DAPI. The LTR staining was performed as follows: the dissected ovaries were first placed into a droplet of a 100 nM LTR solution in 0.01 M PBS for 10 min incubation, washed thrice in PBS, and then fixed in 4 % paraformaldehyde for 20 min; next, the ovaries were washed three times with a 0.1 % Triton X-100 solution in PBS and mounted on a slide with a drop of DAPI-containing *SlowFade* Gold Antifade Mountant (Thermo Fisher Scientific). To make sure the antifade mountant penetrates inner ovarioles, we let the ovaries stay without a cover slip for ~15 min before sealing them under it with nail polish. The samples were stored in the dark at 4 °C until analysis under the LSM 780 confocal microscope (Zeiss) with the Plan-Apochromat 20x/0.8 M27 objective.

Statistical analyses. Survival and recovery curves were compared by the log-rank test. The fecundity, lifespan, number of ovarioles, and number of dying egg chambers per ovariole datasets were tested for normality by the Shapiro–Wilk test; normally distributed data were compared by the heteroscedastic *t* test; data with non-normal distribution were compared by the Mann–Whitney *U* test. Differences in fecundity between the control and mutant fly lines throughout the experiment were evaluated at each point by the heteroscedastic *t* test, followed by the Benjamini–Krieger–Yekutieli method to control the false discovery rate. Analyses of the proportion (%) of LTR-positive germaria obtained in the LTR-staining experiments were performed by the chi-squared test. Differences were considered statistically significant at $p \leq 0.05$.

The Shapiro–Wilk test and the Mann–Whitney *U* test were conducted using Statistics Kingdom statistics calculators (<https://www.statskingdom.com>).

Results

Hsp67Bc-null *D. melanogaster* under normal conditions

To expand our knowledge on the functions of *Hsp67Bc* in the fruit fly we created a *D. melanogaster* line carrying a deletion of almost the entire *Hsp67Bc* gene (described in detail in our recent article (Malkeyeva et al., 2020)). The flies carrying the deletion in the *Hsp67Bc* gene in the homozygous state (*Hsp67Bc*-0) were viable and fertile, and had no visible morphological deviations from the control. The *Hsp67Bc*-0 line had extended longevity under normal conditions (24–25 °C) as compared to the control *Hsp67Bc*-2 line (Fig. 1, *a–c*). The mean lifespan of *Hsp67Bc*-null *D. melanogaster* significantly exceeded that of the control by 35 % in males and by 34 % in females at 24–25 °C (see Fig. 1, *c*). Thus, it was 70.9 ± 1.3 days in the mutant males as compared to 52.4 ± 1.9 days in the control males ($p < 0.001$) and 63.9 ± 2.2 days in *Hsp67Bc*-0 females, compared to 47.8 ± 2.4 days in the control line ($p < 0.001$). On the contrary, the mean fecundity measured during the first month of life was 5.9 % lower ($p = 0.809$) in *Hsp67Bc*-0 females than in the control (see Fig. 1, *d*).

Because under normal conditions the absence of the *Hsp67Bc* gene increased the mean lifespan of the flies while causing only a minor decrease in fecundity, and because no cases of the loss of this gene in wild fruit fly populations have been reported to date, a question arose about the role of the *Hsp67Bc* gene in *D. melanogaster*. It is known that heat shock proteins (which include *Hsp67Bc*) are essential for stress tolerance in all the living organisms (Lindquist, 1986; Sørensen et al., 2003). In our previous study, we discovered involvement of the *Hsp67Bc* gene product in cold stress tolerance in *D. melanogaster* (Malkeyeva et al., 2020). In addition, the *Hsp67Bc* gene expression was shown to increase in response to heat stress (Vos et al., 2016). Therefore, we decided to investigate the impact of the deletion in the *Hsp67Bc* gene on elevated temperature tolerance in the flies.

The effect of heat stress on survival of *Hsp67Bc*-null larvae and pupae

According to FlyBase (<https://flybase.org>), the *Hsp67Bc* gene expression levels are the highest in wandering 3rd instar (late L3) larvae and pupae of *D. melanogaster*, in particular, white prepupae, 12 h pupae, and 48 h pupae. We decided to check how heat stress would affect *Hsp67Bc*-0 flies at those stages of development, in addition to the adult stage.

The larvae and pupae were placed in a 37 °C environment for 2 h, after which they were returned to 24–25 °C to recover and continue development. The survival rates of the larvae and pupae were computed as a proportion (%) of eclosed individuals (Fig. 2, *a*). The mean survival rates to adult stage were similar between the control and *Hsp67Bc*-null pupae, varying between 95.0 ± 2.9 % (12 h *Hsp67Bc*-0 pupae) and 100 % (48 h *Hsp67Bc*-0 pupae). Statistically significant differences were observed between the survival rates of the control and mutant flies at wandering L3 larva stage: the mutant larvae




Fig. 1. The survival, lifespan, and fecundity of *Hsp67Bc*-null (*Hsp67Bc-0*) and control (*Hsp67Bc-2*) *D. melanogaster* under normal conditions (24–25 °C).

Survival curves of males (a) and females (b); the mean lifespan (c); fecundity (eggs per female) dynamics of mutant and control females throughout the first month of life (d). The error bars denote standard error of the mean (SEM). *** $p \leq 0.001$.




Fig. 2. Survival rates of larvae, white prepupae, and pupae of *Hsp67Bc*-null (*Hsp67Bc-0*) and control (*Hsp67Bc-2*) flies after acute heat stress (37 °C, 2 h), and the mean lifespan of adult mutant and control flies under different regimens involving heat treatment.

a, the proportion (%) of wandering L3 larvae (LL3), white prepupae (WPP), 11–13 h pupae (P (12 h)), and of 47–49 h pupae (P (48 h)) surviving to the adult stage after 2 h of heat treatment (37 °C); b, the mean lifespan of the adult males and females constantly kept at 29 °C; c, the mean lifespan of the adult flies kept at 24–25 °C after 2 h heat treatment (37 °C) at 1 day of age; d, the mean lifespan of the adult flies kept at 24–25 °C after 2 h heat treatment (37 °C) at 7 days of age. The error bars represent SEM. * $0.010 < p \leq 0.050$; *** $p \leq 0.001$.

showed higher survival rate as compared to the control line (88.9 ± 3.1 % in the *Hsp67Bc-0* line against 76.6 ± 8.3 % in the control, $p = 0.044$).

The impact of elevated temperature on *Hsp67Bc*-null adult flies

In continuation of the temperature stress experiments on larvae and pupae, *Hsp67Bc-0* and control adult flies were subjected to one of the two variants of elevated temperature treatment. The first variant was life-long maintenance at 29 °C starting from one day of age; the second variant included heat stress (2 h at 37 °C) at either one or seven days of age with subsequent maintenance at 24–25 °C until death of all individuals. The fly ages for heat stress treatment (2 h at 37 °C) were chosen based on FlyBase (<https://flybase.org>) data indicating that the *Hsp67Bc* protein levels are much higher in 1-day-old flies than in 7-day-old flies.

Constant maintenance at 29 °C significantly shortened the lifespan of both control and *Hsp67Bc*-null flies, as compared to maintenance under normal conditions (24–25 °C) without heat treatment (see Fig. 2, b, Fig. 1, c). The mean lifespan of males was 28.1 ± 0.9 days in the control line and 31.3 ± 0.8 days in the mutant line. Still, the mean lifespan of *Hsp67Bc*-null males was 11.5 % higher than that of the control line at 29 °C ($p = 0.010$), and the mutant males passed 50 % survival between days 33 and 34 of the experiment, whereas the control ones had passed it already between days 29 and 30 (Fig. 3). Females of the control line had a mean lifespan of 32.4 ± 0.7 days and *Hsp67Bc-0* females had a mean lifespan of 32.5 ± 1.1 days. Unlike males, females of the control and mutant lines had similar survival dynamics and lifespan at 29 °C (see Fig. 2, b, Fig. 3). Of note, although the mean lifespan of *Hsp67Bc*-null *Drosophila* was exceeding or equal to that of the control flies at the 29 °C environment, the re-




Fig. 3. Survival curves of adult *Hsp67Bc*-null (*Hsp67Bc*-0) and control (*Hsp67Bc*-2) flies kept at 29 °C or at 24–25 °C after 2 h heat treatment (37 °C) at either 1 or 7 days of age.

duction of longevity caused by maintenance at the elevated temperature (29 °C) was more prominent in the mutant flies than in the control ones. Thus, maintenance at 29 °C reduced the lifespan of *Hsp67Bc*-2 males and females 1.9-fold and 1.5-fold, respectively, as compared to normal conditions (24–25 °C) without heat stress, whereas the decline was 2.3-fold in *Hsp67Bc*-null males and 2.0-fold in *Hsp67Bc*-0 females (see Fig. 1, c, Fig. 2, b).

Heat shock (37 °C, 2 h) did not cause death in neither the control nor *Hsp67Bc*-null fly line. The mean lifespan of *Hsp67Bc*-0 flies was higher as compared to the control among both males and females at both variants of heat treatment (at one or seven days of age) (see Fig. 2, c, d). Survival dynamics also significantly differed between the lines with $p \leq 0.010$ (see Fig. 3). The mean lifespan of *Hsp67Bc*-0 males heat-treated at one or seven days of age was higher than that of the control males by ~13 % ($p < 0.001$) and was 67.0 ± 1.7 days in *Hsp67Bc*-0 males heat-shocked at one day of age and 67.9 ± 1.3 days in the mutant males heat-shocked at seven days of age (see Fig. 2, c, d). The mean lifespan of *Hsp67Bc*-null females that underwent heat stress at one day of age exceeded that of the control females by ~16 % (58.9 ± 2.8 days, $p = 0.014$); between the control and mutant females heat-treated at seven days of age, the difference in

the mean lifespan was ~35 % (65.4 ± 2.0 days, whereas that of *Hsp67Bc*-2 females was 48.5 ± 2.4 days, $p < 0.001$) (see Fig. 2, c, d). Of note, the applied heat stress (37 °C, 2 h) had a different impact on the control and *Hsp67Bc*-null flies. In comparison with the maintenance under normal conditions (24–25 °C, without treatment), it increased longevity of the control males and females by 1.5–14.5 % (see Fig. 1, c, Fig. 2, c, d). On the contrary, in the *Hsp67Bc*-0 line, heat stress at 37 °C reduced the mean lifespan of females treated at one day of age by 7.8 %, and the mean lifespan of males heat-shocked at one and seven days of age by 5.5 and 4.2 %, respectively (see Fig. 1, c, Fig. 2, c, d).

These findings may suggest that even though the *Hsp67Bc* gene deletion causes an increase in the lifespan of flies at both normal and elevated temperature, it has a detrimental effect on tolerance to acute heat stress, which normally improves the longevity of flies (Hercus et al., 2003; Le Bourg, 2011; Sarup et al., 2014).

The effect of elevated temperature on *D. melanogaster* fecundity

In parallel with lifespan and survival, we measured fecundity of the control and *Hsp67Bc*-null females as the number of eggs laid in each vial within 24 h divided by the number of




Fig. 4. Fecundity (eggs per female) measured throughout the first month of life of the *Hsp67Bc*-null (*Hsp67Bc*-0) and control (*Hsp67Bc*-2) females kept at 29 °C (a) or at 24–25 °C after 2 h heat treatment (37 °C) at either 1 day (b) or 7 days (c) of age.

The error bars denote SEM. ** 0.001 < p ≤ 0.010; *** p ≤ 0.001.

females kept in those very vials. The mean egg per female ratio calculated throughout the experiment did not statistically differ between the control and mutant lines in any of the heat treatment groups (29 °C, 2 h at 37 °C at one day of age, and 2 h at 37 °C at seven days of age). Nevertheless, *Hsp67Bc*-null females had slightly reduced fecundity as compared to the control flies after being subjected to heat shock (37 °C, 2 h). The differences between the lines were more prominent than at 24–25 °C without treatment. Thus, the mean number of eggs per female was 10.5 % lower in the mutant flies that underwent the heat treatment at one day of age as compared to the control (8.71 eggs/female in *Hsp67Bc*-0 line and 9.73 eggs/female in *Hsp67Bc*-2 line, $p = 0.564$); in the mutant flies subjected to heat stress at seven days of age, this value was 12.8 % (9.77 eggs/female in *Hsp67Bc*-0 line and 11.2 eggs/female in the control, $p = 0.427$).

The egg/female ratio measured each day significantly differed between the mutant and control lines only on some of the days of the experiment (Fig. 4). It is worth mentioning that heat shock (37 °C, 2 h) at one day of age was detrimental for the fecundity of females. The next day after the heat treatment, the number of laid eggs per female was much less than under normal conditions without treatment in both the control and

mutant lines (see Fig. 4, b, Fig. 1, d). The decrease was more prominent in *Hsp67Bc*-null flies (72 % decrease in *Hsp67Bc*-0 line as compared to 40 % reduction in the control line).

In search for the cause of the reduced mean fecundity in *Hsp67Bc*-null females kept at 24–25 °C, we analyzed the morphology of ovaries in the control and mutant lines. In the ovaries of both the control and mutant flies, egg chambers at all stages of oogenesis were present. However, the mutant females had lower number of ovarioles than the control ones. Five- and 15-day-old *Hsp67Bc*-2 females had 16.9–18.4 ovarioles per ovary, whereas *Hsp67Bc*-0 females had 14.6–16.2 ovarioles per ovary ($p = 0.680$ in case of the 5-day-old flies and $p < 0.001$ in case of the 15-day-old flies). This finding may partially explain the difference in the fecundity of the two lines.

The number of ovarioles may be influenced by nutrient deprivation in *D. melanogaster* (Sarıkaya et al., 2012). Dietary restriction stimulates macroautophagy, a process of intracellular component degradation, in regulation of which *Hsp67Bc* was shown to participate (Amano et al., 2006; Carra et al., 2010; Kroemer et al., 2010). Our recent studies on macroautophagy revealed a slight increase in autophagic vacuole number in the brain of adult *Hsp67Bc*-null flies (Malkeyeva et al., 2021). Therefore, we next decided to study the morphology of the control and *Hsp67Bc*-null *D. melanogaster* ovaries under the stress of protein starvation.

The impact of *Hsp67Bc* gene deletion on starvation-induced macroautophagy in *D. melanogaster* ovaries

It is known that starvation, including protein deprivation, induces macroautophagy in *Drosophila* ovaries at two nutrient status checkpoints: germlarium and mid-oogenesis (Hou et al., 2008), which then leads to oogenesis slowdown and to an increase in the number of egg chambers eliminated from oogenesis (Barth et al., 2011). The discarded egg chambers degrade through apoptosis with the participation of autophagy (Bolobolova et al., 2020). To evaluate macroautophagy intensity, we used the LysoTracker Red DND-99 (LTR) dye, which had been shown to label acidic organelles, such as lysosomes and autolysosomes, in *D. melanogaster* (Scott et al., 2004; Klionsky et al., 2007). Massive acidification of the cytoplasm signifies death of forming egg chamber cells. We estimated the percentages of LTR-positive germaria in the control and *Hsp67Bc*-null flies kept on standard food and after five (Fig. 5) and 15 days of protein starvation.

As compared to the control line, in the ovaries of *Hsp67Bc*-null females kept on either the standard or protein-free medium, a higher number of LTR-positive germaria was observed (Fig. 6, see Fig. 5). During early oogenesis, the *Hsp67Bc* gene deletion resulted in a 1.2- to 1.5-fold increase in the LTR-positive-germaria proportion (see Fig. 6). Thus, in 5-day-old *Hsp67Bc*-0 females kept on the standard medium, 32.1 % of germaria were LTR-positive, relative to 21.1 % in the control *Hsp67Bc*-2 line ($p = 0.066$); in 5-day-old starved *Hsp67Bc*-null female ovaries, the percentage of LTR-positive germaria was as high as 77.9 % but was only 59.3 % in *Hsp67Bc*-2 females ($p = 0.045$). In 15-day-old *Hsp67Bc*-null females kept on the standard food, LTR-positive germaria constituted 31.1 %, whereas in the control line, this proportion was 24.3 % ($p = 0.348$); in 15-day-old starved flies, the percentages of LTR-positive germaria in ovaries were 73.3 % in the




Fig. 5. LTR-labelled ovaries of *Hsp67Bc*-null (*Hsp67Bc*-0) and control (*Hsp67Bc*-2) adult *D. melanogaster* females kept on either the standard or protein-free medium for 5 days.

a, an ovary of a control female kept on the standard medium; *b*, *c*, the magnified white rectangle from panel *a* showing DAPI (blue) and LTR (red) channels (*b*) and a separate LTR channel (*c*); *d*–*f*, an ovary (*f*) of an *Hsp67Bc*-2 female kept on the protein-free medium, and its magnified fragment (white rectangle from panel *f*) showing DAPI and LTR channels (*e*) and a separate LTR channel (*d*); *g*–*i*, same as *a*–*f*, for the *Hsp67Bc*-0 line. Yellow asterisks denote egg chambers with highly condensed and/or fragmented nuclei; white arrows indicate LTR-positive germaria; because too many LTR-positive germaria are present in panel *k*, not all of them are indicated by arrows, and three of them are indicated by a brace. Scale bars are 200 µm.

Hsp67Bc-0 line and 60.4 % in the *Hsp67Bc*-2 line ($p = 0.226$). These data reflect an increase in macroautophagy intensity in the germaria of *Hsp67Bc*-null females.

During mid-oogenesis, we noted a decrease in the number of egg chambers with highly condensed chromatin (see Fig. 5), which is a marker of apoptotic cell death, in *Hsp67Bc*-null flies in comparison with the control, except for the 5-day-old females kept on the standard medium. In 5- and 15-day-old starved and in 15-day-old normally fed *Hsp67Bc*-null flies, the mean number of apoptotic egg chambers per ovariole was slightly lower than that in the control flies, the difference being significant only in 5-day-old starved flies (Fig. 6). Thus, the

observed egg chamber apoptosis was 37 % lower in 5-day-old starved mutant females than in the control females ($p = 0.008$). In 15-day-old starved flies, the mean number of apoptotic egg chambers per ovariole was 11 % lower in the *Hsp67Bc*-0 flies than that in the control individuals ($p = 0.528$); in 15-day-old females kept on the standard food, this number was 35 % lower in the mutant flies than in the *Hsp67Bc*-2 line ($p = 0.131$). Although we observed a lower mean number of apoptotically dying mid-oogenesis egg chambers in *Hsp67Bc*-null flies, we can hypothesize that this phenomenon is related to the observed increased apoptosis of forming egg chambers in germaria during early oogenesis in the mutant flies.




Fig. 6. Diagrams depicting ratios of LTR-positive (LTR+) and LTR-negative (LTR-) germaria and mean numbers of mid-oogenesis egg chambers (ECs) with highly condensed and/or fragmented nuclei per ovariole in the ovaries of 5- and 15-day-old mutant (*Hsp67Bc-0*) and control (*Hsp67Bc-2*) females kept on either the standard or protein-free medium.

The numerals written inside columns in the LTR+/- diagrams represent actual numbers of analyzed germaria. The error bars denote SEM. * 0.010 < *p* ≤ 0.050; ** 0.001 < *p* ≤ 0.010.

Discussion

In this study, we investigated the impact of the *Hsp67Bc* gene deletion on *D. melanogaster* fitness under normal conditions and on their heat-stress tolerance. *Hsp67Bc*-null flies showed extended lifespan as compared to the control line under normal conditions (24–25 °C), elevated temperature conditions (29 °C), and after acute heat stress (37 °C, 2 h) (see Fig. 1, Fig. 2). At the same time, the mean fecundity of the mutant females was slightly reduced at 24–25 °C without heat treatment and after the short heat stress (Fig. 1, *d*, Fig. 4, *b*, *c*).

The observed statistically insignificant decrease in *Hsp67Bc*-null female fecundity can be explained by a combination of the following factors. First, the mutants had reduced number of ovarioles, a trait that was reported by other researchers to result in lower egg yield (Yamamoto et al., 2021). Second, the quantity of LTR-positive germaria was higher in *Hsp67Bc*-null females as compared to the control line (Fig. 6), which indicates increased macroautophagy and enhanced death of forming egg chambers resulting in less eggs (Drummond-Barbosa, Spradling, 2001; Nezis et al., 2009). Contrary, in the mutant females, a lower number of mid-oogenesis egg chambers dying via apoptosis was present as compared to the control flies (Fig. 6). This last feature of the mutant ovaries may partially compensate the first two in terms of eventual egg yield making the difference between the lines statistically insignificant. In *D. melanogaster*, ovariole number is determined at the stage of 3rd instar larva and can be influenced by either genetic or environmental factors, such as rearing temperature and diet (Sarıkaya et al., 2012).

Nutrition plays an important role in defining the quantity of ovarioles: larvae kept on medium with reduced nutrient level develop into adult flies with less ovarioles (Sarıkaya et al., 2012). The decreased ovariole number in *Hsp67Bc*-null flies reared on the standard food may be caused by impaired larva nutrition due to reduced food intake or uptake, which was not registered in our studies. Alternatively, the number of ovarioles in mutant flies could be affected by slightly increased macroautophagy, which we detected in *Hsp67Bc*-null fly germaria and, previously, in brain neurons of adult flies

with the *Hsp67Bc* gene deletion (Malkeyeva et al., 2021). It is known that macroautophagy is strongly stimulated in response to starvation (Kroemer et al., 2010); therefore, enhanced macroautophagy on larval stage caused by the absence of the *Hsp67Bc* gene product may mimic nutrient deprivation conditions leading to formation of less ovarioles. The decrease in apoptotic stage 8 egg chambers in *Hsp67Bc*-null females may be a result of increased death of forming egg chambers and, hence, enhanced quality control in the germarium resulting in less defective mid-oogenesis egg chambers in *Hsp67Bc*-null fly ovaries in comparison with the control line.

Extended longevity caused by gene mutations has been reported in *D. melanogaster*. Lifespan is increased in fruit flies carrying hypomorphic mutations in the *InR* (*insulin-like receptor*), *chico*, and *methuselah* genes (Lin et al., 1998; Clansy et al., 2001; Tatar et al., 2001). Notably, products of all these genes are involved in macroautophagy modulation via target of rapamycin (TOR) pathway (Clansy et al., 2001; Wang et al., 2015; Graze et al., 2018; Yamamoto et al., 2021), and their down-regulation leads to macroautophagy stimulation. Similarly, macroautophagy stimulation by dietary restriction or TOR kinase inhibition expands lifespan of animals belonging to various taxa (Masoro, 2000; Kapahi et al., 2004). Moreover, it was shown that longevity extension of *chico*-null *D. melanogaster* is only possible with intact macroautophagy (Bjedov et al., 2020). In this work, we discovered that the number of LTR-positive germaria was slightly higher in *Hsp67Bc*-null *D. melanogaster* ovaries (Fig. 6), which signifies increased macroautophagy. Similarly, our recent study on ultrastructure of neurons in *Wolbachia*-infected *Drosophila* brains revealed an increment in the number of autophagic vacuoles in *Hsp67Bc-0* fly neurons, which, again, points towards enhanced macroautophagy (Malkeyeva et al., 2021). Bjedov et al. (2020) demonstrated that moderate enhancement of macroautophagy in a complex of tissues increases lifespan in *D. melanogaster*, while strong and ubiquitous stimulation of macroautophagy shortens it. Hence, the extension of lifespan we observed in *Hsp67Bc*-null flies may be caused by slight increase in macroautophagy in their tissues.

Although the *Hsp67Bc*-0 flies had an increased lifespan as compared to the control line under all the tested conditions (normal temperature, elevated temperature, and short heat stress), a decline in longevity was present in the mutant flies that were heat treated in relation to untreated *Hsp67Bc*-null flies. Heat shock had the opposite effect on the control line, with acute heat stress improving the longevity of the flies. Generally, mild heat (or other stress) treatment of young adults extends *Drosophila* longevity (Hercus et al., 2003; Le Bourg, 2011; Sarup et al., 2014), as we observed in the control flies. Our experiments with 2–12 h cold treatment of *Hsp67Bc*-null fly line revealed a decreased cold stress tolerance in the mutants (Malkeyeva et al., 2020). Taken together, our results point towards an adverse impact of the *Hsp67Bc* gene deletion on short temperature stress tolerance in adult *D. melanogaster*. While in the laboratory environment flies are rarely exposed to thermal and other stresses, conditions are different in the *D. melanogaster* natural habitat, where fruit flies may experience a wide variety of extreme stresses including heat shock and chill coma. Therefore, an extended lifespan under normal conditions does not guarantee survival in the wild, as was revealed in a study by Wit et al. (2013). It is important for the survival of poikilothermic animals like *Drosophila* to be able to cope with thermal stresses. Hence, the loss of the *Hsp67Bc* gene, the product of which promotes tolerance to acute thermal stresses, though extending lifespan under normal conditions, may be deleterious in a changing environment. Taking into account that *D. melanogaster* overwinter at the adult stage in temperate regions (Izquierdo, 1991), we assume that the *Hsp67Bc* gene was not eliminated from the fruit fly genome because of its prominent role in promoting acute heat and cold tolerance in adult flies.

Conclusion

Here, we studied the effect of the *Hsp67Bc* gene deletion on *D. melanogaster* lifespan and fecundity under normal conditions, and their tolerance to elevated temperature and acute heat stress. We did not detect any difference in survival of heat-shocked (37 °C, 2 h) pupae between the mutant and control lines, and the *Hsp67Bc*-null larvae showed improved survival. Adult *Hsp67Bc*-null flies had a greater lifespan than the control line at all the tested temperature regimes but lower fecundity and decreased acute heat tolerance. We hypothesize that the lifespan extension is caused by slightly increased macroautophagy in the mutant flies, which we observed in ovaries of *Hsp67Bc*-deficient *Drosophila* and – in our earlier work – in the brains of *Hsp67Bc*-null females. At the same time, the enhanced macroautophagy in germlaria, combined with a reduced number of ovarioles, may be the cause of the fecundity reduction in the mutant flies. In conclusion, although the *Hsp67Bc* gene deletion causes the increase in *D. melanogaster* lifespan in a stress-free environment, it has a negative effect on fruit fly acute heat stress tolerance, which may negate the longevity benefits in nature habitat, where stresses like extremely high and low temperature are common.

References

Amano A., Nakagawa I., Yoshimori T. Autophagy in innate immunity against intracellular bacteria. *J. Biochem.* 2006;140(2):161-166. DOI 10.1093/jb/mvj162.

- Bainbridge S.P., Bownes M. Staging the metamorphosis of *Drosophila melanogaster*. *J. Embryol. Exp. Morphol.* 1981;66:57-80.
- Barth J.M.I., Szabad J., Hafen E., Köhler K. Autophagy in *Drosophila* ovaries is induced by starvation and is required for oogenesis. *Cell Death Differ.* 2011;18:915-924. DOI 10.1038/cdd.2010.157.
- Bjedov I., Cochemé H.M., Foley A., Wieser D., Woodling N.S., Castillo-Quan J.I., Norvaisas P., Lujan C., Regan J.C., Toivonen J.M., Murphy M.P., Thornton J., Kinghorn K.J., Neufeld T.P., Cabreiro F., Partridge L. Fine-tuning autophagy maximises lifespan and is associated with changes in mitochondrial gene expression in *Drosophila*. *PLoS Genet.* 2020;16:e1009083. DOI 10.1371/journal.pgen.1009083.
- Bolobolova E.U., Dorogova N.V., Fedorova S.A. Major scenarios of genetically regulated cell death during oogenesis in *Drosophila melanogaster*. *Russ. J. Genet.* 2020;56:655-665. DOI 10.1134/S1022795420060034.
- Carra S., Boncoraglio A., Kanon B., Brunsting J.F., Minoia M., Rana A., Vos M.J., Seidel K., Sibon O.C., Kampinga H.H. Identification of the *Drosophila* ortholog of HSPB8. *J. Biol. Chem.* 2010;285:37811-37822. DOI 10.1074/jbc.M110.127498.
- Clancy D.J., Gems D., Harshman L.G., Oldham S., Stocker H., Hafen E., Leivers S.J., Partridge L. Extension of life-span by loss of CHICO, a *Drosophila* insulin receptor substrate protein. *Science.* 2001;292(5514):104-106. DOI 10.1126/science.1057991.
- Drummond-Barbosa D., Spradling A.C. Stem cells and their progeny respond to nutritional changes during *Drosophila* oogenesis. *Dev. Biol.* 2001;231(1):265-278. DOI 10.1006/dbio.2000.0135.
- Graze R.M., Tzeng R.-Y., Howard T.S., Arbeitman M.N. Perturbation of IIS/TOR signaling alters the landscape of sex-differential gene expression in *Drosophila*. *BMC Genom.* 2018;19:893. DOI 10.1186/s12864-018-5308-3.
- Hercus M.J., Loeschcke V., Rattan S.I.S. Lifespan extension of *Drosophila melanogaster* through hormesis by repeated mild heat stress. *Biogerontology.* 2003;4:149-156. DOI 10.1023/A:1024197806855.
- Hou Y.-C.C., Chittaranjan S., Barbosa S.G., McCall K., Gorski S.M. Effector caspase Dcp-1 and IAP protein Bruce regulate starvation-induced autophagy during *Drosophila melanogaster* oogenesis. *J. Cell Biol.* 2008;182:1127-1139. DOI 10.1083/jcb.200712091.
- Izquierdo J.I. How does *Drosophila melanogaster* overwinter? *Entomol. Exp. Appl.* 1991;59:51-58. DOI 10.1111/j.1570-7458.1991.tb01485.x.
- Jolly C., Morimoto R.I. Role of the heat shock response and molecular chaperones in oncogenesis and cell death. *J. Natl. Cancer. Inst.* 2000;92(19):1564-1572. DOI 10.1093/jnci/92.19.1564.
- Kampinga H.H., Hageman J., Vos M.J., Kubota H., Tanguay R.M., Bruford E.A., Cheetham M.E., Chen B., Hightower L.E. Guidelines for the nomenclature of the human heat shock proteins. *Cell Stress Chaperones.* 2009;14:105-111. DOI 10.1007/s12192-008-0068-7.
- Kapahi P., Zid B.M., Harper T., Koslover D., Sapin V., Benzer S. Regulation of lifespan in *Drosophila* by modulation of genes in the TOR signaling pathway. *Curr. Biol.* 2004;14(10):885-890. DOI 10.1016/j.cub.2004.03.059.
- Klionsky D.J., Cuervo A.M., Seglen P.O. Methods for monitoring autophagy from yeast to human. *Autophagy.* 2007;3(3):181-206. DOI 10.4161/auto.3678.
- Kroemer G., Mariño G., Levine B. Autophagy and the integrated stress response. *Mol. Cell.* 2010;40(2):280-293. DOI 10.1016/j.molcel.2010.09.023.
- Le Bourg É. Using *Drosophila melanogaster* to study the positive effects of mild stress on aging. *Exp. Gerontol.* 2011;46:345-348. DOI 10.1016/j.exger.2010.08.003.
- Lin Y.-J., Seroude L., Benzer S. Extended life-span and stress resistance in the *Drosophila* mutant *methuselah*. *Science.* 1998;282(5390):943-946. DOI 10.1126/science.282.5390.943.
- Lindquist S. The heat-shock response. *Annu. Rev. Biochem.* 1986;55:1151-1191. DOI 10.1146/annurev.bi.55.070186.005443.
- Malkeyeva D., Kiseleva E., Fedorova S. Small heat shock protein *Hsp67Bc* plays a significant role in *Drosophila melanogaster* cold-

- stress tolerance. *J. Exp. Biol.* 2020;223(Pt.21):jeb219592. DOI 10.1242/jeb.219592.
- Malkeyeva D.A., Kiseleva E.V., Fedorova S.A. Loss of *Hsp67Bc* leads to autolysosome enlargement in the *Drosophila* brain. *Cell Biol. Int.* 2021. DOI 10.1002/cbin.11721.
- Masoro E.J. Caloric restriction and aging: an update. *Exp. Gerontol.* 2000;35:299-305. DOI 10.1016/S0531-5565(00)00084-X.
- Nezis I.P., Lamark T., Velentzas A.D., Rusten T.E., Bjørkøy G., Johansen T., Papassideri I.S., Stravopodis D.J., Margaritis L.H., Stenmark H., Brech A. Cell death during *Drosophila melanogaster* early oogenesis is mediated through autophagy. *Autophagy.* 2009;5: 298-302. DOI 10.4161/auto.5.3.7454.
- Parzych K.R., Klionsky D.J. An overview of autophagy: morphology, mechanism, and regulation. *Antioxid. Redox Signal.* 2014;20(3): 460-473. DOI 10.1089/ars.2013.5371.
- Raut S., Mallik B., Parichha A., Amrutha V., Sahi C., Kumar V. RNAi-mediated reverse genetic screen identified *Drosophila* chaperones regulating eye and neuromuscular junction morphology. *G3: Genes Genomes Genetics. (Bethesda).* 2017;7(7):2023-2038. DOI 10.1534/g3.117.041632.
- Sarikaya D.P., Belay A.A., Ahuja A., Dorta A., Green D.A., Extavour C.G. The roles of cell size and cell number in determining ovariole number in *Drosophila*. *Dev. Biol.* 2012;363:279-289. DOI 10.1016/j.ydbio.2011.12.017.
- Sarkar S., Singh M.D., Yadav R., Arunkumar K.P., Pittman G.W. Heat shock proteins: molecules with assorted functions. *Front. Biol. (Beijing).* 2011;6(4):312. DOI 10.1007/s11515-011-1080-3.
- Sarup P., Sørensen P., Loeschcke V. The long-term effects of a life-prolonging heat treatment on the *Drosophila melanogaster* transcriptome suggest that heat shock proteins extend lifespan. *Exp. Gerontol.* 2014;50:34-39. DOI 10.1016/j.exger.2013.11.017.
- Scott R.C., Schuldiner O., Neufeld T.P. Role and regulation of starvation-induced autophagy in the *Drosophila* fat body. *Dev. Cell.* 2004;7(2):167-178. DOI 10.1016/j.devcel.2004.07.009.
- Sørensen J.G., Kristensen T.N., Loeschcke V. The evolutionary and ecological role of heat shock proteins. *Ecol. Lett.* 2003;6:1025-1037. DOI 10.1046/j.1461-0248.2003.00528.x.
- Tatar M., Kopelman A., Epstein D., Tu M.-P., Yin C.-M., Garofalo R.S. A mutant *Drosophila* insulin receptor homolog that extends life-span and impairs neuroendocrine function. *Science.* 2001;292(5514): 107-110. DOI 10.1126/science.1057987.
- Vos M.J., Carra S., Kanon B., Bosveld F., Klauke K., Sibon O.C.M., Kampinga H.H. Specific protein homeostatic functions of small heat-shock proteins increase lifespan. *Aging Cell.* 2016;15:217-226. DOI 10.1111/acel.12422.
- Wang J., Wang Z., Zhang Z., Hua Q., Wang M., Shi C., Xue L., Zhang R., Xie X. *Methuselah* regulates longevity via dTOR: a pathway revealed by small-molecule ligands. *J. Mol. Cell Biol.* 2015; 7:280-282. DOI 10.1093/jmcb/mjv018.
- Wit J., Kristensen T.N., Sarup P., Frydenberg J., Loeschcke V. Laboratory selection for increased longevity in *Drosophila melanogaster* reduces field performance. *Exp. Gerontol.* 2013;48:1189-1195. DOI 10.1016/j.exger.2013.07.012.
- Yamamoto R., Palmer M., Koski H., Curtis-Joseph N., Tatar M. Aging modulated by the *Drosophila* insulin receptor through distinct structure-defined mechanisms. *Genetics.* 2021;217(2):iyaa037. DOI 10.1093/genetics/iyaa037.

ORCID ID

D. Malkeyeva orcid.org/0000-0002-7308-8846
S.A. Fedorova orcid.org/0000-0001-8257-4654
E. Kiseleva orcid.org/0000-0001-6949-182X

Acknowledgements. LTR analyses were performed at the Multiple-access Center for Microscopy of Biological Objects (Institute of Cytology and Genetics SB RAS). This work was supported by budget project No. 0259-2021-0011.

Conflict of interest. The authors declare no conflict of interest.

Received November 19, 2021. Revised December 30, 2021. Accepted January 10, 2022.