

ORIGINAL ARTICLE

Prevalence, antimicrobial susceptibility, and genotyping of *Streptococcus agalactiae* in Tilapia fish (*Oreochromis niloticus*) in Egypt

Asmaa Alazab, Asmaa Sadat , Gamal Younis 

Department of Bacteriology, Mycology and Immunology, Faculty of Veterinary Medicine, Mansoura University, 35516, Egypt

ABSTRACT

Objectives: *Streptococcus agalactiae* is a zoonotic human and animal pathogen that causes global economic losses in aquaculture and fatal outcomes in Tilapia. This study aimed to identify *S. agalactiae* isolated from different fish sources intended for human consumption phenotypically and genotypically and to characterize the virulence-associated genes *fbsA* (fibrinogen-binding protein FbsA), *cfb* (CAMP factor), and *pbp1A/ponA* (penicillin-binding protein 1A).

Materials and Methods: Three hundred Nile Tilapia fish (*Oreochromis niloticus*) were collected from different farms and retail shops in Dakahlia and Damietta, Egypt, during the summer of 2020. The samples were examined using routine phenotypic methods, then characterized using polymerase chain reaction (PCR) targeting *S. agalactiae*-specific *dltS* gene. All *S. agalactiae* isolates were examined for the susceptibility to ten antimicrobial agents by the disc diffusion method. The virulence-associated genes (*fbsA*, *cfb*, and *pbp1A/ponA*) were characterized using multiplex-PCR.

Results: *Streptococcus agalactiae* was detected in 7% ($n = 21/300$) samples. The isolates showed high resistance against ampicillin and erythromycin (20/21; 95%) for each. The most predominant antibiotypes through isolates were P, CN, SXT, CRO, TE, CTX, E, AMP, at 10.5% for each antibiotype. A total of 19 (90.5%) of *S. agalactiae* isolates showed multi-drug resistance (MDR), and those were recovered from market Tilapia fish. The virulence-associated genes (*fbsA*, *cfb*, and *pbp1A/ponA*) were identified in the *S. agalactiae* as 100%, 76%, and 52%, respectively.

Conclusions: The MDR *S. agalactiae* detected in raw Tilapia fish pose a significant health hazard to consumers due to their zoonotic characteristics.

ARTICLE HISTORY

Received November 22, 2021

Revised January 17, 2022

Accepted January 18, 2022

Published March 10, 2022

KEYWORDS

S. agalactiae; Multi-drug resistance; Tilapia; Virulence genes; multiplex-PCR



© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>)

Introduction

Streptococcus agalactiae, a Group B *Streptococcus* (GBS), produces a range of pathological conditions in aquatic animal species [1]. It causes disease in fish and humans [2] and animals, such as dogs, cats, and cattle [3]. GBS is the main causative agent of Streptococcosis outbreaks in freshwater fish aquaculture [4]. *S. agalactiae* has been associated with cases of Streptococcosis outbreaks among humans following fish consumption, indicating its potential as a zoonotic agent [5]. *S. agalactiae* is responsible for meningoencephalitis and septicemia in freshwater fish such as Tilapia [6], and cases of sudden death without detectable symptoms in freshwater, estuarine, and marine fish species [7]. Diseased fish show typical symptoms, including spiral swimming, anorexia, corneal opacity, unilateral or

bilateral exophthalmia, and ulcers and hemorrhages in the skin [8,9]. Affected organs, such as the eye, liver, kidney, and brain, show gross pathological changes, enlarging and showing hemorrhagic and inflammation signs [10]. However, fish may show no signs before sudden death [11].

The severity of the disease produced may be attributable to factors such as its virulence marker and water temperature [12,13]. Virulence genes facilitate the ability of the pathogen to cause disease. They are classified into three classes: adhesin genes (*fbsA*, *fbsB*, *pavA*, *lmb*, and *scpB*), invasin genes (*cylE*, *cfb*, *Spb1*, *hylB*, *rib*, and *bca*), and immune evasion genes (*bac*, *cspA*, and *pbp1A/ponA*). They play a major role in pathogenicity after infection of the host [14]. These genes also help bacteria stay alive,

Correspondence Asmaa Sadat ✉ asmaasadat@mans.edu.eg 📧 Department of Bacteriology, Mycology and Immunology, Faculty of Veterinary Medicine, Mansoura University, 35516, Egypt.

How to cite: Alazab A, Sadat A, Younis G. Prevalence, antimicrobial susceptibility, and genotyping of *Streptococcus agalactiae* in Tilapia fish (*Oreochromis niloticus*) in Egypt. J Adv Vet Anim Res 2022; 9(1):95–103.

make the host less angry, and let the bacteria get around the immune system.

The variation in susceptibility of *S. agalactiae* to antimicrobials could be due to variations in serotypes or to the repeated uncontrolled application of antibiotics in aquaculture [15]. Several antibiotics are used to eliminate streptococcal infection, but these antibiotics' frequent and improper application can produce antimicrobial drug resistance [13,16]. The difference in the effects of antimicrobial treatment and vaccination may be attributed to the ability of streptococci to colonize and survive inside phagocytic cells, preventing their exposure to macrophages [17].

The Nile Tilapia (*Oreochromis niloticus*) is the world's fourth most cultured fish species [18]. It is the main species cultured in Egypt, representing about 65.15% of Egyptian fish production [19], because of its rapid growth rate, good feed conversion, ability to survive in poor environmental conditions, resistance to disease, ease of spawning, palatability, and good consumer acceptance [20]. Large-scale production of farmed Tilapia increases the fish's stress and, consequently, their susceptibility to disease [21]. Streptococcal outbreaks can cause major economic losses and are a major challenge for the development of the tilapia industry worldwide [22]. There is a little consideration about the resistant phenotypes, antimicrobial susceptibility, and virulence mechanisms of *S. agalactiae* recovered from farmed, diseased, and retailed Tilapia fish in Egypt. Thus, this study aimed to detect the prevalence of *S. agalactiae* among Tilapia fish recovered from diseased farmed fish and retail fish, study their antimicrobial susceptibility, and genotype their virulence-associated genes. Finally, to evaluate the hygiene measures taken among retail shop Tilapia fish.

Materials and Methods

Ethical approval

Our study design and fish sampling were approved by the Research Ethics Committee of the Faculty of Veterinary Medicine, Mansoura University, Egypt (Protocol code: M/19).

Fish sampling

From April 2020 to November 2020, 300 Nile Tilapia fish were collected. One hundred fish were sampled from farms located at EL-Manzala Lake, and two hundred fish were collected from retail markets in Dakahlia and Damietta provenances, Egypt. Fish samples were gathered in sterile screw-capped plastic vials, placed in an icebox, and sent for microbiological analysis at the Bacteriology, Mycology, and Immunology Department, Faculty of Veterinary Medicine, Mansoura University.

Isolation of *S. agalactiae* from Tilapia

Under sterile conditions, swabs were sampled from the kidney, brain, liver, eye, spleen, and gills and plated on Edward agar (Oxoid, Basingstoke, UK) and 5% sheep blood agar (Oxoid, Basingstoke, UK), followed by incubation for 24 h at 37°C. Tryptone soy agar (Oxoid, Basingstoke, UK) and blood agar (Oxoid, Basingstoke, UK) were streaked with separate cultured colonies showing morphology similar to Streptococci and then incubated at 37°C for 24 h to obtain pure isolates. Streptococcal isolates were characterized using colonial morphology, Gram staining, motility, catalase tests, cytochrome oxidase tests, hemolysis on 5% sheep blood agar, and esculin hydrolysis on bile esculin slants [23,24]. The CAMP reaction was used as a presumptive diagnosis for *S. agalactiae*, as the CAMP factor was identified first in *S. agalactiae* [25]. All suspected streptococci isolates were preserved in 20% glycerol.

DNA extraction

Genomic DNA was extracted from all strains via homogenizing of a few bacterial colonies (3–5 colonies) in about two hundred milliliters of deionized water, then boiling for fifteen minutes, then centrifugation at 10,000 *g* for three minutes. All the supernatant was collected in a sterile Eppendorf tube and used as a DNA template [26]. The genomic DNA was stored at –20°C until it was used.

Molecular identification of *S. agalactiae* using PCR

All suspected *Streptococcus* spp. were characterized using polymerase chain reaction (PCR) analysis of specific primers targeting the *S. agalactiae*-specific *dltS* gene. The primer sequence and PCR cyclic conditions were mentioned by Poyart et al. [27] (Table 1). The PCR used reaction mix contained 12.5 µl 2× Tag PCR Master Mix (Enzymomics, Daejeon, South Korea), 1 µl of each primer (Metabion International AG, Steinkirchen, Germany), and 3 µl of template DNA in a final volume of 25 µl. The PCR cycle conditions were: 5 min at 94°C, 35 cycles at 94°C for 45 sec, 66°C for 45 sec, and 72°C for 90 sec, and the extension step at 72°C for 5 min using a thermal cycler (Applied Biosystem, 2720 thermal cycler, USA). Then, 10 µl of amplified PCR products were seen using 1% agarose gel through a gel documentation system (Hoefer PS300B life science, USA) and then checked with UV-induced fluorescence (Cleaver Scientific Ltd. UV gel documentation system, USA).

Sequencing of the 16S rRNA of *S. agalactiae*

The primer sequence and PCR cyclic conditions for 16S rRNA used for *S. agalactiae* confirmation and sequencing were performed according to Lagacé et al. [28]. The purification of amplified products for strains was performed using QIAquick PCR Product Extraction Kits (Qiagen Inc.

Table 1. Oligonucleotides and PCR cyclic conditions for PCR amplification of genes used in this study.

Target gene	Primer	Nucleotide sequence (5'-3')	Target gene (bp)	PCR cyclic conditions	Reference
<i>Dlts</i>	<i>dlts-F</i>	AAGTACATGCTGATCAAGT	952	5 min at 94°C, 35 cycles at 94°C for 45 sec, 66°C for 45 sec, 72°C for 1.5 min, and 72°C for 5 min	[27]
	<i>dlts-R</i>	TCTTGATCAACTTGTGTAC			
<i>16S rRNA</i>	<i>F27</i>	AGAGTTTGATCMTGGCTCAG	1485	5 min at 95°C, 35 cycles at 94°C for 30 sec, 56°C for 60 sec, 72°C for 60 sec, and 72°C for 10 min	[28]
	<i>R1492</i>	TACGGYACCTTGTACGACTT			
<i>Pbp1A/ponA</i>	<i>Pbp1A/ponA-F</i>	AGGGGTAGTAGCATTACCAT	939	5 min at 95°C, 38 cycles at	[34]
	<i>Pbp1A/ponA-R</i>	CAACTATGACTGGGATCG			
<i>Cfb</i>	<i>cfb-F</i>	GGATTCAACTGAACTCCAAC	600	95°C for 30 sec, 47°C for 30 sec, 72°C for 30 sec, and 72°C for 10 min	[34]
	<i>cfb-R</i>	GACAACTCCACAAGTGGTAA			
<i>FbsA</i>	<i>fbsA-F</i>	AACCGCAGCGACTTGTTA	278	72°C for 10 min	
	<i>fbsA-R</i>	AAACAAGAGCCAAGTAGGTC			

Valencia, CA) and sequenced using BigDye Terminator V3.1 cycle sequencing kits (Perkin-Elmer, Foster City, CA). According to the manufacturer's instructions, the purification of the sequence reactions (Centrisep, spin-column) used an Applied Biosystems 3130 automated DNA Sequencer (ABI, 3130). BLAST® analysis (Basic Local Alignment Search Tool) was used to identify the sequences and obtain their GenBank accession numbers [29]. Phylogenetic analyses were performed using maximum likelihood, neighbor-joining, and maximum parsimony in MEGA6 (Fig. 3) [30].

Antimicrobial susceptibility test for *S. agalactiae*

Using disc diffusion methods, all *S. agalactiae* strains were tested for susceptibility against 10 antimicrobial agents belonging to seven antimicrobial classes. These antimicrobial agents (Oxoid, Basingstoke, UK) were ciprofloxacin (CIP; 5 µg), penicillin (P; 10 µg), sulfamethoxazole/trimethoprim (SXT; 25 µg), ceftriaxone (CRO; 30 µg), tetracycline (TE; 30 µg), cefotaxime (CTX; 30 µg), erythromycin (E; 15 µg), ampicillin (AMP; 10 µg), imipenem (IPM; 10 µg), and gentamicin (CN; 10 µg) according to CLSI [31]. Tests were performed on Mueller–Hinton agar (Oxoid, Basingstoke, UK) supplemented with 5% sheep blood and incubated aerobically overnight at 37°C. The results were interpreted according to the Clinical and Laboratory Standards Institute [31]. Strains that showed resistance to three or more antimicrobial agents belonging to three different antibiotic classes were considered multi-drug resistance (MDR) bacteria [32]. A multiple antibiotic resistance (MAR) index was calculated according to Krumperman [33].

Genotyping of virulence-associated genes using multiplex PCR

Streptococcus agalactiae strains were genotyped for the standard virulence genes *fbsA* (fibrinogen-binding protein FbsA), *cfb* (CAMP factor), and *pbp1A/ponA* (penicillin-binding protein 1A), which were amplified as described by Kannika et al. [34] using multiplex PCR (Table 1). The reaction mixture contained 12.5 µl of PCR Master Mix, 1 µl of each primer, and 3 µl of template DNA, made up with nuclease-free water to a final reaction volume of 25 µl. The amplification conditions were as follows: an initial denaturation of 5 min at 95°C, followed by 38 cycles of 95°C for 30 sec, 47°C for 30 sec, and 72°C for 30 sec, then a final extension of 10 min at 72°C (Table 1). The amplifications were examined on a thermal cycler (Applied Biosystem, 2720 thermal cycler, USA). The amplified PCR products were visualized on a 1% agarose gel using a gel documentation system (Hoefer PS300B life science, USA) and then checked using UV-induced fluorescence (Cleaver Scientific LTD UV gel documentation system, USA).

Results

Prevalence of *S. agalactiae* isolated from *Tilapia*

The *Streptococcus* species appeared as small, blueish colonies with a smooth, white edge on Edward agar and translucent to slightly opaque, pinpoint colonies surrounded by a beta-hemolytic zone on blood agar. These bacteria were biochemically identified and shown to be Gram-positive cocci, non-motile, oxidase-positive, catalase-negative, and CAMP positive. Of the 300 cultured samples from

Table 2. Prevalence of *Streptococcus spp* isolated from market and farmed Tilapia fish samples using PCR assays.

Source	Farmed Tilapia Fish (No/%)	Market Tilapia Fish (No/%)	Total Number (No/%)
Overall number of Samples	100	200	300
Number of <i>Streptococcus</i> Isolates	34 (34%)	55 (27.5%)	89 (29.6%)
Number of <i>Streptococcus agalactia</i>	13 (13%)	8 (4%)	21 (7%)




Figure 1. PCR analysis of *16S rRNA*. Lanes: 1—MW-DNA ladder (100bp); 2—positive control; 3—negative control; 4,5—*Streptococcus agalactiae* strains.

freshwater fish (*O. niloticus*), 89 isolates (29.6%) were identified as *Streptococcus spp.* using routine phenotypic methods. The suspected *Streptococcus* isolates were subjected to molecular identification (PCR) using species-specific *dlt-s* targeting *S. agalactiae*; almost 23.6% (21/89) of the suspected isolates were confirmed as *S. agalactiae*. The overall prevalence rate of *S. agalactiae* in all examined Tilapia samples in this study was 7% (21/300). *Streptococcus agalactiae* was isolated from 13% (13/100) of diseased farmed Tilapia fish and 4% (8/200) of market Tilapia fish (Table 2, Fig. 1). The *S. agalactiae* strains were isolated from fish samples as follows: 10 (47.6%) from the liver; 6 (28.57%) from the kidney; 2 (9.5%) from each eye and gill samples; 1 (4.85) from spleen samples.

Antimicrobial susceptibility testing of *S. agalactiae*

Susceptibility tests to 10 antimicrobial agents from seven classes of antimicrobial agents showed that the most antimicrobial resistance of *S. agalactiae* strains was detected versus ampicillin at 95% (20/21) and erythromycin at

95%, followed by cefotaxime and trimethoprim-sulfamethoxazole (76% each), ceftriaxone (72%), tetracycline (66%), gentamicin (62%), and penicillin (57%). Less antimicrobial resistance was detected against ciprofloxacin, at 43% (9/21), and imipenem, at 5% (1/21) (Table 3). MDR has been observed in 90.5% of isolates tested. MDR was present in all *S. agalactiae* strains isolated from market Tilapia but in farmed Tilapia at lower levels, of 84.6%. Nineteen antibiotypes were identified in *S. agalactiae*. The most predominant antibiotypes through isolates were P, CN, SXT, CRO, TE, CTX, E, AMP, and CIP, P, CN, SXT, CRO, CTX, E, AMP, at 10.5% for each antibiotype (Table 4).

Virulence associated genes in *S. agalactiae* strains

Three virulence genes were used to predict the virulence of *S. agalactiae*: *fbsA*, *cfb*, and *pbp1A/ponA*. The *fbsA* gene was detected in all isolates, while *cfb* was present in 76% (16/21) of isolates. Around half the isolates (11/21; 52%) carried the *pbp1A/ponA* gene. The *fbsA* and *cfb* genes were recovered from sixteen isolates, and all 3 virulence genes were isolated from 11 isolates (Fig. 2; Table 5).

Sequencing of the *16S rRNA* of *S. agalactiae*

After identifying *S. agalactiae* strains for *16S rRNA* using the PCR method, sequencing of *S. agalactiae* strains was applied by Elim biopharmaceuticals (USA). The nucleotide sequence of the *S. agalactiae* strains was deposited in GenBank under accession no. OL335944. Phylogenetic trees show the genetic relatedness among *S. agalactiae* strains according to nucleotide sequence analysis of the *16S-rRNA* gene (Fig. 3).

Discussion

GBS is a common group of bacteria in humans and fish, causing meningitis in newborns, mastitis in cattle, and sepsis in rabbits [35]. *Streptococcus agalactiae* is the dominant species of streptococci associated with fish disease, especially in Tilapia production [36]. *Streptococcus agalactiae* causes high mortality in susceptible fish species, reaching 50%–70% in intensive farming systems [6]. Some serotypes of *S. agalactiae* from human newborns with meningitis and cattle result in disease and death in infected Nile Tilapia [6], causing severe public health problems. To the best of authors' knowledge, there is a gap in knowledge

Table 3. Antimicrobial susceptibility results of *Streptococcus agalactiae* strains isolated from fish (n = 21).

Antimicrobial Agent	Family	Disc code	Resistant No/(%)	Intermediate No/(%)	Susceptible No/(%)
Ampicillin		Amp	20/(95%)	Zero	1/(5%)
Penicillin	β-lactams	p	12/(57%)	Zero	9/(43%)
Imipenem		Imp	1/(5%)	1/ (5%)	19/(90%)
Erythromycin	Macrolide	E	20/(95%)	Zero	1/(5%)
Cefotaxime		CTX	16/(76%)	Zero	5/(24%)
Ceftriaxone	Cephalosporin	CRO	15/(72%)	Zero	6/(28%)
Trimethoprim/Sulphamethoxazole	Sulphonamide	SXT	16/(76%)	Zero	5/(24%)
Tetracycline	Tetracycline	TE	14/(66%)	5/(24%)	2/(10%)
Gentamicin	Aminoglycoside	CN	13/(62%)	Zero	8/(38%)
Ciprofloxacin	Fluoroquinolone	CIP	9/(43%)	9/(43%)	3/(14%)

Table 4. Antibiogram and MAR of *Streptococcus agalactiae* strains isolated from fish (n = 21).

	Resistance pattern	MAR index	Isolates No. (%)
I	CRO	0.1	1 (5.2)
II	P, E, AMP	0.3	1 (5.2)
III	P, CTX, E, AMP	0.4	1 (5.2)
IV	SXT, TE, CTX, E, AMP	0.5	1 (5.2)
V	P, SXT, TE, E, AMP	0.5	1 (5.2)
VI	P, TE, CTX, E, AMP	0.5	1 (5.2)
VII	CN, CRO, TE, CTX, E, AMP	0.6	1 (5.2)
VIII	CIP, CN, SXT, CRO, E, AMP	0.6	1 (5.2)
IX	CN, SXT, CRO, TE, CTX, E, AMP	0.7	1 (5.2)
X	CIP, SXT, CRO, TE, CTX, E, AMP	0.7	1 (5.2)
XI	CIP, P, CN, SXT, TE, E, AMP	0.7	1 (5.2)
XII	CIP, SXT, CRO, TE, CTX, E, AMP	0.7	1 (5.2)
XIII	CN, SXT, CRO, TE, CTX, E, AMP	0.7	1 (5.2)
XIV	CIP, CN, SXT, CRO, CTX, E, AMP	0.7	1 (5.2)
XV	P, CN, SXT, CRO, TE, CTX, E, AMP	0.8	2 (10.5)
XVI	CIP, P, CN, SXT, CRO, CTX, E, AMP	0.8	2 (10.5)
XVII	CIP, P, CN, SXT, CRO, TE, CTX, E, AMP	0.9	1 (5.2)
XVIII	P, CN, IPM, SXT, CRO, TE, CTX, E, AMP	0.9	1 (5.2)
XIX	CIP, P, CN, SXT, CRO, TE, CTX, E, AMP	0.9	1 (5.2)

considering the resistance mechanisms in *S. agalactiae*; studies from Egypt have not focused on the molecular characterization of GBS from *O. niloticus* to assess its environmental hazard. Our study clarifies the investigation of *S. agalactiae* in both diseased and retail Tilapia fish, antimicrobial resistance, and its role in transmitting virulence determinates between aquatic animals and humans via the food route.

The overall prevalence rate of *S. agalactiae* was 7% (21/300). The prevalence of diseased farmed Tilapia (13/100; 13%) was higher than that of market Tilapia

(8/200; 4%). Previous results were consistent with ours in diseased farmed Tilapia [37]. However, a higher prevalence was found in studies of *S. agalactiae* from diseased fish from Saudi Arabia, Piura, Peru, and Thailand, of 44.4%, 56.25%, and 30.9%, respectively [34,38,39]. In China, *S. agalactiae* has been the causative agent for more than 90% of infected Tilapia since 2009 [40]. In Colombia and Brazil, *S. agalactiae* infection has high mortality in Tilapia farms [41,42]. Higher infection rates have been reported in Latin America and Asia [43]. In previous reports, fresh Nile tilapia showed a lower prevalence of infection than in our




Figure 2. Multiplex PCR analysis of *Streptococcus agalactiae* virulence genes showed MW-DNA ladder (100 bp), *fb*sA gene (278 bp), *cfb* gene (600 bp), and *pbp1A/ponA* gene (939 pb).

Table 5. Prevalence of virulence genes in *Streptococcus agalactiae* isolated from market and farmed Tilapia fish samples.

Virulence genes	Farmed Tilapia Fish No (%)	Market Tilapia Fish No (%)	Total number (%)
<i>fb</i> sA gene	13 (61.9%)	8 (38.09%)	21 (100%)
<i>cfb</i> gene	10 (47.62%)	6 (28.57%)	16 (76%)
<i>Pbp1A/ponA</i> gene	7 (33.33%)	4 (19.05%)	11 (52%)

study [16,44]. In Egypt, a study from Kafr El-Sheikh, Egypt isolated *S. agalactiae* from Tilapia during the summer with a prevalence of 13% [45].

The diversity of GBS prevalence can be attributed to multiple factors. Tilapia under unsuitable conditions is susceptible to several bacterial diseases, especially *S. agalactiae*. The contributing factors are high water temperature, high stocking density, and poor water quality due to high ammonia levels and low dissolved oxygen levels [6,46]. All outbreaks of streptococcosis in Nile Tilapia occurred in summer when the water temperature rose above 27°C [47]. High temperatures are a good environment for expressing the *S. agalactiae* virulence genes, leading to severe damage to fish tissue [47].

The rapidly growing antimicrobial resistance rate is a serious issue worldwide for human and veterinary medicine [43,48–51]. In our study, the susceptibility of *S. agalactiae* to 10 antimicrobials from seven antimicrobial classes was tested. Of these antimicrobials, *S. agalactiae* isolates showed the most potent resistance against ampicillin (95%); however, lower resistance was detected in other reports [38,45]. The isolates studied exhibited greater resistance to tetracycline and penicillin (66% and 57%, respectively) than was reported in previous studies [34,38,39,45,52,53–55]. Similar tetracycline and penicillin resistance result was reported previously [16,22].

The drugs of choice for GBS treatment are penicillin and beta-lactams, followed by macrolides (erythromycin) used in individuals allergic to beta-lactams [56,57]. They are used in aquaculture for prophylactic or treatment purposes [58,59]. Almost all isolates were resistant to erythromycin, as reported by other authors [16,22,60]. The mechanism of resistance against erythromycin in the bacteria includes changes of the ribosomal target by a methylase [49,59]. It is important to search for other options for infection control [50,59].

In our study, a lower rate of sensitivity to trimethoprim/sulfamethoxazole (24%) was detected in our study. However, other studies found higher sensitivity rates [34,39]. Our results showed higher resistance against gentamicin (62%), as reported in several reports [15,21,51,59]. However, other studies found lower levels of resistance [16,22]. Gentamicin affects the survival of bacteria by preventing the binding of its messenger RNA, which causes misreading of its DNA and changes to the proteins it makes.

Antimicrobial resistance against quinolones and third-generation cephalosporins was about 40% and 75%, respectively. This result was different from those of other studies into the antimicrobial sensitivity of *S. agalactiae* to third-generation cephalosporins [38,45]. Fluoroquinolone resistance has increased lately, and this increase has been




Figure 3. Phylogenetic trees viewing the genetic relatedness among *S. agalactiae* strains according to nucleotide sequence analysis of A) *16S-rRNA* gene (OL335944).

attributed to many causes, such as efflux mechanisms or mutations in the quinolone-resistance-determining regions [54,55]. Non-guided and inaccurate use of antibiotics has resulted in many problems, such as resistance, and can affect the safety and quality of food [55,59]. This is a big problem for public health because of the spread of MDR in human communities, hospitals, animal farming, and veterinary medicine. It has become a problem for public health all over the world.

In this study, one of each of the three functional categories of virulence genes—adhesins, invasions, and immune evasions—was characterized in all isolates to determine their pathogenic and invasive abilities. At least two virulence genes were identified in the *S. agalactiae* isolates. The *fbSA* gene, responsible for an adhesion protein on the bacterial surface, promoted adhesion to and invasion of host cells [57,59] and was detected in our isolates. This gene was not expressed in all isolates in a study from Thailand [37]. The *cfb* gene, which encodes CAMP factor, a secreted pore-forming protein causing lysis of red blood cells, was carried by 76% of isolates. Thus, the hemolysis caused by the CAMP factor can be used for the phenotypic characterization of *S. agalactiae*. The penicillin-binding protein A encoding gene (*pbp1A/ponA*), which was detected in almost half of the isolates, leads to enhancement of the resistance of the host to antimicrobial peptides. Previous

studies have seen the three virulence genes in all *S. agalactiae* isolates [34,44]. Understanding pathogenesis in aquatic outbreaks is complex because it is a process that involves both bacterial virulence genes and other related factors. Characterization of the patterns of virulence genes will help determine the genetic diversity of *Streptococcus* spp. and understand the genetic relatedness between bacterial virulence and host adaptations [60].

Conclusions

Intensive cultivation systems with poor environmental conditions are related to the high mortality of fish, causing severe economic losses. Thus, special monitoring programs, efficient sanitary measures, and constant monitoring are essential in controlling and treating GBS. MDR is an emerging global threat, limiting our options for treatment of GBS infections due to uncontrolled use of antibiotics, which must be controlled to limit the emergence of MDR strains in the environment.

List of abbreviations

GBS, Group B Streptococcus; MDR, Multi-drug resistance; DNA, Deoxyribonucleic acid; PCR, Polymerase Chain Reaction; CAMP, Christie Atkins Munch-Petersen; *fbSA*, fibrinogen-binding protein; *fbSB*, fibrinogen-binding protein; *pavA*, fibrinogen-binding protein; *scpB*, Segregation and condensation protein B; *Lmb*, laminin-binding protein; *cylE*, B-haemolysin/cytolysin; Cfb, CAMP factor; *Spb1*, haemolysin III; *hylB*, Hyaluronatylase; *Rib*, the surface protein; *Bca*, C- α protein; *Bac*, C- β protein; *cspA*, the serine protease; *Pbp1A/ponA*, penicillin-binding protein 1A.

Acknowledgment

No Acknowledgment.

Conflicts of interest

The authors declare no conflicts of interest.

Authors' contributions

A.A performed experiments and analyzed the data; A.A and A.S wrote the manuscript. G.Y and A.S designed the experiment and revised the manuscript. All authors approved the final version of the manuscript for publication.

References

- [1] Timoney JF, Gyles CL, Prescott JF, Songer G, Thoen CO. Streptococcus. Gyles CL, Prescott JF, Songer G, Thoen CO (ed.). In: Pathogenesis of bacterial infections in animals. 4th edition, Blackwell, Ames IA, pp 51–74, 2010; <https://doi.org/10.1002/9780470958209.ch4>

- [2] Collin SM, Shetty N, Lamagni T. Invasive Group B *Streptococcus* infections in adults, England, 2015–2016. *Emerg Infect Dis* 2020; 26(6); <https://doi.org/10.3201/eid2606.191141>
- [3] Pang M, Sun L, He T, Bao H, Zhang L, Zhou Y, et al. Molecular and virulence characterization of highly prevalent *Streptococcus agalactiae* circulated in bovine dairy herds. *Vet Res* 2017; 48: Article No. 65; <https://doi.org/10.1186/s13567-017-0461-2>
- [4] Jiménez Leaña Á. Detección de *Streptococcus agalactiae* por PCR en tejidos de *Tilapia rojas* (*Oreochromis* spp.) menores de 20g. [online] 2010, Universidad Nacional de Colombia Sede Bogotá Facultad de Ciencias Posgrado Interfacultades en Microbiología. <https://repositorio.unal.edu.co/handle/unal/6794> [Acceso: 8 de marzo 2022]
- [5] Kalimuddin S, Chen SL, Lim CTK, Koh TH, Tan TY, Kam M, et al. 2015 epidemic of severe *Streptococcus agalactiae* sequence type 283 infections in Singapore associated with the consumption of raw freshwater fish: a detailed analysis of clinical, epidemiological, and bacterial sequencing data. *Clin Infect Dis* 2017; 64(2):S145–52; <https://doi.org/10.1093/cid/cix021>
- [6] Adikesavalu H, Banerjee S, Patra A, Abraham TJ. Meningoencephalitis in farmed monosex Nile Tilapia (*Oreochromis niloticus* L.) caused by *Streptococcus agalactiae*. *Arch Pol Fish* 2017; 25:187–200; <https://doi.org/10.1515/aopf-2017-0018>
- [7] Evans JJ, Bohnsack JF, Klesius, Evans JJ, Klesius PH, Shoemaker CA. An overview of *Streptococcus* in warmwater fish. *Aquacult Health Int* 2006; 7:10–14.
- [8] Laith AA, Ambak MA, Hassan M, Sheriff SM, Nadirah M, Draman AS, et al. Molecular identification and histopathological study of natural *Streptococcus agalactiae* infection in hybrid Tilapia (*Oreochromis niloticus*). *Vet World* 2017; 10(1):101–11; <https://doi.org/10.14202/vetworld.2017.101-111>
- [9] Evans JJ, Pasnik DJ, Klesius PH. Differential pathogenicity of five *Streptococcus agalactiae* isolates of diverse geographic origin in Nile Tilapia (*Oreochromis niloticus* L.). *Aquac Res* 2015; 46:2374–81; <https://doi.org/10.1111/are.12393>
- [10] Pretto-Giordano LG, Müller EE, de Freitas JC, da Silva VG. Evaluation on the pathogenesis of *Streptococcus agalactiae* in Nile Tilapia (*Oreochromis niloticus*). *Braz Arch Biol Technol* 2010; 53(1):87–92; <https://doi.org/10.1590/S1516-89132010000100011>
- [11] Ye X, Li J, Lu M, Deng G, Jiang X, Tian Y, et al. Identification and molecular typing of *Streptococcus agalactiae* isolated from pond-cultured Tilapia in China. *Fish Sci* 2011; 77(4):623–32; <https://doi.org/10.1007/s12562-011-0365-4>
- [12] Liao PC, Tsai YL, Chen YC, Wang PC, Liu SC, Chen SC. Analysis of streptococcal infection and correlation with climatic factors in cultured Tilapia *Oreochromis* spp, Taiwan. *Appl Sci* 2020; 10(11):2076–3417; <https://doi.org/10.3390/app10114018>
- [13] Sudprasert C, Wang PC, Chen SC. Phenotype, genotype and pathogenicity of *Streptococcus agalactiae* isolated from cultured Tilapia (*Oreochromis* spp.) in Taiwan. *J Fish Dis* 2021; 44:747–56; <https://doi.org/10.1111/jfd.13296>
- [14] Lin FP, Lan R, Sintchenko V, Gilbert GL, Kong F, Coiera E. Computational bacterial genome-wide analysis of phylogenetic profiles reveals potential virulence genes of *Streptococcus agalactiae*. *PLoS One* 2011; 6:e17964; <https://doi.org/10.1371/journal.pone.0017964>
- [15] Abuseliana A, Daud H, Aziz S, Bejo S, Alsaid M. *Streptococcus agalactiae* the etiological agent of mass mortality in farmed red Tilapia (*Oreochromis* sp.). *J Anim Vet Adv* 2010; 9:2640–6; <https://doi.org/10.3923/javaa.2010.2640.2646>
- [16] Deng L, Li Y, Geng Y, Zheng L, Rehman T, Zhao R, et al. Molecular serotyping and antimicrobial susceptibility of *Streptococcus agalactiae* isolated from fish in China. *Aquac* 2019; 510:84–9; <https://doi.org/10.1016/j.aquaculture.2019.05.046>
- [17] Zlotkin A, Chilmonczyk S, Eynigor M, Hurvitz A, Ghittino C, Eldar A. Trojan horse effect: phagocyte-mediated *Streptococcus iniae* infection of fish. *Infect Immun* 2003; 71:2318–25; <https://doi.org/10.1128/IAI.71.5.2318-2325.2003>
- [18] FAO yearbook. *Fishery and Aquaculture Statistics 2017/FAO annuaire*, Rome, Italy.
- [19] Gafrd. *Egyptian Fish statistics yearbook*, 2015. 27th edition, General Authority for Fish Resources Development, Cairo, Egypt, 2017.
- [20] Fitzsimmons K, Martinez-Garcia R, Gonzalez-Alanis P. Why Tilapia is becoming the most important food fish on the planet. In: Liu LP, Fitzsimmons K (eds.). *Proceedings of the 9th International Symposium on Tilapia in Aquaculture*, Shanghai Ocean University, Shanghai, China, pp 9–18, 2011 .
- [21] Klesius PH, Shoemaker CA, Evans JJ. *Streptococcus*: a worldwide fish health problem – In: Elghobashy H, Fitzsimmons K, Diab AS (eds.). *8th International Symposium on Tilapia in Aquaculture*, Cairo, Ag. Press Unit, Abbassa, Egypt, pp 83–107, 2008.
- [22] Wang EL, Wang KY, Chen DF, Geng Y, Huang WJ. Molecular cloning and bioinformatic analysis of the *Streptococcus agalactiae* neuA gene isolated from Tilapia. *Genet Mol Res* 2015; 14(2):6003–17; <https://doi.org/10.4238/2015.June.1.18>
- [23] MacFaddin JF. *Media for isolation-cultivation-identification-maintenance of medical bacteria*, vol. I, Williams and Wilkins, Baltimore, MD, 1985.
- [24] Barrow GI, Feltham RKA. *Cowan and steel's manual for the identification of the medical bacteria*. 3rd edition, Cambridge University Press, Cambridge, UK, 2003.
- [25] Jin T, Brefo Mensah E, Fan W, Zeng W, Li Y, Zhang Y, Palmer M. Crystal structure of the *Streptococcus agalactiae* CAMP factor and insights into its membrane-permeabilizing activity. *J Biol Chem* 2018; 293:11867–77; <https://doi.org/10.1074/jbc.RA118.002336>
- [26] Alexopoulou K, Foka A, Petinaki E, Jelastopulu E, Dimitracopoulos G, Spiliopoulou I. Comparison of two commercial methods with PCR restriction fragment length polymorphism of the *tuf* gene in the identification of coagulase-negative *Staphylococci*. *Lett Appl Microbiol* 2006; 43:450–45; <https://doi.org/10.1111/j.1472-765X.2006.01964.x>
- [27] Poyart C, Tazi A, Réglier-Poupet H, Billoët A, Tavares N, Raymond J, et al. Multiplex PCR assay for rapid and accurate capsular typing of group B streptococci. *J Clin Microbiol* 2007; 45(6):1985–8; <https://doi.org/10.1128/JCM.00159-07>
- [28] Lagacé L, Pitre, M, Jacques M, Roy D. Identification of the bacterial community of maple sap by using amplified ribosomal DNA (rDNA) restriction analysis and rDNA sequencing. *Appl Environ Microbiol* 2004; 70(4):2052–60; <https://doi.org/10.1128/AEM.70.4.2052-2060.2004>
- [29] NCBI Resource Coordinators. Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res* 2016;44(D1):D7–19; <https://doi.org/10.1093/nar/gkv1290>
- [30] Tamura K, Stecher G, Peterson D, Filipiski A, Kumar S. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 2013; 30:2725–9; <https://doi.org/10.1093/molbev/mst197>
- [31] CLSI. Clinical and Laboratory Standards Institute. performance standards for antimicrobial susceptibility testing; twenty-seventh informational supplement. Document M100-S27. Wayne, PA, 2017.
- [32] Schwarz S, Silley P, Simjee S, Woodford N, van Duijkeren E, Johnson AP, et al. Editorial: Assessing the antimicrobial susceptibility of bacteria obtained from animals. *J Antimicrob Chemother* 2010; 65:601–4; <https://doi.org/10.1093/jac/dkq037>
- [33] Krumperman PH. Multiple antibiotic resistance indexing of *Escherichia coli* to identify high-risk sources of fecal contamination of foods. *Appl Environ Microbiol* 1983; 46:165–70; <https://doi.org/10.1128/aem.46.1.165-170.1983>
- [34] Kannika K, Pisuttharachai D, Srisapoome P, Wongtavatchai J, Kondo H, Hirono I, et al. Molecular serotyping, virulence gene profiling and pathogenicity of *Streptococcus agalactiae* isolated from Tilapia farms in Thailand by multiplex PCR. *J Appl Microbiol* 2017; 122:1497–507; <https://doi.org/10.1111/jam.13447>

- [35] Ren SY, Geng Y, Wang KY, Zhou ZY, Liu XX, He M, et al. *Streptococcus agalactiae* infection in domestic rabbits, *Oryctolagus cuniculus*. *Transbound Emerg Dis* 2014; 61:92–5; <https://doi.org/10.1111/tbed.12073>
- [36] Liu G, Zhu J, Chen K, Gao T, Yao H, Liu Y, et al. Development of *Streptococcus agalactiae* vaccines for Tilapia. *Dis Aquat Organ* 2016; 21:122(2):163–70; <https://doi.org/10.3354/dao03084>
- [37] Kayansamruaj P, Pirarat N, Katagiri T, Hirono I, Rodkhum C. Molecular characterization and virulence gene profiling of pathogenic *Streptococcus agalactiae* populations from Tilapia (*Oreochromis* sp.) farms in Thailand. *J Vet Diagn Invest* 2014; 26:488–95; <https://doi.org/10.1177/1040638714534237>
- [38] Al-Harbi AH. Phenotypic and genotypic characterization of *Streptococcus agalactiae* isolated from hybrid Tilapia (*Oreochromis niloticus* x *O.aureus*). *Aquaculture* 2016; 464:515–20; <https://doi.org/10.1016/j.aquaculture.2016.07.036>
- [39] Asencios OY, Sanchez BF, Mendizabal BH, Pusari HK, Alfonso OH, Sayan MA, et al. First report of *Streptococcus agalactiae* isolated from *Oreochromis niloticus* in Piura, Peru: molecular identification and histopathological lesions. *Aquac Rep* 2016; 4:74–9; <https://doi.org/10.1016/j.aqrep.2016.06.002>
- [40] Chen M, Li LP, Wang R, Liang WW, Liang W, Huang Y, et al. Screening vaccine candidate strains against *Streptococcus agalactiae* of Tilapia based on PFGE genotype. *Vaccine* 2012; 30: 6088–92; <https://doi.org/10.1016/j.vaccine.2012.07.044>
- [41] Cieslewicz MJ, Chaffin D, Glusman G, Kasper D, Madan A, Rodrigues S, et al. Structural and genetic diversity of group B *Streptococcus* capsular polysaccharides. *Infect Immun* 2005; 73(5):3096–103; <https://doi.org/10.1128/IAI.73.5.3096-3103.2005>
- [42] Marcusso P, Eto S, Claudiano G, Vieira F, Salvador R, Moraes J, et al. Isolamento de *Streptococcus agalactiae* em diferentes órgãos de tilápiasdo Nilo (*Oreochromis niloticus*) criadas em tanques-rede. *J Biosci* 2015; 31:549–54; <https://doi.org/10.14393/BJ-v31n2a2015-22504>
- [43] Xie T, Wu Q, Zhang J, Xu X, Cheng J. Comparison of *Vibrio parahaemolyticus* isolates from aquatic products and clinical by antibiotic susceptibility, virulence, and molecular characterization. *Food Cont* 2017; 71:315–21; <https://doi.org/10.1016/j.foodcont.2016.06.046>
- [44] Legario FS, Casiano H, Turnbull CJF, Crumlish M. Isolation and molecular characterization of Streptococcal species recovered from clinical infections in farmed Nile tilapia (*Oreochromis niloticus*) in the Philippines. *J Fish Dis* 2020; 43:1431–42; <https://doi.org/10.1111/jfd.13247>
- [45] Ghetas HA, Neiana A, Khalil RH, Hussein AM, Mohamed A. Khallaf. *Streptococcus agalactiae* Isolation and Characterization in Nile Tilapia (*Oreochromis niloticus*) with histopathological studies. *JCVR*. 2021; 3:70–9; <https://doi.org/10.21608/jcvr.2021.160254>
- [46] Amal MNA, Saad MZ, Zahrah AS, Zulkafli AR. Water quality influences the presence of *Streptococcus agalactiae* in cage cultured red hybrid Tilapia, *Oreochromis niloticus* x *Oreochromis mossambicus*. *Aquac Res* 2015; 46:313–23; <https://doi.org/10.1111/are.12180>
- [47] Mian F, Godoy DT, Leal CAG, Yuhara TY, Costa GM, Figueiredo HCP. Aspects of the natural history and virulence of *S. agalactiae* infection Nile Tilapia. *Vet Microbiol Aspects*, 2009; 136:180–3; <https://doi.org/10.1016/j.vetmic.2008.10.016>[47].
- [48] Sadat A, El-Sherbiny H, Zakaria A, Ramadan H, Awad A. Prevalence, Antibiogram and virulence characterization of *Vibrio* isolates from fish and shellfish in Egypt: a possible zoonotic hazard to humans. *J Appl Microbiol* 2020; 11:118–43; <https://doi.org/10.1111/jam.14929>
- [49] Sutcliffe J, Tait Kamradt A, Wondrack L. *Streptococcus pneumoniae* and *Streptococcus pyogenes* resistant to macrolides but sensitive to clindamycin: a common resistance pattern mediated by an efflux system. *Antimicrob Agents Chemother* 1996; 40:1817–24; <https://doi.org/10.1128/AAC.40.8.1817>
- [50] Li C, Sapugahawatte DN, Yang Y, Wong KT, Lo NWS, Margaret IP. Multidrug-resistant *Streptococcus agalactiae* strains found in human and fish with high penicillin and cefotaxime non-susceptibilities. *Microorganisms* 2020; 8:1055; <https://doi.org/10.3390/microorganisms8071055>
- [51] Geng Y, Wang KY, Huang XL, Chen DF, Li CW, Ren SY, et al. *Streptococcus agalactiae* an emerging pathogen for cultured ya-fish, *Schizothorax prenanti*, in China. *Transbound Emerg Dis* 2012; 59:369–75; <https://doi.org/10.1111/j.1865-1682.2011.01280.x>
- [52] Amal MNA, Zamri saad M, Iftikhar AR, Siti Zahrah A, Aziel S, Fahmi S. An outbreak of *Streptococcus agalactiae* infection in cage-cultured golden pompano, *Trachinotus blochii* (Lace'pe'de), in Malaysia. *J Fish Dis* 2012; 35:849–52; <https://doi.org/10.1111/j.1365-2761.2012.01443.x>
- [53] Liu L, Li YW, He RZ, Xiao XX, Zhang X, Su YL, et al. Outbreak of *Streptococcus agalactiae* infection in barcoo grunter, *Scortum barcoo* (McCulloch & Waite), in an intensive fish farm in China. *J Fish Dis* 2014; 37:1067–72; <https://doi.org/10.1111/jfd.12187>
- [54] Simoni S, Vincenzi C, Brenciani A, Morroni G, Bagnarelli P, Giovanetti E, et al. Molecular characterization of italian isolates of fluoroquinolone-resistant *Streptococcus agalactiae* and relationships with chloramphenicol resistance. *Microb Drug Resist* 2018; 24:225–31; <https://doi.org/10.1089/mdr.2017.0139>
- [55] Zhang Z, Lan J, Li Y, Hu M, Yu A, Zhang J, et al. The pathogenic and antimicrobial characteristics of an emerging *Streptococcus agalactiae* serotype IX in Tilapia. *Microb Pathog* 2018; 122:39–45; <https://doi.org/10.1016/j.micpath.2018.05.053>
- [56] Doare KL, O'Driscoll M, Turner K, Seedat F, Russell NJ, Seale AC, et al. Intrapartum antibiotic chemoprophylaxis policies for the prevention of group B Streptococcal disease worldwide: systematic review. *Clin Infect Dis* 2017; 65:64; <https://doi.org/10.1093/cid/cix654>
- [57] Yi T, Li Yw, Liu L, Xiao Xx, Li Ax. Protection of Nile Tilapia (*Oreochromis niloticus* L.) against *Streptococcus agalactiae* following immunization with recombinant FbsA and α -enolase. *Aquaculture* 2014; 428–429:35–40; <https://doi.org/10.1016/j.aquaculture.2014.02.027>
- [58] Antimicrobials: OIE—World Organisation for Animal Health. Available via <https://www.oie.int/scientificexpertise/veterinary-products/antimicrobials/> (Accessed 3 June 2020).
- [59] Godoy DT, Carvalho Castro GA, Leal CA, Pereira UP, Leite RC, Figueiredo HC. Genetic diversity and new genotyping scheme for fish pathogenic *Streptococcus agalactiae*. *Lett Appl Microbiol* 2013; 57(6):476–83; <https://doi.org/10.1111/lam.12138>
- [60] Florindo C, Viegas S, Paulino A, Rodrigues E, Gomes JP, Borrego MJ. Molecular characterization and antimicrobial susceptibility profiles in *Streptococcus agalactiae* colonizing strains: association of erythromycin resistance with subtype III-1 genetic clone family. *Clin Microbiol Infect* 2010; 16:1458–63; <https://doi.org/10.1111/j.1469-0691.2010.03106.x>