

CORRECTION

Open Access



Correction to: Patterns and predictors of analgesic use in pregnancy: a longitudinal drug utilization study with special focus on women with migraine

Gerd-Marie Eskerud Harris^{1*}, Mollie Wood¹, Malin Eberhard-Gran^{2,3}, Christofer Lundqvist³ and Hedvig Marie Egeland Nordeng^{1,2}

Correction to: BMC Pregnancy Childbirth 17, 224 (2017)

<https://doi.org/10.1186/s12884-017-1399-0>

In Fig. 3, the names on the y-axis on the two bottom graphs were switched. The correct names are “Use of triptans” on the left bottom graph, and “Use of opioids” on the right bottom graph. The corrected Figure is provided below. This error has no implications for the interpretation of the findings.

Author details

¹Pharmacoepidemiology & Drug Safety Research Group, Department of Pharmacy, School of Pharmacy, Faculty of Mathematics and Natural Sciences, University of Oslo, P.O. Box. 1068, Blindern, 0316 Oslo, Norway. ²Department of Child Health, National Institute of Public Health, Oslo, Norway. ³Health Services Research, Research Department, Akershus University Hospital and University of Oslo, Campus Ahus, Lørenskog, Norway.

Published online: 08 April 2022

The original article can be found online at <https://doi.org/10.1186/s12884-017-1399-0>.

*Correspondence: g.m.e.harris@farmasi.uio.no

¹ Pharmacoepidemiology & Drug Safety Research Group, Department of Pharmacy, School of Pharmacy, Faculty of Mathematics and Natural Sciences, University of Oslo, P.O. Box. 1068, Blindern, 0316 Oslo, Norway
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 3 Patterns of total analgesic use and use of specific analgesics before and during pregnancy among women with no migraine history, previous history of migraine, recent history of migraine (within 1 year prior to pregnancy) and migraine in the past month (% users in each group with 95% confidence intervals). Analgesics include paracetamol, NSAIDs, opioids and triptans, used for either migraine or headache