


Review

Ubiquitin Proteasome System and Microtubules Are Master Regulators of Central and Peripheral Nervous System Axon Degeneration

Daniele Cartelli ^{1,*}, Guido Cavaletti ², Giuseppe Lauria ^{1,3} and Cristina Meregalli ^{2,*}

- ¹ Neuroalgology Unit, Department of Clinical Neurosciences, Fondazione IRCCS Istituto Neurologico Carlo Besta, Via Celoria 11, 20133 Milan, Italy; giuseppe.lauriapinter@istituto-besta.it
- ² Experimental Neurology Unit and Milan Center for Neuroscience, School of Medicine and Surgery, University of Milano-Bicocca, Via Cadore 48, 20900 Monza, Italy; guido.cavaletti@unimib.it
- ³ Department of Medical Biotechnology and Translational Medicine, University of Milan, Via Vanvitelli 32, 20129 Milan, Italy
- * Correspondence: daniele.cartelli@istituto-besta.it (D.C.); cristina.meregalli@unimib.it (C.M.)

Abstract: Axonal degeneration is an active process that differs from neuronal death, and it is the hallmark of many disorders affecting the central and peripheral nervous system. Starting from the analyses of Wallerian degeneration, the simplest experimental model, here we describe how the long projecting neuronal populations affected in Parkinson's disease and chemotherapy-induced peripheral neuropathies share commonalities in the mechanisms and molecular players driving the earliest phase of axon degeneration. Indeed, both dopaminergic and sensory neurons are particularly susceptible to alterations of microtubules and axonal transport as well as to dysfunctions of the ubiquitin proteasome system and protein quality control. Finally, we report an updated review on current knowledge of key molecules able to modulate these targets, blocking the on-going axonal degeneration and inducing neuronal regeneration. These molecules might represent good candidates for disease-modifying treatment, which might expand the window of intervention improving patients' quality of life.

Keywords: axon regeneration; chemotherapy-induced peripheral neuropathies; dying back; microtubules; Parkinson's disease; ubiquitin proteasome system



Citation: Cartelli, D.; Cavaletti, G.; Lauria, G.; Meregalli, C. Ubiquitin Proteasome System and Microtubules Are Master Regulators of Central and Peripheral Nervous System Axon Degeneration. *Cells* **2022**, *11*, 1358. <https://doi.org/10.3390/cells11081358>

Academic Editors: Nicoletta Gagliano, Michele Sommariva, Claudia Moscheni, Francesca Arnaboldi and Francesca Bianchi

Received: 16 March 2022

Accepted: 13 April 2022

Published: 15 April 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Nervous system functioning depends on the intricate arborisation of neurons and the interconnection they establish with each other and with glial cells during development, some of which are continuously re-modelled during life. Axon degeneration causes the detachment of neurons from the targets, with the consequent loss of the regulation of physiological processes, and it underlies many neurological disorders. Axon degeneration is not blocked by genetic or chemical inhibitors, indicating that it is regulated differently from programmed neuronal death [1]. The analyses of programmed cell death identified different intracellular pathways that can lead to neuronal loss, as the increase of oxidative stress or re-activation of factors regulating the cell cycle [2]; nevertheless, a causal relation between these events and the proper neurodegenerative state is still under debate.

Various pathways mediate the removal of axons. One of the simplest to explore experimentally is Wallerian degeneration (WD) which derives from the transection of axons [3], and although it was first described almost two centuries ago, its molecular mechanisms have been the object of clarification during the last three decades [4,5]. Some of the molecular pathways are shared with the Wallerian-like degeneration termed “dying back”, which characterizes the loss of neuronal populations in different neurodegenerative disorders affecting central and peripheral nervous systems like, for example, Parkinson's

disease (PD) and peripheral neuropathies. Typically, dying back degeneration starts distally and progresses toward the soma [6]. Both these conditions share several reasons for axon degeneration, including that axon pathology is length dependent, the extreme vulnerability to metabolic demand for long distance transport, and the fact that they may derive from exposure to environmental neurotoxins and drugs. A further commonality between PD and peripheral neuropathies is the fact that in the central nervous system there is no evidence of long-range axon regeneration and, sometimes, peripheral nerves even fail to regrow [7]. Last but not least, there are hints about the presence of peripheral fiber pathology in PD patients [8], sustaining the hypothesis of common pathological mechanisms, such as the activation of similar intracellular players, among which two deserve particular attention: microtubules (MTs) and the ubiquitin-proteasome system (UPS).

1.1. Microtubules and Axonal Transport

MTs are polar structures constituted of α/β tubulin heterodimers that alternate slow polymerizing and rapid shrinking phases, a behavior called “dynamic instability” which is more pronounced at the so-called plus-end. In axons, MTs seem to be uniformly plus-end out-oriented, meaning the cellular cargo is transported from the cell to the periphery with more ATP-powered growth, an organization that promotes neuronal polarization [9]. Conversely, in dendrites, MTs are non-uniformly aligned and this mixed orientation drives the minus-end directed dyneins to deliver dendrite-specific cargo [10].

In neurons, MTs heterogeneity is further increased by the combination of different tubulin isoforms, post-translational modifications and MT-associated proteins, which have the potential to generate surface patterns associated with diverse functions, correlating with MT stability (i.e., more stable MTs survive for a longer time) and being associated to different neuronal subregions (Figure 1) [11].

For example, the dynamic tyrosinated MTs are enriched in highly mobile neuronal regions, as presynaptic terminals and dendritic spines [12], whereas the axonal shaft is enriched in detyrosinated/ $\Delta 2$ tubulin and acetylated MTs, modifications that are variously related to the recruitment of interacting proteins and change in MT stability [13], to protection from severing enzymes [14] and axonal regeneration. Furthermore, they regulate both kinesins, plus-end oriented molecular motors and dyneins velocities and processivity [15]. Acetylation of lysine 40 on α tubulin deserves particular attention, since it is the only known luminal MT modification, and it acts as a molecular clock regulating the aging of MTs and their self-repairing capability [16], two important features for structures lasting for the entire life of a neuron.

Axonal transport is the bidirectional trafficking of diverse organelles and cargoes. Due to their preferential direction of movement and the uniform orientation of MTs, in axons, kinesins and dyneins are respectively in charge of the anterograde (from soma to tips) or retrograde transport of proteins and organelles [17]. Whether the axonal transport block is the direct consequence of MT defects or other cellular damaging events remains elusive, but it is a crucial event in several neurodegenerative disorders [18].

Therefore, targeting specific pathways which regulate axonal transport could become a potential therapeutic strategy. One of the most promising targets is histone deacetylase 6 (HDAC6), which removes the acetyl groups from the polymerized MTs [19,20], mediating their activity and the regulation of molecular motors and axonal transport. Its genetic or pharmacological modulation shows protective actions in both animal and human experimental models of different neurodegenerative disorders [21].




Figure 1. Confocal micrograph of cultured neurons, showing a conventional distribution of different tubulin post-translational modifications along the axon (arrowhead) and inside an enlarged terminal (arrow). Tyrosinated tubulin (Tyr TUB, red) is present in the axon and constitutes the microtubule network at the terminal; acetylated tubulin (Ac TUB, green) is enriched along the axon and barely present at the tip; $\Delta 2$ tubulin ($\Delta 2$ TUB, white) is detectable only in the axon shaft showing a proximal-to-distal distribution. Scale bar = 15 μ m.

1.2. Ubiquitin-Proteasome System and Protein Quality Control

The UPS and lysosome-mediated protein degradation via aggresome formation/autophagy are two major degradative pathways of proteins and, since about 30% of newly synthesized proteins are misfolded under physiological conditions, the precise co-operation of these systems is fundamental for cell functioning. In the UPS, proteins to be degraded are ubiquitinated by specific enzymes, and the substrate specificity is determined by a family of more than 600 proteins, the E3 ubiquitin ligases. The other two components of the UPS are the deubiquitinating enzymes and the 26S proteasome, a multi-subunit proteinase complex, which ultimately degrades the ubiquitinated substrates (Figure 2, upper panel) [22]. The structural–functional relationships of the UPS components have

not been fully identified yet, albeit recent works have shed light on how the impairment of ubiquitin-mediated degradation can lead to toxic protein aggregation in age-related degenerative diseases [23–25].




Figure 2. In healthy neurons (**upper panel**), ubiquitin-proteasome system is fully active and degrades the unfolded proteins. Not only, the microtubule organization is conventional, as depicted in Figure 1, and sustains the microtubule-dependent functions as the axonal transport of mitochondria or other cargos. In damaged neurons (**lower panel**), the ubiquitin-proteasome system is blocked, promoting the accumulation of misfolded proteins that induce aggregation, toxicity and axonal degeneration. Microtubule-dependent axonal destruction may be derived from two different conditions: microtubule hyperstabilization, which induces microtubule bundling and block of axonal transport, and microtubule destabilization, with consequent release of microtubule interacting proteins and the dismantling of the railways for intracellular trafficking.

Among the three functional components of UPS, the E3 ligase proteins represent the link with the other cell structures and pathways, and gain- or loss-of-function mutations of a number of E3 genes affect normal neuronal functioning and lead to axon degeneration and nervous system disorders. For example, the PAM-Highwire-Rpm-1 (Phr1) E3 ligase is an effector of the axonal destruction pathway and its loss favors the survival of damaged axons in both the central and peripheral nervous systems [26]. In agreement with this observation, increased activity of an E3 ligase limits the availability of the axonal maintenance factor nicotinamide mononucleotide adenylyltransferase 2 (NMNAT2), thereby initiating degeneration [27].

1.3. Axon Degeneration

WD and the dying back process converge on a programmed pathway of injury signaling and cytoskeleton degradation which results in axonal degeneration [28]. Studies have clarified that the loss of one peculiar axonal protective molecular player, the enzyme NMNAT involved in the biosynthesis of NAD⁺ [4], is crucial to axonal degeneration. Since the axonal isoform of NMNAT, NMNAT2, is constantly replenished by anterograde axonal transport, its deficiency due to axonal transport block or other conditions such as increased proteasome activity can induce axonal degeneration.

Conversely, the loss of function of sterile alpha and TIR motif containing 1 (SARM1), an injury-activated NAD⁺ cleavage enzyme, delays axon degeneration and has neuroprotective capabilities through various mechanisms in some [29–31] but not all neurodegenerative disorders [32]. SARM1 function or loss of NMNAT2 enzymatic activity can trigger the activation of dual leucine kinase (DLK)/MAPK signaling that eventually induces calpain-mediated axonal degeneration [33]. It has been reported that the inhibition of DLK activation could delay axonal degeneration after both axotomy and chemotherapeutic treatment [33]. In agreement with this findings, a recent study showed that the *in vivo* overexpression of the DLK-interacting protein FK506-binding protein 8 (FKBP8) delayed the progression of axon degeneration and suppressed neuronal death through the regulation of DLK stability via the UPS and lysosomal protein degradation pathways [34]. Once the axon is damaged, NMNAT2 reduction or SARM1 activation induces calcium influx that starts the calcium-dependent calpain degradation of the axon, in which the active role of SARM1 is confirmed in human derived sensory neurons models of traumatic and neurotoxic degeneration [35]. Other studies revealed that the cytoskeleton, and especially the MTs, are involved in the active phases of axonal destruction (Figure 2, lower panel). For example, one study demonstrated that MT fragmentation is the earliest detectable event during WD, and that both genetic and pharmacological inhibition of UPS delayed degenerative events both *in vitro* and *in vivo* [36]. In agreement with the crucial role of UPS, the E3 ligase Phr1 tunes the levels of NMNAT2 promoting axonal self-destruction [26,27]. Other studies showed the importance of MTs, as disorganized MTs cause the formation of retraction bulbs and the pharmacological inhibition of MTs leads to regeneration failure [37], whereas direct MT stabilization promotes axon regeneration *in vivo* [38].

As an example of how diseases with different etiologies and different types of stress that initiate axon loss can share the same mechanisms of axon degeneration, we will focus on PD and peripheral neuropathies to show some convergences onto common pathways, similar length-dependent axonal pathology and susceptibility to drugs.

2. Parkinson's Disease

Disorders affecting the central nervous system require the activation of widespread processes whose molecular mechanisms are still elusive, and neurodegenerative diseases can be roughly divided into dementias, including Alzheimer's disease, or motor disorders, such as PD, Huntington's disease or Amyotrophic Lateral Sclerosis. This last class usually derive from the degeneration of long-projecting neuronal populations, such as dopaminergic neurons in PD, which is the second most common neurodegenerative disorder, affecting 2–5% of people over 60s worldwide; it is characterized by bradykinesia, resting tremors and rigidity, symptoms that arise from the lack of dopamine in the striatum due to the death of the dopaminergic neurons in the substantia nigra [39]. The identification of druggable pathways leading to disease-modifying treatments remains an unmet clinical need. The inhibition of the mitochondrial respiratory chain and the UPS failure are amongst those acknowledged as playing key roles [40]. Indeed, ubiquitin-immunopositive inclusions in the Lewy bodies, which are a neuropathological hallmark, can be found in the brain of PD patients [41] and impaired proteasomal activity was reported in the substantia nigra [42]. On the other hand, dopaminergic neurons have very long projections whose average arborisation reaches 4.6 m in humans [43], making them particularly dependent on proteins and organelles supply and, thus, on cytoskeletal integrity and axonal transport,

which has been clearly associated to the early phase of PD [44]. The nigrostriatal system displays dying-back degeneration that resembles WD, with MT fragmentation as the first detectable damage [36]. Similarly, MT disorganization has been extensively associated to the driving phase of axon degeneration in PD (for a detailed review see [45]). However, not all experimental models of PD require the participation of actors playing a pivotal role in WD, such as SARM1 or NMNAT2; indeed SARM1-driven axonal degeneration has been reported only in a single neurotoxic model of PD [46].

Studies based on PD-inducing neurotoxins (as rotenone, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine or MPTP, 6-hydroxydopamine and 2,5 hexanedione) demonstrate that dopaminergic neurons are susceptible to MT-targeted drugs more than other neuronal populations [47] and that derangements of MTs are precocious events that precede and may induce trafficking alterations and mitochondrial failure [48,49]. The common defects resulting from these toxins are the early loss of dynamic MTs, the accumulation of disorganized and fragmented acetylated MTs and the block of axonal transport. The early MT destabilization is not only a by-product of the on-going degeneration since the MT-stabilizer epothilone D is neuroprotective in MPTP-treated mice [50]. This neurotoxin likely activates the same pathway involved in WD, as its administration in Wld(S) mice, a strain resistant to Wallerian-like axonal degeneration, did not induce an early axonal loss [51]; similarly, Wld(S) mice treated with 6-hydroxydopamine showed a delayed axonal degeneration [52].

Many pathological mutations have been reported in different genes that are causally linked to PD, but only two of them account for the vast majority: PRKN, encoding for parkin and representing the major cause of juvenile onset PD, and LRRK2 that encodes for leucine-rich repeat kinase 2 (LRRK2). Both proteins are enzymes involved in many intracellular processes and displayed the ability to interact with actin and tubulin, mediating MT acetylation and having an impact on the transport of mitochondria [53–56]. Interestingly, pathological mutants or protein absence corrupts these processes but these could be rescued by the modulation of MT acetylation [53] or by the use of MT-targeted drugs [57].

Genetic evidence suggests the involvement of UPS dysfunction in PD. Pathological mutations have been reported both in the E3 ligase parkin [58], and in the deubiquitinating enzyme ubiquitin C-terminal hydrolase L1 [59], encoded by UCHL1. Loss-of-function parkin mutants fail to interact with α -synuclein, leading to its aggregation, ubiquitination and accumulation into Lewy bodies in humans [60], and to an age-dependent degeneration of the nigrostriatal pathway accompanied by the appearance of motor deficits in mice [61]. Parkin is also involved in the clearance of mitochondria and MTs. Studies demonstrate that the Parkin/phosphatase and tensin homolog (PTEN)-induced kinase 1 (PINK1) pathway regulates both the transport of functional and damaged mitochondria [54] and mitophagy [62], the process through which the dysfunctional mitochondria are selectively removed. Furthermore, parkin interacts with and promotes the degradation of tubulin by UPS [63], highlighting the extent to which MTs and UPS are intertwined and how E3 ligase can represent a point of convergence between these two systems. Thus, the evidence here reported shows how the alteration of similar pathways can lead to axonal degeneration in PD as well as in WD, but that protective mechanisms may be different.

3. Peripheral Neuropathies and Chemotherapy-Induced Peripheral Neuropathy

Neuropathies represent a heterogeneous entity, as they can be classified as motor, sensory or mixed, such as Charcot-Marie-Tooth disease, and are characterized by loss of muscles and weakness, altered touch sensation and sensitivity to warm and cold; symptoms may also include nociceptive and neuropathic pain. Peripheral neuropathies can be either acquired or inherited, whose central abnormality is axon degeneration. Acquired peripheral neuropathies are amongst the most common neurological disorders worldwide [64] and are a major complication of antineoplastic therapies, affecting at least 50% of patients [65]. They experience a number of symptoms including numbness, paresthesia, neuropathic pain and weakness [66]. The most common cause of chemotherapy-induced peripheral neuropathy (CIPN) are platinum-based compounds, anti-tubulin drugs and proteasome

inhibitors [67]. These compounds share a common pathway triggering an early axonal degeneration. Similar to dopaminergic neurons, long sensory nerves are highly susceptible to axonal transport impairment as a causative role contributing to sensory axon loss, which is correlated with the typical dying back and length-dependent presentation of CIPN [68]. Paclitaxel (PTX) is among the most used antineoplastic drugs causing CIPN. The axonal degeneration induced by PTX might be due to interruption of the axonal transport or to the block of regeneration due to the over-stabilization of re-growing neurites [69]. In addition, other studies have demonstrated that PTX-induced post-translational modifications of MTs can interfere with motor proteins binding and motility [70,71]. PTX is a paradigmatic molecule to show how interconnected are the pathways we are discussing. Indeed, its action on the MT system is well described, while the resistance of the mouse mutant *Wld(S)*, characterized by axonal degeneration delay in a variety of disorders, to PTX-induced neuropathy [72] suggests possible common mechanisms between WD and PTX-induced CIPN.

In line with these observations, the involvement of NMNAT2 and SARM1, the two WD actors required for axon survival and degeneration, respectively, has been reported in several models of peripheral neuropathies. For example, reduced transport of NMNAT2 may increase distal axon sensitivity to degeneration due to decreased NMNAT2 levels and is considered common to many peripheral neuropathies [69]. Other studies support the hypothesis of a shared mechanism between WD and CIPN, since both vincristine (VCR) and bortezomib (BTZ) activate a common SARM1-dependent axon degeneration pathway [73]. Interestingly, a recent paper proposed a novel function for SARM1 in sensory neurons, as a negative regulator of axonal cytoskeleton dynamics and collateral branching; indeed, SARM1 depletion leads to increase of MT polymerization dynamics in cultured neurons and to an increased branching of cutaneous sensory innervation in skin of SARM1 *ko* mice [74]. Modifications of cytoskeletal proteins are described in several peripheral neuropathies and they might interfere with cytoskeleton assembly [75] and thus axonal transport [76]. Disruption of MT dynamics can impair the axonal transport of several elements, which is the case in taxanes and vinca alkaloids compounds, resulting in axonal damage [77,78] and CIPN.

The pivotal role of MT system dysfunctions in the development of CIPN is highlighted by the tubulin-directed agents and strengthened by evidence arising from drugs that target other systems, such as BTZ and cisplatin (CDDP). BTZ is the first proteasome inhibitor introduced for the treatment of hematological malignancies and solid tumors, which acts by inhibiting the chymotrypsin-like enzymatic activity of the 26S proteasome (Figure 2, lower panel), inducing an accumulation of the polyubiquitinated proteins [79].

BTZ causes a sensory-predominant peripheral neuropathy and, although it is structurally unrelated to tubulin drugs, the axonopathy is due to MT over-stabilization in tissue culture cells [80] and accumulation of tubulin in DRG neurons with a time-dependent reduction of mitochondria transport [76]. BTZ can also induce alterations in MT dynamics and molecular transport in human induced pluripotent stem cell-derived sensory neurons [81]. Recent studies demonstrated, both *in vitro* and *in vivo*, that BTZ increased tubulin polymerization, suggesting that an alteration of tubulin dynamics and chronic proteasome inhibition contribute to BTZ-induced peripheral neurotoxicity [82]. Furthermore, BTZ was shown to cause *in vitro* and *in vivo* pathological accumulation of $\Delta 2$ tubulin in sensory neurons, while a reduction of $\Delta 2$ tubulin rescued both mitochondria trafficking defects and axon degeneration [83].

As reported for parkin, BTZ affects mitochondria quality inducing mitotoxicity and proteotoxicity [84], and it also promotes the mitochondrial accumulation of the parkin-interactor PINK1 leading to mitophagy [85]. Accordingly, PINK1 expression ameliorates thermal hypersensitivity in PTX-treated *Drosophila* larvae through the suppression of PTX-induced mitophagy [86]. This evidence suggests a close interrelationship between all these systems.

In addition, some *in vitro* studies on CDDP neurotoxicity, a chemotherapeutic drug that interferes with DNA repair inducing apoptosis of cancer cells, showed impaired fast axonal transport associated with light morphologic alterations [87]. Interestingly, CDDP treatment in *Drosophila* resulted in a large number of stationary mitochondria caused by extended pausing during axonal motility [88]. In a mouse model, the HDAC6 inhibitor ACY-1215 (ricolinostat), which acts on acetylated tubulin, was able to prevent mechanical allodynia, rescue mitochondrial bioenergetic contents and restore intraepidermal nerve fiber density in CDDP-treated mice [89]. Therefore, as we already described in PD contexts, the modulation of MT acetylation might be a potential candidate for reverting the clinical and neuropathological defects associated to some CIPN. All this data highlights that axon destruction programs converge on common pathways although the eliciting events might differ, and explain how the study of particular pathologies can be relevant to our understanding of the general process.

4. Regenerative Potentialities

Strategies to manipulate NMNAT2 or SARM1 are particularly attractive for counteracting on-going degeneration. NMNAT2 is an endogenous survival factor able to sustain labile axons. Its expression level needs to be maintained above a certain threshold to protect neuronal processes from degeneration [90], which can be done by its continuous replenishment through active anterograde transport or by limiting its degradation by UPS. In a *Drosophila* injury model, it was demonstrated that the highly conserved E3 ubiquitin ligase, Highwire, strongly inhibits WD through the modulation of its downstream target, NMNAT2 [27]. In agreement with the role of E3 ligase in maintaining and promoting axonal survival, the UPS related neural precursor cell expressed developmentally down-regulated protein 4 (Nedd4) induces the degradation of PTEN and promotes axon branching and regeneration [91]. The crucial role of PTEN in blocking axonal regeneration has been confirmed in a very recent paper showing that PTEN deletion induces neuron hypertrophy and regeneration [92]. Nedd4 is also a critical regulator of excitability of nociceptive neurons [93]; thus, it has the potentiality to induce both axon regeneration and pain relief.

In recent years, SARM1-depletion has gained attention for its ability to protect from axonal degeneration, and the possibility of pre-treating patients with SARM1 inhibitors before they receive chemotherapy might offer a potentially successful strategy to prevent CIPN. Albeit starting from different engaging mechanisms, VCR and BTZ converge toward a core axonal derangement program that can be inhibited by the expression of candidate genes which deplete SARM1 [73]. In keeping with this observation, SARM1-deficient mice are resistant to Wallerian-like neuropathy induced by PTX or a high fat diet [94]. In addition, pharmacological SARM1 inhibition prevented axon structure alterations and provided partial protection of the axonal function in PTX-treated animal models [95].

The modulation of MT functioning for blocking axonal degeneration or inducing their regeneration made MTs a potential druggable target for disease-modifying treatments [96]. The first hypothesis was to use MT-stabilizers like epothilone D in murine models of PD [50] and Alzheimer's disease [97], but their side-effects, i.e., the induction of peripheral neuropathy, limited their use in patients. Recently, new possibilities are under investigation, such as the use of peptides specific for MT lattice and regulation of post-translational MT modifications. Peptides mimicking tubulin sequence at dimer–dimer interface can induce changes in the longitudinal or lateral contacts, thus having the potentiality to regulate MT stability without causing cell resistance or pathological changes. Davunetide (NAPVSIPQ) is one of the first experimentally used peptides and, due to its very high efficiency in inducing recovery from transport defects and axon damage in different neurodegenerative models, it underwent a clinical trial for the treatment of patients affected by Progressive Supranuclear Palsy, which however did not prove to be efficacious [98]. The regulation of MT post-translational modifications has different possible readouts such as the impact on the MT lattice itself or on MT-interacting protein and molecular motors, thus modulating MT stability and axonal transport. For example, the tubulin-tyrosine ligase increases the

level of tyrosinated MTs at axonal injury sites, mediating retrograde organelle transport and starting the signaling essential to axon regeneration [99]. Similarly, the increase of tyrosinated MTs induces axon regeneration in DRG neurons after injury *in vivo* [100] and the inhibition of MT detyrosination by the administration of cnicin or parthenolide accelerates regeneration and functional recovery of axons *in vivo* [101]. Moreover, the accumulation of $\Delta 2$ tubulin has been associated to BTZ-induced neuropathy and its modulation by multiple enzymes reduces axonal degeneration [83]. As already remarked on, a special mention goes to tubulin acetylation and the enzymes that mediate it, because it may represent an accessible target for interventions aimed at interfering with MT functional organization, axonal transport and axon regrowth. For example, sirtuin-2 deacetylates tubulin increasing the resistance to axonal degeneration in *Wld(S)* mice [102] and its active role in the progression of PD and Alzheimer's disease has been suggested [103]. On the other hand, HDAC5 is an injury-regulated enzyme which modulates tubulin acetylation at growth cone promoting axon advancement and regeneration [104].

The most promising target is HDAC6, which proved its ability in rescuing axon degeneration in many experimental models, acting on three interconnected pathways such as tubulin acetylation, axonal transport and autophagy of misfolded proteins [21]. It has been demonstrated that HDAC6 is up-regulated in a spinal cord injury model, and that its inhibition is able to increase tubulin acetylation and organelle transport, thus restoring the autophagic flux and inducing axonal elongation [105]. Through the same mechanisms, its inhibition is neuroprotective in a drug-induced PD animal model [106]. Moreover, both genetic and pharmacological regulation of MT acetylation could rescue axonal transport defects and motor deficits in the LRKK2 mutants-induced *drosophila* model of PD [50] and in the *in vitro* model of Huntington disease [107]. As previously reported, the administration of ricolinostat in CDDP-treated mice could restore axonal defects and induce the regeneration of sensory terminal nerve endings in the skin [89], ameliorate the behavioral effects and improve the cognitive decline associated to Alzheimer's disease transgenic mice [108]. Furthermore, HDAC6 inhibitors demonstrated the beneficial effects in preclinical models of the axonal form of Charcot-Marie-Tooth [109], removing aggregated protein, rescuing organelles transport and promoting axon regeneration. This evidence provides a rationale for the use of HDAC6 inhibitors as a therapeutic strategy for different, although not all, disorders here described.

5. Closing Remarks

We have presented updated results showing how different pathways co-participate in the axon destruction and how their modulation has the potentiality to induce neuronal regeneration. A strong candidate is the pre-treatment of patients that have to undergo chemotherapy with inhibitors of SARM1, or with multiple antineoplastic agents and/or with genetic regulators of convergent pathways (i.e., miRNA modulators), solutions already used in clinical oncology that might widen the window of intervention [110]. Unfortunately, no one can predict if an individual will develop PD or other neurodegenerative disorders; thus, no preventive treatments are allowed. Among all the possible targets that can prompt axon regeneration, HDAC6 is a promising candidate as it is easily accessible to both pharmacological and genetic points of views, and its action impacts multiple systems. Over the next years, researchers and clinicians have to identify which is the best target in the different pathological or therapeutic contexts, as well as how and when to modulate it.

Author Contributions: C.M. and D.C. conceptualized and wrote the review, defined the search strategies and reviewed the literature. G.C. and G.L. reviewed and edited the final manuscript. All authors contributed to the article and approved the submitted version. All authors have read and agreed to the published version of the manuscript.

Funding: This work was funded by Fondazione Cariplo, grant number 2019-1482 to Meregalli C., PRIN, grant number 2017ZFJCS3 to Cavaletti G.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors are particularly thankful to Malacrida Alessio for his help in the preparation of Figure 2.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Whitmore, A.V.; Lindsten, T.; Raff, M.C.; Thompson, C.B. The proapoptotic proteins Bax and Bak are not involved in Wallerian degeneration. *Cell Death Differ.* **2003**, *10*, 260–261. [[CrossRef](#)] [[PubMed](#)]
2. Krantic, S.; Mechawar, N.; Reix, S.; Quirion, R. Molecular basis of programmed cell death involved in neurodegeneration. *Trends Neurosci.* **2005**, *28*, 670–676. [[CrossRef](#)] [[PubMed](#)]
3. Wang, M.S.; Fang, G.; Culver, D.G.; Davis, A.A.; Rich, M.M.; Glass, J.D. The WldS protein protects against axonal degeneration: A model of gene therapy for peripheral neuropathy. *Ann. Neurol.* **2001**, *50*, 773–779. [[CrossRef](#)] [[PubMed](#)]
4. Conforti, L.; Gilley, J.; Coleman, M.P. Wallerian degeneration: An emerging axon death pathway linking injury and disease. *Nat. Rev. Neurosci.* **2014**, *15*, 394–409. [[CrossRef](#)] [[PubMed](#)]
5. Arthur-Farraj, P.; Coleman, M.P. Lessons from injury: How nerve injury studies reveal basic biological mechanisms and therapeutic opportunities for peripheral nerve diseases. *Neurotherapeutics* **2021**, *18*, 2200–2221. [[CrossRef](#)] [[PubMed](#)]
6. Coleman, M. Axon degeneration mechanisms: Commonality amid diversity. *Nat. Rev. Neurosci.* **2005**, *6*, 889–898. [[CrossRef](#)]
7. Coleman, M.P. The challenges of axon survival: Introduction to the special issue on axonal degeneration. *Exp. Neurol.* **2013**, *246*, 1–5. [[CrossRef](#)] [[PubMed](#)]
8. Nolano, M.; Provitera, V.; Manganelli, F.; Iodice, R.; Stancanelli, A.; Caporaso, G.; Saltalamacchia, A.; Califano, F.; Lanzillo, B.; Picillo, M.; et al. Loss of cutaneous large and small fibers in naive and l-dopa-treated PD patients. *Neurology* **2017**, *89*, 776–784. [[CrossRef](#)]
9. Yau, K.W.; Schätzle, P.; Tortosa, E.; Pagès, S.; Holtmaat, A.; Kapitein, L.C.; Hoogenraad, C.C. Dendrites In vitro and in vivo contain microtubules of opposite polarity and axon formation correlates with uniform plus-end-out microtubule orientation. *J. Neurosci.* **2016**, *36*, 1071–1085. [[CrossRef](#)]
10. Kapitein, L.C.; Schlager, M.A.; Kuijpers, M.; Wulf, P.S.; Van Spronsen, M.; MacKintosh, F.C.; Hoogenraad, C.C. Mixed microtubules steer dynein-driven cargo transport into dendrites. *Curr. Biol.* **2010**, *20*, 290–299. [[CrossRef](#)]
11. Janke, C.; Magiera, M.M. The tubulin code and its role in controlling microtubule properties and functions. *Nat. Rev. Mol. Cell Biol.* **2020**, *21*, 307–326. [[CrossRef](#)] [[PubMed](#)]
12. Jaworski, J.; Kapitein, L.C.; Gouveia, S.M.; Dortland, B.R.; Wulf, P.S.; Grigoriev, I.; Camera, P.; Spangler, S.A.; Di Stefano, P.; Demmers, J.; et al. Dynamic microtubules regulate dendritic spine morphology and synaptic plasticity. *Neuron* **2009**, *61*, 85–100. [[CrossRef](#)] [[PubMed](#)]
13. Bär, J.; Popp, Y.; Bucher, M.; Mikhaylova, M. Direct and indirect effects of tubulin post-translational modifications on microtubule stability: Insights and regulations. *Biochim. Biophys. Acta Mol. Cell Res.* **2022**, *1869*, 119241. [[CrossRef](#)] [[PubMed](#)]
14. Sharp, D.J.; Ross, J.L. Microtubule-severing enzymes at the cutting edge. *J. Cell Sci.* **2012**, *125*, 2561–2569. [[CrossRef](#)]
15. Sirajuddin, M.; Rice, L.M.; Vale, R.D. Regulation of microtubule motors by tubulin isoforms and post-translational modifications. *Nat. Cell Biol.* **2014**, *16*, 335–344. [[CrossRef](#)]
16. Portran, D.; Schaedel, L.; Xu, Z.; Théry, M.; Nachury, M.V. Tubulin acetylation protects long-lived microtubules against mechanical ageing. *Nat. Cell Biol.* **2017**, *19*, 391–398. [[CrossRef](#)]
17. Nirschl, J.J.; Ghirelli, A.E.; Holzbaur, E.L.F. The impact of cytoskeletal organization on the local regulation of neuronal transport. *Nat. Rev. Neurosci.* **2017**, *18*, 585–597. [[CrossRef](#)]
18. Sleight, J.N.; Rossor, A.M.; Fellows, A.D.; Tosolini, A.P.; Schiavo, G. Axonal transport and neurological disease. *Nat. Rev. Neurol.* **2019**, *15*, 691–703. [[CrossRef](#)]
19. Hubbert, C.; Guardiola, A.; Shao, R.; Kawaguchi, Y.; Ito, A.; Nixon, A.; Yoshida, M.; Wang, X.F.; Yao, T.P. HDAC6 is a microtubule-associated deacetylase. *Nature* **2002**, *417*, 455–458. [[CrossRef](#)]
20. Zilberman, Y.; Ballestrem, C.; Carramusa, L.; Mazitschek, R.; Khochbin, S.; Bershadsky, A. Regulation of microtubule dynamics by inhibition of the tubulin deacetylase HDAC6. *J. Cell Sci.* **2009**, *122*, 3531–3541. [[CrossRef](#)]
21. Simões-Pires, C.; Zwick, V.; Nurisso, A.; Schenker, E.; Carrupt, P.A.; Cuendet, M. HDAC6 as a target for neurodegenerative diseases: What makes it different from the other HDACs? *Mol. Neurodegener.* **2013**, *8*, 7. [[CrossRef](#)]
22. Park, J.; Cho, J.; Song, E.J. Ubiquitin-proteasome system (UPS) as a target for anticancer treatment. *Arch. Pharm Res.* **2020**, *43*, 1144–1161. [[CrossRef](#)] [[PubMed](#)]
23. Schmidt, M.F.; Gan, Z.Y.; Komander, D.; Dewson, G. Ubiquitin signalling in neurodegeneration: Mechanisms and therapeutic opportunities. *Cell Death Differ.* **2021**, *28*, 570–590. [[CrossRef](#)] [[PubMed](#)]
24. Kumar, D.; Ambasta, R.K.; Kumar, P. Ubiquitin biology in neurodegenerative disorders: From impairment to therapeutic strategies. *Ageing Res. Rev.* **2020**, *61*, 101078. [[CrossRef](#)] [[PubMed](#)]

25. Fernández-Cruz, I.; Reynaud, E. Proteasome subunits involved in neurodegenerative diseases. *Arch. Med. Res.* **2021**, *52*, 1–14. [[CrossRef](#)] [[PubMed](#)]
26. Babetto, E.; Beirowski, B.; Russler, E.V.; Milbrandt, J.; DiAntonio, A. The Phr1 ubiquitin ligase promotes injury-induced axon self-destruction. *Cell Rep.* **2013**, *3*, 1422–1429. [[CrossRef](#)]
27. Xiong, X.; Hao, Y.; Sun, K.; Li, J.; Li, X.; Mishra, B.; Soppina, P.; Wu, C.; Hume, R.I.; Collins, C.A. The highwire ubiquitin ligase promotes axonal degeneration by tuning levels of Nmnat protein. *PLoS Biol.* **2012**, *10*, e1001440. [[CrossRef](#)]
28. Ma, M.; Ferguson, T.A.; Schoch, K.M.; Li, J.; Qian, Y.; Shofer, F.S.; Saatman, K.E.; Neumar, R.W. Calpains mediate axonal cytoskeleton disintegration during Wallerian degeneration. *Neurobiol. Dis.* **2013**, *56*, 34–46. [[CrossRef](#)]
29. Osterloh, J.M.; Yang, J.; Rooney, T.M.; Fox, A.N.; Adalbert, R.; Powell, E.H.; Sheehan, A.E.; Avery, M.A.; Hackett, R.; Logan, M.A.; et al. DSarm/Sarm1 is required for activation of an injury-induced axon death pathway. *Science* **2012**, *337*, 481–484. [[CrossRef](#)]
30. Tian, W.; Czopka, T.; López-Schier, H. Systemic loss of Sarm1 protects Schwann cells from chemotoxicity by delaying axon degeneration. *Commun. Biol.* **2020**, *3*, 49. [[CrossRef](#)]
31. Sarkar, A.; Kumari, N.; Mukherjee, P. The curious case of SARM1: Dr. Jekyll and Mr. Hyde in cell death and immunity? *FEBS J.* **2021**. [[CrossRef](#)] [[PubMed](#)]
32. Peters, O.M.; Lewis, E.A.; Osterloh, J.M.; Weiss, A.; Salameh, J.S.; Metterville, J.; Brown, R.H.; Freeman, M.R. Loss of Sarm1 does not suppress motor neuron degeneration in the SOD1G93A mouse model of amyotrophic lateral sclerosis. *Hum. Mol. Genet.* **2018**, *27*, 3761–3771. [[CrossRef](#)] [[PubMed](#)]
33. Miller, B.R.; Press, C.; Daniels, R.W.; Sasaki, Y.; Milbrandt, J.; DiAntonio, A. A dual leucine kinase-dependent axon self-destruction program promotes Wallerian degeneration. *Nat. Neurosci.* **2009**, *12*, 387–389. [[CrossRef](#)] [[PubMed](#)]
34. Lee, B.; Oh, Y.; Cho, E.; DiAntonio, A.; Cavalli, V.; Shin, J.E.; Choi, H.W.; Cho, Y. FK506-binding protein-like and FK506-binding protein 8 regulate dual leucine zipper kinase degradation and neuronal responses to axon injury. *J. Biol. Chem.* **2022**, *298*, 101647. [[CrossRef](#)]
35. Chen, Y.H.; Sasaki, Y.; DiAntonio, A.; Milbrandt, J. SARM1 is required in human derived sensory neurons for injury-induced and neurotoxic axon degeneration. *Exp. Neurol.* **2021**, *339*, 113636. [[CrossRef](#)]
36. Zhai, Q.; Wang, J.; Kim, A.; Liu, Q.; Watts, R.; Hoopfer, E.; Mitchison, T.; Luo, L.; He, Z. Involvement of the ubiquitin-proteasome system in the early stages of wallerian degeneration. *Neuron* **2003**, *39*, 217–225. [[CrossRef](#)]
37. Ertürk, A.; Hellal, F.; Enes, J.; Bradke, F. Disorganized microtubules underlie the formation of retraction bulbs and the failure of axonal regeneration. *J. Neurosci.* **2007**, *27*, 9169–9180. [[CrossRef](#)]
38. Hellal, F.; Hurtado, A.; Ruschel, J.; Flynn, K.C.; Laskowski, C.J.; Umlauf, M.; Kapitein, L.C.; Strikis, D.; Lemmon, V.; Bixby, J.; et al. Microtubule stabilization reduces scarring and causes axon regeneration after spinal cord injury. *Science* **2011**, *331*, 928–931. [[CrossRef](#)]
39. Obeso, J.A.; Rodriguez-Oroz, M.C.; Goetz, C.G.; Marin, C.; Kordower, J.H.; Rodriguez, M.; Hirsch, E.C.; Farrer, M.; Schapira, A.H.; Halliday, G. Missing pieces in the Parkinson’s disease puzzle. *Nat. Med.* **2010**, *16*, 653–661. [[CrossRef](#)]
40. Dauer, W.; Przedborski, S. Parkinson’s disease: Mechanisms and models. *Neuron* **2003**, *39*, 889–909. [[CrossRef](#)]
41. Kuzuhara, S.; Mori, H.; Izumiyama, N.; Yoshimura, M.; Ihara, Y. Lewy bodies are ubiquitinated. A light and electron microscopic immunocytochemical study. *Acta Neuropathol.* **1988**, *75*, 345–353. [[CrossRef](#)] [[PubMed](#)]
42. McNaught, K.S.; Jenner, P. Proteasomal function is impaired in substantia nigra in Parkinson’s disease. *Neurosci. Lett.* **2001**, *297*, 191–194. [[CrossRef](#)]
43. Pissadaki, E.K.; Bolam, J.P. The energy cost of action potential propagation in dopamine neurons: Clues to susceptibility in Parkinson’s disease. *Front. Comput. Neurosci.* **2013**, *7*, 13. [[CrossRef](#)] [[PubMed](#)]
44. Hunn, B.H.; Cragg, S.J.; Bolam, J.P.; Spillantini, M.G.; Wade-Martins, R. Impaired intracellular trafficking defines early Parkinson’s disease. *Trends Neurosci.* **2015**, *38*, 178–188. [[CrossRef](#)] [[PubMed](#)]
45. Cartelli, D.; Cappelletti, G. Microtubule destabilization paves the way to Parkinson’s disease. *Mol. Neurobiol.* **2017**, *54*, 6762–6774. [[CrossRef](#)]
46. Peters, O.M.; Weiss, A.; Metterville, J.; Song, L.; Logan, R.; Smith, G.A.; Schwarzschild, M.A.; Mueller, C.; Brown, R.H.; Freeman, M. Genetic diversity of axon degenerative mechanisms in models of Parkinson’s disease. *Neurobiol. Dis.* **2021**, *155*, 105368. [[CrossRef](#)]
47. Ren, Y.; Liu, W.; Jiang, H.; Jiang, Q.; Feng, J. Selective vulnerability of dopaminergic neurons to microtubule depolymerization. *J. Biol. Chem.* **2005**, *280*, 34105–34112. [[CrossRef](#)]
48. Cartelli, D.; Ronchi, C.; Maggioni, M.G.; Rodighiero, S.; Giavini, E.; Cappelletti, G. Microtubule dysfunction precedes transport impairment and mitochondria damage in MPP+ -induced neurodegeneration. *J. Neurochem.* **2010**, *115*, 247–258. [[CrossRef](#)]
49. Casagrande, F.V.M.; Amadeo, A.; Cartelli, D.; Calogero, A.M.; Modena, D.; Costa, I.; Cantele, F.; Onelli, E.; Moscatelli, A.; Ascagni, M.; et al. The imbalance between dynamic and stable microtubules underlies neurodegeneration induced by 2,5-hexanedione. *Biochim. Biophys. Acta Mol. Basis Dis.* **2020**, *1866*, 165581. [[CrossRef](#)]
50. Cartelli, D.; Casagrande, F.; Busceti, C.L.; Bucci, D.; Molinaro, G.; Traficante, A.; Passarella, D.; Giavini, E.; Pezzoli, G.; Battaglia, G.; et al. Microtubule alterations occur early in experimental parkinsonism and the microtubule stabilizer epothilone D is neuroprotective. *Sci. Rep.* **2013**, *3*, 1837. [[CrossRef](#)]
51. Hasbani, D.M.; O’Malley, K.L. Wld(S) mice are protected against the Parkinsonian mimetic MPTP. *Exp. Neurol.* **2006**, *202*, 93–99. [[CrossRef](#)] [[PubMed](#)]

52. Sajadi, A.; Schneider, B.L.; Aebischer, P. Wlds-mediated protection of dopaminergic fibers in an animal model of Parkinson disease. *Curr. Biol.* **2004**, *14*, 326–330. [[CrossRef](#)] [[PubMed](#)]
53. Godena, V.K.; Brookes-Hocking, N.; Moller, A.; Shaw, G.; Oswald, M.; Sancho, R.M.; Miller, C.C.; Whitworth, A.J.; De Vos, K.J. Increasing microtubule acetylation rescues axonal transport and locomotor deficits caused by LRRK2 Roc-COR domain mutations. *Nat. Commun.* **2014**, *5*, 5245. [[CrossRef](#)]
54. Wang, X.; Winter, D.; Ashrafi, G.; Schlehe, J.; Wong, Y.L.; Selkoe, D.; Rice, S.; Steen, J.; LaVoie, M.J.; Schwarz, T.L. PINK1 and Parkin target Miro for phosphorylation and degradation to arrest mitochondrial motility. *Cell* **2011**, *147*, 893–906. [[CrossRef](#)] [[PubMed](#)]
55. Ren, Y.; Jiang, H.; Hu, Z.; Fan, K.; Wang, J.; Janoschka, S.; Wang, X.; Ge, S.; Feng, J. Parkin mutations reduce the complexity of neuronal processes in iPSC-derived human neurons. *Stem Cells* **2015**, *33*, 68–78. [[CrossRef](#)]
56. Cartelli, D.; Goldwurm, S.; Casagrande, F.; Pezzoli, G.; Cappelletti, G. Microtubule destabilization is shared by genetic and idiopathic Parkinson’s disease patient fibroblasts. *PLoS ONE* **2012**, *7*, e37467. [[CrossRef](#)]
57. Cartelli, D.; Amadeo, A.; Calogero, A.M.; Casagrande, F.V.M.; De Gregorio, C.; Gioria, M.; Kuzumaki, N.; Costa, I.; Sassone, J.; Ciammola, A.; et al. Parkin absence accelerates microtubule aging in dopaminergic neurons. *Neurobiol. Aging* **2018**, *61*, 66–74. [[CrossRef](#)]
58. Kitada, T.; Asakawa, S.; Hattori, N.; Matsumine, H.; Yamamura, Y.; Minoshima, S.; Yokochi, M.; Mizuno, Y.; Shimizu, N. Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature* **1998**, *392*, 605–608. [[CrossRef](#)]
59. Leroy, E.; Boyer, R.; Auburger, G.; Leube, B.; Ulm, G.; Mezey, E.; Harta, G.; Brownstein, M.J.; Jonnalagada, S.; Chernova, T.; et al. The ubiquitin pathway in Parkinson’s disease. *Nature* **1998**, *395*, 451–452. [[CrossRef](#)]
60. Shimura, H.; Schlossmacher, M.G.; Hattori, N.; Frosch, M.P.; Trockenbacher, A.; Schneider, R.; Mizuno, Y.; Kosik, K.S.; Selkoe, D.J. Ubiquitination of a new form of alpha-synuclein by parkin from human brain: Implications for Parkinson’s disease. *Science* **2001**, *293*, 263–269. [[CrossRef](#)]
61. Lu, X.H.; Fleming, S.M.; Meurers, B.; Ackerson, L.C.; Mortazavi, F.; Lo, V.; Hernandez, D.; Sulzer, D.; Jackson, G.R.; Maidment, N.T.; et al. Bacterial artificial chromosome transgenic mice expressing a truncated mutant parkin exhibit age-dependent hypokinetic motor deficits, dopaminergic neuron degeneration, and accumulation of proteinase K-resistant alpha-synuclein. *J. Neurosci.* **2009**, *29*, 1962–1976. [[CrossRef](#)] [[PubMed](#)]
62. Vives-Bauza, C.; Zhou, C.; Huang, Y.; Cui, M.; De Vries, R.L.; Kim, J.; May, J.; Tocilescu, M.A.; Liu, W.; Ko, H.S.; et al. PINK1-dependent recruitment of Parkin to mitochondria in mitophagy. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 378–383. [[CrossRef](#)]
63. Ren, Y.; Zhao, J.; Feng, J. Parkin binds to alpha/beta tubulin and increases their ubiquitination and degradation. *J. Neurosci.* **2003**, *23*, 3316–3324. [[CrossRef](#)] [[PubMed](#)]
64. Prior, R.; Van Helleputte, L.; Benoy, V.; Van Den Bosch, L. Defective axonal transport: A common pathological mechanism in inherited and acquired peripheral neuropathies. *Neurobiol. Dis.* **2017**, *105*, 300–320. [[CrossRef](#)] [[PubMed](#)]
65. Cavaletti, G. Chemotherapy-induced peripheral neurotoxicity (CIPN): What we need and what we know. *J. Peripher. Nerv. Syst.* **2014**, *19*, 66–76. [[CrossRef](#)] [[PubMed](#)]
66. Toftagen, C. Patient perceptions associated with chemotherapy-induced peripheral neuropathy. *Clin. J. Oncol. Nurs.* **2010**, *14*, E22–E28. [[CrossRef](#)]
67. Argyriou, A.A.; Bruna, J.; Marmiroli, P.; Cavaletti, G. Chemotherapy-induced peripheral neurotoxicity (CIPN): An update. *Crit. Rev. Oncol. Hematol.* **2012**, *82*, 51–77. [[CrossRef](#)]
68. Geisler, S. Vincristine- and bortezomib-induced neuropathies—From bedside to bench and back. *Exp. Neurol.* **2021**, *336*, 113519. [[CrossRef](#)]
69. Cashman, C.R.; Höke, A. Mechanisms of distal axonal degeneration in peripheral neuropathies. *Neurosci. Lett.* **2015**, *596*, 33–50. [[CrossRef](#)]
70. Hammond, J.W.; Huang, C.F.; Kaech, S.; Jacobson, C.; Banker, G.; Verhey, K.J. Posttranslational modifications of tubulin and the polarized transport of kinesin-1 in neurons. *Mol. Biol. Cell* **2010**, *21*, 572–583. [[CrossRef](#)]
71. Gornstein, E.; Schwarz, T.L. The paradox of paclitaxel neurotoxicity: Mechanisms and unanswered questions. *Neuropharmacology* **2014**, *76*, 175–183. [[CrossRef](#)] [[PubMed](#)]
72. Wang, M.S.; Davis, A.A.; Culver, D.G.; Glass, J.D. WldS mice are resistant to paclitaxel (taxol) neuropathy. *Ann. Neurol.* **2002**, *52*, 442–447. [[CrossRef](#)] [[PubMed](#)]
73. Geisler, S.; Doan, R.A.; Cheng, G.C.; Cetinkaya-Fisgin, A.; Huang, S.X.; Höke, A.; Milbrandt, J.; DiAntonio, A. Vincristine and bortezomib use distinct upstream mechanisms to activate a common SARM1-dependent axon degeneration program. *JCI Insight* **2019**, *4*, e129920. [[CrossRef](#)] [[PubMed](#)]
74. Ketschek, A.; Holland, S.M.; Gallo, G. SARM1 suppresses axon branching through attenuation of axonal cytoskeletal dynamics. *Front. Mol. Neurosci.* **2022**, *15*, 726962. [[CrossRef](#)] [[PubMed](#)]
75. Malacrida, A.; Meregalli, C.; Rodriguez-Menendez, V.; Nicolini, G. Chemotherapy-induced peripheral neuropathy and changes in cytoskeleton. *Int. J. Mol. Sci.* **2019**, *20*, 2287. [[CrossRef](#)] [[PubMed](#)]
76. Staff, N.P.; Podratz, J.L.; Grassner, L.; Bader, M.; Paz, J.; Knight, A.M.; Loprinzi, C.L.; Trushina, E.; Windebank, A.J. Bortezomib alters microtubule polymerization and axonal transport in rat dorsal root ganglion neurons. *Neurotoxicology* **2013**, *39*, 124–131. [[CrossRef](#)]

77. LaPointe, N.E.; Morfini, G.; Brady, S.T.; Feinstein, S.C.; Wilson, L.; Jordan, M.A. Effects of eribulin, vincristine, paclitaxel and ixabepilone on fast axonal transport and kinesin-1 driven microtubule gliding: Implications for chemotherapy-induced peripheral neuropathy. *Neurotoxicology* **2013**, *37*, 231–239. [[CrossRef](#)]
78. Topp, K.S.; Tanner, K.D.; Levine, J.D. Damage to the cytoskeleton of large diameter sensory neurons and myelinated axons in vincristine-induced painful peripheral neuropathy in the rat. *J. Comp. Neurol.* **2000**, *424*, 563–576. [[CrossRef](#)]
79. Zhang, L.; Wu, M.; Su, R.; Zhang, D.; Yang, G. The efficacy and mechanism of proteasome inhibitors in solid tumor treatment. *Recent Pat. Anticancer Drug Discov.* **2021**. [[CrossRef](#)]
80. Poruchynsky, M.S.; Sackett, D.L.; Robey, R.W.; Ward, Y.; Annunziata, C.; Fojo, T. Proteasome inhibitors increase tubulin polymerization and stabilization in tissue culture cells: A possible mechanism contributing to peripheral neuropathy and cellular toxicity following proteasome inhibition. *Cell Cycle* **2008**, *7*, 940–949. [[CrossRef](#)]
81. Hrstka, S.C.L.; Ankam, S.; Agac, B.; Klein, J.P.; Moore, R.A.; Narapureddy, B.; Schneider, I.; Hrstka, R.F.; Dasari, S.; Staff, N.P. Proteomic analysis of human iPSC-derived sensory neurons implicates cell stress and microtubule dynamics dysfunction in bortezomib-induced peripheral neurotoxicity. *Exp. Neurol.* **2021**, *335*, 113520. [[CrossRef](#)] [[PubMed](#)]
82. Meregalli, C.; Chiorazzi, A.; Carozzi, V.A.; Canta, A.; Sala, B.; Colombo, M.; Oggioni, N.; Ceresa, C.; Foudah, D.; La Russa, F.; et al. Evaluation of tubulin polymerization and chronic inhibition of proteasome as cytotoxicity mechanisms in bortezomib-induced peripheral neuropathy. *Cell Cycle* **2014**, *13*, 612–621. [[CrossRef](#)] [[PubMed](#)]
83. Pero, M.E.; Meregalli, C.; Qu, X.; Shin, G.J.; Kumar, A.; Shorey, M.; Rolls, M.M.; Tanji, K.; Brannagan, T.H.; Alberti, P.; et al. Pathogenic role of delta 2 tubulin in bortezomib-induced peripheral neuropathy. *Proc. Natl. Acad. Sci. USA* **2021**, *118*, e2012685118. [[CrossRef](#)] [[PubMed](#)]
84. Jannuzzi, A.T.; Arslan, S.; Yilmaz, A.M.; Sari, G.; Beklen, H.; Méndez, L.; Fedorova, M.; Arga, K.Y.; Karademir Yilmaz, B.; Alpertunga, B. Higher proteotoxic stress rather than mitochondrial damage is involved in higher neurotoxicity of bortezomib compared to carfilzomib. *Redox Biol.* **2020**, *32*, 101502. [[CrossRef](#)]
85. Beider, K.; Rosenberg, E.; Dimenshtein-Voevoda, V.; Sirovsky, Y.; Vladimirovsky, J.; Magen, H.; Ostrovsky, O.; Shimoni, A.; Bromberg, Z.; Weiss, L.; et al. Blocking of Transient Receptor Potential Vanilloid 1 (TRPV1) promotes terminal mitophagy in multiple myeloma, disturbing calcium homeostasis and targeting ubiquitin pathway and bortezomib-induced unfolded protein response. *J. Hematol. Oncol.* **2020**, *13*, 158. [[CrossRef](#)]
86. Kim, Y.Y.; Yoon, J.H.; Um, J.H.; Jeong, D.J.; Shin, D.J.; Hong, Y.B.; Kim, J.K.; Kim, D.H.; Kim, C.; Chung, C.G.; et al. PINK1 alleviates thermal hypersensitivity in a paclitaxel-induced Drosophila model of peripheral neuropathy. *PLoS ONE* **2020**, *15*, e0239126. [[CrossRef](#)]
87. Russell, J.W.; Windebank, A.J.; McNiven, M.A.; Brat, D.J.; Brimijoin, W.S. Effect of cisplatin and ACTH4-9 on neural transport in cisplatin induced neurotoxicity. *Brain Res.* **1995**, *676*, 258–267. [[CrossRef](#)]
88. Podratz, J.L.; Lee, H.; Knorr, P.; Koehler, S.; Forsythe, S.; Lambrecht, K.; Arias, S.; Schmidt, K.; Steinhoff, G.; Yudintsev, G.; et al. Cisplatin induces mitochondrial deficits in Drosophila larval segmental nerve. *Neurobiol. Dis.* **2017**, *97*, 60–69. [[CrossRef](#)]
89. Krukowski, K.; Ma, J.; Golonzhka, O.; Laumet, G.O.; Gutti, T.; Van Duzer, J.H.; Mazitschek, R.; Jarpe, M.B.; Heijnen, C.J.; Kavelaars, A. HDAC6 inhibition effectively reverses chemotherapy-induced peripheral neuropathy. *Pain* **2017**, *158*, 1126–1137. [[CrossRef](#)]
90. Gilley, J.; Mayer, P.R.; Yu, G.; Coleman, M.P. Low levels of NMNAT2 compromise axon development and survival. *Hum. Mol. Genet.* **2019**, *28*, 448–458. [[CrossRef](#)]
91. Christie, K.J.; Martinez, J.A.; Zochodne, D.W. Disruption of E3 ligase NEDD4 in peripheral neurons interrupts axon outgrowth: Linkage to PTEN. *Mol. Cell Neurosci.* **2012**, *50*, 179–192. [[CrossRef](#)] [[PubMed](#)]
92. Meyer Zu Reckendorf, S.; Moser, D.; Blechschmidt, A.; Joga, V.N.; Sinske, D.; Hegler, J.; Deininger, S.; Catanese, A.; Vettorazzi, S.; Antoniadis, G.; et al. Motoneuron-specific PTEN deletion in mice induces neuronal hypertrophy and also regeneration after facial nerve injury. *J. Neurosci.* **2022**, *42*, 2474–2491. [[CrossRef](#)]
93. Deftu, A.F.; Chu Sin Chung, P.; Laedermann, C.J.; Gillet, L.; Pertin, M.; Kirschmann, G.; Décosterd, I. The anti-diabetic drug metformin regulates voltage-gated sodium channel Na. *ENEURO* **2022**, *9*, ENEURO.0409-21.2022. [[CrossRef](#)] [[PubMed](#)]
94. Turkiew, E.; Falconer, D.; Reed, N.; Höke, A. Deletion of Sarm1 gene is neuroprotective in two models of peripheral neuropathy. *J. Peripher. Nerv. Syst.* **2017**, *22*, 162–171. [[CrossRef](#)] [[PubMed](#)]
95. Bosanac, T.; Hughes, R.O.; Engber, T.; Devraj, R.; Brearley, A.; Danker, K.; Young, K.; Kopatz, J.; Hermann, M.; Berthemy, A.; et al. Pharmacological SARM1 inhibition protects axon structure and function in paclitaxel-induced peripheral neuropathy. *Brain* **2021**, *144*, 3226–3238. [[CrossRef](#)]
96. Baas, P.W.; Ahmad, F.J. Beyond taxol: Microtubule-based treatment of disease and injury of the nervous system. *Brain* **2013**, *136*, 2937–2951. [[CrossRef](#)]
97. Brunden, K.R.; Zhang, B.; Carroll, J.; Yao, Y.; Potuzak, J.S.; Hogan, A.M.; Iba, M.; James, M.J.; Xie, S.X.; Ballatore, C.; et al. Epothilone D improves microtubule density, axonal integrity, and cognition in a transgenic mouse model of tauopathy. *J. Neurosci.* **2010**, *30*, 13861–13866. [[CrossRef](#)]
98. Boxer, A.L.; Lang, A.E.; Grossman, M.; Knopman, D.S.; Miller, B.L.; Schneider, L.S.; Doody, R.S.; Lees, A.; Golbe, L.I.; Williams, D.R.; et al. Davunetide in patients with progressive supranuclear palsy: A randomised, double-blind, placebo-controlled phase 2/3 trial. *Lancet Neurol.* **2014**, *13*, 676–685. [[CrossRef](#)]
99. Song, W.; Cho, Y.; Watt, D.; Cavalli, V. Tubulin-tyrosine ligase (TTL)-mediated increase in tyrosinated α -tubulin in injured axons is required for retrograde injury signaling and axon regeneration. *J. Biol. Chem.* **2015**, *290*, 14765–14775. [[CrossRef](#)]

100. Wu, D.; Jin, Y.; Shapiro, T.M.; Hinduja, A.; Baas, P.W.; Tom, V.J. Chronic neuronal activation increases dynamic microtubules to enhance functional axon regeneration after dorsal root crush injury. *Nat. Commun.* **2020**, *11*, 6131. [[CrossRef](#)]
101. Gobrecht, P.; Andreadaki, A.; Diekmann, H.; Heskamp, A.; Leibinger, M.; Fischer, D. Promotion of functional nerve regeneration by inhibition of microtubule deetyrosination. *J. Neurosci.* **2016**, *36*, 3890–3902. [[CrossRef](#)] [[PubMed](#)]
102. Suzuki, K.; Koike, T. Mammalian Sir2-related protein (SIRT) 2-mediated modulation of resistance to axonal degeneration in slow Wallerian degeneration mice: A crucial role of tubulin deacetylation. *Neuroscience* **2007**, *147*, 599–612. [[CrossRef](#)] [[PubMed](#)]
103. Esteves, A.R.; Palma, A.M.; Gomes, R.; Santos, D.; Silva, D.F.; Cardoso, S.M. Acetylation as a major determinant to microtubule-dependent autophagy: Relevance to Alzheimer’s and Parkinson disease pathology. *Biochim. Biophys. Acta Mol. Basis Dis.* **2019**, *1865*, 2008–2023. [[CrossRef](#)] [[PubMed](#)]
104. Cho, Y.; Cavalli, V. HDAC5 is a novel injury-regulated tubulin deacetylase controlling axon regeneration. *EMBO J.* **2012**, *31*, 3063–3078. [[CrossRef](#)]
105. Zheng, Z.; Zhou, Y.; Ye, L.; Lu, Q.; Zhang, K.; Zhang, J.; Xie, L.; Wu, Y.; Xu, K.; Zhang, H.; et al. Histone deacetylase 6 inhibition restores autophagic flux to promote functional recovery after spinal cord injury. *Exp. Neurol.* **2020**, *324*, 113138. [[CrossRef](#)]
106. El-Saiy, K.A.; Sayed, R.H.; El-Sahar, A.E.; Kandil, E.A. Modulation of histone deacetylase, the ubiquitin proteasome system, and autophagy underlies the neuroprotective effects of venlafaxine in a rotenone-induced Parkinson’s disease model in rats. *Chem. Biol. Interact.* **2022**, *354*, 109841. [[CrossRef](#)]
107. Dompierre, J.P.; Godin, J.D.; Charrin, B.C.; Cordelières, F.P.; King, S.J.; Humbert, S.; Saudou, F. Histone deacetylase 6 inhibition compensates for the transport deficit in Huntington’s disease by increasing tubulin acetylation. *J. Neurosci.* **2007**, *27*, 3571–3583. [[CrossRef](#)]
108. Zhang, L.; Liu, C.; Wu, J.; Tao, J.J.; Sui, X.L.; Yao, Z.G.; Xu, Y.F.; Huang, L.; Zhu, H.; Sheng, S.L.; et al. Tubastatin A/ACY-1215 improves cognition in Alzheimer’s disease transgenic mice. *J. Alzheimers Dis.* **2014**, *41*, 1193–1205. [[CrossRef](#)]
109. Rossaert, E.; Van Den Bosch, L. HDAC6 inhibitors: Translating genetic and molecular insights into a therapy for axonal CMT. *Brain Res.* **2020**, *1733*, 146692. [[CrossRef](#)]
110. Dai, Z.; Chu, H.; Ma, J.; Yan, Y.; Zhang, X.; Liang, Y. The regulatory mechanisms and therapeutic potential of microRNAs: From chronic pain to morphine tolerance. *Front. Mol. Neurosci.* **2018**, *11*, 80. [[CrossRef](#)]